




Reduction of couplings in the Type-II 2HDM

M. A. May Pech^{1,a}, M. Mondragón^{1,b}, G. Patellis^{2,c} , G. Zoupanos^{3,4,5,6,d}

¹ Instituto de Física, Universidad Nacional Autónoma de México, A.P. 20-364, 01000 Mexico City, Mexico

² Centro de Física Teórica de Partículas-CFTP, Instituto Superior Técnico, Departamento de Física, Universidade de Lisboa, Avenida Rovisco Pais 1, 1049-001 Lisbon, Portugal

³ Physics Department, National Technical University, Zografou, 157 80 Athens, Greece

⁴ Institut für Theoretische Physik der Universität Heidelberg, Philosophenweg 16, 69120 Heidelberg, Germany

⁵ Max-Planck Institut für Physik, Föhringer Ring 6, 80805 Munich, Germany

⁶ Theoretical Physics Department, CERN, Geneva, Switzerland

Received: 3 November 2023 / Accepted: 4 December 2023
© The Author(s) 2023

Abstract The idea of *reduction of couplings* consists in the search for relations between seemingly independent couplings of a renormalizable theory that are renormalization group invariant. In this article, we demonstrate the existence of such 1-loop relations among the top Yukawa, the Higgs quartic and the gauge colour couplings of the Type-II Two Higgs Doublet Model at a high-energy boundary. The phenomenological viability of the reduced theory suggests the value of $\tan \beta$ and the scale in which new physics may appear.

1 Introduction

An essential direction of the last decades in theoretical particle physics is to understand the free parameters of the Standard Model (SM) in terms of a few fundamental ones, i.e. to achieve a *reduction of couplings* (RoC) [1]. However, despite the numerous successes of the SM regarding the description of elementary particles and the interactions among them, there is significantly less progress when it comes to the freedom in the parameter space. The problem of the large number of arbitrary parameters is deeply related to the infinities that emerge at the quantum level. While renormalization succeeds in removing those infinities, it only does so at the cost of introducing counter terms, which leaves the ‘cured’ parameters free to be fixed by the experiment.

Although the success of the SM is undisputed, it is a widespread belief that it is ultimately the low energy limit of a

(more) fundamental theory. Under this light, one of the most popular and efficient ways to reduce that freedom in parameter space is to introduce a symmetry. A well known example of such an idea are the grand unified theories (GUTs) [2–7]. Within the GUT framework, gauge couplings are related and one can even have a unified Yukawa sector. Unfortunately, beyond the minimal $SU(5)$, which was experimentally ruled out a long time ago, theories based on larger groups give rise to new complications regarding the number of free parameters since new degrees of freedom are necessary (i.e. the channels of breaking the symmetry).

The RoC method was proposed as an alternative, systematic way to look for relations among seemingly unrelated parameters [8–10] (see also [11–13]). This technique reduces the number of independent parameters of a theory by relating either all (in its original version) or a number of parameters to a single coupling, which is often called ‘primary coupling’. For this approach to be considered systematic, two conditions should hold. First, both the original and the reduced theory should be renormalizable. Second, the relations among the various parameters involved should be renormalization group invariant (RGI).

This idea was of course first applied on the SM almost four decades ago [14, 15] and, while at the time it produced promising predictions for the top quark and Higgs boson masses, their following respective experimental discoveries ruled them out as too light. However, this work opened the way to a number of theories that extend the SM and are based on the concept of the reduction of couplings, which had significant predictive power and were [16–20] or continue to be [21, 22] successful.

In the present work we choose a minimal extension of the SM, namely the well known Two Higgs Doublet Model

^a e-mail: miguel.maypech@gmail.com

^b e-mail: myriam@fisica.unam.mx

^c e-mail: grigorios.patellis@tecnico.ulisboa.pt (corresponding author)

^d e-mail: george.zoupanos@cern.ch

(2HDM) [23] and, more specifically, its Type-II version (this is just a convenience choice, since similar procedures as the one here described can be applied to all versions). In particular, we use a version of the RoC technique first introduced in [24] in search of a boundary scale at which we have New Physics. The elegance of this approach is that only a second Higgs doublet is needed in order to fit the reduced model with the current experimental constraints and pinpoint the scale above which new field content and/or symmetries could come into effect.

2 Reduction of couplings basics

A brief description of the basic idea of *reduction of couplings* is first in order, as it was introduced in [8] and consequently expanded over the next decades. The goal is to express the couplings of a theory that are considered free in terms of one parameter, that is considered to be more fundamental, called the primary coupling. The basic idea is to search for renormalization group invariant (RGI) relations among parameters that reduce the degree of arbitrariness of the parameter space.

Such relations are in general of the form $F(g_1, \dots, g_A) = \text{const.}$ for A number of parameters, which should satisfy the partial differential equation (PDE):

$$\mu \frac{dF}{d\mu} = \vec{\nabla} F \cdot \vec{\beta} = \sum_{a=1}^A \beta_a \frac{\partial F}{\partial g_a} = 0, \tag{1}$$

where β_a are the β -functions of each coupling g_a , respectively, in order for F to be RGI. This PDE is equivalent to the set of ordinary differential equations below, which are called reduction equations (REs) [8–10],

$$\beta_g \frac{dg_a}{dg} = \beta_a, \quad a = 1, \dots, A - 1, \tag{2}$$

where now g and β_g are the primary coupling and its respective β -function. There are – maximally – $A - 1$ independent RGI constraints in the A -dimensional space of parameters imposed by the F_a 's, thus one could in principle express all parameters in terms of one primary coupling, g .

However, the general solutions of the REs contain as many integration constants as the number of the equations themselves. Thus, so far we have just traded an integration constant for each coupling and these general solutions cannot be considered to have reduced the freedom of the parameter space. The crucial requirement is that the REs admit power series solutions:

$$g_a = \sum_n \rho_a^{(n)} g^{2n+1}, \tag{3}$$

which preserve perturbative renormalizability. Remarkably, the uniqueness of these power series solutions can be already decided at 1-loop level [8–10].

The possibility of a *complete* reduction of couplings described above is without doubt very attractive, as the completely reduced theory features only one independent coupling. However, in many cases this has been proven to be unrealistic. Therefore, fewer RGI constraints are often imposed, leading to a *partial* reduction [14, 15] of the parameter space.

3 Notation and parameters of the 2HDM

For the two Higgs doublets Φ_1, Φ_2 the most general renormalizable scalar potential can be written as [25–27]:

$$\begin{aligned} V_h = & m_{11}^2 \Phi_1^\dagger \Phi_1 + m_{22}^2 \Phi_2^\dagger \Phi_2 - \left(m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h. c.} \right) \\ & + \frac{1}{2} \lambda_1 \left(\Phi_1^\dagger \Phi_1 \right)^2 + \frac{1}{2} \lambda_2 \left(\Phi_2^\dagger \Phi_2 \right)^2 \\ & + \lambda_3 \left(\Phi_1^\dagger \Phi_1 \right) \left(\Phi_2^\dagger \Phi_2 \right) + \lambda_4 \left(\Phi_1^\dagger \Phi_2 \right) \left(\Phi_2^\dagger \Phi_1 \right) \\ & + \left[\frac{1}{2} \lambda_5 \left(\Phi_1^\dagger \Phi_2 \right)^2 + \lambda_6 \left(\Phi_1^\dagger \Phi_1 \right) \left(\Phi_1^\dagger \Phi_2 \right) \right. \\ & \left. + \lambda_7 \left(\Phi_2^\dagger \Phi_2 \right) \left(\Phi_1^\dagger \Phi_2 \right) + \text{h. c.} \right], \end{aligned} \tag{4}$$

where m_{11}^2, m_{22}^2 and $\lambda_{1,2,3,4}$ are always real, while m_{12}^2 and $\lambda_{5,6,7}$ are in general complex. Since in this work we want to demonstrate the simplest possible application of RoC on a 2HDM, we choose to consider all of the above-mentioned parameters to be real.

The discrete symmetries introduced in the context of the Type-II scenario (in which u_R^i couple with Φ_2 , while d_R^i and e_R^i couple with Φ_1) ensure that

$$\lambda_6 = \lambda_7, \tag{5}$$

while in order to conserve the electric charge one needs:

$$\lambda_4 < 0. \tag{6}$$

Furthermore, the potential is bounded from below if [27–29]

$$\lambda_1 > 0, \quad \lambda_2 > 0, \quad \sqrt{\lambda_1 \lambda_2} + \lambda_3 + \lambda_4 - |\lambda_5| > 0. \tag{7}$$

4 A first attempt of reduction

Let us now proceed with the reduction of the parameters of the model. As in past reduced models, the best candidate for ‘primary’ coupling is the strong coupling, g_s . A complete reduc-

tion is not realistic, so we focus on the third fermionic generation Yukawa couplings. Furthermore, reducing in favour of a dimensionless parameter only works for dimensionless parameters, so m_{11}^2, m_{22}^2 and m_{12}^2 will remain free.

The gauge couplings g and g' of the $SU(2)$ and $U(1)$ gauge groups, respectively, will not be considered at the first stage of the reduction, but will be treated as corrections. Since the bottom quark and tau lepton Yukawa couplings are much smaller than the top Yukawa coupling, we do not take them into account in the following work for simplicity. However, they can be straightforwardly incorporated into the following reduction scheme in future studies of the model.

First, we have to specify the 1-loop renormalization group equations (RGEs), which were given (for a general gauge theory and for the specific case of two scalar doublets) in [30–35]. For coherency, we follow the notation of [27]. The gauge β -functions for the model are given by:

$$\mathcal{D}g_s = -7g_s^3 \equiv \beta_3 \tag{8}$$

$$\mathcal{D}g = -3g^3 \equiv \beta_2 \tag{9}$$

$$\mathcal{D}g' = 7g'^3 \equiv \beta_1, \tag{10}$$

where \mathcal{D} is the dimensionless differential operator $16\pi^2 \times \mu(d/d\mu)$. The top Yukawa β -function (with the omission of y_b and y_τ) is:

$$\mathcal{D}y_t = \beta_t = \beta_{t_0} + \beta_{t_c}, \tag{11}$$

where

$$\beta_{t_0} = \left(\frac{9}{2}y_t^2 - 8g_s^2\right)y_t \tag{12}$$

$$\beta_{t_c} = \left(-\frac{9}{4}g^2 - \frac{17}{12}g'^2\right)y_t, \tag{13}$$

and it is understood that β_{t_0} is the top Yukawa β -function without the g, g' contributions, which are notated as β_{t_c} . The λ_i β -functions -without the bottom and tau contributions- are given by:

$$\mathcal{D}\lambda_i = \beta_{\lambda_i} = \beta_{\lambda_{i0}} + \beta_{\lambda_{ic}}, \tag{14}$$

where again $\beta_{\lambda_{i0}}$ are the λ_i β -functions without the g, g' contributions $\beta_{\lambda_{ic}}$ and are given as

$$\begin{aligned} \beta_{\lambda_{10}} = & 12\lambda_1^2 + 4\lambda_3^3 + 4\lambda_3\lambda_4 + 2\lambda_4^2 + 2\lambda_5^2 \\ & + 24\lambda_6^2 + 12\lambda_1\lambda_7^2 - 12\lambda_7^4 \end{aligned} \tag{15}$$

$$\beta_{\lambda_{20}} = 12\lambda_2^2 + 4\lambda_3^3 + 4\lambda_3\lambda_4 + 2\lambda_4^2 + 2\lambda_5^2 + 24\lambda_7^2 \tag{16}$$

$$\begin{aligned} \beta_{\lambda_{30}} = & (\lambda_1 + \lambda_2)(6\lambda_3 + 2\lambda_4) + 4\lambda_3^2 + 2\lambda_4^2 + 2\lambda_5^2 \\ & + 4(\lambda_6^2 + \lambda_7^2) + 16\lambda_6\lambda_7 + 6\lambda_3\lambda_7^2 \end{aligned} \tag{17}$$

$$\beta_{\lambda_{40}} = 2(\lambda_1 + \lambda_2)\lambda_4 + 8\lambda_3\lambda_4 + 4\lambda_4^2 + 8\lambda_5^2$$

$$+ 10(\lambda_6^2 + \lambda_7^2) + 4\lambda_6\lambda_7 + 6\lambda_4\lambda_7^2 \tag{18}$$

$$\begin{aligned} \beta_{\lambda_{50}} = & \lambda_5(2\lambda_1 + 2\lambda_2 + 8\lambda_3 + 12\lambda_4) + 10(\lambda_6^2 + \lambda_7^2) \\ & + 4\lambda_6\lambda_7 + 6\lambda_5\lambda_7^2 \end{aligned} \tag{19}$$

$$\begin{aligned} \beta_{\lambda_{60}} = & (12\lambda_1 + 6\lambda_3 + 8\lambda_4)\lambda_6 + (6\lambda_3 + 4\lambda_4)\lambda_7 \\ & + 10\lambda_5\lambda_6 + 2\lambda_5\lambda_7 + 9\lambda_6\lambda_7^2 \end{aligned} \tag{20}$$

$$\begin{aligned} \beta_{\lambda_{70}} = & (12\lambda_1 + 6\lambda_3 + 8\lambda_4)\lambda_7 + (6\lambda_3 + 4\lambda_4)\lambda_6 \\ & + 10\lambda_5\lambda_7 + 2\lambda_5\lambda_6 + 9\lambda_7\lambda_7^2 \end{aligned} \tag{21}$$

and

$$\beta_{\lambda_{1c}} = \frac{3}{4}(3g^4 + g'^4 + 2g^2g'^2) - 3\lambda_1(3g^2 + g'^2) \tag{22}$$

$$\beta_{\lambda_{2c}} = \frac{3}{4}(3g^4 + g'^4 + 2g^2g'^2) - 3\lambda_2(3g^2 + g'^2) \tag{23}$$

$$\beta_{\lambda_{3c}} = \frac{3}{4}(3g^4 + g'^4 - 2g^2g'^2) - 3\lambda_3(3g^2 + g'^2) \tag{24}$$

$$\beta_{\lambda_{4c}} = 3g^2g'^2 - 3\lambda_4(3g^2 + g'^2) \tag{25}$$

$$\beta_{\lambda_{5c}} = -3\lambda_5(3g^2 + g'^2) \tag{26}$$

$$\beta_{\lambda_{6c}} = -3\lambda_6(3g^2 + g'^2) \tag{27}$$

$$\beta_{\lambda_{7c}} = -3\lambda_7(3g^2 + g'^2), \tag{28}$$

where $\lambda_i = y_i \sin \beta$ and $\tan \beta = v_2/v_1$ is the well known ratio of the two Higgs vacuum expectation values (vevs).

Proceeding with the reduction of λ_i and y_t w.r.t. the primary coupling, g_s and temporarily ‘switching off’ the other two gauge couplings, the power series solutions of Eq. (3) will be:

$$y_t = p_t g_s \tag{29}$$

$$\lambda_i = p_i g_s^2. \tag{30}$$

Substituting the solutions into the REs,

$$\beta_3 \frac{dy_t}{dg_s} = \beta_{t_0} \tag{31}$$

$$\beta_3 \frac{d\lambda_i}{dg_s} = \beta_{\lambda_{i0}}, \tag{32}$$

we get sets of p_i, p_t that depend on $\sin \beta$ and are RGI. Sadly, these solutions do not satisfy Eqs. (6) and (7) and, more importantly, for any choice of $\tan \beta$, the top quark pole mass fails to go above 100 GeV.

One may proceed with taking into account the corrections that come from the other two gauge couplings, in hope that their contributions may ameliorate the above results. Now the full power series solutions of Eq. (3) will be:

$$y_t = p_t g_s + q_t g + r_t g' \tag{33}$$

$$\lambda_i = p_i g_s^2 + q_i g^2 + r_i g'^2, \tag{34}$$

where p_t, p_i are known from the above procedure and the corresponding REs will be

$$\beta_3 \frac{dy_t}{dg_s} = \beta_t \quad (35)$$

$$\beta_3 \frac{d\lambda_i}{dg_s} = \beta_{\lambda_i}. \quad (36)$$

In order for Eqs. (35)–(36) to be solved w.r.t. q_t, q_i, r_t, r_i , one needs a further condition such as [24]

$$\mathcal{D}(q_a g) \sim 0, \quad \mathcal{D}(r_a g') \sim 0, \quad (37)$$

where $a = t, 1, \dots, 7$. However, these conditions only hold for $\mu \geq 10^7$ GeV and are not RGI, thus we cannot have a successful reduction of the theory that way.

An extended discussion about ‘traditional’ (partial) RoC method applied directly to the 2HDM, can be found in [36], where a reduction is performed for the Yukawa and Higgs self-couplings in terms of the strong coupling. Using the bottom quark mass as input the Higgs boson and top quark masses are predicted (with values which are now ruled out by experiment), and also values for the lepton masses and the extra Higgs scalars were found, but already in contradiction with the experimental results at the time. More recently, a similar study of the RoC method specifically applied to the four types of 2HDM with Natural Flavour Conservation with updated data was performed in [37], with similar, albeit not identical, results.

5 A realistic approach to reduction

Since the ‘traditional’ partial RoC method proved to be too restrictive, the next step is to try a reduction at a boundary scale, along the lines of [24] (also explained in [38]). The idea is simple: we solve the REs of Eqs. (35)–(36) at one specific scale, called the boundary scale M_{bdry} , above which a covering theory is assumed, which is supposed to make these solutions RGI. Below M_{bdry} we run the usual 2HDM RGEs, using the reduction solutions as boundary conditions. Thus, using only the experimental values of the gauge couplings and by fixing the $\tan \beta$ value, we obtain the top quark pole mass and all λ_i s at the EW scale and, fixing the mass parameters of the scalar potential, we also obtain the light Higgs boson mass.

The conditions of Eq. (37) demand that $M_{\text{bdry}} \geq 10^7$ GeV. However, since we want to treat g, g' as corrections, we need their values not to be comparable to g_s at the boundary scale. Running the 2HDM gauge RGEs of Eqs. (8)–(10) it becomes clear that, while g' continues to be much smaller than the other gauge couplings until very high energies, the weak coupling starts ‘dangerously’ approaching g_s around

$\mu \sim 10^8$ GeV. This naturally restricts the boundary scale at

$$M_{\text{bdry}} \sim 10^7 \text{ GeV}, \quad (38)$$

which is the value used from now on. The reduction of y_t to g_s does not involve any of the Higgs potential parameters and can be performed independently. As such, we reduce the top Yukawa in favour of the strong coupling, first without the contributions of g, g' , as above. Then, in order to solve the RE of Eq. (35) at the boundary M_{bdry} , we use the values $g(M_{\text{bdry}})$ and $g'(M_{\text{bdry}})$ that we get from Eqs. (9)–(10) using their experimental values at M_Z . From three sets of possible reduction solutions at the boundary scale, the only one that can lead to a phenomenologically viable top mass is

$$y_t = 0.471g_s - 0.119g + 1.228g'. \quad (39)$$

Using the above relation as boundary condition to the top Yukawa RGE, we obtain the EW scale top Yukawa value. In order to satisfy the experimental constraint of [39],

$$m_t = (172.69 \pm 0.30) \text{ GeV}, \quad (40)$$

allowing a theoretical uncertainty of 1 GeV, the ratio of the two Higgs vevs has to be:

$$\tan \beta = 2.2 \pm 0.5. \quad (41)$$

Now, with all the above information, we can perform the same reduction to the full system of g_s, y_t and λ_i at M_{bdry} , including the corrections of g, g' . We get a large number of possible solutions, not all of which are phenomenologically viable. Indeed, once we impose the conditions of Eqs. (5)–(7), there are only four sets of reduction solutions that confirm the light Higgs boson mass measurement [39],

$$m_h^{\text{exp}} = (125.25 \pm 0.17) \text{ GeV}. \quad (42)$$

We have estimated that our theoretical calculations have a 5 GeV uncertainty, due to threshold corrections and higher order contributions. For the calculation of the light Higgs mass we have chosen appropriate mass parameters of the Higgs potential such that the mass of the CP odd scalar, m_A , is 800 GeV, which is allowed by recent LHC searches [40]. However, there is significant freedom in this parameter, since a variation of ± 400 GeV in m_A gives a very small change in the value of m_h , which is covered by the theoretical uncertainty. The four reduction solutions are shown in Table 1, while the results for their respective light Higgs masses are given in Table 2. It is obvious from Table 1 and the $\lambda_i \beta$ -functions that every solution has $\lambda_{6,7} = 0$, while λ_5 vanishes for the latter two solutions.

Table 1 The sets of solutions for $M_{\text{bdry}} = 10^7$ GeV and $\tan \beta = 2.2$ that satisfy the conditions of Eqs. (5)–(7) and yield a light Higgs boson mass within 5 GeV of Eq. (42)

#	p_t	p_1	p_2	p_3	p_4	p_5	p_6	p_7
SET1	0.471	− 1.377	− 1.167	0	0	0	0	0
SET2	0.471	− 1.377	− 1.167	0	0	0	0	0
SET3	0.471	− 1.109	− 0.773	− 0.955	0	0	0	0
SET4	0.471	− 1.109	− 0.773	− 0.955	0	0	0	0
#	q_t	q_1	q_2	q_3	q_4	q_5	q_6	q_7
SET1	− 0.119	4.606	4.198	− 0.087	− 0.060	− 0.060	0	0
SET2	− 0.119	4.598	4.189	0.124	− 0.595	0	0	0
SET3	− 0.119	3.652	2.819	3.317	0	0	0	0
SET4	− 0.119	3.652	2.819	3.317	0	0	0	0
#	r_t	r_1	r_2	r_3	r_4	r_5	r_6	r_7
SET1	1.228	10.022	0.498	− 1.033	− 3.275	0	0	0
SET2	1.228	10.197	0.240	− 0.415	− 1.490	− 1.490	0	0
SET3	1.228	7.929	1.245	5.518	− 9.425	0	0	0
SET4	1.228	− 3.196	0.312	8.017	− 8.394	0	0	0

Table 2 The light Higgs boson mass (in GeV) for each set of solutions of Table 1

#	m_h (GeV)
SET1	127.28
SET2	120.87
SET3	121.20
SET4	122.81

This is the first case in which the RoC method is applied on a non-supersymmetric model and successfully fits the experimentally observed values for both the top quark mass and the (light) Higgs boson mass. It assumes new physics at a specific energy scale -either a covering symmetry or just new field content, or both- with RGI relations among parameters. It is only a simplified example of what the method is capable of, as it can be expanded to either be more restrictive or include other phenomena as well. For example, a 2-loop analysis of the above can rule out some of the solutions, while it can also be applied on a complex Higgs potential. In the latter case, it can result in a realistic description of explicit or spontaneous CP violation with minimal input, and it may single out one of the six symmetries of the Higgs potential [27,41] as the one that naturally occurs from a reduced theory. Lastly, although the extension of the SM with one scalar doublet is one of the simplest and most intuitive ways to tackle questions the SM is unable to, the RoC technique can be applied in more field-rich models (or models with larger symmetries like in [16–18,20–22]). The natural continuation of the present work under this perspective is the application of RoC on models that feature three Higgs doublets (see for instance [42–51]). All the above mentioned cases and extensions are the main subject of our future work.

6 Conclusions

In the present work we described the application of the *reduction of couplings* method on the Two Higgs Doublet Model. In particular, the top Yukawa coupling and the quartic Higgs couplings are expressed in terms of the strong gauge coupling, treating the other two gauge couplings as corrections. The 1-loop reduction is performed at a boundary scale M_{bdry} , over which a covering theory is assumed. The demand for phenomenological viability of the model sets the scale over which new physics appear at $M_{\text{bdry}} \sim 10^7$ GeV. The reduction gives four sets of solutions, which fit the experimental limits for the top quark mass and the (light) Higgs boson mass, using as input only the gauge coupling values at M_Z , while it fixes the value of $\tan \beta \sim 2.2$. Thus, the RoC method provides us with a powerful tool to reduce the number of free parameters of a given theory, guiding the direction of possible viable extensions of the Standard Model.

Acknowledgements GP is supported by the Portuguese Fundação para a Ciência e Tecnologia (FCT) under Contracts UIDB/00777/2020, and UIDP/00777/2020, these projects are partially funded through POCTI (FEDER), COMPETE, QREN, and the EU. GP has a post-doctoral fellowship in the framework of UIDP/00777/2020 with reference BL154/2022_IST_ID. GP and GZ would like to thank CERN-TH for the hospitality and support. GZ would like to thank the MPP-Munich and DFG Exzellenzcluster 2181:STRUCTURES of Heidelberg University for support. MM acknowledge support from DGAPA-UNAM through PAPIIT project IN109321. MAMP acknowledges support from a CONACYT scholarship, and from partial scholarships from IF-UNAM project PRIDIF21-3 and PAPIIT project IN109321.

Data availability This manuscript has no associated data or the data will not be deposited. [Authors’ comment: All relative data and results are displayed in the appropriate tables, there is no extra data to deposit.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- J. Kubo, S. Heinemeyer, M. Mondragon, O. Piguet, K. Sibold, W. Zimmermann, G. Zoupanos, Reduction of couplings and its application in particle physics, finite theories, Higgs and top mass predictions, PoS (Higgs & top)001, Ed. K. Sibold (2014). [arXiv:hep-ph/1411.7155](https://arxiv.org/abs/hep-ph/1411.7155)
- J.C. Pati, A. Salam, Is baryon number conserved? *Phys. Rev. Lett.* **31**, 661–664 (1973)
- H. Georgi, S.L. Glashow, Unity of all elementary particle forces. *Phys. Rev. Lett.* **32**, 438–441 (1974)
- H. Georgi, H.R. Quinn, S. Weinberg, Hierarchy of interactions in unified gauge theories. *Phys. Rev. Lett.* **33**, 451–454 (1974)
- H. Fritzsch, P. Minkowski, Unified interactions of leptons and hadrons. *Ann. Phys.* **93**, 193–266 (1975)
- F. Gursev, P. Ramond, P. Sikivie, A universal gauge theory model based on E₆. *Phys. Lett. B* **60**, 177–180 (1976)
- Y. Achiman, B. Stech, Quark lepton symmetry and mass scales in an E₆ unified gauge model. *Phys. Lett. B* **77**, 389–393 (1978)
- W. Zimmermann, Reduction in the number of coupling parameters. *Commun. Math. Phys.* **97**, 211 (1985)
- R. Oehme, W. Zimmermann, Relation between effective couplings for asymptotically free models. *Commun. Math. Phys.* **97**, 569 (1985)
- R. Oehme, Reduction and reparametrization of quantum field theories. *Prog. Theor. Phys. Suppl.* **86**, 215 (1986)
- E. Ma, Modified quantum chromodynamics. I. Exact global color symmetry and asymptotic freedom. *Phys. Rev. D* **17**, 623 (1978)
- N.-P. Chang, Eigenvalue conditions and asymptotic freedom for Higgs scalar gauge theories. *Phys. Rev. D* **10**, 2706 (1974)
- S. Nandi, W.-C. Ng, Can coupling constants be related? *Phys. Rev. D* **20**, 972 (1979)
- J. Kubo, K. Sibold, W. Zimmermann, Higgs and top mass from reduction of couplings. *Nucl. Phys. B* **259**, 331 (1985)
- J. Kubo, K. Sibold, W. Zimmermann, New results in the reduction of the Standard Model. *Phys. Lett. B* **220**, 185 (1989)
- D. Kapetanakis, M. Mondragon, G. Zoupanos, Finite unified models. *Z. Phys. C* **60**, 181–186 (1993). [arXiv:hep-ph/9210218](https://arxiv.org/abs/hep-ph/9210218)
- J. Kubo, M. Mondragon, G. Zoupanos, Reduction of couplings and heavy top quark in the minimal SUSY GUT. *Nucl. Phys. B* **424**, 291–307 (1994)
- M. Mondragon, G. Zoupanos, Finite unified theories and the top quark mass. *Nucl. Phys. Proc. Suppl.* **37C**, 98–105 (1995)
- T. Kobayashi, J. Kubo, M. Mondragon, G. Zoupanos, Constraints on finite soft supersymmetry-breaking terms. *Nucl. Phys. B* **511**, 45–68 (1998). [arXiv:hep-ph/9707425](https://arxiv.org/abs/hep-ph/9707425)
- S. Heinemeyer, M. Mondragon, G. Zoupanos, Confronting finite unified theories with low-energy phenomenology. *JHEP* **07**, 135 (2008). [arXiv:0712.3630](https://arxiv.org/abs/0712.3630)
- S. Heinemeyer, M. Mondragon, G. Patellis, N. Tracas, G. Zoupanos, Updates and new results in models with reduced couplings. *Fortsch. Phys.* **68**(6), 2000028 (2020). [arXiv:2002.10983](https://arxiv.org/abs/2002.10983)
- S. Heinemeyer, J. Kalinowski, W. Kotlarski, M. Mondragon, G. Patellis, N. Tracas, G. Zoupanos, Probing unified theories with reduced couplings at future hadron colliders. *Eur. Phys. J. C* **81**(2), 185 (2021). [arXiv:2011.07900](https://arxiv.org/abs/2011.07900)
- T.D. Lee, A theory of spontaneous T violation. *Phys. Rev. D* **8**, 1226–1239 (1973)
- S. Heinemeyer, M. Mondragon, N. Tracas, G. Zoupanos, Reduction of the parameters in MSSM. *JHEP* **08**, 150 (2018). [arXiv:1712.02729](https://arxiv.org/abs/1712.02729)
- Y.L. Wu, L. Wolfenstein, Sources of CP violation in the two Higgs doublet model. *Phys. Rev. Lett.* **73**, 1762–1764 (1994). [arXiv:hep-ph/9409421](https://arxiv.org/abs/hep-ph/9409421)
- S. Davidson, H.E. Haber, Basis-independent methods for the two-Higgs-doublet model. *Phys. Rev. D* **72**, 035004 (2005). [Erratum: *Phys. Rev. D* **72**, 099902 (2005)]. [arXiv:hep-ph/0504050](https://arxiv.org/abs/hep-ph/0504050)
- G.C. Branco, P.M. Ferreira, L. Lavoura, M.N. Rebelo, M. Sher, J.P. Silva, Theory and phenomenology of two-Higgs-doublet models. *Phys. Rep.* **516**, 1–102 (2012). [arXiv:1106.0034](https://arxiv.org/abs/1106.0034)
- C.T. Hill, C.N. Leung, S. Rao, Renormalization group fixed points and the Higgs boson spectrum. *Nucl. Phys. B* **262**(3), 517–537 (1985)
- I.P. Ivanov, Building and testing models with extended Higgs sectors. *Prog. Part. Nucl. Phys.* **95**, 160–208 (2017)
- T.P. Cheng, E. Eichten, L.-F. Li, Higgs phenomena in asymptotically free gauge theories. *Phys. Rev. D* **9**, 2259 (1974)
- D.R.T. Jones, The two loop beta function for a G(1) x G(2) gauge theory. *Phys. Rev. D* **25**, 581 (1982)
- M.E. Machacek, M.T. Vaughn, Fermion and Higgs masses as probes of unified theories. *Phys. Lett. B* **103**, 427–432 (1981)
- H.E. Haber, R. Hempfling, The renormalization group improved Higgs sector of the minimal supersymmetric model. *Phys. Rev. D* **48**, 4280–4309 (1993). [arXiv:hep-ph/9307201](https://arxiv.org/abs/hep-ph/9307201)
- W. Grimus, L. Lavoura, Renormalization of the neutrino mass operators in the multi-Higgs-doublet standard model. *Eur. Phys. J. C* **39**, 219–227 (2005). [arXiv:hep-ph/0409231](https://arxiv.org/abs/hep-ph/0409231)
- P.M. Ferreira, L. Lavoura, J.P. Silva, Renormalization-group constraints on Yukawa alignment in multi-Higgs-doublet models. *Phys. Lett. B* **688**, 341–344 (2010). [arXiv:1001.2561](https://arxiv.org/abs/1001.2561)
- A. Denner, Reduction of couplings in the two Higgs doublet extension of the electroweak standard model. *Nucl. Phys. B* **347**(1–2), 184–202 (1990)
- M.A. May Pech, Reducción de acoplamientos en modelos multi-Higgs. Master thesis, Universidad Nacional Autónoma de México, Ciudad de México (2023). <http://132.248.9.195/ptd2023/mayo/0840422/Index.html>
- S. Heinemeyer, M. Mondragon, N. Tracas, G. Zoupanos, Reduction of couplings and its application in particle physics. *Phys. Rep.* **814**, 1–43 (2019). [arXiv:1904.00410](https://arxiv.org/abs/1904.00410)
- Particle Data Group** Collaboration, R.L. Workman et al., Review of particle physics. *PTEP* **2022**, 083C01 (2022)
- L. Wang, J.M. Yang, Y. Zhang, Two-Higgs-doublet models in light of current experiments: a brief review. *Commun. Theor. Phys.* **74**(9), 097202 (2022). [arXiv:2203.07244](https://arxiv.org/abs/2203.07244)
- P.M. Ferreira, H.E. Haber, J.P. Silva, Generalized CP symmetries and special regions of parameter space in the two-Higgs-doublet model. *Phys. Rev. D* **79**, 116004 (2009). [arXiv:0902.1537](https://arxiv.org/abs/0902.1537)
- V. Keus, S.F. King, S. Moretti, Three-Higgs-doublet models: symmetries, potentials and Higgs boson masses. *JHEP* **01**, 052 (2014). [arXiv:1310.8253](https://arxiv.org/abs/1310.8253)
- J. Kubo, A. Mondragon, M. Mondragon, E. Rodriguez-Jauregui, The flavor symmetry. *Prog. Theor. Phys.* **109**, 795–807 (2003). [Erratum: *Prog. Theor. Phys.* **114**, 287 (2005)]. [arXiv:hep-ph/0302196](https://arxiv.org/abs/hep-ph/0302196)

44. M. Gómez-Bock, M. Mondragón, A. Pérez-Martínez, Scalar and gauge sectors in the 3-Higgs Doublet Model under the S_3 symmetry. *Eur. Phys. J. C* **81**(10), 942 (2021). [arXiv:2102.02800](#)
45. D. Das, U.K. Dey, Analysis of an extended scalar sector with S_3 symmetry. *Phys. Rev. D* **89**(9), 095025 (2014). [Erratum: *Phys. Rev. D* **91**(3), 039905 (2015)]. [arXiv:1404.2491](#)
46. A. Kunčinas, O.M. Ogreid, P. Osland, M.N. Rebelo, S_3 -inspired three-Higgs-doublet models: a class with a complex vacuum. *Phys. Rev. D* **101**(7), 075052 (2020). [arXiv:2001.01994](#)
47. J. Kalinowski, W. Kotlarski, M.N. Rebelo, I. de Medeiros Varzielas, 3HDM with Δ (27) symmetry and its phenomenological consequences. *JHEP* **02**, 231 (2023). [arXiv:2112.12699](#)
48. A.E. Cárcamo Hernández, C. Espinoza, J.C. Gómez-Izquierdo, J.M. González, M. Mondragón, Predictive extended 3HDM with S_4 family symmetry. [arXiv:2212.12000](#)
49. P.S. Bhupal Dev, A. Pilaftsis, Natural alignment in the two Higgs doublet model. *J. Phys. Conf. Ser.* **873**(1), 012008 (2017). [arXiv:1703.05730](#)
50. N. Darvishi, A. Pilaftsis, Quartic coupling unification in the maximally symmetric 2HDM. *Phys. Rev. D* **99**(11), 115014 (2019). [arXiv:1904.06723](#)
51. N. Darvishi, M.R. Masouminia, A. Pilaftsis, Maximally symmetric three-Higgs-doublet model. *Phys. Rev. D* **104**(11), 115017 (2021). [arXiv:2106.03159](#)