

LHC HIGGS WORKING GROUP^{*}
PUBLIC NOTE

Ad interim recommendations for the Higgs boson production cross sections at $\sqrt{s} = 13.6$ TeV

WG1 conveners: Alexander Karlberg¹, Julie Malcles², Bernhard Mistlberger³, Roberto Di Nardo⁴

ggF conveners: Syed Haider Abidi⁵, Robin Hayes⁶, Alexander Huss¹, Stephen Jones⁷,

VBF conveners: Gaetano Barone⁸, Jiayi Chen⁹, Stephane Cooperstein¹⁰, Silvia Ferrario Ravasio¹, Mathieu Pellen¹¹

VH conveners: Hannah Arnold⁶, Alessandro Calandri¹², Suman Chatterjee¹³, Giancarlo Ferrera¹⁴, Ciaran Williams¹⁵

t \bar{t} H/tH conveners: Małgorzata Worek¹⁶, Marco Zaro¹⁴

b \bar{b} H/bH conveners: Chayanit Asawatangtrakuldee¹⁷, Tim Barklow³, Michael Spira¹⁸, Marius Wiesemann¹⁹

¹ CERN, Theoretical Physics Department, CH-1211 Geneva 23, Switzerland

² IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, F-91191, France

³ SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94039, USA

⁴ Università degli Studi and INFN Roma Tre, Via della Vasca Navale 84, I-00146, Rome, Italy

⁵ Brookhaven National Laboratory, PO Box 5000, Upton, NY 11973, USA

⁶ Nikhef, Science Park 105, 1098 XG Amsterdam, The Netherlands

⁷ Institute for Particle Physics Phenomenology, Durham University, Durham DH1 3LE, UK

⁸ Brown University, Providence, RI 02912, USA

⁹ Department of Physics, Simon Fraser University, Burnaby, BC V5H0G9, Canada

¹⁰ University of California San Diego, 500 Gilman Dr, La Jolla, CA 92093, USA

¹¹ Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, D-79104 Freiburg, Germany

¹² ETH Zurich—Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

¹³ Institute of High Energy Physics (HEPHY), Austrian Academy of Sciences (ÖAW), Nikolsdorfer Gasse 18, 1050 Vienna, Austria

¹⁴ Università degli Studi di Milano & INFN, Sezione di Milano, Via Celoria 16, 20133 Milano, Italy

* <https://twiki.cern.ch/twiki/bin/view/LHCPhysics/LHCWG>

¹⁵*Department of Physics, University at Buffalo, The State University of New York, Buffalo, 14260, USA*

¹⁶*Institute for Theoretical Particle Physics and Cosmology, RWTH Aachen University,
D-52056 Aachen, Germany*

¹⁷*Department of Physics, Faculty of Science, Chulalongkorn University, 10330 Bangkok, Thailand*

¹⁸*Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland*

¹⁹*Max-Planck-Institut für Physik, Föhringer Ring 6, 80805 München, Germany*

Abstract This note documents predictions for the inclusive production cross sections of the Standard Model Higgs boson at the Large Hadron Collider at a centre of mass energy of 13.6 TeV. The predictions here are based on simple extrapolations of previously documented predictions published in the CERN Yellow Report "Deciphering the Nature of the Higgs Sector". The predictions documented in this note should serve as a reference while a more complete and update-to-date derivation of cross section predictions is in progress.

Higgs boson production cross sections at $\sqrt{s} = 13.6$ TeV

This document collects the ad interim recommendation for the Higgs boson inclusive production cross section at $\sqrt{s} = 13.6$ TeV. These are obtained from extrapolations based on the cross-section values computed at different value of m_H and centre of mass energy in the CERN Yellow Report "Deciphering the Nature of the Higgs Sector" (YR4) (CERN-2017-002) [1]. The values have been computed within the relevant subgroups of the WG1 LHC Higgs Working Group. Consequently, the derived numerical predictions are a direct product of the work of YR4 and the references cited therein, and it is therefore recommended both this note and the YR4 are cited whenever the interim predictions are used in experimental analyses.

ggF

The ggF cross sections and uncertainties at different values of m_H are reported in Table 1 and obtained with a linear interpolation of the YR4 [1] cross sections available at $\sqrt{s} = 13$ TeV and $\sqrt{s} = 14$ TeV. The interpolation formula utilised is,

$$\sigma(13.6 \text{ TeV}) = 0.4 \times \sigma(13 \text{ TeV}) + 0.6 \times \sigma(14 \text{ TeV}), \quad (1)$$

where $\sigma(E)$ is the production cross section at centre of mass energy E . We have checked explicitly by computing a few cross sections exactly, that the error introduced by the linear interpolation is below the permille level.

The YR4 numbers incorporate results from:

- QCD corrections in the heavy-top limit at NLO [2, 3, 4], NNLO [6, 7, 8] and N³LO [9, 10, 11].
- Exact finite quark mass effects at NLO [3, 5].
- Approximate top quark mass effects obtained in a $1/m_t$ expansion at NNLO [12, 13, 14, 15].
- Electroweak corrections at NLO [16, 17, 18]
- Approximate mixed QCD-electroweak corrections obtained in the $m_t, m_W, m_Z \gg m_H$ limit [19].

The combination procedure and error prescription are described in Ref. [20], and numbers are produced using iHIXS [21].

VBF

In Ref. [1], the state-of-the-art predictions were obtained at NNLO QCD + NLO EW accuracy. The NNLO QCD corrections were obtained from PROVBFH [22] while the EW corrections have been computed with the help of HAWK [23, 24, 25]. Both types of corrections are obtained in the VBF approximation and do not include s -channel contributions. They are combined using the following formula:

$$\sigma^{\text{VBF}} = \sigma_{\text{NNLO QCD}}^{\text{DIS}}(1 + \delta_{\text{EW}}) + \sigma_\gamma, \quad (2)$$

where δ_{EW} is the relative EW correction and σ_γ the photon-induced contributions. The VBF cross sections at 13.6 TeV for different values of m_H are reported in Table 2. These are obtained upon applying a cubic spline interpolation based on numbers computed in Ref. [1] at various centre of mass energies.

VH

The WH and ZH cross sections at different values of m_H are reported in Table 3 and Table 4 respectively. These are obtained by first performing a quadratic fit to the available YR4 cross-sections at various centre of mass energies and then extracting the value for $\sqrt{s} = 13.6$ TeV. The NNLO-QCD corrections were obtained with VH@NNLO [66, 67, 64, 65, 68] and the NLO-EW with HAWK [63, 25].

$t\bar{t}H$ and tH

The $t\bar{t}H$ and tH (t -channel, s -channel, tWH) cross sections at different values of m_H are reported in Table tables 5 to 8 and are obtained with a linear interpolation of the YR4 cross sections. In YR4, cross-sections for $t\bar{t}H$ and tH have been computed with MADGRAPH5_AMC@NLO [26, 27], as follows:

- $t\bar{t}H$ production is computed at NLO QCD+EW accuracy [28, 29, 30, 31, 32, 33, 34, 35, 27].
- t -channel tH is computed at NLO QCD, as documented in Ref. [36]. In particular, the central value of the cross section is computed in a five-flavour scheme, but the quoted scale uncertainty are the envelope of the scale uncertainty obtained both in the five- and four-flavour scheme; hence, they account also for the flavor-scheme ambiguity.
- t -channel tH is computed at NLO QCD, as documented in Ref. [36].
- tWH is computed at NLO QCD, following Ref. [37]. In particular, the “diagram removal with interference” (DR2) approach, as implemented in the MADSTR plugin [38], is employed to subtract contributions from resonant top quarks which appear at NLO.

It is worth to notice that, since YR4, important theoretical progress has been achieved for these processes, in particular: soft-gluon resummation has been performed for $t\bar{t}H$ at NNLL accuracy [39, 40, 41, 42], and it has been subsequently combined with the EW corrections [43, 44, 45]. More recently, the NNLO cross section has been computed [46], relying on an approximation solely for the two-loop amplitude; finally, EW corrections have been computed for tH production [47]; their effect is rather small (3%), and furthermore they cannot be computed for the s -, t - and tW channels separately.

$b\bar{b}H$

The $b\bar{b}H$ cross sections at different values of m_H are reported in Table 9 and obtained with a linear interpolation of the YR4 cross sections available at $\sqrt{s} = 13.5$ TeV and $\sqrt{s} = 14$ TeV (For $M_H = 125.4$ GeV we used the numbers for 13 TeV.). These are obtained for both the YR4 values and for the NLO+NNLLpart+ $y_b y_t$ predictions that are based on the following results:

- QCD corrections within the 4-flavour scheme (4FS) at NLO including the interference between top- and bottom-Yukawa induced contributions [48, 49, 50].
- QCD corrections within the 5-flavour scheme at NLO [52, 51] and NNLO [53].
- Santander matching of the 4FS and 5FS [54].
- Fully matched results of the 4FS and 5FS using the FONLL [55, 56] and NLO+NNLLpart+ $y_b y_t$ [57, 58] methods.

Since YR4 there have been additional work on the determination of the N³LO corrections to bottom-induced Higgs-boson production in the 5FS that has been matched to the 4FS consistently [59]. Another study pointed out the relevance or dominance of non bottom-Yukawa induced processes as $gg \rightarrow Hg^* \rightarrow Hb\bar{b}$ [60] and several other competing processes that overwhelm the SM bottom-Yukawa induced contribution [61]. The disentanglement of the bottom-Yukawa induced contribution from the other competing processes, however, might be possible by machine-learning techniques [62]. In general, the bottom-Yukawa induced cross sections are relevant for BSM scenarios with an enhanced bottom Yukawa coupling as e.g. supersymmetric extensions or a 2HDM of type II.

Acknowledgments

This work was done on behalf of the LHCHWG.

Table 1: ggF cross sections at the LHC at 13.6 TeV and corresponding scale and PDF+ α_s uncertainties computed according to the PDF4LHC recommendation.

ggF (N3LO QCD + NLO EW)							
m_H [GeV]	σ [pb]	+Th %	-Th %	TH Gaussian %	$\pm(\text{PDF}+\alpha_s)\%$	$\pm\text{PDF}\%$	$\pm\alpha_s\%$
120.00	5.611E+01	+4.7	-6.9	± 4.0	± 3.2	± 1.9	± 2.6
120.50	5.571E+01	+4.7	-6.9	± 4.0	± 3.2	± 1.9	± 2.6
121.00	5.531E+01	+4.7	-6.8	± 3.9	± 3.2	± 1.9	± 2.6
121.50	5.490E+01	+4.7	-6.8	± 3.9	± 3.2	± 1.9	± 2.6
122.00	5.451E+01	+4.6	-6.8	± 3.9	± 3.2	± 1.9	± 2.6
122.50	5.412E+01	+4.6	-6.8	± 3.9	± 3.2	± 1.9	± 2.6
123.00	5.374E+01	+4.6	-6.8	± 3.9	± 3.2	± 1.9	± 2.6
123.50	5.336E+01	+4.6	-6.8	± 3.9	± 3.2	± 1.9	± 2.6
124.00	5.298E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
124.10	5.290E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
124.20	5.283E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
124.30	5.275E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
124.40	5.268E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
124.50	5.260E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
124.60	5.253E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
124.70	5.245E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
124.80	5.238E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
124.90	5.231E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.00	5.223E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.09	5.217E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.10	5.216E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.20	5.209E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.30	5.202E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.38	5.196E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.40	5.194E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.50	5.187E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.60	5.180E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.70	5.172E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.80	5.165E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
125.90	5.158E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
126.00	5.151E+01	+4.6	-6.7	± 3.9	± 3.2	± 1.9	± 2.6
126.50	5.115E+01	+4.5	-6.7	± 3.8	± 3.2	± 1.9	± 2.6
127.00	5.080E+01	+4.5	-6.6	± 3.8	± 3.2	± 1.9	± 2.6
127.50	5.045E+01	+4.5	-6.6	± 3.8	± 3.2	± 1.9	± 2.6
128.00	5.011E+01	+4.5	-6.6	± 3.8	± 3.2	± 1.9	± 2.6
128.50	4.976E+01	+4.5	-6.6	± 3.8	± 3.2	± 1.9	± 2.6
129.00	4.943E+01	+4.5	-6.6	± 3.8	± 3.2	± 1.9	± 2.6
129.50	4.909E+01	+4.5	-6.6	± 3.8	± 3.2	± 1.9	± 2.6
130.00	4.875E+01	+4.5	-6.6	± 3.8	± 3.2	± 1.8	± 2.6

Table 2: VBF cross sections at the LHC at 13.6 TeV and corresponding scale and PDF+ α_s uncertainties computed according to the PDF4LHC recommendation.

VBF ((approx.) NNLO QCD + NLO EW)						
m_H [GeV]	σ [pb]	+Scale %	-Scale %	$\pm(\text{PDF}+\alpha_s)\%$	$\pm\text{PDF}\%$	$\pm\alpha_s\%$
125	4.078	+0.5	-0.3	± 2.1	± 2.1	± 0.5
125.09	4.075	+0.5	-0.3	± 2.1	± 2.1	± 0.5
125.38	4.067	+0.5	-0.3	± 2.1	± 2.1	± 0.5

Table 3: WH cross sections at the LHC at 13.6 TeV and corresponding scale and PDF+ α_s uncertainties computed according to the PDF4LHC recommendation.

$pp \rightarrow WH$ (NNLO QCD + NLO EW)								
m_H [GeV]	σ [pb]	+Scale %	-Scale %	$\pm(\text{PDF}+\alpha_s)\%$	$\pm\text{PDF}\%$	$\pm\alpha_s\%$	W^+H [pb]	W^-H [pb]
125	1.457	+0.4	-0.7	± 1.8	± 1.6	± 0.9	0.8889	0.5677
125.09	1.453	+0.4	-0.7	± 1.8	± 1.6	± 0.9	0.8870	0.5664
125.38	1.442	+0.4	-0.7	± 1.8	± 1.6	± 0.9	0.8801	0.5620

Table 4: ZH cross sections at the LHC at 13.6 TeV and corresponding scale and PDF+ α_s uncertainties computed according to the PDF4LHC recommendation.

$pp \rightarrow ZH$ (NNLO QCD + NLO EW)							
m_H [GeV]	σ [pb]	+Scale %	-Scale %	$\pm(\text{PDF}+\alpha_s)\%$	$\pm\text{PDF}\%$	$\pm\alpha_s\%$	$\sigma(gg \rightarrow ZH)$ [pb]
125	9.439E-01	+3.7	-3.2	± 1.6	± 1.3	± 0.9	1.360E-01
125.09	9.422E-01	+3.8	-3.2	± 1.6	± 1.3	± 0.9	1.359E-01
125.38	9.361E-01	+3.8	-3.2	± 1.6	± 1.3	± 0.9	1.347E-01

Table 5: $t\bar{t}H$ cross sections at the LHC at 13.6 TeV and corresponding scale and PDF+ α_s uncertainties computed according to the PDF4LHC recommendation.

$t\bar{t}H$ (NLO QCD + NLO EW)						
m_H [GeV]	σ [pb]	+Scale %	-Scale %	$\pm(\text{PDF}+\alpha_s)\%$	$\pm\text{PDF}\%$	$\pm\alpha_s\%$
125	$5.700 \cdot 10^{-1}$	+6.0	-9.3	± 3.5	± 3	± 2
125.09	$5.688 \cdot 10^{-1}$	+6.0	-9.3	± 3.5	± 3	± 2
125.38	$5.638 \cdot 10^{-1}$	+6.0	-9.3	± 3.5	± 3	± 2
125.40	$5.634 \cdot 10^{-1}$	+6.0	-9.3	± 3.5	± 3	± 2

Table 6: tH t -channel cross sections at the LHC at 13.6 TeV and corresponding scale and PDF+ α_s uncertainties computed according to the PDF4LHC recommendation. Sum of both tH and $\bar{t}H$ cross sections is given.

tH t-chan (NLO QCD)						
m_H [GeV]	σ [pb]	+Scale %	-Scale %	$\pm(\text{PDF}+\alpha_s)\%$	$\pm\text{PDF}\%$	$\pm\alpha_s\%$
125	$8.362 \cdot 10^{-2}$	+6.5	-14.8	± 3.7	± 3.5	± 1.2
125.09	$8.353 \cdot 10^{-2}$	+6.5	-14.8	± 3.7	± 3.5	± 1.2
125.38	$8.320 \cdot 10^{-2}$	+6.5	-14.8	± 3.7	± 3.5	± 1.2
125.40	$8.317 \cdot 10^{-2}$	+6.5	-14.8	± 3.7	± 3.5	± 1.2

Table 7: tH s -channel cross sections at the LHC at 13.6 TeV and corresponding scale and PDF+ α_s uncertainties computed according to the PDF4LHC recommendation. Sum of both tH and $\bar{t}H$ cross sections is given.

tH s-chan (NLO QCD)						
m_H [GeV]	σ [pb]	+Scale %	-Scale %	$\pm(\text{PDF}+\alpha_s)\%$	$\pm\text{PDF}\%$	$\pm\alpha_s\%$
125	$3.068 \cdot 10^{-3}$	+2.4	-1.7	± 2.2	± 2.2	± 0.3
125.09	$3.064 \cdot 10^{-3}$	+2.4	-1.7	± 2.2	± 2.2	± 0.3
125.38	$3.044 \cdot 10^{-3}$	+2.4	-1.7	± 2.2	± 2.2	± 0.3
125.40	$3.040 \cdot 10^{-3}$	+2.4	-1.7	± 2.2	± 2.2	± 0.3

Table 8: tH W -associated cross sections at the LHC at 13.6 TeV and corresponding scale and PDF+ α_s uncertainties computed according to the PDF4LHC recommendation. Sum of both tH and $\bar{t}H$ cross sections is given.

tWH (NLO QCD)						
m_H [GeV]	σ [pb]	+Scale %	-Scale %	$\pm(\text{PDF}+\alpha_s)\%$	$\pm\text{PDF}\%$	$\pm\alpha_s\%$
125	$1.720 \cdot 10^{-2}$	+5.0	-6.8	± 6.3	± 6.1	± 1.5

Table 9: $b\bar{b}H$ cross sections at the LHC at 13.6 TeV and corresponding scale and PDF+ α_s uncertainties computed according to the PDF4LHC recommendation.

$b\bar{b}H$ (NNLO QCD in 5FS, NLO QCD in 4FS, Santander matching)			
m_H [GeV]	σ [pb]	$+(Scale+PDF+\alpha_s)\%$	$-(Scale+PDF+\alpha_s)\%$
125.0	0.5269	+20.1	-24.0
125.09	0.5257	+20.3	-24.0
125.40	0.5213	+20.4	-24.1

$b\bar{b}H$ (NLO+NNLLpart+ybyt matching)				
m_H [GeV]	σ [pb]	$\pm(Ren.+Fact.+Matching\ Scales)\%$	$+(PDF+\alpha_s+m_b)\%$	$-(PDF+\alpha_s+m_b)\%$
125	0.568	± 5.0	+2.0	-2.1
125.09	0.566	± 5.1	+2.0	-2.0
125.40	0.563	± 4.9	+1.9	-1.9

Bibliography

- [1] D. de Florian *et al.* [LHC Higgs Cross Section Working Group], doi:10.23731/CYRM-2017-002 [arXiv:1610.07922 [hep-ph]].
- [2] S. Dawson, Nucl. Phys. B **359** (1991), 283-300 doi:10.1016/0550-3213(91)90061-2
- [3] D. Graudenz, M. Spira and P. M. Zerwas, Phys. Rev. Lett. **70** (1993), 1372-1375 doi:10.1103/PhysRevLett.70.1372
- [4] A. Djouadi, M. Spira and P. M. Zerwas, Phys. Lett. B **264** (1991), 440-446 doi:10.1016/0370-2693(91)90375-Z
- [5] M. Spira, A. Djouadi, D. Graudenz and P. M. Zerwas, Nucl. Phys. B **453** (1995), 17-82 doi:10.1016/0550-3213(95)00379-7
- [6] C. Anastasiou and K. Melnikov, Nucl. Phys. B **646** (2002), 220-256 doi:10.1016/S0550-3213(02)00837-4 [arXiv:hep-ph/0207004 [hep-ph]].
- [7] R. V. Harlander and W. B. Kilgore, Phys. Rev. Lett. **88** (2002), 201801 doi:10.1103/PhysRevLett.88.201801 [arXiv:hep-ph/0201206 [hep-ph]].
- [8] V. Ravindran, J. Smith and W. L. van Neerven, Nucl. Phys. B **665** (2003), 325-366 doi:10.1016/S0550-3213(03)00457-7 [arXiv:hep-ph/0302135 [hep-ph]].
- [9] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog and B. Mistlberger, JHEP **03** (2015), 091 doi:10.1007/JHEP03(2015)091 [arXiv:1411.3584 [hep-ph]].
- [10] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog and B. Mistlberger, Phys. Lett. B **737** (2014), 325-328 doi:10.1016/j.physletb.2014.08.067 [arXiv:1403.4616 [hep-ph]].
- [11] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog and B. Mistlberger, Phys. Rev. Lett. **114** (2015), 212001 doi:10.1103/PhysRevLett.114.212001 [arXiv:1503.06056 [hep-ph]].
- [12] R. V. Harlander and K. J. Ozeren, JHEP **11** (2009), 088 doi:10.1088/1126-6708/2009/11/088 [arXiv:0909.3420 [hep-ph]].
- [13] A. Pak, M. Rogal and M. Steinhauser, JHEP **02** (2010), 025 doi:10.1007/JHEP02(2010)025 [arXiv:0911.4662 [hep-ph]].
- [14] R. V. Harlander and K. J. Ozeren, Phys. Lett. B **679** (2009), 467-472 doi:10.1016/j.physletb.2009.08.012 [arXiv:0907.2997 [hep-ph]].
- [15] R. V. Harlander, H. Mantler, S. Marzani and K. J. Ozeren, Eur. Phys. J. C **66** (2010), 359-372 doi:10.1140/epjc/s10052-010-1258-x [arXiv:0912.2104 [hep-ph]].
- [16] U. Aglietti, R. Bonciani, G. Degrassi and A. Vicini, Phys. Lett. B **595** (2004), 432-441 doi:10.1016/j.physletb.2004.06.063 [arXiv:hep-ph/0404071 [hep-ph]].
- [17] S. Actis, G. Passarino, C. Sturm and S. Uccirati, Phys. Lett. B **670** (2008), 12-17 doi:10.1016/j.physletb.2008.10.018 [arXiv:0809.1301 [hep-ph]].

- [18] S. Actis, G. Passarino, C. Sturm and S. Uccirati, Nucl. Phys. B **811** (2009), 182-273 doi:10.1016/j.nuclphysb.2008.11.024 [arXiv:0809.3667 [hep-ph]].
- [19] C. Anastasiou, R. Boughezal and F. Petriello, JHEP **04** (2009), 003 doi:10.1088/1126-6708/2009/04/003 [arXiv:0811.3458 [hep-ph]].
- [20] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog, A. Lazopoulos and B. Mistlberger, JHEP **05** (2016), 058 doi:10.1007/JHEP05(2016)058 [arXiv:1602.00695 [hep-ph]].
- [21] F. Dulat, A. Lazopoulos and B. Mistlberger, Comput. Phys. Commun. **233** (2018), 243-260 doi:10.1016/j.cpc.2018.06.025 [arXiv:1802.00827 [hep-ph]].
- [22] M. Cacciari, F. A. Dreyer, A. Karlberg, G. P. Salam and G. Zanderighi, Phys. Rev. Lett. **115** (2015) no.8, 082002 [erratum: Phys. Rev. Lett. **120** (2018) no.13, 139901] doi:10.1103/PhysRevLett.115.082002 [arXiv:1506.02660 [hep-ph]].
- [23] M. Ciccolini, A. Denner and S. Dittmaier, Phys. Rev. D **77** (2008), 013002 doi:10.1103/PhysRevD.77.013002 [arXiv:0710.4749 [hep-ph]].
- [24] M. Ciccolini, A. Denner and S. Dittmaier, Phys. Rev. Lett. **99** (2007), 161803 doi:10.1103/PhysRevLett.99.161803 [arXiv:0707.0381 [hep-ph]].
- [25] A. Denner, S. Dittmaier, S. Kallweit and A. Mück, Comput. Phys. Commun. **195** (2015), 161-171 doi:10.1016/j.cpc.2015.04.021 [arXiv:1412.5390 [hep-ph]].
- [26] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli and M. Zaro, JHEP **07** (2014), 079 doi:10.1007/JHEP07(2014)079 [arXiv:1405.0301 [hep-ph]].
- [27] R. Frederix, S. Frixione, V. Hirschi, D. Pagani, H. S. Shao and M. Zaro, JHEP **07** (2018), 185 [erratum: JHEP **11** (2021), 085] doi:10.1007/JHEP11(2021)085 [arXiv:1804.10017 [hep-ph]].
- [28] W. Beenakker, S. Dittmaier, M. Kramer, B. Plumper, M. Spira and P. M. Zerwas, Phys. Rev. Lett. **87** (2001), 201805 doi:10.1103/PhysRevLett.87.201805 [arXiv:hep-ph/0107081 [hep-ph]].
- [29] L. Reina and S. Dawson, Phys. Rev. Lett. **87** (2001), 201804 doi:10.1103/PhysRevLett.87.201804 [arXiv:hep-ph/0107101 [hep-ph]].
- [30] W. Beenakker, S. Dittmaier, M. Krämer, B. Plumper, M. Spira and P. M. Zerwas, Nucl. Phys. B **653** (2003), 151-203 doi:10.1016/S0550-3213(03)00044-0 [arXiv:hep-ph/0211352 [hep-ph]].
- [31] L. Reina, S. Dawson and D. Wackerth, Phys. Rev. D **65** (2002), 053017 doi:10.1103/PhysRevD.65.053017 [arXiv:hep-ph/0109066 [hep-ph]].
- [32] S. Dawson, L. H. Orr, L. Reina and D. Wackerth, Phys. Rev. D **67** (2003), 071503 doi:10.1103/PhysRevD.67.071503 [arXiv:hep-ph/0211438 [hep-ph]].
- [33] S. Frixione, V. Hirschi, D. Pagani, H. S. Shao and M. Zaro, JHEP **09** (2014), 065 doi:10.1007/JHEP09(2014)065 [arXiv:1407.0823 [hep-ph]].
- [34] Y. Zhang, W. G. Ma, R. Y. Zhang, C. Chen and L. Guo, Phys. Lett. B **738** (2014), 1-5 doi:10.1016/j.physletb.2014.09.022 [arXiv:1407.1110 [hep-ph]].
- [35] S. Frixione, V. Hirschi, D. Pagani, H. S. Shao and M. Zaro, JHEP **06** (2015), 184 doi:10.1007/JHEP06(2015)184 [arXiv:1504.03446 [hep-ph]].
- [36] F. Demartin, F. Maltoni, K. Mawatari and M. Zaro, Eur. Phys. J. C **75** (2015) no.6, 267 doi:10.1140/epjc/s10052-015-3475-9 [arXiv:1504.00611 [hep-ph]].

- [37] F. Demartin, B. Maier, F. Maltoni, K. Mawatari and M. Zaro, Eur. Phys. J. C **77** (2017) no.1, 34 doi:10.1140/epjc/s10052-017-4601-7 [arXiv:1607.05862 [hep-ph]].
- [38] S. Frixione, B. Fuks, V. Hirschi, K. Mawatari, H. S. Shao, P. A. Sunder and M. Zaro, JHEP **12** (2019), 008 doi:10.1007/JHEP12(2019)008 [arXiv:1907.04898 [hep-ph]].
- [39] A. Broggio, A. Ferroglio, B. D. Pecjak, A. Signer and L. L. Yang, JHEP **03** (2016), 124 doi:10.1007/JHEP03(2016)124 [arXiv:1510.01914 [hep-ph]].
- [40] A. Kulesza, L. Motyka, T. Stebel and V. Theeuwes, JHEP **03** (2016), 065 doi:10.1007/JHEP03(2016)065 [arXiv:1509.02780 [hep-ph]].
- [41] A. Broggio, A. Ferroglio, B. D. Pecjak and L. L. Yang, JHEP **02** (2017), 126 doi:10.1007/JHEP02(2017)126 [arXiv:1611.00049 [hep-ph]].
- [42] A. Kulesza, L. Motyka, T. Stebel and V. Theeuwes, Phys. Rev. D **97** (2018) no.11, 114007 doi:10.1103/PhysRevD.97.114007 [arXiv:1704.03363 [hep-ph]].
- [43] A. Kulesza, L. Motyka, D. Schwartländer, T. Stebel and V. Theeuwes, Eur. Phys. J. C **79** (2019) no.3, 249 doi:10.1140/epjc/s10052-019-6746-z [arXiv:1812.08622 [hep-ph]].
- [44] A. Broggio, A. Ferroglio, R. Frederix, D. Pagani, B. D. Pecjak and I. Tsinikos, JHEP **08** (2019), 039 doi:10.1007/JHEP08(2019)039 [arXiv:1907.04343 [hep-ph]].
- [45] A. Kulesza, L. Motyka, D. Schwartländer, T. Stebel and V. Theeuwes, Eur. Phys. J. C **80** (2020) no.5, 428 doi:10.1140/epjc/s10052-020-7987-6 [arXiv:2001.03031 [hep-ph]].
- [46] S. Catani, S. Devoto, M. Grazzini, S. Kallweit, J. Mazzitelli and C. Savoini, Phys. Rev. Lett. **130** (2023) no.11, 111902 doi:10.1103/PhysRevLett.130.111902 [arXiv:2210.07846 [hep-ph]].
- [47] D. Pagani, I. Tsinikos and E. Vryonidou, JHEP **08** (2020), 082 doi:10.1007/JHEP08(2020)082 [arXiv:2006.10086 [hep-ph]].
- [48] S. Dittmaier, M. Krämer and M. Spira, Phys. Rev. D **70** (2004), 074010 doi:10.1103/PhysRevD.70.074010 [arXiv:hep-ph/0309204 [hep-ph]].
- [49] S. Dawson, C. B. Jackson, L. Reina and D. Wackerlo, Phys. Rev. D **69** (2004), 074027 doi:10.1103/PhysRevD.69.074027 [arXiv:hep-ph/0311067 [hep-ph]].
- [50] M. Wiesemann, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni and P. Torrielli, JHEP **02** (2015), 132 doi:10.1007/JHEP02(2015)132 [arXiv:1409.5301 [hep-ph]].
- [51] D. Dicus, T. Stelzer, Z. Sullivan and S. Willenbrock, Phys. Rev. D **59** (1999), 094016 doi:10.1103/PhysRevD.59.094016 [arXiv:hep-ph/9811492 [hep-ph]].
- [52] C. Balazs, H. J. He and C. P. Yuan, Phys. Rev. D **60** (1999), 114001 doi:10.1103/PhysRevD.60.114001 [arXiv:hep-ph/9812263 [hep-ph]].
- [53] R. V. Harlander and W. B. Kilgore, Phys. Rev. D **68** (2003), 013001 doi:10.1103/PhysRevD.68.013001 [arXiv:hep-ph/0304035 [hep-ph]].
- [54] R. Harlander, M. Kramer and M. Schumacher, [arXiv:1112.3478 [hep-ph]].
- [55] S. Forte, D. Napoletano and M. Ubiali, Phys. Lett. B **751** (2015), 331-337 doi:10.1016/j.physletb.2015.10.051 [arXiv:1508.01529 [hep-ph]].
- [56] S. Forte, D. Napoletano and M. Ubiali, Phys. Lett. B **763** (2016), 190-196 doi:10.1016/j.physletb.2016.10.040 [arXiv:1607.00389 [hep-ph]].

- [57] M. Bonvini, A. S. Papanastasiou and F. J. Tackmann, JHEP **11** (2015), 196 doi:10.1007/JHEP11(2015)196 [arXiv:1508.03288 [hep-ph]].
- [58] M. Bonvini, A. S. Papanastasiou and F. J. Tackmann, JHEP **10** (2016), 053 doi:10.1007/JHEP10(2016)053 [arXiv:1605.01733 [hep-ph]].
- [59] C. Duhr, F. Dulat, V. Hirschi and B. Mistlberger, JHEP **08** (2020) no.08, 017 doi:10.1007/JHEP08(2020)017 [arXiv:2004.04752 [hep-ph]].
- [60] N. Deutschmann, F. Maltoni, M. Wiesemann and M. Zaro, JHEP **07** (2019), 054 doi:10.1007/JHEP07(2019)054 [arXiv:1808.01660 [hep-ph]].
- [61] D. Pagani, H. S. Shao and M. Zaro, JHEP **11** (2020), 036 doi:10.1007/JHEP11(2020)036 [arXiv:2005.10277 [hep-ph]].
- [62] C. Grojean, A. Paul and Z. Qian, JHEP **04** (2021), 139 doi:10.1007/JHEP04(2021)139 [arXiv:2011.13945 [hep-ph]].
- [63] A. Denner, S. Dittmaier, S. Kallweit and A. Muck, JHEP **03** (2012), 075 doi:10.1007/JHEP03(2012)075 [arXiv:1112.5142 [hep-ph]].
- [64] O. Brein, R. V. Harlander and T. J. E. Zirke, Comput. Phys. Commun. **184** (2013), 998-1003 doi:10.1016/j.cpc.2012.11.002 [arXiv:1210.5347 [hep-ph]].
- [65] R. V. Harlander, S. Liebler and T. Zirke, JHEP **02** (2014), 023 doi:10.1007/JHEP02(2014)023 [arXiv:1307.8122 [hep-ph]].
- [66] O. Brein, A. Djouadi and R. Harlander, Phys. Lett. B **579** (2004), 149-156 doi:10.1016/j.physletb.2003.10.112 [arXiv:hep-ph/0307206 [hep-ph]].
- [67] L. Altenkamp, S. Dittmaier, R. V. Harlander, H. Rzehak and T. J. E. Zirke, JHEP **02** (2013), 078 doi:10.1007/JHEP02(2013)078 [arXiv:1211.5015 [hep-ph]].
- [68] R. V. Harlander, A. Kulesza, V. Theeuwes and T. Zirke, JHEP **11** (2014), 082 doi:10.1007/JHEP11(2014)082 [arXiv:1410.0217 [hep-ph]].