

CERN-EP-2024-006  
2024/01/29

CMS-BPH-22-002

# Observation of the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ decay

The CMS Collaboration\*

## Abstract

Using proton-proton collision data corresponding to an integrated luminosity of  $140 \text{ fb}^{-1}$  collected by the CMS experiment at  $\sqrt{s} = 13 \text{ TeV}$ , the  $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$  decay is observed for the first time, with a statistical significance exceeding 5 standard deviations. The relative branching fraction, with respect to the  $\Lambda_b^0 \rightarrow \psi(2S) \Lambda$  decay, is measured to be  $\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+) / \mathcal{B}(\Lambda_b^0 \rightarrow \psi(2S) \Lambda) = [3.38 \pm 1.02 \pm 0.61 \pm 0.03]\%$ , where the first uncertainty is statistical, the second is systematic, and the third is related to the uncertainties in  $\mathcal{B}(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$  and  $\mathcal{B}(\Xi^- \rightarrow \Lambda \pi^-)$ .

*Submitted to the European Physical Journal C*



## 1 Introduction

Multibody decays of beauty hadrons present a rich laboratory to search for intermediate resonances in the decay products. When decay products contain a charmonium state, such intermediate resonances could decay into a charmonium meson and a hadron, which could be a manifestation of their exotic nature. An important turning point in exotic spectroscopy was achieved at the LHC, when the LHCb Collaboration reported the observation of statistically significant  $J/\psi p$  pentaquark-like structures in the decay of the lightest beauty baryon  $\Lambda_b^0 \rightarrow J/\psi p K^-$  [1]. Various interpretations of these structures have been proposed, including tightly bound hidden-charm  $[c\bar{c}uud]$  pentaquark states [2, 3], loosely bound molecular baryon-meson states [4–6], or being due to a double triangle singularity [7]. More recently, additional exotic states have been reported by LHCb in the decays  $\Lambda_b^0 \rightarrow J/\psi p K^-$  [8],  $\Xi_b^- \rightarrow J/\psi \Lambda K^-$  [9],  $B_s^0 \rightarrow J/\psi p \bar{p}$  [10], and  $B^- \rightarrow J/\psi \Lambda \bar{p}$  [11]. Up to now, the hidden-charm pentaquark candidates have been reported only in  $J/\psi p$  and  $J/\psi \Lambda$  systems. Investigation of other channels with heavier baryons in the decay products, such as  $\Xi^-$  and  $\Omega^-$ , could unveil the existence of doubly or triply strange pentaquarks [12, 13].

In this paper, we report on the search for the  $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$  decay, where the  $J/\psi \rightarrow \mu^+ \mu^-$ ,  $\Xi^- \rightarrow \Lambda \pi^-$ , and  $\Lambda \rightarrow p \pi^-$  channels are used to reconstruct the intermediate decay products. Charge-conjugate states are implied throughout the text. The measurement of the ratio of branching fractions

$$\mathcal{R} \equiv \frac{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+)}{\mathcal{B}(\Lambda_b^0 \rightarrow \psi(2S) \Lambda)} = \frac{N(\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+)}{N(\Lambda_b^0 \rightarrow \psi(2S) \Lambda)} \frac{\epsilon_{\psi(2S)\Lambda}}{\epsilon_{J/\psi \Xi^- K^+}} \frac{\mathcal{B}(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)}{\mathcal{B}(\Xi^- \rightarrow \Lambda \pi^-)} \quad (1)$$

is also reported, where  $N$  is the measured  $\Lambda_b^0$  yield and  $\epsilon$  is the total efficiency. The normalization channel is chosen to be  $\Lambda_b^0 \rightarrow \psi(2S) \Lambda$ , with the subsequent  $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$  and  $J/\psi \rightarrow \mu^+ \mu^-$  decays, because of its similar decay topology and kinematics to the signal decay, leading to the reduction of many systematic uncertainties. The branching fractions of the intermediate decays  $\mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)$  and  $\mathcal{B}(\Lambda \rightarrow p \pi^-)$  cancel in the ratio. Invariant mass distributions of the three two-body combinations for the signal channel are also presented in order to look for intermediate resonances.

The analysis uses proton-proton (pp) collision data recorded by the CMS experiment in 2016–2018, at  $\sqrt{s} = 13$  TeV, corresponding to an integrated luminosity of  $140 \text{ fb}^{-1}$  [14–16]. Tabulated results are provided in the HEPData record for this analysis [17].

## 2 The CMS detector and simulated event samples

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter, and a brass and scintillator hadron calorimeter, each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity coverage provided by the barrel and endcap detectors. Muons are measured in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Ref. [18].

Muons are measured in the pseudorapidity range  $|\eta| < 2.4$ , with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive-plate chambers. Matching muons to tracks measured in the silicon tracker results in a transverse momentum ( $p_T$ ) resolution for muons with  $p_T$  up to 100 GeV of 1% in the barrel and 3% in the endcaps. The silicon

tracker used in 2016 measured charged particles within the range  $|\eta| < 2.5$ . For nonisolated particles of  $1 < p_T < 10$  GeV and  $|\eta| < 1.4$ , the track resolutions were typically 1.5% in  $p_T$  and 25–90  $\mu\text{m}$  in the transverse impact parameter [19]. At the start of 2017, a new pixel detector was installed [20]; the upgraded tracker measured nonisolated particles of  $1 < p_T < 10$  GeV up to  $|\eta| < 3$  with typical resolutions of 1.5% in  $p_T$  and 20–75  $\mu\text{m}$  in the transverse impact parameter [21].

Events of interest are selected using a two-tiered trigger system [22]. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about 4  $\mu\text{s}$  [23]. The second level, known as the high-level trigger (HLT), consists of a farm of computing processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage. All events used in this analysis are selected by a set of triggers requiring two identified muons of opposite charge plus an additional track to form a secondary vertex, displaced from the region of the pp interactions. The trigger demanded for each muon to have  $p_T > 4$  GeV and to pass within 2 cm of the beam axis. The dimuon system was required to have  $p_T > 6.9$  GeV, invariant mass between 2.9 and 3.3 GeV, a vertex fit probability greater than 10%, a separation of the secondary vertex relative to the beam axis in the transverse plane larger than 3 standard deviations (s.d.), and a cosine of the angle in the transverse plane between the dimuon momentum vector and the vector joining the beam axis and the dimuon vertex greater than 0.9. The additional track was required to have  $p_T > 0.8$  (1.2) GeV and an impact parameter with respect to the beam axis greater than 0 (2) s.d., for data collected in 2016 (2017–2018). Finally, the two muons and the additional track were required to originate from the same vertex with a  $\chi^2$  per degree of freedom (dof) less than 10.

Monte Carlo (MC) simulated event samples are generated with PYTHIA v8.240 [24] using the CP5 underlying event tune [25]. The EVTGEN 1.6.0 [26] program models the beauty baryon decays  $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$  and  $\Lambda_b^0 \rightarrow \psi(2S) \Lambda$  with a phase space decay model, followed by the  $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$  and  $J/\psi \rightarrow \mu^+ \mu^-$  decays. Final-state radiation is included in EVTGEN using PHOTOS 3.61 [27]. The events are then passed through a detailed GEANT4-based simulation [28] of the CMS detector, including also the decays of long-lived hyperons  $\Xi^- \rightarrow \Lambda \pi^-$  and  $\Lambda \rightarrow p \pi^-$ , followed by the trigger and reconstruction algorithms identical to those used for the collision data. The simulation includes additional interactions due to multiple pp collisions in each bunch crossing, with the same distribution as observed in the experiment.

### 3 Event reconstruction and selection

The reconstruction for all the decays considered in this analysis starts by finding two muons of opposite charge, which must match those that triggered the event readout and pass the soft-muon identification criteria [29]. The offline selection for both muons requires  $p_T(\mu^\pm) > 3$  GeV,  $|\eta(\mu^\pm)| < 2.4$ ,  $\chi^2$  fit probability to a common dimuon vertex  $P_{\text{vtx}}(\mu^+ \mu^-) > 1\%$ , and dimuon invariant mass  $2.9 < m(\mu^+ \mu^-) < 3.3$  GeV.

The  $\Lambda \rightarrow p \pi^-$  candidates are selected from displaced two-prong vertices as described in Ref. [30]. The track with the higher momentum is assumed to be the proton one, and together with the pion track it is fit to a common vertex with their invariant mass constrained to the known  $\Lambda$  hyperon mass of  $m_{\text{PDG}}(\Lambda) = 1115.683$  MeV [31]. The  $\chi^2$  fit probability for the  $\Lambda$  vertex is required to be  $P_{\text{vtx}}(p \pi^-) > 1\%$ .

For the signal channel, to form the  $\Xi^- \rightarrow \Lambda \pi^-$  candidates, an additional high-purity [19] track

assumed to be a pion is selected with  $p_T > 0.2 \text{ GeV}$ . This track and the selected  $\Lambda$  candidate are then fit to a common vertex with the  $\Lambda\pi^-$  mass constrained to the known  $\Xi^-$  hyperon mass of  $m_{\text{PDG}}(\Xi^-) = 1321.71 \text{ MeV}$  [31]. To form the  $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$  candidate, a high-purity track is chosen with an assigned kaon mass and  $p_T(K^+) > 1.2 \text{ GeV}$ , which aligns with the HLT  $p_T$  requirement. The final reconstruction step in the signal channel is the  $\mu^+\mu^-\Xi^-K^+$  vertex fit with a  $\chi^2$  probability above 1%, where the dimuon mass is constrained to the world-average  $J/\psi$  meson mass of  $3096.9 \text{ MeV}$  [31].

For the normalization channel, two high-purity tracks of opposite charges with  $p_T > 0.4 \text{ GeV}$ , assumed to be pions from the  $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$  decay, are selected. One of them is required to have  $p_T > 1.2 \text{ GeV}$  to match the HLT  $p_T$  requirement. The  $\Lambda_b^0$  candidates are obtained by a vertex fit of the  $\mu^+\mu^-\pi^+\pi^-\Lambda$  system with a  $J/\psi \rightarrow \mu^+\mu^-$  mass constraint, as for the signal channel. The invariant mass of the  $J/\psi\pi^+\pi^-$  candidates is required to be in the range  $3.60 < m(J/\psi\pi^+\pi^-) < 3.95 \text{ GeV}$ .

From all reconstructed pp collision points in each event, the primary vertex (PV) is chosen as the one with the smallest  $\Lambda_b^0$  pointing angle, which is the angle between the momentum of the  $\Lambda_b^0$  candidate and the vector from the PV to the reconstructed  $\Lambda_b^0$  candidate vertex. If any of the tracks used in the  $\Lambda_b^0$  candidate reconstruction were included in the fit of the chosen PV, they are removed, and the PV is refitted.

Selection criteria for the signal channel  $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$  are optimized using the Punzi figure of merit [32]. The signal efficiency is evaluated using simulated event samples. Estimation of the background yield involves combining the collision data from the  $\Lambda_b^0$  mass sideband, excluding the signal region which spans twice the mass resolution around the known  $\Lambda_b^0$  mass. Additionally, the wrong-sign candidates ( $J/\psi\Xi^-K^-$  and  $J/\psi\Xi^+K^+$ ) from the full mass range are included, after ensuring that the shape of the mass distribution of the wrong-sign candidates matches that of the correct-sign ones. Combining these two background sources reduces the impact of the statistical uncertainty in the optimization procedure. The variables used in the optimization include the  $p_T$  of all decay products; the flight length significance in the transverse plane of the  $\Lambda_b^0$ ,  $\Lambda$ , and  $\Xi^-$  baryon candidates and the corresponding pointing angles; the vertex fit probabilities; and the mass windows for hyperon candidates. The resulting criteria are summarized in Table 1. The background is reduced by a factor of 15 after the optimization, whereas the signal efficiency is 70% of the initial selection described above. It has been verified that very similar performance is obtained when reshuffling the order in which the selection criteria are optimized. The selection criteria in the normalization channel  $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$  are chosen to be the same, wherever possible, as in the signal channel, to reduce the systematic uncertainties. The  $J/\psi\pi^+\pi^-$  mass is required to be within  $11.1 \text{ MeV}$  of the known  $\psi(2S)$  meson mass of  $3686.1 \text{ MeV}$  [31], which corresponds to approximately 2.5 times the mass resolution.

For the measurement of  $\mathcal{R}$  defined in Eq. (1), the pion from the  $\Xi^-$  decay is required to have  $p_T > 0.4 \text{ GeV}$ . Additionally, the HLT requirements are repeated offline by requiring  $p_T(\mu) > 4 \text{ GeV}$ ,  $p_T(J/\psi) > 6.9 \text{ GeV}$ ,  $P_{\text{vtx}}(\mu^+\mu^-) > 5\%$ , and track (kaon for the signal channel, the harder of the two pions in the normalization channel) impact parameter above 2 s.d. with respect to the PV. These extra criteria ensure that events from potentially inadequately modeled phase space regions are avoided, as the reliability of the efficiency evaluation from simulated samples in those regions is questionable. Nevertheless, the reconstruction algorithm works reliably in those regions, and thus the corresponding events are used to study the mass distribution, as discussed in the following section.

In less than 5% of the events, multiple  $\Lambda_b^0$  candidates in the same channel are found; all of them are used in the analysis.

Table 1: Optimized selection criteria for the signal decay mode  $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ . The first two requirements are applied using the momenta before the corresponding mass constraint.

Variable	Selection
$ m(p\pi^-) - m_{\text{PDG}}(\Lambda) $	$< 8 \text{ MeV}$
$ m(\Lambda\pi^-) - m_{\text{PDG}}(\Xi^-) $	$< 6 \text{ MeV}$
$p_T(\Lambda_b^0)$	$> 11.5 \text{ GeV}$
$p_T(J/\psi)$	$> 6.5 \text{ GeV}$
$p_T(\Xi^-)$	$> 2.6 \text{ GeV}$
$p_T(\Lambda)$	$> 2.2 \text{ GeV}$
$p_T(K^+)$	$> 1.2 \text{ GeV}$
$\mu^+\mu^-\Xi^-K^+$ vertex fit probability	$> 5\%$
$\Lambda\pi^-$ vertex fit probability	$> 5\%$
$p\pi^-$ vertex fit probability	$> 1\%$
$\Xi^-$ vertex displacement from $\Lambda_b^0$ vertex	$> 3 \text{ s.d.}$
$\Lambda$ vertex displacement from $\Xi^-$ vertex	$> 0 \text{ s.d.}$
$\Lambda_b^0$ vertex displacement from PV	$> 3 \text{ s.d.}$
Angle between $\Xi^-$ momentum and displacement	$< 0.0447 \text{ rad}$
Angle between $\Lambda$ momentum and displacement	$< 0.14 \text{ rad}$
Angle between $\Lambda_b^0$ momentum and displacement	$< 0.0447 \text{ rad}$
PV impact parameter for pion from $\Xi^-$ decay	$> 0.4 \text{ s.d.}$
PV impact parameter for kaon	$> 0.4 \text{ s.d.}$

## 4 Invariant mass distributions

The measured mass distribution of the  $\psi(2S)\Lambda$  candidates is shown in Fig. 1 (left) together with the results of an unbinned maximum likelihood fit. We model the signal with a Student's  $t$ -distribution [33] with all parameters (mean,  $\sigma$ ,  $n$ ) free. The combinatorial background is described by an exponential function with a free slope parameter and normalization. The fitted mass of  $5619.3 \pm 0.3 \text{ MeV}$  is in agreement with the world-average  $\Lambda_b^0$  mass of  $5619.60 \pm 0.17 \text{ MeV}$  [31], and the mass resolution of  $8.90 \pm 0.40 \text{ MeV}$  is slightly larger than, yet in agreement with, its value of  $8.52 \text{ MeV}$  found in simulation. The measured yield is  $N(\Lambda_b^0 \rightarrow \psi(2S)\Lambda) = 1744 \pm 63$ . The  $\chi^2$  between the binned distribution and the fit function is 76.6 for 94 degrees of freedom, demonstrating the good quality of the fit.

The measured invariant mass distribution of the selected  $J/\psi\Xi^-K^+$  candidates is shown in




Figure 1: Measured  $\psi(2S)\Lambda$  (left) and  $J/\psi\Xi^-K^+$  (right) invariant mass distributions and overlaid fit results.

Fig. 1 (right). A narrow peak at the  $\Lambda_b^0$  mass is seen on top of a smooth background. The  $\Lambda_b^0$  signal is modeled with a Student's  $t$ -distribution with mean and  $\sigma$  floating, but the  $n$  parameter fixed to the value found by fitting the simulated distribution, because of the limited signal yield of  $N(\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+) = 46 \pm 11$ . The background is fitted with an exponential function. The  $\Lambda_b^0$  mass returned by the fit ( $5625.9 \pm 3.2$  MeV) is within 2 s.d. of the world-average value [31]. The width of the signal peak ( $\sigma$ ) is found to be  $10.4 \pm 3.3$  MeV, consistent within 1.2 s.d. with the value found in simulation,  $6.6 \pm 0.2$  MeV. The fit quality is good, as demonstrated by the  $\chi^2/\text{dof} = 30.1/45$  for the binned distribution.

The signal significance is evaluated using the likelihood ratio technique by applying the background-only and signal-plus-background hypotheses. In these two fits, a Gaussian constraint is applied on the background shape parameter to the one obtained from a fit to the wrong-sign data. Similarly, a Gaussian constraint is applied to the signal shape parameter  $n$  (from simulation) and the resolution  $\sigma = \sigma_{\text{MC}} (8.90/8.52)$ . The correction factor is extracted from the normalization channel and accounts for the difference in the widths of the peak between the measured and simulated event samples. The mean value of the peak is also Gaussian-constrained with a central value and uncertainty equal to the known  $\Lambda_b^0$  mass and its uncertainty [31], respectively. The fit with the signal-plus-background model with these constraints returns a signal yield of  $36 \pm 8$  and is presented in Appendix A. Since the conditions of Wilks' theorem [34] are satisfied, the asymptotic formulae of Ref. [35] (Eqs. (12) and (52)) are used to determine the  $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$  signal significance, which is found to be 5.8 standard deviations. To evaluate the effect of the choice of the model for fitting the signal significance, several alternative models of signal and background were tested, including double-Gaussian or Johnson [36] functions for the signal and a second-degree polynomial or a modified threshold function for the background. An alternative without a constraint on the background shape was also tested. The significance obtained with the alternative models varies in the range from 5.3 to 5.9 standard deviations. This allows us to claim the first observation of the  $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$  decay.

The sensitivity of this analysis to potential pentaquark signals in the intermediate invariant mass distributions of the  $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$  decay is limited by the low signal yield. The background-subtracted two-body invariant mass distributions, obtained with the  $s\mathcal{P}$ Plot technique [37], are shown in Fig. 2. The distributions do not show any narrow peaks and agree, within uncertainties, with the predictions from the phase space simulation. The distributions are also consistent with the results of extracting the yields by fitting the  $\Lambda_b^0$  signal in each of the five intermediate invariant mass bins.




Figure 2: Intermediate invariant mass distributions of the  $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$  decay. The filled circles and empty squares show the measured background-subtracted distributions and the results from the simulation with a phase-space model, respectively.

For the measurement of  $\mathcal{R}$  (Eq. (1)), more stringent requirements are used, as explained at the end of Section 3, and the measured signal yields decrease to  $1179 \pm 47$  and  $23 \pm 7$  for the




Figure 3: Measured  $\psi(2S)\Lambda$  (left) and  $J/\psi\Xi^-K^+$  (right) invariant mass distributions and corresponding fits used for the measurement of  $\mathcal{R}$ .

$\Lambda_b^0 \rightarrow \psi(2S)\Lambda$  and  $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$  channels, respectively, using unconstrained fits as for Fig 1. These are the baseline results referred to in Section 6. The corresponding mass distributions and fits are presented in Fig. 3.

## 5 Efficiencies

Efficiencies for the signal and normalization channels are calculated using simulated event samples. The total efficiency is calculated by factorizing into two components: detector acceptance and a combined trigger, reconstruction, and selection efficiency.

As only the ratio of the total efficiencies is needed to measure  $\mathcal{R}$ , the systematic uncertainties associated with the muon and track reconstruction are reduced. The obtained efficiency ratio is  $\epsilon_{\psi(2S)\Lambda} / \epsilon_{J/\psi\Xi^-K^+} = 5.06 \pm 0.29$ , where the uncertainty reflects the limited size of the simulated samples. Efficiencies for different years of data-taking are combined with weights corresponding to the integrated luminosity collected in each year. The efficiency for the  $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$  channel is significantly lower than that for the  $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$  channel for several reasons including the low energy release in the  $\Xi^- \rightarrow \Lambda\pi^-$  decay, resulting in a low-momentum pion track.

## 6 Systematic uncertainties

Many systematic uncertainties, related to the muon reconstruction and identification as well as to the trigger efficiency, partially cancel in the measured ratios. Since the signal and normalization channels have the same number of tracks in the final state, most uncertainties related to track reconstruction also cancel in the measured ratio  $\mathcal{R}$ . However, the  $p_T$  spectrum of kaons from the  $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$  decay is observed to differ from that of the highest- $p_T$  pion in the  $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$  channel used for normalization. Despite the signal and normalization channels having the same number of final-state tracks, an uncertainty of 2.3% [38] is included, which reflects the difference in tracking efficiency between the measured and simulated event samples. The MC event samples are validated using the normalization channel by comparing the measured distributions of variables used in the event selection, after background subtraction, to those found in simulation; no significant discrepancies are found in most of the distributions. A small discrepancy was observed in the  $p_T(\Lambda_b^0)$  distribution, and the MC event samples for both channels were reweighted using  $p_T(\Lambda_b^0)$ -dependent weights so that the  $p_T(\Lambda_b^0)$  distribution in



the weighted simulation sample matches the background-subtracted distribution measured in the  $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$  channel. The efficiency ratio evaluated using these weighted MC samples is found to be  $\epsilon_{\psi(2S)\Lambda}/\epsilon_{J/\psi\Xi^-K^+} = 4.82 \pm 0.39$ , which is lower than 4.7%, yet still in agreement with the value reported in the previous section. An uncertainty of 4.7% is assigned to account for potential mismodeling of the  $p_T(\Lambda_b^0)$  spectrum.

The systematic uncertainty related to the choice of the signal model is evaluated by testing three different models. For the normalization channel  $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$  the signal shape parameters are floating, while for the signal channel  $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$  the mass resolution parameters are fixed to those found in simulation, after correcting the width of the peak for the ratio between the two resolutions in the measured and simulated event samples evaluated in the normalization channel. The tested models simultaneously vary the signal and normalization channels and use a Student's  $t$ -distribution, a double-Gaussian, and a Johnson function [36] to model the signal. The largest deviation in the ratio of the  $\Lambda_b^0$  signal yields from the baseline value is taken as the systematic uncertainty.

The systematic uncertainty related to the choice of the background model is estimated in a similar way, with three alternative models: a second-degree polynomial, a threshold function [39, 40] multiplied by an exponential, and a threshold function multiplied by a first-degree polynomial.

The decay  $\Lambda_b^0 \rightarrow J/\psi\pi^+\pi^-\Lambda$  can proceed through the  $\psi(2S)$  resonance as well as via other intermediate resonances, or as a nonresonant four-body decay, even if  $m(J/\psi\pi^+\pi^-)$  is required to be in a narrow mass window around the  $\psi(2S)$  meson mass [31]. In order to look for a possible non- $\psi(2S)$  contribution to the  $\Lambda_b^0 \rightarrow J/\psi\pi^+\pi^-\Lambda$  decays, we have studied the background-subtracted mass distribution of  $J/\psi\pi^+\pi^-$ , obtained by the  $s\mathcal{P}$ lot technique, corresponding to the signal  $\Lambda_b^0$  peak. The non- $\psi(2S)$  background contribution in the baseline  $J/\psi\pi^+\pi^-$  mass region window is evaluated to be no more than 2.5%, which is taken as a systematic uncertainty.

The uncertainty in the efficiency ratio due to the limited size of the simulated samples, calculated to be 5.6% in Section 5, is considered as a systematic uncertainty.

In order to assess the reliability of the efficiency evaluation from the simulated samples, the selection criteria on muon and  $J/\psi$   $p_T$ , dimuon vertex probability, track impact parameter, and  $p_T$  of the soft pion from  $\Xi^-$  decay are tightened, one at a time, until the signal efficiency decreases by 10 or 20% with respect to that obtained with the selection used for the  $\mathcal{R}$  measurement. The analysis is repeated each time, and the value of  $\mathcal{R}$  is re-calculated and compared to the baseline  $\mathcal{R}$  value. The differences ( $d$ ) between the two values and their uncertainties ( $\delta d$ ), which also account for the correlation between the two values, are evaluated. The largest value of  $\sqrt{d^2 - (\delta d)^2}$  among the different variations of the selection criteria is found to be 14.3% and is used as the systematic uncertainty in the efficiency ratio.

Table 2 summarizes the previously discussed systematic uncertainties in the ratio  $\mathcal{R}$ . The total uncertainty is calculated as the sum in quadrature of the individual sources.

## 7 Branching fraction ratio measurement

The branching fraction of the newly observed  $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$  decay, with respect to the  $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$  one, is measured using Eq. (1) to be

$$\mathcal{R} \equiv \frac{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+)}{\mathcal{B}(\Lambda_b^0 \rightarrow \psi(2S)\Lambda)} = [3.38 \pm 1.02 \text{ (stat)} \pm 0.61 \text{ (syst)} \pm 0.03 \text{ (}\mathcal{B}\text{)}] \%,$$

Table 2: The relative systematic uncertainties in the measurement of  $\mathcal{R}$ .

Source	Uncertainty (%)
Tracking efficiency	2.3
$p_T(\Lambda_b^0)$ spectrum	4.7
Signal model	3.9
Background model	6.7
Non- $\psi(2S)$ contribution	2.5
Limited size of MC samples	5.6
Selection efficiency	14.3
Total	18.2

where the last uncertainty is related to the uncertainties in the branching fractions  $\mathcal{B}(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 34.68 \pm 0.30\%$  and  $\mathcal{B}(\Xi^- \rightarrow \Lambda\pi^-) = 99.887 \pm 0.035\%$  [31].

## 8 Summary

The  $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$  decay is observed with a significance exceeding 5 standard deviations using  $\sqrt{s} = 13$  TeV proton-proton collision data corresponding to an integrated luminosity of  $140 \text{ fb}^{-1}$  collected by the CMS experiment. The branching fraction is measured with respect to the  $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$  decay to be  $\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+)/\mathcal{B}(\Lambda_b^0 \rightarrow \psi(2S)\Lambda) = [3.38 \pm 1.02 (\text{stat}) \pm 0.61 (\text{syst}) \pm 0.03 (\mathcal{B})]\%$ . The distributions of intermediate invariant masses  $m(J/\psi\Xi^-)$ ,  $m(J/\psi K^+)$ , and  $m(\Xi^-K^+)$  from the  $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$  decay are also presented. This is the first discovered multibody decay containing the  $J/\psi\Xi^-$  system, which opens the possibility to search for doubly-strange hidden-charm pentaquarks when more data are collected. The new results are important for understanding the strong interaction processes in hadronic decays of beauty baryons and the possible formation of exotic multi-quark states.

## Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TEN-

MAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the "Excellence of Science – EOS" – be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), under Germany's Excellence Strategy – EXC 2121 "Quantum Universe" – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 124845, K 124850, K 128713, K 128786, K 129058, K 131991, K 133046, K 138136, K 143460, K 143477, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Center for High Performance Computing, Big Data and Quantum Computing, funded by the EU NexGeneration program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF "a way of making Europe", and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B37G660013 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

## References

- [1] LHCb Collaboration, "Observation of  $J/\psi p$  resonances consistent with pentaquark states in  $\Lambda_b^0 \rightarrow J/\psi p K^-$  decays", *Phys. Rev. Lett.* **115** (2015) 072001, doi:10.1103/PhysRevLett.115.072001, arXiv:1507.03414.
- [2] L. Maiani, A. D. Polosa, and V. Riquer, "The new pentaquarks in the diquark model", *Phys. Lett. B* **749** (2015) 289, doi:10.1016/j.physletb.2015.08.008, arXiv:1507.04980.
- [3] E. Santopinto and A. Giachino, "Compact pentaquark structures", *Phys. Rev. D* **96** (2017) 014014, doi:10.1103/PhysRevD.96.014014, arXiv:1604.03769.
- [4] L. Roca, J. Nieves, and E. Oset, "LHCb pentaquark as a  $\bar{D}^* \Sigma_c - \bar{D}^* \Sigma_c^*$  molecular state", *Phys. Rev. D* **92** (2015) 094003, doi:10.1103/PhysRevD.92.094003, arXiv:1507.04249.

- 
- [5] M. I. Eides, V. Y. Petrov, and M. V. Polyakov, “Pentaquarks with hidden charm as hadroquarkonia”, *Eur. Phys. J. C* **78** (2018) 36, doi:10.1140/epjc/s10052-018-5530-9, arXiv:1709.09523.
- [6] M.-L. Du et al., “Interpretation of the LHCb  $P_c$  states as hadronic molecules and hints of a narrow  $P_c(4380)$ ”, *Phys. Rev. Lett.* **124** (2020) 072001, doi:10.1103/PhysRevLett.124.072001, arXiv:1910.11846.
- [7] S. X. Nakamura, “ $P_c(4312)^+$ ,  $P_c(4380)^+$ , and  $P_c(4457)^+$  as double triangle cusps”, *Phys. Rev. D* **103** (2021) 111503, doi:10.1103/PhysRevD.103.L111503, arXiv:2103.06817.
- [8] LHCb Collaboration, “Observation of a narrow pentaquark state,  $P_c(4312)^+$ , and of two-peak structure of the  $P_c(4450)^+$ ”, *Phys. Rev. Lett.* **122** (2019) 222001, doi:10.1103/PhysRevLett.122.222001, arXiv:1904.03947.
- [9] LHCb Collaboration, “Evidence of a  $J/\psi\Lambda$  structure and observation of excited  $\Xi^-$  states in the  $\Xi_b^- \rightarrow J/\psi\Lambda K^-$  decay”, *Sci. Bull.* **66** (2021) 1278, doi:10.1016/j.scib.2021.02.030, arXiv:2012.10380.
- [10] LHCb Collaboration, “Evidence for a new structure in the  $J/\psi p$  and  $J/\psi\bar{p}$  systems in  $B_s^0 \rightarrow J/\psi p\bar{p}$  decays”, *Phys. Rev. Lett.* **128** (2022) 062001, doi:10.1103/PhysRevLett.128.062001, arXiv:2108.04720.
- [11] LHCb Collaboration, “Observation of a  $J/\psi\Lambda$  resonance consistent with a strange pentaquark candidate in  $B^- \rightarrow J/\psi\Lambda\bar{p}$  decays”, *Phys. Rev. Lett.* **131** (2023) 031901, doi:10.1103/PhysRevLett.131.031901, arXiv:2210.10346.
- [12] K. Azizi, Y. Sarac, and H. Sundu, “Investigation of hidden-charm double strange pentaquark candidate  $P_{css}$  via its mass and strong decays”, *Eur. Phys. J. C* **82** (2022) 543, doi:10.1140/epjc/s10052-022-10495-7, arXiv:2112.15543.
- [13] F.-L. Wang, X.-D. Yang, R. Chen, and X. Liu, “Hidden-charm pentaquarks with triple strangeness due to the  $\Omega_c^{(*)}\bar{D}_s^{(*)}$  interactions”, *Phys. Rev. D* **103** (2021) 054025, doi:10.1103/PhysRevD.103.054025, arXiv:2101.11200.
- [14] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at  $\sqrt{s} = 13$  TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [15] CMS Collaboration, “CMS luminosity measurement for the 2017 data-taking period at  $\sqrt{s} = 13$  TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-17-004, 2018.
- [16] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at  $\sqrt{s} = 13$  TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
- [17] HEPData record for this analysis, 2024. doi:10.17182/hepdata.145642.
- [18] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [19] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.

- [20] CMS Tracker Collaboration, “The CMS Phase-1 pixel detector upgrade”, *JINST* **16** (2021) P02027, doi:10.1088/1748-0221/16/02/P02027, arXiv:2012.14304.
- [21] CMS Collaboration, “Track impact parameter resolution for the full pseudo rapidity coverage in the 2017 dataset with the CMS Phase-1 pixel detector”, CMS Detector Performance Note CMS-DP-2020-049, 2020.
- [22] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [23] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at  $\sqrt{s} = 13$  TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [24] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [25] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
- [26] D. J. Lange, “The EvtGen particle decay simulation package”, *Nucl. Instrum. Meth. A* **462** (2001) 152, doi:10.1016/S0168-9002(01)00089-4.
- [27] E. Barberio and Z. Waş, “PHOTOS — a universal Monte Carlo for QED radiative corrections: version 2.0”, *Comput. Phys. Commun.* **79** (1994) 291, doi:10.1016/0010-4655(94)90074-4.
- [28] GEANT4 Collaboration, “GEANT4 — a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [29] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at  $\sqrt{s} = 13$  TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [30] CMS Collaboration, “CMS tracking performance results from early LHC operation”, *Eur. Phys. J. C* **70** (2010) 1165, doi:10.1140/epjc/s10052-010-1491-3, arXiv:1007.1988.
- [31] Particle Data Group Collaboration, “Review of particle physics”, *Prog. Theor. Exp. Phys.* **2022** (2022) 083C01, doi:10.1093/ptep/ptac097.
- [32] G. Punzi, “Sensitivity of searches for new signals and its optimization”, in *Proceedings of PHYSTAT 2003, Statistical problems in particle physics, astrophysics and cosmology*, p. MODT002. 2003. arXiv:physics/0308063. eConf C030908.
- [33] S. Jackman, “Bayesian analysis for the social sciences”. John Wiley & Sons, New Jersey, USA, 2009. doi:10.1002/9780470686621.
- [34] S. S. Wilks, “The large-sample distribution of the likelihood ratio for testing composite hypotheses”, *Annals Math. Statist.* **9** (1938) 60, doi:10.1214/aoms/1177732360.
- [35] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* **71** (2011) 1554, doi:10.1140/epjc/s10052-011-1554-0, arXiv:1007.1727. [Erratum: doi:10.1140/epjc/s10052-013-2501-z].

- [36] N. L. Johnson, “Systems of frequency curves generated by methods of translation”, *Biometrika* **36** (1949) 149, doi:10.2307/2332539.
- [37] M. Pivk and F. R. Le Diberder, “ $\mathcal{P}$ lot: a statistical tool to unfold data distributions”, *Nucl. Instrum. Meth. A* **555** (2005) 356, doi:10.1016/j.nima.2005.08.106, arXiv:physics/0402083.
- [38] CMS Collaboration, “Tracking performances for charged pions with Run2 legacy data”, CMS Detector Performance Note CMS-DP-2022-012, 2022.
- [39] CMS Collaboration, “Observation of a new excited beauty strange baryon decaying to  $\Xi_b^- \pi^+ \pi^-$ ”, *Phys. Rev. Lett.* **126** (2021) 252003, doi:10.1103/PhysRevLett.126.252003, arXiv:2102.04524.
- [40] CMS Collaboration, “Study of excited  $\Lambda_b^0$  states decaying to  $\Lambda_b^0 \pi^+ \pi^-$  in proton-proton collisions at  $\sqrt{s} = 13$  TeV”, *Phys. Lett. B* **803** (2020) 135345, doi:10.1016/j.physletb.2020.135345, arXiv:2001.06533.

## A Invariant mass distribution fit with constraints

The fit of the measured  $J/\psi \Xi^- K^+$  invariant mass distribution with constraints on the background shape parameter, signal shape parameter, resolution, and the mean value of the peak is presented in Fig. 4. The fit quality is good, as demonstrated by  $\chi^2/\text{dof} = 35.6/44$  for the binned distribution.






















Figure 4: Measured  $J/\psi \Xi^- K^+$  invariant mass distribution and overlaid constrained fit result.

## B The CMS Collaboration




### Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan<sup>1</sup> 

### Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , A. Escalante Del Valle , P.S. Hussain , M. Jeitler<sup>2</sup> , N. Krammer , D. Liko , I. Mikulec , J. Schieck<sup>2</sup> , R. Schöffbeck , D. Schwarz , M. Sonawane , S. Templ , W. Waltenberger , C.-E. Wulz<sup>2</sup> 















### Universiteit Antwerpen, Antwerpen, Belgium

M.R. Darwish<sup>3</sup> , T. Janssen , P. Van Mechelen 

### Vrije Universiteit Brussel, Brussel, Belgium

E.S. Bols , J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , H. El Faham , S. Lowette , I. Makarenko , D. Müller , A.R. Sahasransu , S. Tavernier , M. Tytgat<sup>4</sup> , S. Van Putte , D. Vannerom 

### Université Libre de Bruxelles, Bruxelles, Belgium

B. Clerbaux , G. De Lentdecker , L. Favart , D. Hohov , J. Jaramillo , A. Khalilzadeh, K. Lee , M. Mahdavihorrani , A. Malara , S. Paredes , L. Pétré , N. Postiau, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 






### Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , Y. Hong , J. Knolle , L. Lambrecht , G. Mestdach, C. Rendón, A. Samalan, K. Skovpen , N. Van Den Bossche , L. Wezenbeek 



















### Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , G. Bruno , C. Caputo , C. Delaere , I.S. Donertas , A. Giammanco , K. Jaffel , Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , K. Mondal , T.T. Tran , S. Wertz 

### Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , C. Hensel , T. Menezes De Oliveira, A. Moraes , P. Rebello Teles , M. Soeiro

### Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Alves Gallo Pereira , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato<sup>5</sup>, E.M. Da Costa , G.G. Da Silveira<sup>6</sup> , D. De Jesus Damiao , S. Fonseca De Souza , J. Martins<sup>7</sup> , C. Mora Herrera , K. Mota Amarilo , L. Mundim , H. Nogima , A. Santoro , A. Sznajder , M. Thiel , A. Vilela Pereira 

### Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes<sup>6</sup> , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

### Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 





### University of Sofia, Sofia, Bulgaria

A. Dimitrov , T. Ivanov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

**Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile**

S. Keshri , S. Thakur 













**Beihang University, Beijing, China**

T. Cheng , Q. Guo, T. Javaid , M. Mittal , L. Yuan 

**Department of Physics, Tsinghua University, Beijing, China**

G. Bauer<sup>8,9</sup>, Z. Hu , J. Liu, K. Yi<sup>8,10</sup> 

**Institute of High Energy Physics, Beijing, China**

G.M. Chen<sup>11</sup> , H.S. Chen<sup>11</sup> , M. Chen<sup>11</sup> , F. Iemmi , C.H. Jiang, A. Kapoor<sup>12</sup> , H. Liao , Z.-A. Liu<sup>13</sup> , F. Monti , M.A. Shahzad<sup>11</sup>, R. Sharma<sup>14</sup> , J.N. Song<sup>13</sup>, J. Tao , C. Wang<sup>11</sup>, J. Wang , Z. Wang<sup>11</sup>, H. Zhang 

**State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China**

A. Agapitos , Y. Ban , A. Levin , C. Li , Q. Li , Y. Mao, S.J. Qian , X. Sun , D. Wang , H. Yang, L. Zhang , C. Zhou 




**Sun Yat-Sen University, Guangzhou, China**

Z. You 

**University of Science and Technology of China, Hefei, China**

N. Lu 

**Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China**

X. Gao<sup>15</sup> , D. Leggat, H. Okawa , Y. Zhang 





**Zhejiang University, Hangzhou, Zhejiang, China**

Z. Lin , C. Lu , M. Xiao 





**Universidad de Los Andes, Bogota, Colombia**

C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

**Universidad de Antioquia, Medellin, Colombia**

J. Mejia Guisao , F. Ramirez , M. Rodriguez , J.D. Ruiz Alvarez 

**University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia**

D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 









**University of Split, Faculty of Science, Split, Croatia**

M. Kovac , T. Sculac 




**Institute Rudjer Boskovic, Zagreb, Croatia**

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , S. Mishra , A. Starodumov<sup>16</sup> , T. Susa 

**University of Cyprus, Nicosia, Cyprus**

A. Attikis , K. Christoforou , S. Konstantinou , J. Mousa , C. Nicolaou, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 








































































































**Charles University, Prague, Czech Republic**

M. Finger , M. Finger Jr. , A. Kveton 

**Escuela Politecnica Nacional, Quito, Ecuador**











E. Ayala 



**Universidad San Francisco de Quito, Quito, Ecuador**E. Carrera Jarrin **Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**H. Abdalla<sup>17</sup> , Y. Assran<sup>18,19</sup>**Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt**M. Abdullah Al-Mashad , M.A. Mahmoud **National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**R.K. Dewanjee<sup>20</sup> , K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken **Department of Physics, University of Helsinki, Helsinki, Finland**H. Kirschenmann , K. Osterberg , M. Voutilainen **Helsinki Institute of Physics, Helsinki, Finland**S. Bharthuar , E. Brücken , F. Garcia , J. Havukainen , K.T.S. Kallonen , M.S. Kim , R. Kinnunen, T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Lotti, L. Martikainen , M. Myllymäki , M.m. Rantanen , H. Siikonen , E. Tuominen , J. Tuominiemi **Lappeenranta-Lahti University of Technology, Lappeenranta, Finland**P. Luukka , H. Petrow , T. Tuuva<sup>†</sup>**IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France**M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , V. Lohezic , J. Malcles , J. Rander, A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro<sup>21</sup> , P. Simkina , M. Titov , M. Tornago **Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France**C. Baldenegro Barrera , F. Beaudette , A. Buchot Perraguin , P. Busson , A. Cappati , C. Charlot , F. Damas , O. Davignon , A. De Wit , G. Falmagne , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , J. Motta , M. Nguyen , C. Ochando , L. Portales , R. Salerno , U. Sarkar , J.B. Sauvan , Y. Sirois , A. Tarabini , E. Vernazza , A. Zabi , A. Zghiche **Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France**J.-L. Agram<sup>22</sup> , J. Andrea , D. Apparou , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , C. Grimault, R. Haeberle , A.-C. Le Bihan , M.A. Sessini , P. Van Hove **Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France**S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , J. Choi , D. Contardo , P. Depasse , C. Dozen<sup>23</sup> , H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg, G. Grenier , B. Ille , I.B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries, A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao **Georgian Technical University, Tbilisi, Georgia**G. Adamov, I. Lomidze , Z. Tsamalaidze<sup>16</sup> **RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**

V. Botta , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , N. Röwert , M. Teroerde 

### **RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany**

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , M.y. Lee , L. Mastrolorenzo, M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , A. Novak , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , A. Sharma , A. Stein , F. Torres Da Silva De Araujo<sup>24</sup> , L. Vigilante, S. Wiedenbeck , S. Zaleski











### **RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany**

C. Dziwok , G. Flügge , W. Haj Ahmad<sup>25</sup> , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 











### **Deutsches Elektronen-Synchrotron, Hamburg, Germany**

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , S. Bhattacharya , F. Blekman<sup>26</sup> , K. Borras<sup>27</sup> , D. Brunner , A. Campbell , A. Cardini , C. Cheng, F. Colombina , S. Consuegra Rodríguez , G. Correia Silva , M. De Silva , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo<sup>26</sup> , A. Geiser , A. Giraldi , G. Greau, V. Guglielmi , M. Guthoff , A. Hinzmann , A. Jafari<sup>28</sup> , L. Jeppe , N.Z. Jomhari , B. Kaech , M. Kasemann , H. Kaveh , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka<sup>29</sup> , W. Lohmann<sup>30</sup> , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , J. Metwally, A.B. Meyer , G. Milella , A. Mussgiller , A. Nürnberg , Y. Otari, D. Pérez Adán , E. Ranken , A. Raspereza , B. Ribeiro Lopes , J. Rübenach, A. Saggio , M. Scham<sup>31,27</sup> , S. Schnake<sup>27</sup> , P. Schütze , C. Schwanenberger<sup>26</sup> , D. Selivanova , M. Shchedrolosiev , R.E. Sosa Ricardo , L.P. Sreelatha Pramod , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , Q. Wang , Y. Wen , K. Wichmann, L. Wiens<sup>27</sup> , C. Wissing , S. Wuchterl , Y. Yang , A. Zimmermann Castro Santos 

### **University of Hamburg, Hamburg, Germany**

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , M. Bonanomi , P. Connor , M. Eich, K. El Morabit , Y. Fischer , A. Fröhlich, C. Garbers , E. Garutti , A. Grohsjean , M. Hajheidari, J. Haller , H.R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcari , T. Kramer , V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , A. Mehta , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoiu , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf 

### **Karlsruher Institut fuer Technologie, Karlsruhe, Germany**

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann<sup>32</sup> , R. Hofsaess , M. Horzela , U. Husemann , M. Klute , R. Koppenhöfer , M. Link, A. Lintuluoto , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H.J. Simonis , N. Trevisani , R. Ulrich , J. van der Linden , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , S. Wunsch, X. Zuo 

**Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi,**



**Greece**

G. Anagnostou, P. Assiouras , G. Daskalakis , A. Kyriakis, A. Papadopoulos<sup>32</sup>, A. Stakia 








**National and Kapodistrian University of Athens, Athens, Greece**

P. Kontaxakis , G. Melachroinos, A. Panagiotou, I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 






**National Technical University of Athens, Athens, Greece**

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsiopolitis, A. Zacharopoulou

**University of Ioánnina, Ioánnina, Greece**

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, P. Gianneios , C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 



**HUN-REN Wigner Research Centre for Physics, Budapest, Hungary**

M. Bartók<sup>33</sup> , C. Hajdu , D. Horvath<sup>34,35</sup> , F. Sikler , V. Veszpremi 

**MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary**

M. Csanád , K. Farkas , M.M.A. Gadallah<sup>36</sup> , Á. Kadlecik , P. Major , K. Mandal , G. Pásztor , A.J. Rádl<sup>37</sup> , G.I. Veres 




**Faculty of Informatics, University of Debrecen, Debrecen, Hungary**

P. Raics, B. Ujvari<sup>38</sup> , G. Zilizi 
















**Institute of Nuclear Research ATOMKI, Debrecen, Hungary**

G. Bencze, S. Czellar, J. Karancsi<sup>33</sup> , J. Molnar, Z. Szillasi

**Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary**

T. Csorgo<sup>37</sup> , F. Nemes<sup>37</sup> , T. Novak 

**Panjab University, Chandigarh, India**

J. Babbar , S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra<sup>39</sup> , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , M. Meena , K. Sandeep , T. Sheokand, J.B. Singh , A. Singla 
















**University of Delhi, Delhi, India**

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , M. Naimuddin , K. Ranjan , S. Saumya 




**Saha Institute of Nuclear Physics, HBNI, Kolkata, India**

S. Acharya<sup>40</sup> , S. Baradia , S. Barman<sup>41</sup> , S. Bhattacharya , D. Bhowmik, S. Dutta , S. Dutta, B. Gomber<sup>40</sup> , P. Palit , G. Saha , B. Sahu<sup>40</sup> , S. Sarkar






**Indian Institute of Technology Madras, Madras, India**

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , P. Jana , P. Kalbhor , J.R. Komaragiri<sup>42</sup> , D. Kumar<sup>42</sup> , L. Panwar<sup>42</sup> , R. Pradhan , P.R. Pujahari , N.R. Saha , A. Sharma , A.K. Sikdar , S. Verma 

**Tata Institute of Fundamental Research-A, Mumbai, India**















T. Aziz, I. Das , S. Dugad, M. Kumar , G.B. Mohanty , P. Suryadevara

**Tata Institute of Fundamental Research-B, Mumbai, India**

A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , Sh. Jain , S. Karmakar 

S. Kumar , G. Majumder , K. Mazumdar , S. Mukherjee , S. Parolia , A. Thachayath 




**National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India**

S. Bahinipati<sup>43</sup> , A.K. Das , C. Kar , D. Maity<sup>44</sup> , P. Mal , T. Mishra , V.K. Muraleedharan Nair Bindhu<sup>44</sup> , K. Naskar<sup>44</sup> , A. Nayak<sup>44</sup> , P. Sadangi , P. Saha , S.K. Swain , S. Varghese<sup>44</sup> , D. Vats<sup>44</sup> 





**Indian Institute of Science Education and Research (IISER), Pune, India**

A. Alpana , S. Dube , B. Kansal , A. Laha , A. Rastogi , S. Sharma 


**Isfahan University of Technology, Isfahan, Iran**

H. Bakhshiansohi<sup>45</sup> , E. Khazaie<sup>46</sup> , M. Zeinali<sup>47</sup> 
































**Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**

S. Chenarani<sup>48</sup> , S.M. Etesami , M. Khakzad , M. Mohammadi Najafabadi 




























**University College Dublin, Dublin, Ireland**

M. Grunewald 

**INFN Sezione di Bari<sup>a</sup>, Università di Bari<sup>b</sup>, Politecnico di Bari<sup>c</sup>, Bari, Italy**

M. Abbrescia<sup>a,b</sup> , R. Aly<sup>a,c,49</sup> , A. Colaleo<sup>a,b</sup> , D. Creanza<sup>a,c</sup> , B. D'Anzi<sup>a,b</sup> , N. De Filippis<sup>a,c</sup> , M. De Palma<sup>a,b</sup> , A. Di Florio<sup>a,c</sup> , W. Elmetenawee<sup>a,b,49</sup> , L. Fiore<sup>a</sup> , G. Iaselli<sup>a,c</sup> , G. Maggi<sup>a,c</sup> , M. Maggi<sup>a</sup> , I. Margjeka<sup>a,b</sup> , V. Mastrapasqua<sup>a,b</sup> , S. My<sup>a,b</sup> , S. Nuzzo<sup>a,b</sup> , A. Pellecchia<sup>a,b</sup> , A. Pompili<sup>a,b</sup> , G. Pugliese<sup>a,c</sup> , R. Radogna<sup>a</sup> , G. Ramirez-Sanchez<sup>a,c</sup> , D. Ramos<sup>a</sup> , A. Ranieri<sup>a</sup> , L. Silvestris<sup>a</sup> , F.M. Simone<sup>a,b</sup> , Ü. Sözbilir<sup>a</sup> , A. Stamerra<sup>a</sup> , R. Venditti<sup>a</sup> , P. Verwilligen<sup>a</sup> , A. Zaza<sup>a,b</sup> 



















**INFN Sezione di Bologna<sup>a</sup>, Università di Bologna<sup>b</sup>, Bologna, Italy**

G. Abbiendi<sup>a</sup> , C. Battilana<sup>a,b</sup> , D. Bonacorsi<sup>a,b</sup> , L. Borgonovi<sup>a</sup> , R. Campanini<sup>a,b</sup> , P. Capiluppi<sup>a,b</sup> , F.R. Cavallo<sup>a</sup> , M. Cuffiani<sup>a,b</sup> , G.M. Dallavalle<sup>a</sup> , T. Diotallevi<sup>a,b</sup> , F. Fabbri<sup>a</sup> , A. Fanfani<sup>a,b</sup> , D. Fasanella<sup>a,b</sup> , P. Giacomelli<sup>a</sup> , L. Giommi<sup>a,b</sup> , C. Grandi<sup>a</sup> , L. Guiducci<sup>a,b</sup> , S. Lo Meo<sup>a,50</sup> , L. Lunerti<sup>a,b</sup> , S. Marcellini<sup>a</sup> , G. Masetti<sup>a</sup> , F.L. Navarria<sup>a,b</sup> , A. Perrotta<sup>a</sup> , F. Primavera<sup>a,b</sup> , A.M. Rossi<sup>a,b</sup> , T. Rovelli<sup>a,b</sup> , G.P. Siroli<sup>a,b</sup> 

**INFN Sezione di Catania<sup>a</sup>, Università di Catania<sup>b</sup>, Catania, Italy**

S. Costa<sup>a,b,51</sup> , A. Di Mattia<sup>a</sup> , R. Potenza<sup>a,b</sup> , A. Tricomi<sup>a,b,51</sup> , C. Tuve<sup>a,b</sup> 

**INFN Sezione di Firenze<sup>a</sup>, Università di Firenze<sup>b</sup>, Firenze, Italy**

G. Barbagli<sup>a</sup> , G. Bardelli<sup>a,b</sup> , B. Camaiani<sup>a,b</sup> , A. Cassese<sup>a</sup> , R. Ceccarelli<sup>a</sup> , V. Ciulli<sup>a,b</sup> , C. Civinini<sup>a</sup> , R. D'Alessandro<sup>a,b</sup> , E. Focardi<sup>a,b</sup> , T. Kello<sup>a</sup> , G. Latino<sup>a,b</sup> , P. Lenzi<sup>a,b</sup> , M. Lizzo<sup>a</sup> , M. Meschini<sup>a</sup> , S. Paoletti<sup>a</sup> , A. Papanastassiou<sup>a,b</sup> , G. Sguazzoni<sup>a</sup> , L. Viliani<sup>a</sup> 

**INFN Laboratori Nazionali di Frascati, Frascati, Italy**













L. Benussi , S. Bianco , S. Meola<sup>52</sup> , D. Piccolo 

**INFN Sezione di Genova<sup>a</sup>, Università di Genova<sup>b</sup>, Genova, Italy**











P. Chatagnon<sup>a</sup> , F. Ferro<sup>a</sup> , E. Robutti<sup>a</sup> , S. Tosi<sup>a,b</sup> 

**INFN Sezione di Milano-Bicocca<sup>a</sup>, Università di Milano-Bicocca<sup>b</sup>, Milano, Italy**




























A. Benaglia<sup>a</sup> , G. Boldrini<sup>a,b</sup> , F. Brivio<sup>a</sup> , F. Cetorelli<sup>a</sup> , F. De Guio<sup>a,b</sup> , M.E. Dinardo<sup>a,b</sup> , P. Dini<sup>a</sup> , S. Gennai<sup>a</sup> , R. Gerosa<sup>a,b</sup> , A. Ghezzi<sup>a,b</sup> , P. Govoni<sup>a,b</sup> 

L. Guzzi<sup>a</sup> , M.T. Lucchini<sup>a,b</sup> , M. Malberti<sup>a</sup> , S. Malvezzi<sup>a</sup> , A. Massironi<sup>a</sup> , D. Menasce<sup>a</sup> , L. Moroni<sup>a</sup> , M. Paganoni<sup>a,b</sup> , D. Pedrini<sup>a</sup> , B.S. Pinolini<sup>a</sup>, S. Ragazzi<sup>a,b</sup> , T. Tabarelli de Fatis<sup>a,b</sup> , D. Zuolo<sup>a</sup> 












**INFN Sezione di Napoli<sup>a</sup>, Università di Napoli 'Federico II'<sup>b</sup>, Napoli, Italy; Università della Basilicata<sup>c</sup>, Potenza, Italy; Scuola Superiore Meridionale (SSM)<sup>d</sup>, Napoli, Italy**

S. Buontempo<sup>a</sup> , A. Cagnotta<sup>a,b</sup> , F. Carnevali<sup>a,b</sup>, N. Cavallo<sup>a,c</sup> , A. De Iorio<sup>a,b</sup> , F. Fabozzi<sup>a,c</sup> , A.O.M. Iorio<sup>a,b</sup> , L. Lista<sup>a,b,53</sup> , P. Paolucci<sup>a,32</sup> , B. Rossi<sup>a</sup> , C. Sciacca<sup>a,b</sup> 
















**INFN Sezione di Padova<sup>a</sup>, Università di Padova<sup>b</sup>, Padova, Italy; Università di Trento<sup>c</sup>, Trento, Italy**

R. Ardino<sup>a</sup> , P. Azzi<sup>a</sup> , N. Bacchetta<sup>a,54</sup> , D. Bisello<sup>a,b</sup> , P. Bortignon<sup>a</sup> , A. Bragagnolo<sup>a,b</sup> , R. Carlin<sup>a,b</sup> , T. Dorigo<sup>a</sup> , F. Gasparini<sup>a,b</sup> , U. Gasparini<sup>a,b</sup> , A. Gozzelino<sup>a</sup> , G. Grosso<sup>a</sup>, L. Layer<sup>a,55</sup>, E. Lusiani<sup>a</sup> , M. Margoni<sup>a,b</sup> , A.T. Meneguzzo<sup>a,b</sup> , M. Migliorini<sup>a,b</sup> , J. Pazzini<sup>a,b</sup> , P. Ronchese<sup>a,b</sup> , R. Rossin<sup>a,b</sup> , F. Simonetto<sup>a,b</sup> , G. Strong<sup>a</sup> , M. Tosi<sup>a,b</sup> , A. Triossi<sup>a,b</sup> , S. Ventura<sup>a</sup> , H. Yarar<sup>a,b</sup>, M. Zanetti<sup>a,b</sup> , P. Zotto<sup>a,b</sup> , A. Zucchetta<sup>a,b</sup> , G. Zumerle<sup>a,b</sup> 































**INFN Sezione di Pavia<sup>a</sup>, Università di Pavia<sup>b</sup>, Pavia, Italy**

S. Abu Zeid<sup>a,56</sup> , C. Aimè<sup>a,b</sup> , A. Braghieri<sup>a</sup> , S. Calzaferri<sup>a,b</sup> , D. Fiorina<sup>a,b</sup> , P. Montagna<sup>a,b</sup> , V. Re<sup>a</sup> , C. Riccardi<sup>a,b</sup> , P. Salvini<sup>a</sup> , I. Vai<sup>a,b</sup> , P. Vitulo<sup>a,b</sup> 




















**INFN Sezione di Perugia<sup>a</sup>, Università di Perugia<sup>b</sup>, Perugia, Italy**

S. Ajmal<sup>a,b</sup> , P. Asenov<sup>a,57</sup> , G.M. Bilei<sup>a</sup> , D. Ciangottini<sup>a,b</sup> , L. Fanò<sup>a,b</sup> , M. Magherini<sup>a,b</sup> , G. Mantovani<sup>a,b</sup>, V. Mariani<sup>a,b</sup> , M. Menichelli<sup>a</sup> , F. Moscatelli<sup>a,57</sup> , A. Piccinelli<sup>a,b</sup> , M. Presilla<sup>a,b</sup> , A. Rossi<sup>a,b</sup> , A. Santocchia<sup>a,b</sup> , D. Spiga<sup>a</sup> , T. Tedeschi<sup>a,b</sup> 







































**INFN Sezione di Pisa<sup>a</sup>, Università di Pisa<sup>b</sup>, Scuola Normale Superiore di Pisa<sup>c</sup>, Pisa, Italy; Università di Siena<sup>d</sup>, Siena, Italy**

P. Azzurri<sup>a</sup> , G. Bagliesi<sup>a</sup> , R. Bhattacharya<sup>a</sup> , L. Bianchini<sup>a,b</sup> , T. Boccali<sup>a</sup> , E. Bossini<sup>a</sup> , D. Bruschini<sup>a,c</sup> , R. Castaldi<sup>a</sup> , M.A. Ciocci<sup>a,b</sup> , M. Cipriani<sup>a,b</sup> , V. D'Amante<sup>a,d</sup> , R. Dell'Orso<sup>a</sup> , S. Donato<sup>a</sup> , A. Giassi<sup>a</sup> , F. Ligabue<sup>a,c</sup> , D. Matos Figueiredo<sup>a</sup> , A. Messineo<sup>a,b</sup> , M. Musich<sup>a,b</sup> , F. Palla<sup>a</sup> , A. Rizzi<sup>a,b</sup> , G. Rolandi<sup>a,c</sup> , S. Roy Chowdhury<sup>a</sup> , T. Sarkar<sup>a</sup> , A. Scribano<sup>a</sup> , P. Spagnolo<sup>a</sup> , R. Tenchini<sup>a</sup> , G. Tonelli<sup>a,b</sup> , N. Turini<sup>a,d</sup> , A. Venturi<sup>a</sup> , P.G. Verdini<sup>a</sup> 

**INFN Sezione di Roma<sup>a</sup>, Sapienza Università di Roma<sup>b</sup>, Roma, Italy**







P. Barria<sup>a</sup> , M. Campana<sup>a,b</sup> , F. Cavallari<sup>a</sup> , L. Cunqueiro Mendez<sup>a,b</sup> , D. Del Re<sup>a,b</sup> , E. Di Marco<sup>a</sup> , M. Diemoz<sup>a</sup> , F. Errico<sup>a,b</sup> , E. Longo<sup>a,b</sup> , P. Meridiani<sup>a</sup> , J. Mijuskovic<sup>a,b</sup> , G. Organtini<sup>a,b</sup> , F. Pandolfi<sup>a</sup> , R. Paramatti<sup>a,b</sup> , C. Quaranta<sup>a,b</sup> , S. Rahatlou<sup>a,b</sup> , C. Rovelli<sup>a</sup> , F. Santanastasio<sup>a,b</sup> , L. Soffi<sup>a</sup> 

**INFN Sezione di Torino<sup>a</sup>, Università di Torino<sup>b</sup>, Torino, Italy; Università del Piemonte Orientale<sup>c</sup>, Novara, Italy**








N. Amapane<sup>a,b</sup> , R. Arcidiacono<sup>a,c</sup> , S. Argiro<sup>a,b</sup> , M. Arneodo<sup>a,c</sup> , N. Bartosik<sup>a</sup> , R. Bellan<sup>a,b</sup> , A. Bellora<sup>a,b</sup> , C. Biino<sup>a</sup> , N. Cartiglia<sup>a</sup> , M. Costa<sup>a,b</sup> , R. Covarelli<sup>a,b</sup> , N. Demaria<sup>a</sup> , L. Finco<sup>a</sup> , M. Grippo<sup>a,b</sup> , B. Kiani<sup>a,b</sup> , F. Legger<sup>a</sup> , F. Luongo<sup>a,b</sup> , C. Mariotti<sup>a</sup> , S. Maselli<sup>a</sup> , A. Mecca<sup>a,b</sup> , E. Migliore<sup>a,b</sup> , M. Monteno<sup>a</sup> , R. Mulargia<sup>a</sup> , M.M. Obertino<sup>a,b</sup> , G. Ortona<sup>a</sup> , L. Pacher<sup>a,b</sup> , N. Pastrone<sup>a</sup> , M. Pelliccioni<sup>a</sup> , M. Ruspa<sup>a,c</sup> , F. Siviero<sup>a,b</sup> , V. Sola<sup>a,b</sup> , A. Solano<sup>a,b</sup> , D. Soldi<sup>a,b</sup> , A. Staiano<sup>a</sup> , C. Tarricone<sup>a,b</sup> , D. Trocino<sup>a</sup> , G. Umoret<sup>a,b</sup> , E. Vlasov<sup>a,b</sup> 



**INFN Sezione di Trieste<sup>a</sup>, Università di Trieste<sup>b</sup>, Trieste, Italy**

S. Belforte<sup>a</sup> , V. Candelise<sup>a,b</sup> , M. Casarsa<sup>a</sup> , F. Cossutti<sup>a</sup> , K. De Leo<sup>a,b</sup> ,  
G. Della Ricca<sup>a,b</sup> 





**Kyungpook National University, Daegu, Korea**

S. Dogra , J. Hong , C. Huh , B. Kim , D.H. Kim , J. Kim, H. Lee, S.W. Lee ,  
C.S. Moon , Y.D. Oh , M.S. Ryu , S. Sekmen , Y.C. Yang 

**Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea**

G. Bak , P. Gwak , H. Kim , D.H. Moon 

**Hanyang University, Seoul, Korea**

E. Asilar , D. Kim , T.J. Kim , J.A. Merlin, J. Park 

**Korea University, Seoul, Korea**

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , S. Lee , J. Park, S.K. Park, J. Yoo 

**Kyung Hee University, Department of Physics, Seoul, Korea**

J. Goh 









**Sejong University, Seoul, Korea**

H. S. Kim , Y. Kim, S. Lee



**Seoul National University, Seoul, Korea**

J. Almond, J.H. Bhyun, J. Choi , W. Jun , J. Kim , J.S. Kim, S. Ko , H. Kwon , H. Lee ,  
J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

**University of Seoul, Seoul, Korea**

W. Jang , D.Y. Kang, Y. Kang , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh,  
I.J. Watson , S. Yang 

**Yonsei University, Department of Physics, Seoul, Korea**

S. Ha , H.D. Yoo 

**Sungkyunkwan University, Suwon, Korea**

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 


**College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait**

T. Beyrouthy, Y. Maghrbi 

**Riga Technical University, Riga, Latvia**

K. Dreimanis , A. Gaile , G. Pikurs, A. Potrebko , M. Seidel , V. Veckalns<sup>58</sup> 

**University of Latvia (LU), Riga, Latvia**

N.R. Strautnieks 







**Vilnius University, Vilnius, Lithuania**

M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 






**National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia**

N. Bin Norjoharuddeen , I. Yusuff<sup>59</sup> , Z. Zolkapli

**Universidad de Sonora (UNISON), Hermosillo, Mexico**

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,  
M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 




**Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico**

G. Ayala , H. Castilla-Valdez , E. De La Cruz-Burelo , I. Heredia-De La Cruz<sup>60</sup> ,  
R. Lopez-Fernandez , C.A. Mondragon Herrera, A. Sánchez Hernández 

**Universidad Iberoamericana, Mexico City, Mexico**

C. Oropeza Barrera , M. Ramírez García 


**Benemerita Universidad Autonoma de Puebla, Puebla, Mexico**

I. Bautista , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 





**University of Montenegro, Podgorica, Montenegro**

I. Bubanja, N. Raicevic 

**University of Canterbury, Christchurch, New Zealand**

P.H. Butler 

**National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan 







**AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland**

V. Avati, L. Grzanka , M. Malawski 

**National Centre for Nuclear Research, Swierk, Poland**

H. Bialkowska , M. Bluj , B. Boimska , M. Górski , M. Kazana , M. Szeleper ,  
P. Zalewski 















**Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski ,  
A. Muhammad 



**Warsaw University of Technology, Warsaw, Poland**

K. Pozniak , W. Zabolotny 

**Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , P. Faccioli ,  
M. Gallinaro , J. Hollar , N. Leonardo , T. Niknejad , A. Petrilli , M. Pisano ,  
J. Seixas , J. Varela , J.W. Wulff




























**Faculty of Physics, University of Belgrade, Belgrade, Serbia**

P. Adzic , P. Milenovic 

**VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**

M. Dordevic , J. Milosevic , V. Rekovic














**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**

M. Aguilar-Benitez, J. Alcaraz Maestre , Cristina F. Bedoya , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , D. Fernández Del Val ,  
J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez ,  
J.M. Hernandez , M.I. Josa , J. León Holgado , D. Moran , C. M. Morcillo Perez ,  
Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo ,  
I. Redondo , D.D. Redondo Ferrero , L. Romero, S. Sánchez Navas , L. Urda Gómez ,  
J. Vazquez Escobar , C. Willmott




















**Universidad Autónoma de Madrid, Madrid, Spain**

J.F. de Trocóniz 

**Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain**

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , J.R. González Fernández , E. Palencia Cortezon , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

**Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain**

S. Bhowmik , S. Blanco Fernández , J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , C. Fernandez Madrazo , G. Gomez , C. Lasaos García , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizán García 

**University of Colombo, Colombo, Sri Lanka**

M.K. Jayananda , B. Kailasapathy<sup>61</sup> , D.U.J. Sonnadara , D.D.C. Wickramarathna 





**University of Ruhuna, Department of Physics, Matara, Sri Lanka**

W.G.D. Dharmaratna<sup>62</sup> , K. Liyanage , N. Perera , N. Wickramage 



































**CERN, European Organization for Nuclear Research, Geneva, Switzerland**

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , B. Bilin , A.A. Bin Anuar , A. Bocci , E. Brondolin , C. Caillol , T. Camporesi , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , F. Fallavollita<sup>63</sup>, L. Forthomme , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , L. Gouskos , M. Haranko , J. Hegeman , B. Huber, V. Innocente , T. James , P. Janot , J. Kieseler , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , B. Maier , L. Malgeri , M. Mannelli , A.C. Marini , M. Matthewman, F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Moortgat , M. Mulders , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo , H. Qu , D. Rabadý , G. Reales Gutiérrez, M. Rovere , H. Sakulin , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas<sup>64</sup> , A.G. Stahl Leitner , A. Steen , S. Summers , D. Treille , P. Tropea , A. Tsiroú, D. Walter , J. Wanczyk<sup>65</sup> , K.A. Wozniak<sup>66</sup> , P. Zehetner , P. Zejdl , W.D. Zeuner

**Paul Scherrer Institut, Villigen, Switzerland**

T. Bevilacqua<sup>67</sup> , L. Caminada<sup>67</sup> , A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli<sup>67</sup> , L. Noehte<sup>67</sup> , T. Rohe 



















**ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland**

T.K. Aarrestad , K. Androsov<sup>65</sup> , M. Backhaus , A. Calandri , C. Cazzaniga , K. Datta , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani , L. Marchese , C. Martin Perez , A. Mascellani<sup>65</sup> , E. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , M.G. Ratti , M. Reichmann , C. Reissel , T. Reitenspiess , B. Ristic , F. Riti , D. Ruini, D.A. Sanz Becerra , R. Seidita , J. Steggemann<sup>65</sup> , D. Valsecchi , R. Wallny 




**Universität Zürich, Zurich, Switzerland**

C. Amsler<sup>68</sup> , P. Bäertschi , C. Botta , D. Brzhechko, M.F. Canelli , K. Cormier 












R. Del Burgo, J.K. Heikkilä , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechti , A. Macchiolo , P. Meiring , V.M. Mikuni , U. Molinatti , I. Neutelings , A. Reimers , P. Robmann, S. Sanchez Cruz , K. Schweiger , M. Senger , Y. Takahashi , R. Tramontano 




#### National Central University, Chung-Li, Taiwan

C. Adloff<sup>69</sup>, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari<sup>42</sup> , S.S. Yu 




















#### National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, Y. Chao , K.F. Chen , P.s. Chen, Z.g. Chen, W.-S. Hou , T.h. Hsu, Y.w. Kao, R. Khurana, G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , A. Psallidas, X.f. Su, J. Thomas-Wilsker , L.s. Tsai, H.y. Wu, E. Yazgan 


#### High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatrangkuldee , N. Srimanobhas , V. Wachirapusanand 

#### Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , Z.S. Demiroglu , F. Dolek , I. Dumanoglu<sup>70</sup> , E. Eskut , Y. Guler<sup>71</sup> , E. Gurpinar Guler<sup>71</sup> , C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir<sup>72</sup> , A. Polatoz , B. Tali<sup>73</sup> , U.G. Tok , S. Turkcapar , E. Uslan , I.S. Zorbakir 

#### Middle East Technical University, Physics Department, Ankara, Turkey

M. Yalvac<sup>74</sup> 














#### Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya<sup>75</sup> , O. Kaya<sup>76</sup> , S. Tekten<sup>77</sup> 

#### Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak<sup>70,78</sup> , Y. Komurcu , S. Sen<sup>79</sup> 

#### Istanbul University, Istanbul, Turkey

O. Aydilek , S. Cerci<sup>73</sup> , V. Epshteyn , B. Haciasahinoglu , I. Hos<sup>80</sup> , B. Isildak<sup>81</sup> , B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , D. Sunar Cerci<sup>73</sup> , C. Zorbilmez 
















#### Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , B. Grynyov 


















#### National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 
































#### University of Bristol, Bristol, United Kingdom

D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki, J. Goldstein , H.F. Heath , L. Kreczko , B. Krikler , S. Paramesvaran , S. Seif El Nasr-Storey, V.J. Smith , N. Stylianou<sup>82</sup> , K. Walkingshaw Pass, R. White 




#### Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev<sup>83</sup> , C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , K.V. Ellis, K. Harder , S. Harper , M.-L. Holmberg<sup>84</sup> , J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , T. Williams 














**Imperial College, London, United Kingdom**

R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller, V. Cacchio, C.A. Carrillo Montoya , G.S. Chahal<sup>85</sup> , D. Colling , J.S. Dancu, P. Dauncey , G. Davies , J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M.H. Hassanshahi , A. Howard, G. Iles , M. Knight , J. Langford , L. Lyons , A.-M. Magnan , S. Malik, A. Martelli , M. Mieskolainen , J. Nash<sup>86</sup> , M. Pesaresi, B.C. Radburn-Smith , A. Richards, A. Rose , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , L.H. Vage, T. Virdee<sup>32</sup> , M. Vojinovic , N. Wardle , D. Winterbottom 

**Brunel University, Uxbridge, United Kingdom**

K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

**Baylor University, Waco, Texas, USA**

S. Abdullin , A. Brinkerhoff , B. Caraway , J. Dittmann , K. Hatakeyama , J. Hiltbrand , A.R. Kanuganti , B. McMaster , M. Saunders , S. Sawant , C. Sutantawibul , M. Toms<sup>87</sup> , J. Wilson 

















**Catholic University of America, Washington, DC, USA**

R. Bartek , A. Dominguez , C. Huerta Escamilla, A.E. Simsek , R. Uniyal , A.M. Vargas Hernandez 


















**The University of Alabama, Tuscaloosa, Alabama, USA**

R. Chudasama , S.I. Cooper , S.V. Gleyzer , C.U. Perez , P. Rumerio<sup>88</sup> , E. Usai , C. West , R. Yi 



















**Boston University, Boston, Massachusetts, USA**

A. Akpinar , A. Albert , D. Arcaro , C. Cosby , Z. Demiragli , C. Erice , E. Fontanesi , D. Gastler , S. Jeon , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , S. Yuan 

**Brown University, Providence, Rhode Island, USA**

G. Benelli , X. Coubez<sup>27</sup>, D. Cutts , M. Hadley , U. Heintz , J.M. Hogan<sup>89</sup> , T. Kwon , G. Landsberg , K.T. Lau , D. Li , J. Luo , S. Mondal , M. Narain<sup>†</sup> , N. Pervan , S. Sagir<sup>90</sup> , F. Simpson , M. Stamenkovic , W.Y. Wong, X. Yan , W. Zhang





**University of California, Davis, Davis, California, USA**

S. Abbott , J. Bonilla , C. Brainerd , R. Breedon , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , D. Pellett , W. Wei , Y. Yao , F. Zhang 























**University of California, Los Angeles, California, USA**

M. Bachtis , R. Cousins , A. Datta , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , E. Manca , D. Saltzberg , V. Valuev 




















**University of California, Riverside, Riverside, California, USA**

R. Clare , M. Gordon, G. Hanson , W. Si , S. Wimpenny<sup>†</sup> 











**University of California, San Diego, La Jolla, California, USA**

J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , L. Giannini , J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , M. Pieri , M. Quinnan , B.V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil 















**University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA**

A. Barzdukas , L. Brennan , C. Campagnari , G. Collura , A. Dorsett , J. Incandela , M. Kilpatrick , J. Kim , A.J. Li , P. Masterson , H. Mei , M. Oshiro , J. Richman , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , S. Wang 














#### California Institute of Technology, Pasadena, California, USA

A. Bornheim , O. Cerri, A. Latorre, J.M. Lawhorn , J. Mao , H.B. Newman , T. Q. Nguyen , M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu 

#### Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , M.B. Andrews , P. Bryant , V. Dutta , T. Ferguson , A. Harilal , C. Liu , T. Mudholkar , S. Murthy , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 










































































#### University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hassani , G. Karathanasis , E. MacDonald, N. Manganello , F. Marini , A. Perloff , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper 





















#### Cornell University, Ithaca, New York, USA

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan , D. Gadkari , S. Hogan , J. Monroy , J.R. Patterson , J. Reichert , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 















#### Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V.D. Elvira , Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig , J. Hirschauer , L. Horyn , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K.H.M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , V. Papadimitriou , N. Pastika , K. Pedro , C. Pena<sup>91</sup> , F. Ravera , A. Reinsvold Hall<sup>92</sup> , L. Ristori , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , I. Zoi 






#### University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , L. Cadamuro , P. Chang , V. Cherepanov , R.D. Field, E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K.H. Lo, K. Matchev , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , D. Rosenzweig , S. Rosenzweig , K. Shi , J. Wang 

#### Florida State University, Tallahassee, Florida, USA


















T. Adams , A. Al Kadhimi , A. Askew , N. Bower , R. Habibullah , V. Hagopian , R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, O. Viazlo , M. Wulansatiti , R. Yohay , J. Zhang

#### Florida Institute of Technology, Melbourne, Florida, USA
















B. Alsufyani, M.M. Baarmand , S. Butalla , T. Elkafrawy<sup>56</sup> , M. Hohlmann , R. Kumar Verma , M. Rahmani

#### University of Illinois Chicago, Chicago, USA, Chicago, USA













M.R. Adams , C. Bennett, R. Cavanaugh , S. Dittmer , R. Escobar Franco , O. Evdoki-

mov , C.E. Gerber , D.J. Hofman , J.h. Lee , D. S. Lemos , A.H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , T. Roy , S. Rudrabhatla , M.B. Tonjes , N. Varelas , X. Wang , Z. Ye , J. Yoo 


























#### **The University of Iowa, Iowa City, Iowa, USA**

M. Alhousseini , D. Blend , K. Dilsiz<sup>93</sup> , L. Emediato , G. Karaman , O.K. Köseyan , J.-P. Merlo , A. Mestvirishvili<sup>94</sup> , J. Nachtman , O. Neogi , H. Ogul<sup>95</sup> , Y. Onel , A. Penzo , C. Snyder , E. Tiras<sup>96</sup> 











#### **Johns Hopkins University, Baltimore, Maryland, USA**

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz , T.Á. Vámi 

#### **The University of Kansas, Lawrence, Kansas, USA**

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , C. Lindsey , J. Marquez , N. Minafra , M. Murray , M. Nickel , M. Pitt , S. Popescu<sup>97</sup> , C. Rogan , C. Royon , R. Salvatico , S. Sanders , C. Smith , Q. Wang , G. Wilson 


















#### **Kansas State University, Manhattan, Kansas, USA**

B. Allmond , A. Ivanov , K. Kaadze , A. Kalogeropoulos , D. Kim , Y. Maravin , K. Nam , J. Natoli , D. Roy , G. Sorrentino 





























#### **Lawrence Livermore National Laboratory, Livermore, California, USA**

F. Rebassoo , D. Wright 















#### **University of Maryland, College Park, Maryland, USA**

A. Baden , A. Belloni , A. Bethani , Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M.M. Paranjpe , L. Wang 


#### **Massachusetts Institute of Technology, Cambridge, Massachusetts, USA**

J. Bendavid , W. Busza , I.A. Cali , Y. Chen , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov , P. Harris , D. Hoang , D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. Mironov , C. Paus , D. Rankin , C. Roland , G. Roland , S. Rothman , Z. Shi , G.S.F. Stephans , J. Wang , Z. Wang , B. Wyslouch , T. J. Yang 

#### **University of Minnesota, Minneapolis, Minnesota, USA**

B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , S. Pandey , M. Revering , R. Rusack , R. Saradhy , N. Schroeder , N. Strobbe , M.A. Wadud 

#### **University of Mississippi, Oxford, Mississippi, USA**

L.M. Cremaldi 

















#### **University of Nebraska-Lincoln, Lincoln, Nebraska, USA**

K. Bloom , M. Bryson , D.R. Claes , C. Fangmeier , F. Golf , G. Haza , J. Hossain , C. Joo , I. Kravchenko , I. Reed , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu , A.G. Zecchinelli 











#### **State University of New York at Buffalo, Buffalo, New York, USA**

G. Agarwal , H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar , A. Williams 






























**Northeastern University, Boston, Massachusetts, USA**

E. Barberis , Y. Haddad , Y. Han , A. Krishna , J. Li , M. Lu , G. Madigan , R. Mccarthy , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , A. Tishelman-Charny , B. Wang , D. Wood 












**Northwestern University, Evanston, Illinois, USA**

S. Bhattacharya , J. Bueghly , Z. Chen , K.A. Hahn , Y. Liu , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliercio , M. Velasco 




















**University of Notre Dame, Notre Dame, Indiana, USA**

R. Band , R. Bucci , S. Castells , M. Cremonesi , A. Das , R. Goldouzian , M. Hildreth , K.W. Ho , K. Hurtado Anampa , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano , N. Marinelli , I. Mcalister , T. McCauley , C. Mcgrady , C. Moore , Y. Musienko<sup>16</sup> , H. Nelson , M. Osherson , R. Ruchti , A. Townsend , M. Wayne , H. Yockey , M. Zarucki , L. Zygala 

**The Ohio State University, Columbus, Ohio, USA**

A. Basnet , B. Bylsma , M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , A. Lesauvage , M. Nunez Ornelas , K. Wei , B.L. Winer , B. R. Yates 
























**Princeton University, Princeton, New Jersey, USA**

F.M. Addesa , H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , S. Higginbotham , G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , A. Shevelev , D. Stickland , C. Tully 




**University of Puerto Rico, Mayaguez, Puerto Rico, USA**

S. Malik 

















**Purdue University, West Lafayette, Indiana, USA**

A.S. Bakshi , V.E. Barnes , S. Chandra , R. Chawla , S. Das , A. Gu , L. Gutay , M. Jones , A.W. Jung , D. Kondratyev , A.M. Koshy , M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , V. Scheurer , J.F. Schulte , M. Stojanovic , J. Thieman , A. K. Viridi , F. Wang , W. Xie 













**Purdue University Northwest, Hammond, Indiana, USA**

J. Dolen , N. Parashar , A. Pathak 

**Rice University, Houston, Texas, USA**

D. Acosta , A. Baty , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed , P. Gardner , F.J.M. Geurts , A. Kumar , W. Li , O. Miguel Colin , B.P. Padley , R. Redjimi , J. Rotter , E. Yigitbasi , Y. Zhang 






















**University of Rochester, Rochester, New York, USA**

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , C. Fallon , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , P. Parygin<sup>87</sup> , E. Popova<sup>87</sup> , R. Taus , G.P. Van Onsem 

**The Rockefeller University, New York, New York, USA**










K. Goulianos 

**Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA**















B. Chiarito , J.P. Chou , Y. Gershtein , E. Halkiadakis , A. Hart , M. Heindl , D. Jaroslowski , O. Karacheban<sup>30</sup> , I. Laflotte , A. Lath , R. Montalvo , K. Nash , H. Routray , S. Salur , S. Schnetzer , S. Somalwar , R. Stone , S.A. Thayil , S. Thomas , J. Vora , H. Wang 














**University of Tennessee, Knoxville, Tennessee, USA**

H. Acharya , D. Ally , A.G. Delannoy , S. Fiorendi , T. Holmes , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 

**Texas A&M University, College Station, Texas, USA**

D. Aebi , M. Ahmad , O. Bouhali<sup>98</sup> , M. Dalchenko , R. Eusebi , J. Gilmore , T. Huang , T. Kamon<sup>99</sup> , H. Kim , S. Luo , S. Malhotra, R. Mueller , D. Overton , D. Rathjens , A. Safonov 









**Texas Tech University, Lubbock, Texas, USA**

N. Akchurin , J. Damgov , V. Hegde , A. Hussain , Y. Kazhykarim, K. Lamichhane , S.W. Lee , A. Mankel , T. Mengke, S. Muthumuni , T. Peltola , I. Volobouev , A. Whitbeck 

**Vanderbilt University, Nashville, Tennessee, USA**

E. Appelt , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 






















**University of Virginia, Charlottesville, Virginia, USA**

B. Cardwell , B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , A. Li , C. Neu , C.E. Perez Lara 

**Wayne State University, Detroit, Michigan, USA**

P.E. Karchin 

**University of Wisconsin - Madison, Madison, Wisconsin, USA**

A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

**Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN**

S. Afanasiev , V. Andreev , Yu. Andreev , T. Aushev , M. Azarkin , A. Babaev , A. Belyaev , V. Blinov<sup>100</sup>, E. Boos , V. Borshch , D. Budkouski , V. Chekhovsky, R. Chistov<sup>100</sup> , M. Danilov<sup>100</sup> , A. Dermenev , T. Dimova<sup>100</sup> , D. Druzhkin<sup>101</sup> , M. Dubinin<sup>91</sup> , L. Dudko , A. Ershov , G. Gavrilo , V. Gavrilo , S. Gninenko , V. Golovtsov , N. Golubev , I. Golutvin , I. Gorbunov , A. Gribushin , Y. Ivanov , V. Kachanov , L. Kardapoltsev<sup>100</sup> , V. Karjavine , A. Karneyeu , V. Kim<sup>100</sup> , M. Kirakosyan, D. Kirpichnikov , M. Kirsanov , V. Klyukhin , O. Kodolova<sup>102</sup> , D. Konstantinov , V. Korenkov , A. Kozyrev<sup>100</sup> , N. Krasnikov , A. Lanev , P. Levchenko<sup>103</sup> , N. Lychkovskaya , V. Makarenko , A. Malakhov , V. Matveev<sup>100</sup> , V. Murzin , A. Nikitenko<sup>104,102</sup> , S. Obraztsov , V. Oreshkin , V. Palichik , V. Perelygin , S. Petrushanko , S. Polikarpov<sup>100</sup> , V. Popov , O. Radchenko<sup>100</sup> , M. Savina , V. Savrin , M. Sergeev , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen<sup>100</sup> , S. Slabospitskii , V. Smirnov , A. Snigirev , D. Sosnov , V. Sulimov , E. Tcherniaev , A. Terkulov , O. Teryaev , I. Tlisova , A. Toropin , L. Uvarov , A. Uzunian , A. Vorobyev<sup>†</sup>, N. Voytishin , B.S. Yuldashev<sup>105</sup>, A. Zarubin , I. Zhizhin , A. Zhokin 

†: Deceased

<sup>1</sup>Also at Yerevan State University, Yerevan, Armenia

<sup>2</sup>Also at TU Wien, Vienna, Austria

- 
- <sup>3</sup>Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt
- <sup>4</sup>Also at Ghent University, Ghent, Belgium
- <sup>5</sup>Also at Universidade Estadual de Campinas, Campinas, Brazil
- <sup>6</sup>Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil
- <sup>7</sup>Also at UFMS, Nova Andradina, Brazil
- <sup>8</sup>Also at Nanjing Normal University, Nanjing, China
- <sup>9</sup>Now at Henan Normal University, Xinxiang, China
- <sup>10</sup>Now at The University of Iowa, Iowa City, Iowa, USA
- <sup>11</sup>Also at University of Chinese Academy of Sciences, Beijing, China
- <sup>12</sup>Also at China Center of Advanced Science and Technology, Beijing, China
- <sup>13</sup>Also at University of Chinese Academy of Sciences, Beijing, China
- <sup>14</sup>Also at China Spallation Neutron Source, Guangdong, China
- <sup>15</sup>Also at Université Libre de Bruxelles, Bruxelles, Belgium
- <sup>16</sup>Also at an institute or an international laboratory covered by a cooperation agreement with CERN
- <sup>17</sup>Also at Cairo University, Cairo, Egypt
- <sup>18</sup>Also at Suez University, Suez, Egypt
- <sup>19</sup>Now at British University in Egypt, Cairo, Egypt
- <sup>20</sup>Also at Birla Institute of Technology, Mesra, Mesra, India
- <sup>21</sup>Also at Purdue University, West Lafayette, Indiana, USA
- <sup>22</sup>Also at Université de Haute Alsace, Mulhouse, France
- <sup>23</sup>Also at Department of Physics, Tsinghua University, Beijing, China
- <sup>24</sup>Also at The University of the State of Amazonas, Manaus, Brazil
- <sup>25</sup>Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- <sup>26</sup>Also at University of Hamburg, Hamburg, Germany
- <sup>27</sup>Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- <sup>28</sup>Also at Isfahan University of Technology, Isfahan, Iran
- <sup>29</sup>Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- <sup>30</sup>Also at Brandenburg University of Technology, Cottbus, Germany
- <sup>31</sup>Also at Forschungszentrum Jülich, Juelich, Germany
- <sup>32</sup>Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- <sup>33</sup>Also at Institute of Physics, University of Debrecen, Debrecen, Hungary
- <sup>34</sup>Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- <sup>35</sup>Now at Universitatea Babeş-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania
- <sup>36</sup>Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- <sup>37</sup>Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- <sup>38</sup>Also at Faculty of Informatics, University of Debrecen, Debrecen, Hungary
- <sup>39</sup>Also at Punjab Agricultural University, Ludhiana, India
- <sup>40</sup>Also at University of Hyderabad, Hyderabad, India
- <sup>41</sup>Also at University of Visva-Bharati, Santiniketan, India
- <sup>42</sup>Also at Indian Institute of Science (IISc), Bangalore, India
- <sup>43</sup>Also at IIT Bhubaneswar, Bhubaneswar, India
- <sup>44</sup>Also at Institute of Physics, Bhubaneswar, India
- <sup>45</sup>Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- <sup>46</sup>Also at Department of Physics, Isfahan University of Technology, Isfahan, Iran
- <sup>47</sup>Also at Sharif University of Technology, Tehran, Iran
- <sup>48</sup>Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran

<sup>49</sup>Also at Helwan University, Cairo, Egypt

<sup>50</sup>Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy

<sup>51</sup>Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy

<sup>52</sup>Also at Università degli Studi Guglielmo Marconi, Roma, Italy

<sup>53</sup>Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy

<sup>54</sup>Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA

<sup>55</sup>Also at Università di Napoli 'Federico II', Napoli, Italy

<sup>56</sup>Also at Ain Shams University, Cairo, Egypt

<sup>57</sup>Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy

<sup>58</sup>Also at Riga Technical University, Riga, Latvia

<sup>59</sup>Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia

<sup>60</sup>Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico

<sup>61</sup>Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka

<sup>62</sup>Also at Saegis Campus, Nugegoda, Sri Lanka

<sup>63</sup>Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy

<sup>64</sup>Also at National and Kapodistrian University of Athens, Athens, Greece

<sup>65</sup>Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland

<sup>66</sup>Also at University of Vienna Faculty of Computer Science, Vienna, Austria

<sup>67</sup>Also at Universität Zürich, Zurich, Switzerland

<sup>68</sup>Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria

<sup>69</sup>Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France

<sup>70</sup>Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey

<sup>71</sup>Also at Konya Technical University, Konya, Turkey

<sup>72</sup>Also at Izmir Bakircay University, Izmir, Turkey

<sup>73</sup>Also at Adiyaman University, Adiyaman, Turkey

<sup>74</sup>Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey

<sup>75</sup>Also at Marmara University, Istanbul, Turkey

<sup>76</sup>Also at Milli Savunma University, Istanbul, Turkey

<sup>77</sup>Also at Kafkas University, Kars, Turkey

<sup>78</sup>Now at Istanbul Okan University, Istanbul, Turkey

<sup>79</sup>Also at Hacettepe University, Ankara, Turkey

<sup>80</sup>Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey

<sup>81</sup>Also at Yildiz Technical University, Istanbul, Turkey

<sup>82</sup>Also at Vrije Universiteit Brussel, Brussel, Belgium

<sup>83</sup>Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom

<sup>84</sup>Also at University of Bristol, Bristol, United Kingdom

<sup>85</sup>Also at IPPP Durham University, Durham, United Kingdom

<sup>86</sup>Also at Monash University, Faculty of Science, Clayton, Australia

<sup>87</sup>Now at an institute or an international laboratory covered by a cooperation agreement with CERN

<sup>88</sup>Also at Università di Torino, Torino, Italy

<sup>89</sup>Also at Bethel University, St. Paul, Minnesota, USA

<sup>90</sup>Also at Karamanoğlu Mehmetbey University, Karaman, Turkey

<sup>91</sup>Also at California Institute of Technology, Pasadena, California, USA



---

<sup>92</sup>Also at United States Naval Academy, Annapolis, Maryland, USA

<sup>93</sup>Also at Bingol University, Bingol, Turkey

<sup>94</sup>Also at Georgian Technical University, Tbilisi, Georgia

<sup>95</sup>Also at Sinop University, Sinop, Turkey

<sup>96</sup>Also at Erciyes University, Kayseri, Turkey

<sup>97</sup>Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania

<sup>98</sup>Also at Texas A&M University at Qatar, Doha, Qatar

<sup>99</sup>Also at Kyungpook National University, Daegu, Korea

<sup>100</sup>Also at another institute or international laboratory covered by a cooperation agreement with CERN

<sup>101</sup>Also at Universiteit Antwerpen, Antwerpen, Belgium

<sup>102</sup>Also at Yerevan Physics Institute, Yerevan, Armenia

<sup>103</sup>Also at Northeastern University, Boston, Massachusetts, USA

<sup>104</sup>Also at Imperial College, London, United Kingdom

<sup>105</sup>Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan