



Interpretations of the ATLAS measurements of Higgs boson production and decay rates and differential cross-sections in pp collisions at $\sqrt{s} = 13$ TeV

The ATLAS Collaboration

Measurements of the Higgs boson production times decay rates and differential cross-sections have recently been performed by the ATLAS experiment in several decay channels using up to 139 fb^{-1} of proton–proton collision data at $\sqrt{s} = 13$ TeV recorded at the Large Hadron Collider. This paper presents multiple interpretations of these Higgs boson measurements. Measurements of production-mode cross-sections, simplified template cross-sections and fiducial differential cross-sections in different decay channels are reparameterised in terms of the impact of Standard Model effective field theory operators, and constraints are reported on the corresponding Wilson coefficients. Production and decay rate measurements are interpreted in UV-complete extensions of the Standard Model, namely the two-Higgs-doublet model (2HDM) near the alignment limit and the Minimal Supersymmetric Standard Model (MSSM) for various MSSM benchmark scenarios. The constraints on the 2HDM parameters $(\cos(\beta - \alpha), \tan\beta)$ and the MSSM parameters $(m_A, \tan\beta)$ are complementary to those obtained from direct searches for additional Higgs bosons.

Contents

1	Introduction	3
2	Measurements of Higgs boson cross-sections	5
2.1	Data samples	5
2.2	Simulation of the Standard Model signal	5
2.3	Signal yield parameterisation	7
2.3.1	Fiducial differential cross-section measurements	8
2.3.2	Production and decay rate and simplified template cross-section measurements	9
3	Interpretations based on SM Effective Field Theory	14
3.1	Methodology of effective field theory interpretations	14
3.1.1	Simulation of the impact of SMEFT operators	14
3.1.2	Cross-section calculation with linear terms	18
3.1.3	Cross-section calculation including quadratic terms	19
3.1.4	Assumptions on SM Higgs boson kinematics	19
3.2	Constraints from STXS measurements	22
3.2.1	Sensitivity estimate and choice of parameters	23
3.2.2	Results	29
3.2.3	Comparison between the linearised model and the model including quadratic terms	36
3.3	Constraints from fiducial differential measurements	40
3.3.1	Sensitivity estimate and choice of parameters	40
3.3.2	Results	41
4	Interpretations based on UV-complete BSM models	43
4.1	Constraints on two-Higgs-doublet models	43
4.1.1	Constraints based on the κ -framework	44
4.1.2	Constraints using an EFT-based approach	45
4.2	Constraints on the MSSM	50
4.2.1	MSSM benchmark scenarios	50
4.2.2	Interpretation procedure	52
4.2.3	Results	53
5	Conclusions	57
Appendix: Validity of Gaussian approximation of STXS measurements in SMEFT interpretation		59

1 Introduction

Since a particle (H) with attributes consistent with those of the Higgs boson [1–6] of the Standard Model (SM) of particle physics was discovered [7, 8] by the ATLAS and CMS experiments, its properties have been measured with proton–proton (pp) collision data produced by the Large Hadron Collider (LHC) at CERN. The ATLAS and CMS Collaborations recently presented their first results on the combination of Higgs boson production and decay rate measurements with up to 139 fb^{-1} of 13 TeV pp collision data [9, 10], collected during the full LHC Run 2 in the years 2015 through 2018. Two interpretations of the ATLAS measurements are presented here. The first one is based on an effective field theory (EFT) framework of the Standard Model. The second interpretation relies on specific theories that provide ultraviolet (UV) completions of the SM, specifically several benchmark scenarios within the two-Higgs doublet model (2HDM) [11–14] or within a minimal supersymmetric extension of the Standard Model (MSSM) [15–19].

In addition to the production and decay rates mentioned above, the fiducial differential cross-sections for Higgs boson production as a function of selected event kinematics have been measured by the ATLAS Collaboration with the same data sample in the $H \rightarrow ZZ^* \rightarrow 4\ell$ ($\ell = e, \mu$) [20] and $H \rightarrow \gamma\gamma$ [21] decay channels, among others [22, 23]. The fiducial differential cross-sections measured as a function of the total transverse momentum¹ of the Higgs boson decay products in the two decay channels, $p_{\text{T}}^{4\ell}$ and $p_{\text{T}}^{\gamma\gamma}$, are used in this paper to constrain anomalous Higgs boson couplings to gluons and top quarks, and an anomalous Higgs-gluon-top-antitop quark coupling, within the same EFT framework. These couplings are also probed using the EFT interpretation of Higgs boson production and decay rate measurements; the sensitivities obtained with the two approaches are compared.

Effective field theories provide a model-independent approach to describe new physics at high mass scales, systematically improvable with higher-order perturbative calculations, to parametrise the effects of candidate theories beyond the Standard Model (BSM) that reduce to the Standard Model at low energies. In the SM Effective Field Theory (SMEFT), the effects of BSM dynamics at energy scales Λ that are large in comparison to the vacuum-expectation-value (vev) v of the SM Higgs field ($\Lambda \gg v$) can be parameterised at low energies, $E \ll \Lambda$, in terms of higher-dimensional operators that are built up from the Standard Model fields and respect its symmetries. Measurements of (fiducial) cross-sections can be used to place constraints on the Wilson coefficients associated with the SMEFT operators, corresponding to limits on BSM physics at some fixed scale Λ .

The methodology employed here for the EFT interpretation of the combined measurements of Higgs boson production and decay rates is similar to the one used by the individual analyses. The interpretation is applied to the combined measurements of the Higgs boson production cross-sections times branching ratios in the following decay channels: $H \rightarrow ZZ^* (\rightarrow 4\ell)$, $H \rightarrow \gamma\gamma$, $H \rightarrow bb$, $H \rightarrow \tau\tau$, $H \rightarrow WW^* (\rightarrow e\nu\mu\nu)$, $H \rightarrow Z\gamma$, and $H \rightarrow \mu\mu$ [24–34].

A similar approach is used for the interpretation of the fiducial p_{T} -differential cross-sections in the $H \rightarrow ZZ^* \rightarrow 4\ell$ and $H \rightarrow \gamma\gamma$ channels. For this study the p_{T} distribution was chosen as it was previously measured in both channels and is sensitive to anomalous Higgs boson couplings to top-quarks and gluons.

¹ ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the centre of the LHC ring, and the y -axis points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$. The transverse momentum p_{T} is the momentum component in the transverse plane.

The method resembles closely that described in Ref. [21] for the joint interpretation of five differential cross-sections in the $H \rightarrow \gamma\gamma$ decay channel in terms of a few dimension-6 SMEFT operators, but only the cross-section measurements as a function of $p_T^{4\ell}$ and $p_T^{\gamma\gamma}$ are used, and a smaller number of operators is probed. On the other hand, the joint interpretation with a second decay channel with similar statistical precision increases the sensitivity to the probed BSM operators.

In the interpretations presented in the second part of this paper, performed in the context of UV-complete BSM theories, two additional measurements of production and decay rates, based on a partial Run 2 data sample collected in the years 2015 and 2016, are considered: $H \rightarrow WW^*$ in the WH and ZH production modes [35], and $H \rightarrow$ multileptons in the $t\bar{t}H$ production mode [36]. In the latter, the selection requirement of two prompt leptons primarily selects $H \rightarrow WW^*$ and $H \rightarrow \tau\tau$ decays, with a small additional contribution of $H \rightarrow ZZ^*$ decays.

In a previous publication [37], measurements were interpreted within the hMSSM model [38] in terms of constraints on the mass of the neutral CP-odd boson A (m_A) and on the ratio of the vacuum expectation values of the Higgs doublets ($\tan\beta$). The use of this model as a benchmark MSSM scenario suffers from theoretical limitations in regions with small m_A or large $\tan\beta$ because its assumptions may not hold in these regions of parameter space [39]. In this paper, in addition to the hMSSM model, seven more recent benchmark MSSM scenarios [39, 40] are tested. These include two new scenarios [40] that were recently designed to be compatible with a low value of $\tan\beta$ (< 10). This parameter space region is otherwise excluded in most MSSM scenarios, which predict for $\tan\beta \lesssim 10$ a mass of the lightest neutral Higgs boson significantly lower than the observed Higgs boson mass of 125.09 ± 0.24 GeV [41]. In addition, an interpretation in terms of the parameters $\tan\beta$ and $\cos(\beta - \alpha)$ of the 2HDM near the alignment limit, in which the light neutral Higgs boson of the 2HDM is SM-like², is also reported. These constraints are compared with those inferred from the limits on the SMEFT Wilson coefficients after matching the 2HDM parameters to the SMEFT operators.

The paper is structured as follows. Section 2 gives an overview of the Higgs boson production-mode cross-sections and decay rate measurements and of the fiducial differential cross-section measurements that are used to derive the results presented. Section 3, devoted to the EFT interpretation of the measurements, starts with a description of the general approach of deriving limits on Wilson coefficients (Section 3.1). This is followed by the details of the specific choices of Wilson coefficients probed in the interpretations of the combined measurements of Higgs boson production-mode cross-sections and decay rates (Section 3.2) or of the measurements of the fiducial differential cross-sections as a function of p_T^H (Section 3.3), together with the corresponding constraints. The interpretations in UV-complete extensions of the SM are detailed in Section 4. The approach taken to interpret the Higgs boson production-mode cross-sections and decay rate measurements in terms of constraints on the parameters of the 2HDM or the MSSM, and the derived results, are described in Sections 4.1 and 4.2, respectively. Finally, Section 5 presents the conclusions.

² The terms SM-like, wino-like etc are used throughout this paper to denote particles with similar properties to those of the named particle.

2 Measurements of Higgs boson cross-sections

2.1 Data samples

The results presented in this article are based on up to 139 fb^{-1} of pp collision data collected by the ATLAS detector [42–44] in the years 2015 to 2018, with the LHC operating at a centre-of-mass energy of 13 TeV. The decay channels, targeted production modes and integrated luminosity of the data sample used in each input analysis are summarised in Table 1, where the simplified template cross-section measurements labelled ‘STXS’ are identical to those used in Ref. [9]. The analyses used for this interpretation were published while the uncertainty in the combined 2015–2018 integrated luminosity was 1.7%. It was derived from the calibration of the luminosity scale using x – y beam-separation scans, following a methodology similar to that detailed in Ref. [45], and using the LUCID-2 detector [46] for the baseline luminosity measurements. Subsequently an improved estimate of the integrated luminosity was made [47], with a slightly higher central value (+0.8%) and half the uncertainty (0.8%). The results described in the following sections assume the older, less precise luminosity calibration; they would not be significantly affected if recomputed using the latest one.

2.2 Simulation of the Standard Model signal

Monte Carlo (MC) simulated event samples of SM Higgs bosons produced in pp collisions and decaying into the final states considered in this study were generated and normalised to SM predictions calculated for a Higgs boson mass $m_H = 125.09 \text{ GeV}$. For each Higgs boson decay mode, the branching ratio that was used corresponds to the theoretical calculation at the highest available order of accuracy in perturbation theory [49].

Higgs boson production via gluon–gluon fusion was simulated using the POWHEG BOX v2 [50–53] NNLOPS implementation [54, 55]. The event generator uses HNNLO [56] to reweight the inclusive Higgs boson rapidity distribution produced by the next-to-leading-order (NLO) generation of $pp \rightarrow H + \text{parton}$ events, with the scale of each parton emission determined using the MiNLO procedure [57]. The PDF4LHC15 [58] next-to-next-to-leading-order (NNLO) parton distribution functions (PDFs) were used for the central prediction and uncertainty, except for the $H \rightarrow b\bar{b}$ and $H \rightarrow Z\gamma$ measurements, which used the NNPDF3.0 [59] NNLO PDFs. The sample is normalised such that it reproduces the total cross-section predicted by a next-to-next-to-next-to-leading-order (N³LO) quantum chromodynamics (QCD) calculation with NLO electroweak (EW) corrections applied [49, 60–63]. The NNLOPS generator reproduces the Higgs boson p_T distribution predicted by the NNLO plus next-to-next-to-leading logarithm (NNLL) calculation of HRES2.3 [64], which includes the effects of top- and bottom-quark masses and uses dynamical renormalisation and factorisation scales.

The VBF and $q\bar{q} \rightarrow VH$ ($gg \rightarrow ZH$) production processes were simulated at NLO (LO) accuracy in QCD using the POWHEG BOX v2 [65] generator where the simulation of $q\bar{q} \rightarrow VH$ relies on improved NLO calculations [66]. The PDF4LHC15 NLO set of PDFs was used, except for the $H \rightarrow b\bar{b}$ and $H \rightarrow Z\gamma$ measurements, which used the NNPDF3.0 NLO PDFs. In the vector-boson fusion (VBF) $H \rightarrow b\bar{b}$ analysis, NLO EW corrections were calculated using MADGRAPH5_AMC@NLO 3.0 and applied as a function of the generated Higgs boson transverse momentum. The VBF sample is normalised to an approximate-NNLO QCD cross-section with NLO EW corrections applied [49, 67–69]. The VH samples are normalised to

Table 1: The decay channels (1st column), targeted production modes (2nd column) and integrated luminosity \mathcal{L} (3rd column) of the data sample used for each analysis included in the combined measurements of Higgs boson production and decay rates, couplings and simplified template cross-sections (labelled as ‘STXS’ in the ‘Binning’ column), or in the measurements of fiducial p_T -differential cross-sections (labelled as ‘differential’ in the ‘Binning’ column). The references for the input analyses (5th column) and information about which input analyses are considered for the SMEFT-based interpretation (6th column) and the interpretations in terms of UV-complete models (7th column) are also provided. The STXS-0* binning only distinguishes inclusive production processes, whereas the STXS-1.2 and differential binnings partition production processes in kinematic regions at a varying level of granularity. The full definitions of these binning schemes for the measurements and the corresponding signal yield parameterisations are detailed in Section 2.3. In all the analyses, the tiny expected contribution from associated $b\bar{b}H$ production is applied as a correction to ggF as described in the text.

Decay channel	Analysis Production mode	\mathcal{L} [fb ⁻¹]	Reference	Binning	SMEFT	2HDM and (h)MSSM
$H \rightarrow \gamma\gamma$	(ggF, VBF, WH , ZH , ttH , tH)	139	[25] [21]	STXS-1.2 differential	✓ ✓(subset)	✓
$H \rightarrow ZZ^*$	($ZZ^* \rightarrow 4\ell$: ggF, VBF, $WH + ZH$, $ttH + tH$)	139	[24] [20]	STXS-1.2 differential	✓ ✓(subset)	✓
	($ZZ^* \rightarrow \ell\ell\nu\bar{\nu}/\ell\ell q\bar{q}$: ttH multileptons)	36.1	[36]	STXS-0*		✓
$H \rightarrow \tau\tau$	(ggF, VBF, $WH + ZH$, $ttH + tH$)	139	[31]	STXS-1.2	✓	✓
	(ttH multileptons)	36.1	[36]	STXS-0*		✓
$H \rightarrow WW^*$	(ggF, VBF)	139	[32]	STXS-1.2	✓	✓
	(WH , ZH)	36.1	[48]	STXS-0*		✓
	(ttH multileptons)	36.1	[36]	STXS-0*		✓
$H \rightarrow bb$	(WH , ZH)	139	[26, 27]	STXS-1.2	✓	✓
	(VBF)	126	[28]	STXS-1.2	✓	✓
	($ttH + tH$)	139	[30]	STXS-1.2	✓	✓
	(boosted Higgs bosons: inclusive production)	139	[29]	STXS-1.2	✓	✓
$H \rightarrow Z\gamma$	(inclusive production)	139	[33]	STXS-0*	✓	✓
$H \rightarrow \mu\mu$	(ggF + $ttH + tH$, VBF + $WH + ZH$)	139	[34]	STXS-0*	✓	✓

cross-sections calculated at NNLO in QCD with NLO EW corrections [70, 71] and additional NLO QCD corrections [72] for the $gg \rightarrow ZH$ subprocess [49].

Higgs boson production in association with a top–antitop quark pair ($t\bar{t}H$) was simulated at NLO accuracy in QCD using the POWHEG Box v2 generator, except for the $H \rightarrow \mu\mu$ analysis and the measurements based on 36.1 fb^{-1} of data, which used the MADGRAPH5_AMC@NLO 2.6 [73] generator. The PDF4LHC15 NLO set of PDFs was used in the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ analyses, while the NNPDF3.0 (N)NLO set of PDFs was used in the other ones. In all cases, the sample is normalised to a calculation with NLO QCD and EW corrections [49, 74–77]. Higgs boson production in association with a single top quark in the t -channel (tHq) or with an additional W boson (tHW) were simulated at NLO accuracy using MADGRAPH5_AMC@NLO 2.6 and either the NNPDF3.0(N)NLO PDF set or the CT10 NLO PDF set. The two tH samples are normalised to NLO QCD calculations [49, 78].

All parton-level events were input to PYTHIA 8.2 [79] to model the Higgs boson decay, parton showering, hadronisation, and multiple parton interactions. The generators were interfaced to PYTHIA 8.2 using the AZNLO [80] or A14 [81] sets of tuned parameters (tunes).

Particle-level events were passed through a GEANT 4 [82] simulation of the ATLAS detector [83] and reconstructed using the same analysis software as used for the data. Event pile-up was included in the simulation by overlaying inelastic pp collisions, such that the distribution of the average number of interactions per bunch crossing reproduces that observed in the data. The inelastic pp collisions were simulated with PYTHIA 8.1 using the MSTW2008LO [84] set of PDFs with the A2 tune [85], or using the NNPDF2.3LO [86] set of PDFs with the A3 tune [87].

2.3 Signal yield parameterisation

In all analyses listed in Table 1, the likelihood function for each signal region k , with one or more bins r , is modelled as

$$L(N_k | \mu, \theta) = \prod_r \text{Poisson} \left(N_{k,r} | s_k(\mu, \theta) \cdot f_s^{k,r}(\theta) + b_{k,r}(\theta) \right), \quad (1)$$

where $N_{k,r}$ is the observed event count of bin r in region k , s_k is the expected signal count in region k , $f_s^{k,r}$ is the expected fraction of the signal in region k that is contained in bin r , and $b_{k,r}$ represents the expected event count from background processes. The ensemble of *parameters of interest* μ describes the Higgs boson signal normalisation, while θ represents the set of *nuisance parameters* taking into account the systematic uncertainties that originate from theoretical and experimental sources, as well as additional degrees of freedom without prior constraints such as background yields or normalisations in some of the input channels. The full likelihood function is then the product of the likelihood functions for each signal region k and of Gaussian or log-normal probability density functions that constrain the nuisance parameters.

Depending on the level of detail implemented in each analysis, the signal yield parameters μ can be indexed by Higgs boson production process (i), decay mode (X), and fiducial phase space region defined at the particle level (k'). Analysis region k , defined at the reconstruction level, is typically chosen to match the particle-level region k' as closely as possible, to reduce the extrapolation uncertainty. As

reconstruction-level selections do not generally correspond exactly to particle-level regions, multiple particle-level regions will contribute to the signal yield s_k .

For the measurements listed in Table 1, two distinct signal parameterisation strategies are followed as defined in Eq. (2) and Eq. (3). For analyses labelled as ‘STXS-0*’ or ‘STXS-1.2’, the signal yield for region k is modelled as a scale factor $\mu_k^{i,k',X}$ applied to the SM Higgs boson production cross-section times branching ratio, for each Higgs boson production process i and decay X , in a fiducial region k' defined at the particle level. Alternatively, for analyses labelled as ‘differential’, the signal yield is modelled as a cross-section $\sigma_{\text{fid.}}^{k',X}$ describing the sum of all production processes, separately for each Higgs boson decay mode X and fiducial region k' defined at the particle level. The corresponding parameterisations of the signal yield s_k in terms of the parameters of interest $\boldsymbol{\mu}_k = \{\mu_k^{i,k',X}\}$ or $\{\mu_k^{k',X}\}$ and of the nuisance parameters $\boldsymbol{\theta}$ are:

$$s_k^{\text{STXS}}(\boldsymbol{\mu}_k, \boldsymbol{\theta}) = \mathcal{L} \times \sum_{i,k',X} \mu_k^{i,k',X} \times (\sigma \times \mathcal{B})_{\text{SM,(N(N))NLO}}^{i,k',X}(\boldsymbol{\theta}) \times \epsilon_{\text{STXS},k}^{i,k',X}(\boldsymbol{\theta}), \quad (2)$$

$$s_k^{\text{diff.}}(\boldsymbol{\mu}_k, \boldsymbol{\theta}) = \mathcal{L} \times \sum_{k',X} \mu_k^{k',X} \times \sigma_{\text{SM, fid.}}^{k',X}(\boldsymbol{\theta}) \times \epsilon_{\text{diff.},k}^{k',X}(\boldsymbol{\theta}), \quad (3)$$

where \mathcal{L} is the integrated luminosity, $(\sigma \times \mathcal{B})_{\text{SM,(N(N))NLO}}^{i,k',X}$ is the calculation, at the highest available order, of the SM Higgs boson cross-section for the production process i in particle-level region k' multiplied by the SM Higgs boson branching ratio to the final state X , and $\sigma_{\text{SM, fid.}}^{k',X}$ is the SM Higgs boson production cross-section for the fiducial region k' , summed over all production modes. The factors $\epsilon_{\text{STXS},k}^{i,k',X}$ and $\epsilon_{\text{diff.},k}^{k',X}$ represent the products of acceptance times efficiency of the reconstruction-level region k for the particle-level fiducial phase space region k' and Higgs boson decay X (in production mode i for the STXS interpretation).

For each interpretation based on a particular model (SMEFT, 2HDM, or MSSM) with a vector of model parameters $\boldsymbol{\alpha}$, the original signal parameters $\boldsymbol{\mu}$ are replaced with expressions that parameterise the model predictions, i.e., $\boldsymbol{\mu} \rightarrow \boldsymbol{\mu}(\boldsymbol{\alpha})$, so that the likelihood of Eq. (1) is directly expressed in terms of the parameters $\boldsymbol{\alpha}$. Then, constraints on these parameters can be directly inferred from the modified likelihood expression. The model-specific re-parameterisations of the signal parameters are detailed in Sections 3 and 4.

The acceptance factors $\epsilon_{\text{STXS},k}^{i,k',X}$ and $\epsilon_{\text{diff.},k}^{k',X}$, as well as the signal bin fractions $f_s^{k,r}$, are derived assuming SM Higgs boson kinematics. For interpretations of the measurements in physics models that significantly alter kinematic distributions, additional correction factors may be needed to account for changes in the acceptance and signal shape as a function of BSM model parameters. These are discussed when applicable in Sections 3 and 4.

2.3.1 Fiducial differential cross-section measurements

For the interpretation of fiducial differential cross-section measurements in the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ analyses, the differential distributions in the transverse momentum of the Higgs boson decay products are considered. The $H \rightarrow ZZ^* \rightarrow 4\ell$ analysis defines nine analysis regions with boundaries in $p_T^{4\ell}$ at

$$\{0, 10, 20, 30, 45, 60, 80, 120, 200, 300\} \text{ GeV},$$

whereas the $H \rightarrow \gamma\gamma$ analysis defines a finer granularity with 20 analysis regions with boundaries in $p_T^{\gamma\gamma}$ at

$$\{0, 5, 10, 15, 20, 25, 30, 35, 45, 60, 80, 100, 120, 140, 170, 200, 250, 300, 450, 650, 13000\} \text{ GeV},$$

resulting in a total of 29 fiducial cross-section measurements that are interpreted in Section 3.3. Further details of these analyses, including the full definitions of the fiducial regions, can be found in Refs. [20, 21].

2.3.2 Production and decay rate and simplified template cross-section measurements

For the interpretations based on the SMEFT framework presented in Section 3, measurements of ‘simplified template cross-sections’ of various Higgs boson production processes in the regions of phase space defined within the STXS framework [49, 88–90] are used. The inclusive production process classes defined at ‘Stage-0’ in the STXS scheme are bbH production, ttH and tH processes, $qq \rightarrow Hqq$ processes (with contributions from both VBF production and quark-initiated VH production with a hadronic decay of the gauge boson), WH and ZH production (including $gg \rightarrow ZH$, followed by any leptonic decay of the vector boson $V(\text{lep})H$) and finally the $gg \rightarrow H$ process, consisting of ggF and $gg \rightarrow ZH, Z \rightarrow q\bar{q}$ production. Since the acceptances for $gg \rightarrow H$ and bbH production are similar for all input analyses, the bbH production mode is modelled as a 1% increase of the $gg \rightarrow H$ yield in each region of production phase space [49]. Theory uncertainties for the $gg \rightarrow H$, $qq \rightarrow Hqq$, and ttH processes are defined as in Refs. [24, 25], while those of the $V(\text{lep})H$ process follow the scheme described in Ref. [91].

The analyses listed in Table 1 as using the ‘STXS-0*’ binning provide measurements of the scale factors $\mu^{i,X}$ for the product of the inclusive cross-section of one or more production modes i , with $i = \text{ggF, VBF, } WH, ZH, ttH, tH^3$, and branching ratio of the Higgs boson decay mode X . Analyses of the most sensitive production modes performed with the full Run 2 data sample, labelled ‘STXS-1.2’ in Table 1, report measurements of signal yield scale factors in the more fine-grained ‘Stage-1.2’ definition of Higgs boson production cross-sections. At Stage-1.2, the Stage-0 regions are partitioned in particle-level kinematic volumes, such as in a few coarse intervals of Higgs boson transverse momentum (p_T^H). The ensemble of parameters $\mu_{\text{STXS-1.2}}^{i,k',X}$ describes deviations in differential distributions, with the level of detail controlled by the number of particle-level regions that are defined. The precision with which the more fine-grained set of scale factors $\mu_{\text{STXS-1.2}}^{i,k',X}$ can be measured depends on the design of the analysis and the amount of available data. As individual analyses provide only limited sensitivity to some of the Stage-1.2 categories with the current data sample, some of these categories are merged compared to the Stage-1.2 definitions given in Ref. [49]. Details of the merged category definitions in each analysis are provided in Refs. [24–32]. The grouping of the production modes for STXS-0* and STXS-1.2 regions is also shown in the second column of Table 1.

After merging, a total of 78 signal yield parameters are extracted from the STXS measurements listed in Table 1. The kinematic definitions of these 78 STXS signal yield parameters, along with their measured signal strengths (i.e., their ratios to the corresponding SM predictions) and uncertainties, are shown in Figures 1–3. While the data underlying the measurements in these figures is the same as that used for the interpretation of the combined Higgs boson measurements shown in Ref. [9], the granularity of the STXS

³ The classification of production modes used here differs in one respect from the Stage-0 STXS definition: in the STXS definition $qq \rightarrow VH(qq)$ production and VBF production are both labelled as a common $qq \rightarrow Hqq$ production mode, whereas these production modes are labelled separately here, based on the corresponding definition/settings of the MC event generators.

model used here is substantially larger, providing additional constraining power for the parameters of the SMEFT interpretation of the data. The χ^2 for the compatibility between the observed values and the SM hypothesis is 50 for 78 degrees of freedom, corresponding to a p -value of 99.4%.

For the MSSM and tree-level 2HDM interpretations of Section 4, no predictions are available for all production modes for the full STXS Stage-1.2 phase space partitioning. In this case, the interpretation is performed based on the STXS-0* production mode scale factors measured by each input analysis.




Figure 1: Measured signal strength for each of the $H \rightarrow WW^*$ and $H \rightarrow ZZ^*$ STXS categories included in the combination. In the $H \rightarrow ZZ^*$ analysis, VH production events with the vector boson V decaying into leptons are assigned to the $VH\text{-lep}$ category. In one category the negative uncertainty is truncated, as indicated by the hatched region, which corresponds to cross-section values for which the total yield prediction in an analysis region is negative. In that case, the negative uncertainty and its breakdown in terms of statistical and systematic components are not reported, and replaced with a long dash (-).




Figure 2: Measured signal strength for each of the $H \rightarrow \gamma\gamma$ and $H \rightarrow Z\gamma$ STXS categories included in the combination. In the $H \rightarrow \gamma\gamma$ analysis, events with two or more particle-level jets are assigned to the VH -veto (VH -had) categories if the leading dijet invariant mass m_{jj} is in the range $m_{jj} < 60 \text{ GeV}$ or $120 \leq m_{jj} < 350 \text{ GeV}$ ($60 \leq m_{jj} < 120 \text{ GeV}$). In one category the uncertainty is truncated, as indicated by the hatched region, which corresponds to cross-section values for which the total yield prediction in an analysis region is negative. In that case, the negative uncertainty and its breakdown in terms of statistical and systematic components are not reported, and replaced with a long dash (-).




Figure 3: Measured signal strength for each of the $H \rightarrow \tau\tau$, $b\bar{b}$, and $\mu\mu$ STXS categories implemented.

3 Interpretations based on SM Effective Field Theory

3.1 Methodology of effective field theory interpretations

The SM effective field theory provides an elegant language to encode the modifications of the Higgs boson properties induced by a wide class of BSM theories. Within the mathematical language of the SMEFT, the effects of BSM dynamics at high energies $\Lambda \gg v$, i.e., well above the electroweak scale $v = 246$ GeV, can be parameterised at low energies, $E \ll \Lambda$, in terms of higher-dimensional operators built up from the SM fields and respecting its symmetries such as gauge invariance. This yields an effective Lagrangian:

$$\mathcal{L}_{\text{SMEFT}} = \mathcal{L}_{\text{SM}} + \sum_i^{N_{d=6}} \frac{c_i}{\Lambda^2} \mathcal{O}_i^{(6)} + \sum_j^{N_{d=8}} \frac{b_j}{\Lambda^4} \mathcal{O}_j^{(8)} + \dots, \quad (4)$$

where \mathcal{L}_{SM} is the SM Lagrangian, $\mathcal{O}_i^{(6)}$ and $\mathcal{O}_j^{(8)}$ represent a complete set of operators of mass-dimensions $d = 6$ and $d = 8$, and c_i, b_j are the corresponding dimensionless Wilson coefficients. Operators with $d = 5$ and $d = 7$ violate lepton and/or baryon number conservation and are not considered in this study. The effective theory expansion in Eq. (4) is robust, fully general, and can be systematically matched to explicit UV-complete BSM scenarios.

The goal of the analysis is to constrain the $d = 6$ Wilson coefficients that correspond to operators that either directly or indirectly impact Higgs boson couplings to SM particles [92, 93]. Contributions of operators of mass-dimension $d = 8$, which are suppressed by $1/\Lambda^2$ relative to the leading effects from $d = 6$ operators and whose impact on Higgs boson production and decay in the kinematic regions of interest are not fully calculated, are not considered. The ‘Warsaw’ basis [94] is used, which forms a complete set of all $\mathcal{O}_i^{(6)}$ operators in Eq. (4) allowed for by the SM gauge symmetries. This basis is widely used in EFT constraints in various fields of particle physics.

The ‘top’ flavour symmetry scheme for EFT operators is assumed, following the recommendation of Ref. [95]. In this scheme, quarks of the first two generations and quarks of the 3rd generation are described by independent fields, a $U(2)^3$ symmetry is imposed on the light quarks, and the quark sector of the EFT is assumed to have no mixing; all lepton generations are modelled independently. Table 2 lists the operators considered in the analysis of STXS data, and their corresponding Wilson coefficients c_i . Only CP-even $d = 6$ operators for which the Λ^{-2} -suppressed contribution to any of the STXS measurements shown in Figures 1–3 exceeds 0.1% relative to the SM prediction at $c_i/\Lambda^2 = 1/\text{TeV}^2$ are listed. The EFT analysis of the differential data only considers a subset of the coefficients listed, c_{HG}, c_{tH} and c_{tG} , due to the more restricted sensitivity of the differential analyses.

A value of $\Lambda = 1$ TeV is assumed, coefficients for alternative values of $\Lambda = X$ can be obtained through a scaling of the results presented in this article by a factor $(X/1 \text{ TeV})^2$. All Wilson coefficients are assumed to be real.

3.1.1 Simulation of the impact of SMEFT operators

The impact of the $d = 6$ SMEFT operators listed in Table 2 are computed with two universal FEYNRULES output (UFO) models of MADGRAPH [73] normalised to theoretical calculations of production cross-sections and decay branching ratios as follows.

Table 2: Wilson coefficients c and corresponding $d = 6$ SMEFT operators O_j used in this analysis. The operator and coefficient notation follows the convention of Ref. [96].

Wilson coefficient	Operator	Wilson coefficient	Operator
c_H	$(H^\dagger H)^3$	$c_{Qq}^{(1,1)}$	$(\bar{Q}\gamma_\mu Q)(\bar{q}\gamma^\mu q)$
$c_{H\Box}$	$(H^\dagger H)\Box(H^\dagger H)$	$c_{Qq}^{(1,8)}$	$(\bar{Q}T^a\gamma_\mu Q)(\bar{q}T^a\gamma^\mu q)$
c_G	$f^{abc}G_\mu^{av}G_\nu^{b\rho}G_\rho^{c\mu}$	$c_{Qq}^{(3,1)}$	$(\bar{Q}\sigma^i\gamma_\mu Q)(\bar{q}\sigma^i\gamma^\mu q)$
c_W	$\epsilon^{IJK}W_\mu^{I\nu}W_\nu^{J\rho}W_\rho^{K\mu}$	$c_{Qq}^{(3,8)}$	$(\bar{Q}\sigma^iT^a\gamma_\mu Q)(\bar{q}\sigma^iT^a\gamma^\mu q)$
c_{HDD}	$(H^\dagger D^\mu H)^*(H^\dagger D_\mu H)$	$c_{qq}^{(3,1)}$	$(\bar{q}\sigma^i\gamma_\mu q)(\bar{q}\sigma^i\gamma^\mu q)$
c_{HG}	$H^\dagger H G_{\mu\nu}^A G^{A\mu\nu}$	$c_{tu}^{(1)}$	$(\bar{t}\gamma_\mu t)(\bar{u}\gamma^\mu u)$
c_{HB}	$H^\dagger H B_{\mu\nu} B^{\mu\nu}$	$c_{tu}^{(8)}$	$(\bar{t}T^a\gamma_\mu t)(\bar{u}T^a\gamma^\mu u)$
c_{HW}	$H^\dagger H W_{\mu\nu}^I W^{I\mu\nu}$	$c_{td}^{(1)}$	$(\bar{t}\gamma_\mu t)(\bar{d}\gamma^\mu d)$
c_{HWB}	$H^\dagger \tau^I H W_{\mu\nu}^I B^{\mu\nu}$	$c_{td}^{(8)}$	$(\bar{t}T^a\gamma_\mu t)(\bar{d}T^a\gamma^\mu d)$
$c_{Hl,11}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_1\gamma^\mu l_1)$	$c_{Qu}^{(1)}$	$(\bar{Q}\gamma_\mu Q)(\bar{u}\gamma^\mu u)$
$c_{Hl,22}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_2\gamma^\mu l_2)$	$c_{Qu}^{(8)}$	$(\bar{Q}T^a\gamma_\mu Q)(\bar{u}T^a\gamma^\mu u)$
$c_{Hl,33}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_3\gamma^\mu l_3)$	$c_{Qd}^{(1)}$	$(\bar{Q}\gamma_\mu Q)(\bar{d}\gamma^\mu d)$
$c_{Hl,11}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{l}_1\tau^I\gamma^\mu l_1)$	$c_{Qd}^{(8)}$	$(\bar{Q}T^a\gamma_\mu Q)(\bar{d}T^a\gamma^\mu d)$
$c_{Hl,22}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{l}_2\tau^I\gamma^\mu l_2)$	$c_{tq}^{(1)}$	$(\bar{q}\gamma_\mu q)(\bar{t}\gamma^\mu t)$
$c_{Hl,33}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{l}_3\tau^I\gamma^\mu l_3)$	$c_{tq}^{(8)}$	$(\bar{q}T^a\gamma_\mu q)(\bar{t}T^a\gamma^\mu t)$
$c_{He,11}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_1\gamma^\mu e_1)$	$c_{eH,22}$	$(H^\dagger H)(\bar{l}_2 e_2 H)$
$c_{He,22}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_2\gamma^\mu e_2)$	$c_{eH,33}$	$(H^\dagger H)(\bar{l}_3 e_3 H)$
$c_{He,33}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_3\gamma^\mu e_3)$	c_{uH}	$(H^\dagger H)(\bar{q}Y_u^\dagger u \tilde{H})$
$c_{Hq}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}\gamma^\mu q)$	c_{tH}	$(H^\dagger H)(\bar{Q}\tilde{H}t)$
$c_{Hq}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{q}\tau^I\gamma^\mu q)$	c_{bH}	$(H^\dagger H)(\bar{Q}Hb)$
c_{Hu}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p\gamma^\mu u_r)$	c_{tG}	$(\bar{Q}\sigma^{\mu\nu}T^A t)\tilde{H}G_{\mu\nu}^A$
c_{Hd}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p\gamma^\mu d_r)$	c_{tW}	$(\bar{Q}\sigma^{\mu\nu}t)\tau^I\tilde{H}W_{\mu\nu}^I$
$c_{HQ}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{Q}\gamma^\mu Q)$	c_{tB}	$(\bar{Q}\sigma^{\mu\nu}t)\tilde{H}B_{\mu\nu}$
$c_{HQ}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{Q}\tau^I\gamma^\mu Q)$	$c_{ll,1221}$	$(\bar{l}_1\gamma_\mu l_2)(\bar{l}_2\gamma^\mu l_1)$
c_{Ht}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{t}\gamma^\mu t)$		
c_{Hb}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{b}\gamma^\mu b)$		

Calculations for Higgs boson production modes with tree-level diagrams are performed with SMEFT_{SM} [96] assuming the ‘top’ flavour symmetry scheme, following the recommendation of Ref. [95] and providing as input $G_F = 1.1663787 \times 10^{-5} \text{ GeV}^{-2}$, $m_Z = 91.1876 \text{ GeV}$, and $m_W = 80.387 \text{ GeV}$, where G_F is the Fermi constant and m_Z and m_W are the Z and W boson masses respectively. Cross-sections are calculated at NLO accuracy in QCD for ggF , $gg \rightarrow ZH$ and $H \rightarrow gg$ with SMEFTatNLO [97] and at NLO EW accuracy for SMEFT-SM interference terms in $H \rightarrow \gamma\gamma$ [98] and $H \rightarrow Z\gamma$ [99]. Lowest-order calculations in QCD for $m_H = 125.09 \text{ GeV}$ are used for all other production and decay modes. SMEFT modifications to the background processes in the included analyses are not considered, but their impact on the SMEFT measurements of the signal is expected to be small as most background distributions are constrained by control regions.

In the simulation, the following requirements are applied at particle level: the minimal jet transverse momentum is $p_T > 20 \text{ GeV}$ and an angular separation $\Delta R > 0.05$ between two jets or two leptons is required, to avoid divergences in the matrix element calculation. For all events, PYTHIA8 was used for the simulation of parton showering, where the Higgs boson decay is based on the width calculated by MADGRAPH for the EFT parameters in question. In ggF events a matching was performed with the MLM algorithm [100, 101] to remove phase space overlap between the jets from the matrix element and the shower.

For analyses based on the STXS methodology, the RIVET program [102] is used to analyse the simulated events, where the computation of high-level kinematic quantities and classification of the events according to their STXS region is performed by the HiggsTemplateCrossSections in accordance with definitions in [49].⁴

The cross-section predictions for a specific process, calculated as described above, are estimated as the sum of three terms:

$$\sigma_{\text{SMEFT}} = \sigma_{\text{SM}} + \sigma_{\text{int}} + \sigma_{\text{BSM}}, \quad (5)$$

where σ_{SM} is the SM cross-section, σ_{int} describes the interference between the SMEFT operators (BSM processes) and SM operators, and σ_{BSM} is the cross-section involving exclusively SMEFT operators. When considering only $d = 6$ SMEFT operators, it follows from Eq. (4) that σ_{int} consists of terms involving a single $d = 6$ SMEFT operator, suppressing each term by a factor Λ^{-2} , and that σ_{BSM} contains terms involving products of two $d = 6$ SMEFT operators, suppressing each term by a factor Λ^{-4} . For this reason, the impact of the σ_{BSM} term is generally expected to be small, though its impact may still be non-negligible in certain regions of phase space, e.g., when energy scales are of order Λ .

Additional terms suppressed by a factor Λ^{-4} occur in the general SMEFT expansion of Eq. (4), i.e., the lowest-order cross-section terms generated by $d = 8$ operators that involve interference between SM and these operators. As only a subset of Higgs boson processes have been calculated with $d = 8$ operators [103–105], the list of terms suppressed by a factor Λ^{-4} considered is incomplete and the effect of the missing $d = 8$ terms relative to σ_{BSM} is not known. However, their effect may be of a magnitude comparable to that of products of two $d = 6$ SMEFT operators, as the suppression factor is at the same order. For this reason, SMEFT interpretations are presented both with and without the σ_{BSM} contribution to give a general indication of the sensitivity of the analysis to Λ^{-4} terms.

⁴ The RIVET algorithm has been modified to classify events in which a Higgs boson and two leptons arise from the same production vertex as VH production events. This change ensures the proper classification of events with leptons from off-shell V decays, since MADGRAPH only saves on-shell intermediate particles. Contributions from off-shell V bosons are small in the SM, but can be enhanced by SMEFT operators.

To reduce the perturbative QCD uncertainty in the extraction of the SMEFT Wilson coefficients, the parameterisation of the predicted cross-section of Eq. (5) is computed as a relative correction to the SM prediction computed at the highest available order for each process:

$$\sigma_{\text{SMEFT}}^i = \sigma_{\text{SM}}^{i,((\text{N})\text{N})\text{NLO}} \times \left(1 + \frac{\sigma_{\text{int}}^{i,(\text{N})\text{LO}}}{\sigma_{\text{SM}}^{i,(\text{N})\text{LO}}} + \frac{\sigma_{\text{BSM}}^{i,(\text{N})\text{LO}}}{\sigma_{\text{SM}}^{i,(\text{N})\text{LO}}} \right). \quad (6)$$

This calculation strategy assumes that higher-order terms have the same relative effect on σ_{int} and σ_{BSM} as on σ_{SM} [106] and that the SMEFT operator predominantly modifies the production process i under consideration.

The predictions are further modified by the impact of SMEFT operators on Higgs boson decay branching ratios. Since the Higgs boson is a narrow, scalar particle, and only on-shell production is considered in this analysis, its production cross-section and decay width factorise. The impact of SMEFT operators on production and decay therefore also factorises and can be derived independently. Thus, the cross-section for a given Higgs boson production process i in particle-level region k' and for a given decay mode $H \rightarrow X$ is

$$(\sigma \times \mathcal{B})_{\text{SMEFT}}^{i,k',H \rightarrow X} = \sigma_{\text{SMEFT}}^{i,k'} \times \mathcal{B}_{\text{SMEFT}}^{H \rightarrow X} = \left(\sigma_{\text{SM}}^{i,k'} + \sigma_{\text{int}}^{i,k'} + \sigma_{\text{BSM}}^{i,k'} \right) \times \left(\frac{\Gamma_{\text{SM}}^{H \rightarrow X} + \Gamma_{\text{int}}^{H \rightarrow X} + \Gamma_{\text{BSM}}^{H \rightarrow X}}{\Gamma_{\text{SM}}^H + \Gamma_{\text{int}}^H + \Gamma_{\text{BSM}}^H} \right).$$

The factorised SMEFT prediction is calculated with ratios as in Eq. (6) to utilise the SM prediction at the highest available order:

$$(\sigma \times \mathcal{B})_{\text{SMEFT}}^{i,k',H \rightarrow X} = (\sigma \times \mathcal{B})_{\text{SM},(\text{N}(\text{N}))\text{NLO}}^{i,k',H \rightarrow X} \left(1 + \frac{\sigma_{\text{int},(\text{N})\text{LO}}^{i,k'}}{\sigma_{\text{SM},(\text{N})\text{LO}}^{i,k'}} + \frac{\sigma_{\text{BSM},(\text{N})\text{LO}}^{i,k'}}{\sigma_{\text{SM},(\text{N})\text{LO}}^{i,k'}} \right) \left(\frac{1 + \frac{\Gamma_{\text{int}}^{H \rightarrow X}}{\Gamma_{\text{SM}}^{H \rightarrow X}} + \frac{\Gamma_{\text{BSM}}^{H \rightarrow X}}{\Gamma_{\text{SM}}^{H \rightarrow X}}}{1 + \frac{\Gamma_{\text{int}}^H}{\Gamma_{\text{SM}}^H} + \frac{\Gamma_{\text{BSM}}^H}{\Gamma_{\text{SM}}^H}} \right), \quad (7)$$

where the ratios $\sigma_{\text{int}}/\sigma_{\text{SM}}$ and $\Gamma_{\text{int}}/\Gamma_{\text{SM}}$ have a linear dependence on SMEFT operators and are suppressed by a factor Λ^{-2} , and the ratios $\sigma_{\text{BSM}}/\sigma_{\text{SM}}$ and $\Gamma_{\text{BSM}}/\Gamma_{\text{SM}}$ have a quadratic dependence on SMEFT operators and are suppressed by a factor Λ^{-4} . In the analysis, these ratios are parameterised as

$$\frac{\sigma_{\text{int}}^{i,k'}}{\sigma_{\text{SM}}^{i,k'}} = \sum_j A_j^{\sigma_{i,k'}} c_j \quad \frac{\sigma_{\text{BSM}}^{i,k'}}{\sigma_{\text{SM}}^{i,k'}} = \sum_{j,l \geq j} B_{jl}^{\sigma_{i,k'}} c_j c_l \quad (8)$$

$$\frac{\Gamma_{\text{int}}^{H \rightarrow X}}{\Gamma_{\text{SM}}^{H \rightarrow X}} = \sum_j A_j^{\Gamma^{H \rightarrow X}} c_j \quad \frac{\Gamma_{\text{BSM}}^{H \rightarrow X}}{\Gamma_{\text{SM}}^{H \rightarrow X}} = \sum_{j,l \geq j} B_{jl}^{\Gamma^{H \rightarrow X}} c_j c_l \quad (9)$$

$$\frac{\Gamma_{\text{int}}^H}{\Gamma_{\text{SM}}^H} = \sum_j A_j^{\Gamma^H} c_j \quad \frac{\Gamma_{\text{BSM}}^H}{\Gamma_{\text{SM}}^H} = \sum_{j,l \geq j} B_{jl}^{\Gamma^H} c_j c_l, \quad (10)$$

with

$$A_j^{\Gamma^H} = \frac{\sum_X \Gamma_{\text{SM}}^{H \rightarrow X} A_j^{\Gamma^{H \rightarrow X}}}{\sum_X \Gamma_{\text{SM}}^{H \rightarrow X}} \quad B_{jl}^{\Gamma^H} = \frac{\sum_X \Gamma_{\text{SM}}^{H \rightarrow X} B_{jl}^{\Gamma^{H \rightarrow X}}}{\sum_X \Gamma_{\text{SM}}^{H \rightarrow X}}. \quad (11)$$

In Eq. (11), the sum includes all possible final states X with up to four elementary particles produced by one of the following Higgs boson decays: $H \rightarrow f\bar{f}$ ($f = c, b, \mu, \tau$), $H \rightarrow \gamma\gamma$, $H \rightarrow Z\gamma$, and $H \rightarrow VV^*$ ($V = W, Z$). All $A_j^{\sigma_{i,k'}}$, $A_j^{\Gamma^{H \rightarrow X}}$, $B_{jl}^{\sigma_{i,k'}}$ and $B_{jl}^{\Gamma^{H \rightarrow X}}$ coefficients are constant factors obtained from simulation that express the sensitivity of the process to the operators \mathcal{O}_j and \mathcal{O}_l that correspond to the Wilson coefficients c_j and c_l , where the indices j, l run over all non-negligible operators. The coefficients A_j and B_{jl} are proportional to Λ^{-2} and Λ^{-4} respectively. The values of all SM Higgs boson branching ratios that enter Eq. (11) are taken from Ref. [49].

For the interpretation of the differential analyses, where the measured parameters $\mu^{k',X}$ do not label individual production processes, the relative fractions of the different processes are predicted according to the SMEFT calculations as in Eq. (6).

Using Eqs. (7)–(10), two statistical models are constructed for the interpretation of the data: a linearised variant that only considers terms suppressed by up to a factor Λ^{-2} , and a variant including quadratic terms with suppression factor Λ^{-4} .

3.1.2 Cross-section calculation with linear terms

In a scenario where Λ^{-4} -suppressed contributions are ignored, the predicted deviation of the cross-section, partial width and total width from their SM values can each be explicitly linearised as a function of the Wilson coefficients c . Ignoring all Λ^{-4} -suppressed BSM terms in Eq. (7), and using the parameterisation of Eqs. (8)–(10), the expression for the cross-section times branching ratio reduces to

$$\begin{aligned}
(\sigma \times \mathcal{B})_{\text{SMEFT}}^{i,k',H \rightarrow X} &= (\sigma \times \mathcal{B})_{\text{SM},((N)N)\text{NLO}}^{i,k',H \rightarrow X} \times \left(1 + \frac{\sigma_{\text{int},(N)\text{LO}}^{i,k'}}{\sigma_{\text{SM},(N)\text{LO}}^{i,k'}} \right) \times \left(\frac{1 + \frac{\Gamma_{\text{int}}^{H \rightarrow X}}{\Gamma_{\text{SM}}^{H \rightarrow X}}}{1 + \frac{\Gamma_{\text{int}}^H}{\Gamma_{\text{SM}}^H}} \right) \\
&= (\sigma \times \mathcal{B})_{\text{SM},((N)N)\text{NLO}}^{i,k',H \rightarrow X} \times \left(1 + \sum_j A_j^{\sigma_{i,k'}} c_j \right) \times \left(\frac{1 + \sum_j A_j^{\Gamma^{H \rightarrow X}} c_j}{1 + \sum_j A_j^{\Gamma^H} c_j} \right), \\
&= (\sigma \times \mathcal{B})_{\text{SM},((N)N)\text{NLO}}^{i,k',H \rightarrow X} \times \left(\frac{1 + \sum_j \left(A_j^{\sigma_{i,k'}} + A_j^{\Gamma^{H \rightarrow X}} \right) c_j + O(\Lambda^{-4})}{1 + \sum_j A_j^{\Gamma^H} c_j + O(\Lambda^{-4})} \right), \quad (12)
\end{aligned}$$

where all higher order terms in the expansion are suppressed by power Λ^{-4} or beyond.

A subsequent Taylor expansion of the width ratio expression of Eq. (12) and truncation of terms beyond Λ^{-2} would result in a completely linearised expression for $\sigma \times \mathcal{B}$. However, as the parametric dependence of a Wilson coefficient in a branching fraction f is effectively of the form $c/(1 + f \cdot c)$, the linearity in c of this expression assumed for small values of c does not hold for large values of c . This effect is particularly pronounced for operators affecting Higgs boson decays with a large branching fraction, e.g., c_{bH} , and operators that have a large measured uncertainty that allows large values of c in the error propagation. For this reason, a full Taylor expansion of the ratio is not performed.

3.1.3 Cross-section calculation including quadratic terms

The SMEFT prediction including the available terms proportional to Λ^{-4} is:

$$\begin{aligned}
(\sigma \times \mathcal{B})_{\text{SMEFT}}^{i,k',H \rightarrow X} &= (\sigma \times \mathcal{B})_{\text{SM},((\text{N})\text{N})\text{NLO}}^{i,k',H \rightarrow X} \left(1 + \sum_j A_j^{\sigma_{i,k'}} c_j + \sum_{j,l \geq j} B_{jl}^{\sigma_{i,k'}} c_j c_l \right) \left(\frac{1 + \sum_j A_j^{\Gamma_{H \rightarrow X}} c_j + \sum_{j,l \geq j} B_{jl}^{\Gamma_{H \rightarrow X}} c_j c_l}{1 + \sum_j A_j^{\Gamma_H} c_j + \sum_{j,l \geq j} B_{jl}^{\Gamma_H} c_j c_l} \right), \\
&= (\sigma \times \mathcal{B})_{\text{SM},((\text{N})\text{N})\text{NLO}}^{i,k',H \rightarrow X} \cdot \\
&\quad \left(\frac{1 + \sum_j (A_j^{\sigma_{i,k'}} + A_j^{\Gamma_{H \rightarrow X}}) c_j + \sum_{j,l} (A_j^{\sigma_{i,k'}} A_l^{\Gamma_{H \rightarrow X}}) c_j c_l + \sum_{j,l \geq j} (B_{jl}^{\sigma_{i,k'}} + B_{jl}^{\Gamma_{H \rightarrow X}}) c_j c_l + \mathcal{O}(\Lambda^{-6})}{1 + \sum_j (A_j^{\Gamma_H}) c_j + \sum_{j,l \geq j} (B_{jl}^{\Gamma_H}) c_j c_l + \mathcal{O}(\Lambda^{-6})} \right) \quad (13)
\end{aligned}$$

where both the numerator and denominator are a second-order Taylor expansion resulting in a linearised expression for terms of order Λ^{-2} and a quadratic expression for terms of order Λ^{-4} . Similarly to Eq. (12), and for the same reason, the width ratio expression in Eq. (13) is not subjected to a further Taylor expansion. While the set of operators proportional to Λ^{-4} considered in Eq. (13) is not complete due to missing $d = 8$ operators, as noted earlier a comparison of results obtained with the linear statistical model and the model including quadratic terms is indicative of the sensitivity of the measurements to terms suppressed by Λ^{-4} .

Figure 4 illustrates the impact of the linear terms A_j for the three SMEFT operators considered in the interpretation of the fiducial differential cross-section measurements, whereas Figure 5 shows the impact of linear terms and the impact including quadratic terms on production ($A_j^{\sigma_{i,k'}}$) and decay ($\frac{1+A_j^{\Gamma_{H \rightarrow X}}}{1+A_j^{\Gamma_H}} - 1$) rates.

The relative importance of the quadratic term increases linearly with the considered variation of the corresponding Wilson coefficient. Figure 5 shows that the quadratic terms have the highest impact in the high- p_T regions of VH production for the coefficients $c_{Hq}^{(3)}$, $c_{Hq}^{(1)}$, c_{Hd} , and c_{Hu} , on high- p_T (inclusive) $t\bar{t}H$ production for the coefficients c_{tG} , $c_{Qq}^{(1,8)}$, $c_{Qq}^{(3,1)}$, $c_{tq}^{(8)}$, $c_{Qu}^{(8)}$, $c_{tu}^{(8)}$, (c_G) and in the tH channel for the coefficients $c_{HQ}^{(3)}$ and $c_{Qq}^{(3,1)}$.

3.1.4 Assumptions on SM Higgs boson kinematics

With the insertion of the SMEFT cross-section predictions of Eqs. (12) or (13) in Eqs. (2) and (3) describing the expected signal yield s_k for each analysis region k , i.e.,

$$\begin{aligned}
\mu^{i,k',X} &\rightarrow \frac{(\sigma \times \mathcal{B})_{\text{SM}+\Lambda^{-2,-4}}^{i,k',H \rightarrow X}}{(\sigma \times \mathcal{B})_{\text{SM},((\text{N})\text{N})\text{NLO}}^{i,k',H \rightarrow X}} \text{ in Eq. (2),} \\
\mu^{k',X} &\rightarrow \frac{\sigma_{\text{SM}+\Lambda^{-2,-4},\text{fid.}}^{k',H \rightarrow X}}{\sigma_{\text{SM},\text{fid.}}^{k',H \rightarrow X}} \text{ in Eq. (3),}
\end{aligned}$$

the effect of SMEFT cross-section modifications on the inclusive Higgs boson signal yield for each particle-level region k' is fully taken into account for every reconstruction-level region k . However, the




Figure 4: Expected impact of the SMEFT operators on the fiducial differential distributions of the transverse momentum of the Higgs boson decay products in $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^*$ decays, relative to the SM cross-section, assuming the linearised SMEFT model. The values of the Wilson coefficients, specified in the legend, are chosen to show the distribution of the operator impact in the same range as the typical uncertainty of the measurement. To judge the experimental sensitivity to constrain the operators from the data in the listed fiducial regions, the total uncertainty in the measurement in each region is shown in the top panel. For presentational clarity, the uncertainty of low precision regions is clipped off in the plot. The impact of these three operators in the STXS analysis is shown in matching colours in the bottom panel of Figure 5 for coefficient variations of identical magnitude.

SMEFT operators may also affect other terms of the likelihood, notably the efficiency times acceptance factors $\epsilon_k^{i,X,k'}$ occurring in Eqs. (2) and (3), as well as the signal yield distribution inside each region k represented by the factor $f_s^{r,k}$ occurring in Eq. (1).

In both the linear SMEFT model and the model including quadratic terms, it is assumed that the theoretical systematic uncertainties assigned to the acceptance factors $\epsilon_k^{i,X,k'}$, which are fully taken into account in the SM cross-section prediction, are large compared to the effect of the possible acceptance changes induced by SMEFT operators in Higgs boson production through the full validity range of the SMEFT model, and that this additional SMEFT related theory uncertainty can therefore be neglected.

This assumption is motivated by the similarity of reconstruction-level analysis regions to STXS and differential fiducial regions, which are designed to be relatively insensitive to acceptance changes induced by SMEFT operators. In addition, the rather fine binning of the Stage-1.2 STXS regions reduces further the possibility of significant variations of the acceptance factors inside each region. The effect of SMEFT operators on other observables used in the definition of the analysis regions, e.g., through multivariate discriminants, is assumed to be negligible.

Conversely, the effect of SMEFT operators on Higgs boson decays can strongly affect the acceptance factors $\epsilon_k^{i,X}$, since decays are not limited to a restricted fiducial phase space in the STXS framework. For 2-body decays, such acceptance effects are generally small [25]. However, the four-body $H \rightarrow ZZ^* \rightarrow 4\ell$ and $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ decays are significantly impacted by acceptance effects, hence for these decays the impact factors $A_j^{H \rightarrow X}$ and $B_{jl}^{H \rightarrow X}$ are recalculated including an approximate implementation of the reconstruction-level requirements at the particle level. The impact of acceptance effects is evaluated for all operators in Table 2 that can affect $\Gamma^{H \rightarrow ZZ^*}$ or $\Gamma^{H \rightarrow WW^*}$, i.e., $c_{H\Box}$, c_{HDD} , c_{HW} , c_{HB} , c_{HWB} , $c_{HI,11}^{(1)}$, $c_{HI,22}^{(1)}$,




Figure 6: Illustration of the effect of a variation of the SMEFT coefficient c_{HW} on kinematic observables used in the event selection: (a) impact on the dilepton invariant mass in the $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ analysis, and (b) impact on the invariant mass of the 3rd and 4th p_T -ranked lepton in the $H \rightarrow ZZ^* \rightarrow 4\ell$ analysis. The vertical dotted lines indicate the selection criteria applied to the corresponding variables by the $H \rightarrow WW^*$ analysis ($10 < m_{\ell\ell} < 55$ GeV) and by the $H \rightarrow ZZ^*$ analysis ($m_{Z\ell} > 12$ GeV). The shaded bands show the statistical uncertainty of the SMEFT simulation. The value of the Wilson coefficient ($c_{HW} = 1$) is significantly larger than the sensitivity of this study, and is chosen for illustration purposes.

Finally, if the signal yield distribution inside an analysis region k deviates from the SM distribution encoded in $f_s^{r,k}$ in Eq. (1), the measured yield s_k may be biased. This effect is expected to be negligible for all regions where the discriminant observable inside the region is the reconstructed invariant mass of the Higgs boson, which is largely unaffected by SMEFT operators. Regions that feature a multivariate discriminant as the observable can potentially be more affected by SMEFT induced shape differences, but are only used in regions with low signal yields where it is assumed that any effect induced by signal shape deviations is small compared to the statistical uncertainties. For the high-yield $H \rightarrow WW^*$ channel, where the transverse mass m_T is used as discriminant observable, which is sensitive to SMEFT operators, it is verified that the effect of shape variations induced by the SMEFT operators is negligible for the range of coefficient values that are consistent with the observed measurement.

3.2 Constraints from STXS measurements

The aim of the EFT analysis is to obtain constraints on the Wilson coefficients c_j through a maximum likelihood analysis of the Higgs boson STXS measurements shown in Figures 1–3. All signal strength

modifiers $\mu^{i,k',X}$ of Eq. (2) are directly expressed in terms of the coefficients c_j :

$$s_k(\mathbf{c}, \boldsymbol{\theta}) = \sum_{i,k',X} \left(\mu^{i,k',X} \equiv \frac{(\sigma \times \mathcal{B})_{\text{SMEFT}}^{i,k',H \rightarrow X}(\mathbf{c})}{(\sigma \times \mathcal{B})_{\text{SM},(\text{N}(\text{N}))\text{NLO}}^{i,k',H \rightarrow X}} \right) \times \mathcal{L} \times (\sigma \times \mathcal{B})_{\text{SM},(\text{N}(\text{N}))\text{NLO}}^{i,k',H \rightarrow X}(\boldsymbol{\theta}) \times \epsilon_k^{i,k',X}(\boldsymbol{\theta}),$$

with $\mathcal{L}, \epsilon_k^{i,k',X}(\boldsymbol{\theta})$ and $\boldsymbol{\theta}$ as defined in Eq. (2), and where the signal cross-section $(\sigma \times \mathcal{B})_{\text{SMEFT}}^{i,k',H \rightarrow X}(\mathbf{c})$ is either taken from the linear model of Eq. (12) or the model including quadratic terms of Eq. (13).

The available data samples contain insufficient information to constrain all coefficients \mathbf{c} listed in Table 2 simultaneously. A standard numerical joint maximum likelihood estimation of this set of parameters \mathbf{c} will not converge due to the presence of flat directions in the parameter space, corresponding to the unconstrained degrees of freedom.

As degrees of freedom left unconstrained by the data do not necessarily correspond to individual coefficients c_j , but may also be linear combinations $\sum_j a_j c_j$, a modified basis is defined and used in the following, to ensure the convergence of the numerical likelihood maximisation procedure for the largest number of degrees of freedom that can be probed by the current set of measurements.

3.2.1 Sensitivity estimate and choice of parameters

To determine a modified basis \mathbf{c}' that can be estimated from the data, the SM expected covariance matrix V_{STXS} of the measurement, expressed in the STXS basis $\{\mu^{i,k',X}\}$, is analysed. From V_{STXS} , the Hessian matrix V_{STXS}^{-1} is obtained, and then rotated to the SMEFT basis $\{c_j\}$:

$$V_{\text{SMEFT}}^{-1} = P_{(i,k',X) \rightarrow (j)}^T V_{\text{STXS}}^{-1} P_{(i,k',X) \rightarrow (j)}. \quad (14)$$

The expected Jacobian matrix $P_{(i,k',X) \rightarrow (j)}$ is based on the linearised SMEFT model of Eq. (12) assuming small c_j :

$$P_{(i,k',X) \rightarrow (j)} = A_j^{\sigma_{i,k'}} + A_j^{\Gamma^{H \rightarrow X}} - A_j^{\Gamma^H}, \quad (15)$$

where $A_j^{\sigma_{i,k'}}$, $A_j^{\Gamma^{H \rightarrow X}}$ and $A_j^{\Gamma^H}$ are the constant factors obtained from the simulation.

In the limit of Gaussian STXS measurements, the matrix V_{SMEFT}^{-1} represents the Fisher information matrix of its linearised SMEFT model re-parameterisation, with the additional caveat that for large coefficient variations the representation of V_{SMEFT}^{-1} may be suboptimal, as a consequence of the fully linearized form of Eq. (15) that is only valid in the limit of small coefficient values. An eigenvalue decomposition of the Fisher information matrix yields the eigenvectors $ev^{[i]}$ and their corresponding eigenvalues λ_i and variances $1/\lambda_i$.

Figure 7 lists the eigenvectors obtained from the expected measurements accounting for the observed values of nuisance parameters, ranked by eigenvalue and truncated to eigenvalues $\lambda_i \geq 0.01$, corresponding to a truncation at an estimated uncertainty of $\sigma(ev^{[i]}) \leq 10$, well beyond the validity range of the Wilson coefficients of $\mathcal{O}(1)$.

From the ranking shown in Figure 7 and a survey of the sensitivity of the STXS regions to the Wilson coefficients in the linearised model, as shown in Figure 5, the following observations are made:




Figure 7: Eigenvectors of the inverse EFT covariance matrix obtained by propagating the SMEFT parameterisation to the covariance matrix V_{STXS} and requiring a significant (≥ 0.01) eigenvalue λ . The corresponding expected uncertainty $\sigma_{\text{exp}} = 1/\sqrt{\lambda}$ for each eigenvector is also shown.

- The coefficients $c_{eH,33}$, $c_{eH,22}$, and c_{bH} , representing the Yukawa coupling modifiers of the $H \rightarrow \tau\tau$, $H \rightarrow \mu\mu$, and $H \rightarrow bb$ decays respectively, can be individually measured from the corresponding Higgs boson channels that enter the combination. There are weak correlations of these operators with other SMEFT operators, primarily for c_{bH} , through their contribution to the total Higgs boson width.
- The coefficient $c_{Hq}^{(3)}$ affects both the WH and ZH production modes with an increasing impact as a function of the transverse momentum p_T^V of the W and Z bosons, and is constrained almost exclusively by the $VH, H \rightarrow b\bar{b}$ analysis.
- The coefficients c_{HG} , c_{tG} and c_{tH} are constrained by ggF and $t\bar{t}H$ production. As the uncertainties of the constrained directions span more than two orders of magnitude, it is beneficial to represent them by decorrelated parameters, even though the rotation matrix required to decorrelate them is close to an identity matrix.
- The coefficients c_{HB} , c_{HW} , c_{HWB} , c_{tB} and c_{tW} are constrained in two directions through their impacts on the branching ratios of the $H \rightarrow \gamma\gamma$ and $H \rightarrow Z\gamma$ decays. There is an additional small impact of these operators on VBF and VH production, mildly constraining a third direction in this parameter group.
- The coefficients c_{Hu} , c_{Hd} , c_{Ht} , c_{Hb} and $c_{Hq}^{(1)}$ mainly affect the neutral current interactions with quarks, while $c_{He,33}$, $c_{Hl,33}^{(1)}$, $c_{Hl,33}^{(3)}$ mainly affect the W and Z vertices with third generation leptons. Both sets of coefficients are primarily constrained by the $VH, H \rightarrow b\bar{b}$ analysis. The small sensitivity to c_{Ht} and c_{Hb} arises from the p_T -dependent corrections that they induce in the subleading production mode $gg \rightarrow ZH$.
- The coefficients of the four-fermion operators involving the top-quark, i.e., $c_{Qq}^{(1,8)}$, $c_{Qq}^{(3,1)}$, $c_{tq}^{(8)}$, $c_{Qu}^{(8)}$, $c_{tu}^{(8)}$, $c_{td}^{(8)}$, $c_{Qd}^{(8)}$, $c_{Qq}^{(3,8)}$, $c_{Qq}^{(1,1)}$, $c_{tu}^{(1)}$, $c_{tq}^{(1)}$, $c_{Qu}^{(1)}$, and $c_{Qd}^{(1)}$, as well as the trilinear gluon coupling c_G , affect the $t\bar{t}H$ and tH production modes and are largely degenerate. The p_T^H spectrum measured in the $H \rightarrow \gamma\gamma$ and $H \rightarrow bb$ channels constrains two linear combinations of these 14 coefficients. The separate constraint on tH production in the $H \rightarrow \gamma\gamma$ analysis weakly constrains a third direction in this group.
- The coefficients $c_{H,1221}$, $c_{Hl,11}^{(3)}$, and $c_{Hl,22}^{(3)}$ primarily cause a shift in the Fermi constant, resulting in an overall normalisation factor across different production modes. The coefficient $c_{H\Box}$ only affects the measurements through a Higgs boson propagator correction and thus acts similarly as an overall scale factor in the observed cross-sections. Only a single linear combination of these four operators can be constrained from the data.
- The coefficients $c_{Hl,11}^{(1)}$, $c_{Hl,22}^{(1)}$, $c_{He,11}$ and $c_{He,22}$ introduce anomalous $HZee$ and $HZ\mu\mu$ vertices and are mainly constrained by the branching ratio of the Higgs boson decay into four leptons (electrons or muons), together with c_{HDD} (responsible for an anomalous HZZ vertex) and $c_{HQ}^{(3)}$, $c_{HQ}^{(1)}$ (leading to $HZtt$ and $HZbb$ couplings that modify the Z boson propagator, but to which the Higgs boson measurements included here are not sensitive). The $ZH, H \rightarrow b\bar{b}$ analysis also constrains the couplings to first- and second-generation leptons such as $c_{He,11}$, $c_{Hl,11}^{(1)}$, $c_{He,22}$, $c_{Hl,22}^{(1)}$, but the constraints are significantly looser than those from the $H \rightarrow ZZ^* \rightarrow 4\ell$ decay.

Based on these observations, a new fit basis \mathbf{c}' is defined that achieves both fit stability and fit-parameter interpretability. This basis \mathbf{c}' is expressed in terms of single Warsaw basis coefficients c_j whenever possible, and in terms of linear combinations (denoted by e) of coefficients when necessary:

$$\begin{aligned}
\mathbf{c} &= \{c_{eH,22}\} \cup & \mathbf{c}' &= \{c_{eH,22}\} \cup \\
&\{c_{eH,33}\} \cup & &\{c_{eH,33}\} \cup \\
&\{c_{Hq}^{(3)}\} \cup & &\{c_{Hq}^{(3)}\} \cup \\
&\{c_{bH}\} \cup & &\{c_{bH}\} \cup \\
&\{c_{HG}, c_{tG}, c_{tH}\} \cup & \rightarrow &\{e_{\text{ggF}}^{[1]}, e_{\text{ggF}}^{[2]}, e_{\text{ggF}}^{[3]}\} \cup \\
&\{c_{HB}, c_{HW}, c_{HWB}, c_{tB}, c_{tW}\} \cup & \rightarrow &\{e_{H\gamma\gamma, Z\gamma}^{[1]}, e_{H\gamma\gamma, Z\gamma}^{[2]}, e_{H\gamma\gamma, Z\gamma}^{[3]}\} \cup \\
&\{c_{Hu}, c_{Hq}^{(1)}, c_{Hd}, c_{Hl,33}^{(3)}, & & \\
&\quad c_{Ht}, c_{He,33}, c_{Hl,33}^{(1)}, c_{Hb}\} \cup & \rightarrow &\{e_{ZH}^{[1]}, e_{ZH}^{[2]}, e_{ZH}^{[3]}, e_{ZH}^{[4]}\} \cup \\
&\{c_G, c_{Qq}^{(1,8)}, c_{Qq}^{(3,1)}, c_{tq}^{(8)}, c_{Qu}^{(8)}, c_{tu}^{(8)}, c_{td}^{(8)}, & & \\
&\quad c_{Qd}^{(8)}, c_{Qq}^{(3,8)}, c_{Qq}^{(1,1)}, c_{tu}^{(1)}, c_{tq}^{(1)}, c_{Qu}^{(1)}, c_{Qd}^{(1)}\} \cup & \rightarrow &\{e_{\text{tH}}^{[1]}, e_{\text{tH}}^{[2]}, e_{\text{tH}}^{[3]}\} \cup \\
&\{c_{H\Box}, c_{Hl,11}^{(3)}, c_{Hl,22}^{(3)}, c_{ll,1221}\} \cup & \rightarrow &\{e_{\text{glob}}^{[1]}\} \cup \\
&\{c_{Hl,11}^{(1)}, c_{Hl,22}^{(1)}, c_{He,11}, c_{He,22}, c_{HDD}, c_{HQ}^{(3)}, c_{HQ}^{(1)}\} & \rightarrow &\{e_{Hllll}^{[1]}\}.
\end{aligned}$$

Only subgroup eigenvectors with an expected uncertainty ≤ 10 are retained as model parameters. The remaining coefficients, to which the data are not really sensitive, are fixed to zero in the analysis. The subgroup eigenvector directions relative to the Warsaw basis are shown in Figure 8. The names of the eigenvectors are chosen to reflect the production or decay process that dominates the experimental sensitivity of the parameter group, as shown in Figure 5, although single parameters within some of the groups sometimes have a dominant sensitivity to another process.

No separate optimisation of the parameter basis \mathbf{c}' is performed for the SMEFT model including quadratic terms of Eq. (13) as the non-linear effects of this model are expected to vanish for small c_j , and thus to asymptotically yield the same rotation matrix as Eq. (15).

The impacts of the variations of the coefficients of the rotated basis \mathbf{c}' on the STXS regions and Higgs boson branching ratios are shown in Figure 9, with the same parameter grouping as shown in Figure 5. The figure clearly demonstrates that the basis \mathbf{c}' represents impact variations across regions that are much less correlated than those represented by the Warsaw basis \mathbf{c} . The magnitude of the parameter variations shown in Figure 9 are set to the *expected uncertainty* for each parameter in the linear SMEFT model, thus giving a realistic indication of the magnitude of variations that can be constrained from the data. For completeness, the impact of the quadratic terms for the same values of the parameter basis is overlaid, indicating the relative magnitude of the quadratic terms at the expected sensitivity level of the linear model. The impact of these expected sensitivities for the measured SMEFT parameters is discussed in Section 3.2.3.




Figure 8: Definition of the fit basis coefficients c' in terms of the Warsaw basis coefficients c .




Figure 9: Impact of coefficients of the rotated basis c' on the STXS regions, relative to the SM cross-section, under the assumption of the linearised SMEFT model (shaded histogram) and the SMEFT model including quadratic terms (open histogram). For all coefficients, a variation equal to the expected uncertainty when using the linearised model is shown. Groupings in this figure are consistent with those in Figure 5, but vertical axis scales differ in order to completely show all quadratic impact terms, which can occasionally be very large. To judge the experimental sensitivity to constrain the operators from the data in the listed STXS regions, the total uncertainty in the corresponding regions is shown in the top panel. For presentational clarity, the uncertainty of low precision STXS regions is clipped off in the plot.

3.2.2 Results

Linear model Figure 10 compares the expected and observed results obtained using the linearised SMEFT model, showing good agreement of the observed data with the SM expectation, corresponding to a p -value of 94.5%. The parameter value ranges shown in the bottom panel correspond to 68% and 95% confidence level (CL) intervals, where all other coefficients and nuisance parameters are profiled. The observed uncertainties are generally about 10% smaller than the expected uncertainties, since the assumed Higgs boson width, when computed with the observed parameter values, is smaller than its SM expectation value. This effect is mostly driven by a high observed value of c_{bH} , which corresponds to a reduced value of $\Gamma^{H \rightarrow b\bar{b}}$. This reduction of the observed linear model uncertainties is comparable to that of the reduced observed uncertainties in the κ model analysis of the same data reported in Ref. [9], due to an equivalently low measured value (< 1) of κ_b .

Figure 10 also shows the contribution of each measured Higgs boson decay or production mode to the sensitivity of each measurement of the coefficients of the rotated basis. The contribution of a measurement i to the sensitivity to a coefficient c'_j is determined from the ratio of the Fisher information $\mathcal{I}_i(c'_j)$ of that measurement to the sum of the Fisher information of all contributing components, i.e., $\sum_k \mathcal{I}_k(c'_j)$, where in all cases the Fisher information is estimated as the inverse of the covariance matrix V_{SMEFT} , as defined in Eq. (14). This breakdown reveals that six parameters are (almost) exclusively measured by a single decay mode: the Yukawa coefficients $c_{He,22}$ and $c_{He,33}$ by the decay modes $H \rightarrow \mu\mu$ and $H \rightarrow \tau\tau$ respectively, the coefficient $c_{Hq}^{(3)}$ by $H \rightarrow b\bar{b}$; coefficients $e_{H\gamma\gamma,Z\gamma}^{[1]}$ and $e_{H\gamma\gamma,Z\gamma}^{[2]}$ by $H \rightarrow \gamma\gamma$ and $H \rightarrow Z\gamma$ respectively and the coefficient $e_{Hllll}^{[1]}$ by $H \rightarrow ZZ^*$. Due to its large effect on the Higgs boson width, the measurement of the Yukawa coefficient c_{bH} is not dominated by $H \rightarrow b\bar{b}$, but instead constrained by a combination of measured decays. Similarly, five parameters are measured (almost) exclusively by a single production mode: coefficient $e_{\text{ggF}}^{[1]}$ by ggF production, $e_{ZH}^{[1]}$ by ZH production, $e_{\text{ttH}}^{[1]}$ and $e_{\text{ttH}}^{[2]}$ by $t\bar{t}H$ production at high- p_T and medium- p_T respectively, and $e_{\text{ttH}}^{[3]}$ by tH production. No measured coefficient is predominantly sensitive to $H \rightarrow WW^*$ decays. The VBF or WH production modes each only dominate a single parameter measurement, $c_{He,33}$ and $c_{Hq}^{(3)}$, respectively.

Generally, the tightest constraints on SMEFT coefficients are observed for processes where the SM amplitudes are suppressed by factors that do not enter in SMEFT operators contributing to the same measured final states. The most prominent example of these is the $O_{He,22}$ operator, for which the corresponding SM $H \rightarrow \mu\mu$ amplitude is so strongly suppressed by the Yukawa factor that it is the most precisely measured coefficient, despite being constrained only by the very low-yield $H \rightarrow \mu\mu$ channel. Along similar lines, the precisely measured coefficients $e_{\text{ggF}}^{[1]}$, $e_{H\gamma\gamma,Z\gamma}^{[1]}$ and $e_{H\gamma\gamma,Z\gamma}^{[2]}$ benefit from suppressed SM amplitudes that require loops at lowest order.

Analysis of parameter uncertainty sources Figure 11 illustrates the relative importance of the various uncertainty components on the coefficients measured in the linearised model, where the top panel shows the relative contributions of the statistical and systematic components, and the bottom panel shows the relative contributions of the experimental, signal theory and background theory contributions to the total systematic uncertainty, where signal theory systematic uncertainties include both total cross-section uncertainties and acceptance uncertainties. The breakdown of the total uncertainty into statistical and systematic components is performed as follows. The statistical uncertainty in each parameter is estimated from the 68% confidence interval obtained when fixing to their best-fit values all nuisance parameters associated with systematic uncertainties, and leaving unconstrained all other parameters. The total systematic uncertainty is calculated




Figure 10: Comparison of the expected (gray) and observed (blue) parameters of the rotated basis c' with the SMEFT linearised model, where all other coefficients and nuisance parameters are profiled. The middle panel shows the symmetrised 68% CL uncertainty σ of each parameter measurement (left vertical axis) and the corresponding energy scale $\Lambda/\sqrt{\sigma}$ that is probed (right vertical axis). The bottom panel shows the measured parameter value (dot) and 68% (solid line) and 95% (dashed line) CL intervals, divided by the symmetrised uncertainty shown in the middle panel. The p -value for the compatibility of the data with the Standard Model expectation (all coefficients vanishing) is 94.5%. The top panel shows the expected breakdown of contributions to the sensitivity of each measurement from the various measured Higgs boson decay and production modes. Green shades represent production and decay modes with Higgs boson couplings to vector bosons, red shades represent modes with Higgs boson couplings to fermions, and blue shades represent modes with loop couplings. The production category labelled ‘inc’ collects final states for which the breakdown into production modes is not available ($H \rightarrow \mu\mu$ and $H \rightarrow Z\gamma$). The 95% CL interval for $e_{\text{tH}}^{[3]}$ is truncated at the boundary at which the logarithm of the likelihood function becomes undefined, resulting in a small undercoverage.

as the square root of the difference of the squares of the total uncertainty and the statistical uncertainty. The contribution from a group of systematic uncertainties is estimated as the difference of the squares of the total uncertainty and the uncertainty from a fit in which the associated nuisance parameters are fixed to their best-fit values.




Figure 11: Expected fractional contributions of the statistical (orange) and systematic (blue) uncertainties to the total uncertainty in the measurements of the parameters of the rotated basis \mathbf{c}' with the SMEFT linearised model (top panel), and the corresponding expected fractional contributions of experimental (cyan), signal theory (light blue) and background theory (dark blue) uncertainties to the total systematic uncertainty (bottom panel).

For parameters $c_{He,33}$, $e_{\text{glob}}^{[1]}$ and $e_{Hlll}^{[1]}$, the fractional contribution of systematic uncertainties is close to 50%; for parameters $e_{\text{ggF}}^{[1]}$, $e_{\text{ggF}}^{[2]}$, $e_{\text{ttH}}^{[1]}$ and $e_{\text{ttH}}^{[2]}$, the fractional contribution of systematic uncertainties is around 40%; the remaining parameters have smaller relative systematic uncertainties. In particular, the uncertainties in the $c_{He,22}$ and $e_{\text{ttH}}^{[3]}$ parameters, probed by the measurements of the rare processes $H \rightarrow \mu\mu$ decay and tH production, respectively, are almost completely dominated by the statistical component. The total systematic uncertainty in most parameters is dominated by experimental systematic uncertainties, except $e_{\text{ggF}}^{[1]}$ and $e_{\text{glob}}^{[1]}$, where signal theory systematic uncertainties dominate, and $e_{\text{ggF}}^{[2]}$, $e_{\text{ttH}}^{[1]}$ and $e_{\text{ttH}}^{[2]}$, where background theory systematic uncertainties dominate. The breakdown of the uncertainty components, including a breakdown of the systematic uncertainty into experimental, signal theory and background theory components, is also given in Tables 3 and 4.

The expected and observed correlation matrices are shown in Figure 12, and show a reasonably good decorrelation achieved between the fit basis parameters: over half of the off-diagonal elements are smaller than 0.1 and over 85% are smaller than 0.3 in both the observed and expected matrix. A few exceptions stand out, notably the correlation between $e_{\text{ggF}}^{[2]}$ and $e_{\text{ttH}}^{[1]}$ and between $e_{\text{ggF}}^{[3]}$ and $e_{H\gamma\gamma, Z\gamma}^{[1]}$, which are caused by a common sensitivity to ttH production and ggF $H \rightarrow \gamma\gamma$, respectively.

Model including quadratic terms Figure 13 compares the expected and observed results obtained using the SMEFT model including quadratic terms, again showing good agreement of the observed data with the SM expectation, corresponding to a p -value of 98.2%. For most parameters, the observed uncertainty is noticeably smaller than the expected uncertainty. The cause of this discrepancy is related to appearance of

Table 3: Summary of the observed measurements of the parameters of the rotated basis \mathbf{c}' in the SMEFT linearised model, with a breakdown of the total uncertainty into statistical and systematic components. A further breakdown of the systematic uncertainty into experimental and theoretical uncertainties in the signal and background is also listed. All MC statistical uncertainties are classified as experimental systematic uncertainties in the breakdown. All uncertainties correspond to 68% CL intervals. The sum in quadrature of the systematic uncertainty components may differ from the total systematic uncertainty due to correlations.

Parameter	Value	σ_{tot}	σ_{stat}	σ_{syst}	$\sigma_{\text{syst}}^{\text{exp}}$	$\sigma_{\text{syst}}^{\text{th.sig}}$	$\sigma_{\text{syst}}^{\text{th.bkg}}$
$c_{eH,22}$	-0.0002	0.0028	0.0026	0.0009	0.0006	0.00049	+0.0003 -0.0005
$c_{eH,33}$	0.019	0.016	0.011	0.011	0.008	0.006	0.005
$c_{Hq}^{(3)}$	0.039	0.040	0.034	0.021	0.015	+0.011 -0.005	0.012
c_{bH}	0.055	+0.039 -0.058	+0.032 -0.045	+0.023 -0.037	+0.014 -0.022	0.015	+0.013 -0.025
$e_{\text{ggF}}^{[1]}$	0.0003	0.0039	0.0029	0.0026	0.0013	0.0021	+0.0013 -0.0008
$e_{\text{ggF}}^{[2]}$	-0.07	0.38	0.29	0.25	0.13	0.11	0.18
$e_{\text{ggF}}^{[3]}$	5.5	7.1	5.8	4.00	2.71	+2.8 -2.1	1.6
$e_{H\gamma\gamma,Z\gamma}^{[1]}$	0.0048	0.0043	0.0034	0.0026	0.0017	0.0015	0.0012
$e_{H\gamma\gamma,Z\gamma}^{[2]}$	-0.042	0.041	0.037	+0.015 -0.022	0.014	+0.006 -0.011	+0.006 -0.010
$e_{H\gamma\gamma,Z\gamma}^{[3]}$	-0.47	0.62	0.51	0.35	0.24	0.20	0.16
$e_{ZH}^{[1]}$	-0.14	0.34	0.29	0.19	0.16	0.09	+0.08 -0.11
$e_{ZH}^{[2]}$	3.7	3.9	3.3	2.1	1.6	0.9	1.0
$e_{ZH}^{[3]}$	1.8	8.5	7.2	4.5	3.4	2.2	2.0
$e_{ZH}^{[4]}$	4.8	9.8	8.2	5.2	4.0	2.5	2.2
$e_{\text{ttH}}^{[1]}$	-0.65	0.84	0.64	0.54	0.31	0.17	0.41
$e_{\text{ttH}}^{[2]}$	-0.05	3.7	2.8	2.4	0.94	+0.3 -0.6	2.2
$e_{\text{ttH}}^{[3]}$	8.6	+13.4 -9.2	+12.1 -8.8	+5.73 -2.91	+3.9 -2.1	+3.8 -1.5	+2.0 -1.2
$e_{\text{glob}}^{[1]}$	0.13	+0.83 -0.57	+0.57 -0.42	+0.60 -0.38	+0.30 -0.20	+0.42 -0.29	+0.29 -0.17
$e_{Hllll}^{[1]}$	-0.349	1.1	0.76	0.74	0.44	0.38	0.46

Table 4: Summary of the expected measurements of the parameters of the rotated basis c' with the SMEFT linearised model, with a breakdown of the total uncertainty into a statistical and systematic component. A further breakdown of the systematic uncertainty into experimental and theoretical uncertainties in the signal and background is also listed. All MC statistical uncertainties are classified as experimental systematic uncertainties in the breakdown. All uncertainties correspond to 68% CL intervals. The sum in quadrature of the systematic uncertainty components may differ from the total systematic uncertainty due to correlations.

Parameter	Value	σ_{tot}	σ_{stat}	σ_{syst}	$\sigma_{\text{syst}}^{\text{exp}}$	$\sigma_{\text{syst}}^{\text{th.sig}}$	$\sigma_{\text{syst}}^{\text{th.bkg}}$
$c_{eH,22}$	0.0	0.0031	0.0030	0.0010	0.0007	+0.0004 -0.0005	+0.0004 -0.0006
$c_{eH,33}$	0.0	0.019	0.013	0.014	0.010	0.007	0.007
$c_{Hq}^{(3)}$	0.0	0.044	0.037	0.023	0.017	0.007	0.015
c_{bH}	0.0	+0.05 -0.09	+0.04 -0.07	+0.03 -0.06	+0.02 -0.03	+0.01 -0.02	+0.02 -0.04
$e_{\text{ggF}}^{[1]}$	0.0	0.0046	0.0035	0.0029	0.0015	0.0022	+0.0015 -0.0011
$e_{\text{ggF}}^{[2]}$	0.0	0.46	0.36	0.29	0.15	0.13	0.20
$e_{\text{ggF}}^{[3]}$	0.0	8.4	7.1	4.4	3.1	2.5	1.9
$e_{H\gamma\gamma,Z\gamma}^{[1]}$	0.0	0.0057	0.0043	0.0030	0.0021	0.0016	+0.0012 -0.0018
$e_{H\gamma\gamma,Z\gamma}^{[2]}$	0.0	0.044	0.040	0.019	0.016	+0.005 -0.007	0.009
$e_{H\gamma\gamma,Z\gamma}^{[3]}$	0.0	0.68	0.56	0.37	0.25	+0.25 -0.15	0.17
$e_{ZH}^{[1]}$	0.0	0.40	0.34	0.21	0.15	0.10	0.10
$e_{ZH}^{[2]}$	0.0	4.4	3.7	2.3	1.8	0.96	1.10
$e_{ZH}^{[3]}$	0.0	9.8	8.4	5.0	3.7	2.5	2.3
$e_{ZH}^{[4]}$	0.0	11	9.9	5.7	4.5	2.6	2.5
$e_{\text{ttH}}^{[1]}$	0.0	1.0	0.79	0.68	0.40	0.22	0.49
$e_{\text{ttH}}^{[2]}$	0.0	4.7	3.5	3.1	1.2	0.61	2.8
$e_{\text{ttH}}^{[3]}$	0.0	+10.4 -7.6	+10.0 -7.0	3.02	2.25	+1.79 -1.30	+0.97 -1.51
$e_{\text{glob}}^{[1]}$	0.0	+0.93 -0.60	+0.64 -0.46	+0.67 -0.39	0.28	+0.49 -0.23	+0.35 -0.18
$e_{Hllll}^{[1]}$	0.0	1.23	0.89	0.85	0.54	0.39	0.53





Figure 12: The (a) expected and (b) observed correlation matrix for the parameters of the rotated basis c' obtained from a fit to the linearised SMEFT model.

quadratic parameter terms in the cross-section predictions, which cause the likelihood function to generally have multiple minima in each parameter. In the expected result, most nearby minima that are exactly degenerate with the global minimum merge into a single wider uncertainty interval. Conversely, in the observed result, the generally non-zero observed coefficient values lift some of these minima to become local minima, thereby resulting in a narrower interval around the remaining global minimum. This effect can be clearly seen in more detail in the profile likelihood scans of each parameter, shown in Figures 14 and 15. Most secondary minima are raised by 1–2 units in $-2 \log L$, thus narrowing the 68% confidence intervals, which are based on a unit threshold, but less so the 95% confidence intervals based on a 3.84 unit threshold. For all parameters with multiple solutions, the global minimum in the observed result resolves to the minimum closest to a coefficient value of zero, except parameter $e_{H\gamma\gamma,Z\gamma}^{[1]}$, where the minimum furthest from zero becomes the global minimum. For the parameter $c_{eH,22}$, there are two exactly degenerate minima. In that case, the minimum closest to that of the interpretation with the linear model is quoted as the best-fit result.




Figure 13: Comparison of the expected (gray) and observed (blue) parameters of the rotated basis c' with the SMEFT model with quadratic terms, where all other coefficients and nuisance parameters are profiled. The top panel shows the symmetrised 68% uncertainty σ of each parameter measurement (left vertical axis) and the corresponding energy scale $\Lambda/\sqrt{\sigma}$ that is probed (right vertical axis). The bottom panel shows the measured parameter value and 68% (solid) and 95% (dashed) CL intervals, divided by the symmetrised uncertainty shown in the top panel. Some of the observed intervals shown are clipped off. Due to multiple minima, the likelihood curves from which these CL intervals are derived are non parabolic. For the parameter $c_{eH,22}$, for which there are two degenerate minima in the interpretation with the model including quadratic terms, the point corresponds to the best-fit value that is closest to that of the interpretation with the linear model.

Almost all profile likelihood functions show Gaussian behaviour to a good approximation within the 95% CL range of each parameter, although this is masked by the appearance of degenerate solutions. The

only exception is the expected profile likelihood for the $e_{ttH}^{[3]}$ parameter in the linearised model, where a boundary occurs at $e_{ttH}^{[3]} = -10$, just before the likelihood threshold of the 95% CL is reached. The yield prediction in the high- p_T region of the ttH , $H \rightarrow \gamma\gamma$ channel becomes negative for very negative values of $e_{ttH}^{[3]}$, rendering the likelihood undefined beyond that point. The 95% CL interval for $e_{ttH}^{[3]}$ shown in Figures 10 and 16 is truncated at this boundary, resulting in a small undercoverage.

3.2.3 Comparison between the linearised model and the model including quadratic terms

Figure 16 compares the results for the linearised model and the model with quadratic terms for the observed data and shows that for several of the coefficients the constraints from the model with the quadratic terms are significantly stronger than those observed for the linear model. In most cases, the stronger constraints arise from the relatively weak impact of the BSM–SM interference term on the cross-section compared to the quadratic BSM terms in specific production or decay modes.

Figure 9 illustrates the impact of both the linear model and the model with quadratic terms, where the impact strength visualised for *both* models corresponds to the expected 68% uncertainty of each fit basis parameter c' under the linear model. The comparatively larger impact of the model including quadratic terms at these chosen parameter values identifies STXS regions where quadratic terms outweigh the linear terms at the measured scale and thus identifies regions that predominantly provide the extra constraining power of the models including quadratic terms. The strongest observed sensitivity enhancements occur in $e_{ggF}^{[3]}$, $e_{ttH}^{[2]}$ and $e_{ttH}^{[3]}$. For coefficient $e_{ggF}^{[3]}$, the quadratic term significantly increases sensitivity in the ggF, ZH and ttH production modes as well as in the $H \rightarrow \gamma\gamma$ decay mode. For coefficients $e_{ttH}^{[2]}$ and $e_{ttH}^{[3]}$, the quadratic term enhances sensitivity in high- p_T ttH production and tH production, respectively. Conversely, for the coefficients $e_{ggF}^{[2]}$ and $e_{H\gamma\gamma,Z\gamma}^{[3]}$, the effect of the increased impact of the quadratic terms is limited, and the strong reduction in uncertainties is mostly driven by the lifting of degenerated solutions in the observed case compared with expected, as visible in the profile likelihood scans in Figure 14.

Operators in Warsaw basis: $c_{eH,22}$, $c_{eH,33}$, $c_{Hq}^{(3)}$ and c_{bH}






Figure 14: Profile likelihood scans of the fitted coefficients of the rotated basis c' obtained from fits to the SM expectation (dashed lines) and to the observed data (solid lines), based on the SMEFT linearised model (blue) and the SMEFT model with quadratic terms (orange). The horizontal dashed lines in each plot correspond to the asymptotic threshold values for 68% and 95% CL intervals. The profile likelihood scans of the SMEFT model with quadratic terms can exhibit two minima in the scanned parameter, whereas the scans of the linearised model have one minimum per parameter. Multiple minima in the coefficients that are profiled in each scan may furthermore result in discontinuous derivatives in the profile likelihood (e.g., prominently visible in the observed data in (e) and (l)).

Eigenvector group top



Eigenvector group $overall\ normalisation$



Eigenvector group $H \rightarrow ZZ^*$




Figure 15: Profile likelihood scans of the fitted coefficients of the rotated basis c' obtained from fits to the SM expected (dashed lines) and to the observed data (solid lines), based on the SMEFT linearised model (blue) and the SMEFT model with quadratic terms (orange). The horizontal dashed lines in each plot correspond to the asymptotic threshold values for 68% and 95% CL intervals. The profile likelihood scans of the SMEFT model with quadratic terms can exhibit two minima in the scanned parameter, whereas the scans of the linearised model can only have one minimum per parameter. Multiple minima in the coefficients that are profiled in each scan may furthermore result in discontinuous derivatives in the profile likelihood (e.g., prominently visible in the observed data in (d)).




Figure 16: Comparison of the observed parameters of the rotated basis c' with the SMEFT linearised model (blue) and the model including quadratic terms (orange), where all other coefficients and nuisance parameters are profiled. The top panel shows the symmetrised 68% CL uncertainty σ of each parameter measurement (left vertical axis) and the corresponding energy scale $\Lambda/\sqrt{\sigma}$ that is probed (right vertical axis). The bottom panel shows the measured parameter value and 68% (solid) and 95% (dashed) CL intervals, divided by the symmetrised uncertainty shown in the top panel. Some of the intervals shown for the model with quadratic terms are clipped off. Due to multiple minima the likelihood curves from which these CL intervals are derived are non-parabolic. The 95% CL interval for $e_{\text{tH}}^{[3]}$ is truncated at the boundary at which the log likelihood becomes undefined, resulting in a small undercoverage. For the parameter $c_{eH,22}$, for which there are two degenerate minima in the interpretation with the model including quadratic terms, the point corresponds to the best-fit value that is closest to that of the interpretation with the linear model.

3.3 Constraints from fiducial differential measurements

Coefficients of SMEFT operators can alternatively be constrained from fiducial cross-section measurements. Fiducial differential cross-section measurements are, unlike STXS measurements, explicitly unfolded to the particle level. The finer granularity of kinematic regions defined in differential measurements, compared to STXS measurements, offers an advantage in the analysing power of differential distributions, but is counteracted by the inclusive treatment of Higgs boson production processes, which are modelled separately in the STXS approach.

For a comparison to the STXS approach, differential measurements in the p_T of the Higgs boson are considered. Anomalous couplings of the Higgs boson to gluons and top quarks, as well as between gluons and top quarks, can affect the total Higgs boson production cross-section and its dependence on the Higgs boson transverse momentum.

The expected deviations from the SM predictions due to these anomalous couplings can be relatively large in high Higgs boson p_T regions, which are also characterised by a better signal-to-background ratio, making the p_T^H -differential cross-section measurement more sensitive to these effects compared to a measurement of the inclusive rate.

3.3.1 Sensitivity estimate and choice of parameters

In this study, constraints on anomalous Higgs boson couplings to gluons and top quarks are set from the observed transverse momentum differential spectra of the Higgs boson decay products in the $H \rightarrow ZZ^* \rightarrow 4\ell$ and $H \rightarrow \gamma\gamma$ decay channels. The constraints are inferred using an EFT approach in which the SM Lagrangian is augmented with the three dimension-6 SMEFT operators O_{HG} (Higgs-gluon point-like contact term), O_{tH} (top quark Yukawa coupling modifier) and O_{tG} (chromomagnetic dipole operator), that are defined in Table 2. The operator O_{HG} introduces an Hgg contact interaction that modifies the value and p_T -dependence of the ggF production cross-section, as well as the $H \rightarrow gg$ partial decay width, thus affecting indirectly the $H \rightarrow ZZ^* \rightarrow 4\ell$ and $H \rightarrow \gamma\gamma$ branching ratios. The operator O_{tH} modifies the $t\bar{t}H$ vertex and thus affects Higgs boson production through top-quark-loop mediated gluon-gluon fusion or in association with a $t\bar{t}$ pair, as well as the top-quark-loop amplitude contributing to the $H \rightarrow \gamma\gamma$ partial width. The operator O_{tG} introduces a $t\bar{t}Hg$ vertex that leads to additional contributions to the amplitude for ggF or $t\bar{t}H$ Higgs boson production, as well as for $H \rightarrow gg$ decays, affecting in turn the branching ratios for both $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$.

The study presented in this section is performed in the scenario in which only the SM–BSM interference effects, linear in the Wilson coefficients, are considered. For simplicity, due to the statistical precision of the data sample and the very large correlations between the estimators for the three Wilson coefficients, the constraints are initially set on one Wilson coefficient at a time, while the values of the remaining coefficients are assumed to be equal to zero. Subsequently, similarly to the approach presented in Section 3.2.1, a rotation in the parameter space is performed to define a new set of coefficients that are decorrelated and can be probed simultaneously. The new basis is formed by the three eigenvectors $ev^{[i]}$ ($i = 1..3$) of the Fisher information matrix, which are related to the three Wilson coefficients by the following rotation:

$$\begin{aligned} ev^{[1]} &= 0.999c_{HG} - 0.035c_{tG} - 0.003c_{tH}, \\ ev^{[2]} &= 0.035c_{HG} + 0.978c_{tG} + 0.205c_{tH}, \\ ev^{[3]} &= -0.005c_{HG} - 0.205c_{tG} + 0.979c_{tH}. \end{aligned}$$

The impact of the operators defining the new basis on the fiducial differential cross-sections is shown in Figure 17.




Figure 17: Impact of the rotated SMEFT operators on the fiducial cross-section differential in p_T^H in $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^*$ decays, relative to the SM cross-section, under the assumption of the linearised SMEFT model. To judge the experimental sensitivity to constrain the operators from the data in the listed fiducial regions, the total uncertainty in the corresponding regions is shown in the top panel. For presentational clarity, the uncertainty of low precision fiducial regions is clipped off in the plot.

3.3.2 Results

The statistical interpretation is performed using a joint likelihood model of the data in the two decay channels, built, as described in Section 2.3, from the observed and expected values of the fiducial differential cross-sections in each bin of the transverse momentum of the Higgs boson decay products for both $H \rightarrow 4\ell$ and $H \rightarrow \gamma\gamma$ decays. In the likelihood function, the theoretical prediction for each decay channel and transverse momentum interval is parameterised as a function of the Wilson coefficients, as summarised in Section 3.1. The procedure assumes common higher-order corrections for the SMEFT and SM processes. The theoretical uncertainties are taken from the SM calculations in Section 2.2, with the uncertainty associated with the SMEFT scaling assumed to be negligible.

Figure 18 shows the 68% CL one-dimensional intervals in the linearised model for the rotated parameters $ev^{[i]}$ with the other two parameters profiled. The constraint on the first rotated parameter, which is almost aligned with c_{HG} , is of the order of 1%, while the constraints on the other two rotated parameters are $O(1)$. Figure 18 compares these results to those obtained from the STXS measurements performed with the same decay channels using a linearised model in which all Wilson coefficients except for c_{tH} , c_{tG} and c_{HG} are set to zero. The comparison illustrates that the differential cross-section measurements, which have a finer granularity compared to the STXS measurements but probe the distribution of a single observable inclusively in production mode, have less constraining power than the STXS measurements, which use information on the particles produced with the Higgs boson decay products to separate the different production modes whose cross-sections are affected in different ways by the different operators, as

shown in Figure 5. In particular, the large sensitivity difference observed for $ev^{[2]}$ originates from the separate measurements of ggF and $t\bar{t}H$ production in the STXS framework.




Figure 18: The (a) observed and (b) expected 68% CL intervals on the three rotated parameters $ev^{[i]}$ obtained with the SMEFT linearised model using either STXS (blue) or fiducial p_T -differential cross-section measurements (green) in the $H \rightarrow \gamma\gamma$ and $H \rightarrow 4\ell$ decay channels. The parameter $ev^{[1]}$ is almost aligned with c_{HG} , which mainly affects ggF production, while the parameters $ev^{[2]}$ and $ev^{[3]}$ are close to c_{tG} and c_{tH} respectively, which impact both ggF and $t\bar{t}H$ production.

4 Interpretations based on UV-complete BSM models

In this section, the measurements of Higgs boson cross-sections and decay rates are interpreted in two UV-complete extensions of the SM: a two-Higgs-doublet model near the alignment limit (Section 4.1), and eight benchmark scenarios of the minimal supersymmetric extension of the SM (Section 4.2).

4.1 Constraints on two-Higgs-doublet models

In two-Higgs-doublet models, the SM Higgs sector with one doublet of scalar complex fields Φ_1 is extended by introducing a second doublet Φ_2 [11–13]. The models considered here assume CP conservation and a (softly broken) Z_2 discrete symmetry that forbids quartic terms of the scalar field potential $V(\Phi_1, \Phi_2)$ that contain odd powers of either Φ_1 or Φ_2 . The vacuum expectation values $v_{1,2}$ of the scalar doublets $\Phi_{1,2}$ that minimise V are related by the sum rule $v_1^2 + v_2^2 = v^2$. These can be assumed to be both real and non-negative without loss of generality. Electroweak symmetry breaking leads to five physical scalar Higgs fields: two neutral CP-even Higgs bosons h and H , one neutral CP-odd Higgs boson A , and two charged Higgs bosons H^\pm . For the results presented in this section, the observed Higgs boson is identified with the lightest CP-even neutral scalar particle h .

The Z_2 symmetry of the potential forbids tree-level flavour-changing neutral currents [107, 108], which are strongly constrained by existing data, and implies that all fermions with the same quantum numbers couple to only one Higgs doublet. Depending on which Higgs doublet couples to the three groups of elementary fermions (up-type and down-type quarks, and leptons), four types of 2HDM are defined:

- Type I: All fermions couple to the same Higgs doublet.
- Type II: One Higgs doublet couples to up-type quarks while the other one couples to down-type quarks and charged leptons.
- Lepton-specific: One Higgs doublet couples to leptons while the other one couples to up- and down-type quarks.
- Flipped: One Higgs doublet couples to down-type quarks while the other one couples to up-type quarks and leptons.

The Higgs sector of these models contains seven free physical parameters: four Higgs boson masses (m_h , m_H , m_A and m_{H^\pm}), two mixing angles α and β , and the coefficient m_{12}^2 of the $(\Phi_1^\dagger \Phi_2 + \Phi_2^\dagger \Phi_1)$ term in the Higgs field potential that softly breaks the Z_2 symmetry. The angle β is defined as $\tan \beta = \frac{v_2}{v_1}$ and can be assumed to be in the first quadrant. The mixing angle α of the neutral CP-even Higgs sector is defined, modulo a phase, equal to π . It can thus be assumed that $0 \leq \beta - \alpha \leq \pi$.

The values of α and β determine the couplings of the Higgs fields to the vector bosons and to the fermions. In this section, the *decoupling limit* is assumed, in which $m_H \gg v$. This in turn implies the *alignment limit*, $|\cos(\beta - \alpha)| \ll 1$, in which the light scalar Higgs boson h has SM-like couplings. The mass of the light Higgs boson m_h is assumed to be close to 125 GeV, while all other Higgs bosons are assumed to be significantly heavier (and their masses are assumed to be degenerate, $m_A \sim m_{H^\pm} \sim m_H \sim M$). Near the alignment limit, i.e., for $|\cos(\beta - \alpha)|$ small, the modifications of the tree-level couplings of the h boson with respect to the SM predictions follow the expressions summarised in Table 5 [13].

Table 5: Multiplicative factors predicted in the four 2HDM scenarios near the alignment limit, as a function of $\tan\beta$ and $\cos(\beta - \alpha)$, for the Higgs boson couplings to up-type quarks (1st row), down-type quarks (2nd row), charged leptons (3rd row), vector bosons (4th row), and to itself (5th row). The symbol $c_{\beta-\alpha}$ stands for $\cos(\beta - \alpha)$, while $s_{\beta-\alpha}$ stands for $\sin(\beta - \alpha)$. The definition of the parameter \bar{m} is given in the text.

Coupling	Type I	Type II	Lepton-specific	Flipped
u, c, t			$s_{\beta-\alpha} + c_{\beta-\alpha}/\tan\beta$	
d, s, b	$s_{\beta-\alpha} + c_{\beta-\alpha}/\tan\beta$	$s_{\beta-\alpha} - c_{\beta-\alpha} \times \tan\beta$	$s_{\beta-\alpha} + c_{\beta-\alpha}/\tan\beta$	$s_{\beta-\alpha} - c_{\beta-\alpha} \times \tan\beta$
e, μ, τ	$s_{\beta-\alpha} + c_{\beta-\alpha}/\tan\beta$	$s_{\beta-\alpha} - c_{\beta-\alpha} \times \tan\beta$	$s_{\beta-\alpha} - c_{\beta-\alpha} \times \tan\beta$	$s_{\beta-\alpha} + c_{\beta-\alpha}/\tan\beta$
W, Z			$s_{\beta-\alpha}$	
h		$s_{\beta-\alpha}^3 + \left(3 - 2\frac{\bar{m}^2}{m_h^2}\right) c_{\beta-\alpha}^2 s_{\beta-\alpha} + 2 \cot(2\beta) \left(1 - \frac{\bar{m}^2}{m_h^2}\right) c_{\beta-\alpha}^3$		

In addition to the impact of the tree-level coupling modifications, the production and decay rates of the h boson are modified through NLO EW corrections involving the trilinear hhh coupling λ [109]. In the 2HDM framework, this self-coupling is modified with respect to the SM expectation λ_{SM} by the scale factor (denoted by κ_λ) shown in the last row of Table 5 [110]. The parameter \bar{m} is $\bar{m}^2 = \frac{m_{12}^2}{\sin\beta\cos\beta} = m_A^2 + \lambda_5 v^2$, where λ_5 is the coefficient of the $(\Phi_1^\dagger \Phi_2)^2$ term of the Higgs potential. Near the alignment limit considered, the value of \bar{m} is close to that of m_A ($\lambda_5 v^2 \ll m_A^2$), and a value $\bar{m} = m_A = 1$ TeV ($\lambda_5 = 0$) is assumed in this section in all scenarios in which the effect of the self-coupling is considered in the calculation of limits on $\cos(\beta - \alpha)$ and $\tan\beta$.

In the following sections, limits on the 2HDM parameters $\cos(\beta - \alpha)$ and $\tan\beta$ are inferred from studies of the ensemble of Higgs boson production and decay rate measurements in two distinct scenarios: first by comparing the measured rates for each production mode with inclusive 2HDM predictions expressed in the ‘ κ -framework’, and then by comparing the more fine-grained measurements described in Section 2 with the 2HDM predictions expressed in the linearised statistical model of the EFT-based parameterisation described in Section 3.

4.1.1 Constraints based on the κ -framework

In the κ -framework [14], the Higgs boson production-mode cross-sections and decay branching ratios are parameterised in terms of multiplicative coupling strength modifiers κ [9]. In the model considered in this study, the loop-induced processes (gluon–gluon fusion $gg \rightarrow H$ and $gg \rightarrow ZH$ production, as well as $H \rightarrow \gamma\gamma$, $H \rightarrow Z\gamma$ and $H \rightarrow gg$ decays) are expressed in terms of the strength factors for the couplings to the SM particles inside the loop. The model also assumes that there are no invisible or undetected Higgs boson decays beyond the SM. The impact of the trilinear hhh coupling modifier κ_λ on the Higgs boson production and decay rates via NLO EW corrections to the LO amplitudes is also included in the parameterisations. Constraints on κ_λ from searches for Higgs boson pair production [111] are not taken into account. In the likelihood function in Eq. (1), the signal strengths $\mu_k^{i,X}$ are re-parametrised as $\mu_k^{i,X} = \mu^{i,X}(\{\kappa(\tan\beta, \cos(\beta - \alpha))\})$ using the relations listed in Table 5. Confidence regions for $\tan\beta$ and $\cos(\beta - \alpha)$ can thus be inferred from the combined measurements of Higgs boson production and decay rates.

The resulting 95% CL contours in the $(\tan\beta, \cos(\beta - \alpha))$ plane for $0.1 < \tan\beta < 10$ are shown in Figure 19. All models exhibit similar exclusion regions in the $(\tan\beta, \cos(\beta - \alpha))$ plane at low values ($\lesssim 1$) of $\tan\beta$,

where only a narrow region of $\cos(\beta - \alpha)$ around zero is consistent with the measured values of the Higgs boson production and decay rates. The interval of allowed values of $\cos(\beta - \alpha)$ increases in size with increasing $\tan\beta$, up to a total width of about 0.1–0.2 for $\tan\beta = 1$, depending on the model. For higher values of $\tan\beta$, in the models in which at least one of the coupling strength modifiers is enhanced by a factor $\tan\beta$, i.e., all models except Type-I, the allowed range of $\cos(\beta - \alpha)$ around zero shrinks as $\tan\beta$ gets larger than one. This trend is observed also for the region $\tan\beta > 10$, not shown in Figure 19. For the Type-I model, the allowed range of $\cos(\beta - \alpha)$ further increases with $\tan\beta$, reaching a width of around 0.5 for $\tan\beta = 10$. For $\tan\beta > 10$, the allowed range of $\cos(\beta - \alpha)$ remains approximately constant. This includes regions of the parameter space that deviate significantly from the alignment-limit hypothesis. When the constraint from the trilinear hhh coupling is included, the width of the 95% CL interval for $\cos(\beta - \alpha)$ at large $\tan\beta$ for the Type-I model is reduced by about 50%. An additional feature of the Type-I model result is that in the large $\tan\beta$ region, for positive $\cos(\beta - \alpha)$, the observed exclusion region is significantly larger than the expected one. This derives from the fact that the ATLAS measurement of Higgs boson production and decay rates favours values of the coupling strength modifiers to b -quarks, t -quarks and τ -leptons smaller than one and of the couplings to W and Z bosons larger than one ($\kappa_F = 0.95 \pm 0.05$ and $\kappa_V = 1.035 \pm 0.031$ when assuming that all fermions share the same modifier κ_F and that $\kappa_W = \kappa_Z = \kappa_V$ [9]). In this scenario, the model predicts $\sin(\beta - \alpha) = \kappa_V \approx 1$ and $\cos(\beta - \alpha) = \tan\beta(\kappa_F - \sin(\beta - \alpha)) \approx \tan\beta(\kappa_F - 1)$, disfavouring positive values of $\cos(\beta - \alpha)$ for $\kappa_F < 1$.

A second small region of allowed $\tan\beta$ and $\cos(\beta - \alpha)$ values for large $\tan\beta$ and $\cos(\beta - \alpha) \approx 0.2$ is present in the Type-II, Lepton-specific, and Flipped models. This corresponds to regions with $\cos(\beta + \alpha) \approx 0$, for which some of the fermion couplings have the same magnitude as in the SM but the opposite sign. In particular, this corresponds to a negative sign of the lepton couplings in the Lepton-specific model, of the down-type quark couplings in the Flipped model, or both in the Type-II model. Since the sign of these couplings is not sufficiently well constrained by current experimental measurements, this region is not excluded. The same region is however not allowed in the Type-I model, in which the coupling strength modifiers of down-type quarks and leptons have the same sign (and value) as for up-type quarks. In that case, a negative sign of the top-quark coupling strength modifier κ_t is experimentally excluded by the measurement of $H \rightarrow \gamma\gamma$ decays, whose branching ratio depends linearly on $\kappa_t\kappa_W$ because of the interference between W boson- and top-quark-mediated loop amplitudes.

4.1.2 Constraints using an EFT-based approach

The modifications to the tree-level Higgs boson couplings to fermions near the alignment limit described in Section 4.1 can be generated by the dimension-6 operators \mathcal{O}_{uH} , \mathcal{O}_{bH} , \mathcal{O}_{tH} and \mathcal{O}_{eH} of the SMEFT Lagrangian in the ‘top’ flavour scheme, with the following matching between the Wilson coefficients and the 2HDM parameters [112]:

$$\frac{v^2 c_{iH}}{\Lambda^2} = -Y_i \eta_i \frac{\cos(\beta - \alpha)}{\tan\beta}, \quad (16)$$

with $i = u, b, t, \tau, \mu$, and $c_{\mu H}$ and $c_{\tau H}$ referring to the $c_{eH,22}$ and $c_{eH,33}$ coefficients, respectively. In these formulae, computed to first order in $\cos(\beta - \alpha)$, $Y_i = \sqrt{2}m_i/v$ are the SM Higgs boson couplings and the values of η_i depend on the type of model ($\eta_b = -\tan^2\beta$ in the Type-II and Flipped models, as are η_μ and η_τ in the Type-II and the Lepton-specific models; $\eta_{b,t,\mu,\tau} = 1$ otherwise). The corrections to the HVV




Figure 19: Regions of the 2HDM ($\tan\beta$, $\cos(\beta - \alpha)$) parameter plane excluded at 95% CL (light yellow filled regions) in the κ -framework-based approach by the measured rates of Higgs boson production and decays in the (a) Type-I, (b) Type-II, (c) Lepton-specific and (d) Flipped models. The dark yellow dashed lines show the borders of the corresponding expected exclusion regions for the SM hypothesis. For the Type-I model, the observed and expected regions excluded at 95% CL when the κ_λ constraint is considered are also shown (solid and dashed blue lines). The red dashed lines correspond to the exact SM alignment limit ($\cos(\beta - \alpha) = 0$). Results derived in the κ -framework assume the decoupling limit, i.e., that the masses of the non-SM-like Higgs bosons are large compared to the SM vev, which implies the assumption of the alignment limit, i.e., $|\cos(\beta - \alpha)| \ll 1$.

vertices are quadratic in $\cos(\beta - \alpha)$ and are neglected. For the EFT to be valid, $\Lambda \gg v$ and therefore $|\cos(\beta - \alpha)| \propto \frac{v^2}{\Lambda^2}$ should be small, close to the alignment limit.

The operator O_H is also generated in the models under study, with a coefficient that scales with $\cos^2(\beta - \alpha)$ but which can be significantly enhanced if the other scalar states are much heavier than the SM-like Higgs boson. In that case, c_H is proportional to $(M/v)^2$ [112]:

$$\frac{v^2 c_H}{\Lambda^2} = \cos(\beta - \alpha)^2 \left(\frac{M}{v} \right)^2.$$

Non-zero values of c_H modify the trilinear hhh coupling λ by a scale factor [113]:

$$\kappa_\lambda = 1 - \frac{v^2 c_H}{\lambda_{\text{SM}} \Lambda^2} = 1 - \frac{\cos(\beta - \alpha)^2}{\lambda_{\text{SM}}} \left(\frac{M}{v} \right)^2.$$

This in turn affects Higgs boson production and decay rates via NLO EW corrections to the LO amplitudes, whose effect is included in the parameterisations used to obtain the results. As in Section 4.1.1, constraints on κ_λ from searches for Higgs boson pair production are not taken into account. In the following, a value of $M = 1$ TeV is assumed for the masses of the heavy scalar particles, and thus $\kappa_\lambda \approx 1 - (11 \cos(\beta - \alpha))^2$. Given the large predicted deviations of κ_λ from one for $\cos(\beta - \alpha)$ different from zero, including κ_λ in the interpretation is expected to affect significantly, for $\tan \beta \gtrsim 1$, the results for the Type-I model, for which the impact of $\cos(\beta - \alpha) \neq 0$ on the other Wilson coefficients, c_{bH} , c_{tH} and $c_{\tau H}$, is suppressed by a $1/\tan \beta$ factor. Exclusion limits for all other types of 2HDM models are expected to be largely unaffected.

Constraints on the values of $\tan \beta$ and $\cos(\beta - \alpha)$ can thus be inferred from those set on c_{bH} , c_{tH} , $c_{\tau H}$ and c_H using the SMEFT interpretation of the combined measurement of production and decay rates in STXS bins. The likelihood model defined in Section 3.2 is used, expressing the four Wilson coefficients c_{bH} , c_{tH} , $c_{\tau H}$ and c_H in terms of the 2HDM parameters as in Eq. (16) and fixing all the other Wilson coefficients to zero. The results are shown in Figure 20.

A comparison between the excluded regions from the two approaches is shown in Figure 21. In the regions where the assumptions used in this study are valid, the excluded regions are very similar in the two approaches. In the Type-I model for large values of $\tan \beta$, the EFT-based approach leads to looser constraints on $\cos(\beta - \alpha)$ than the κ -framework-based approach. This difference stems from the fact that the EFT-based approach (i) does not exploit the constraints from the HVV couplings, that only enter at dimension-8 in the SMEFT expansion and are not considered here, and (ii) retains only terms of $O(\cos(\beta - \alpha))$ in the expansion of κ_λ , while in the κ -framework-based approach the constraint $\kappa_V = \sin(\beta - \alpha)$ and the full dependence of κ_λ on $\cos(\beta - \alpha)$ are considered. However, part of the region of Type-I model parameter space allowed in the EFT-based approach but not in that based on the κ -framework is inconsistent with the alignment limit hypothesis of $|\cos(\beta - \alpha)| \ll 1$.

A significant difference between the $\tan \beta$, $\cos(\beta - \alpha)$ constraints from the approach presented in this section compared to those obtained using the κ -framework-based approach is the absence of the second small allowed region around $\tan \beta \approx 10$, $\cos(\beta - \alpha) \approx 0.2$. This difference follows from the fact that in the EFT-based approach only dimension-6 terms are considered in the SMEFT Lagrangian and a linear expansion of $\sigma \times B$, which does not include terms of second order in the Wilson coefficients, is performed. The likelihood function in the EFT-based approach is thus Gaussian and has a single maximum [114]. As a consequence, the region with a flipped coupling sign does not appear.




Figure 20: Regions of the 2HDM $(\tan\beta, \cos(\beta - \alpha))$ parameter plane excluded at 95% CL (blue filled regions) in the EFT-based approach by the measured rates of Higgs boson production and decays in the (a) Type-I, (b) Type-II, (c) Lepton-specific and (d) Flipped models. The dashed black lines show the borders of the corresponding expected exclusion regions for the SM hypothesis. For the Type-I model, the observed and expected regions excluded at 95% CL when the c_H constraint is considered are also shown (solid and dashed green lines). Results are derived assuming that $|\cos(\beta - \alpha)| \ll 1$, near the alignment limit represented by the red dashed lines, and that the masses of the non-SM-like Higgs bosons are large compared to the SM vev. In particular, the limits on the Type-1 model at high $\tan\beta$ are likely outside the validity range of the assumption made in the model.




Figure 21: Comparison of the constraints from the approaches based on the κ - and EFT-frameworks in the $\tan\beta$, $\cos(\beta - \alpha)$ plane in the (a) Type-I, (b) Type-II, (c) Lepton-specific and (d) Flipped models. The κ_λ constraint is included in the Type-I model interpretation. Solid (dashed) lines indicate the observed (expected) constraints. Results are derived assuming that $|\cos(\beta - \alpha)| \ll 1$, near the alignment limit represented by the red dashed lines, and that the masses of the non-SM-like Higgs bosons are large compared to the SM vev. In particular, the EFT limits on the Type-1 model at high $\tan\beta$ are likely outside the validity range of the assumption made in the model.

4.2 Constraints on the MSSM

Supersymmetry (SUSY) [115–123], a theoretically motivated framework for extending the SM, was conceived to address some of the SM’s unanswered questions. In particular, SUSY offers a solution to the hierarchy problem [124–127], which is related to the fine tuning needed to obtain the relatively small value of the Higgs boson mass compared to the Planck scale. Supersymmetry can also provide credible dark matter candidates [128, 129] and can improve the unification of the electroweak and strong interactions [130–138].

Measurements of Higgs production and decay are interpreted in the context of the minimal supersymmetric extension of the SM (MSSM) [15–19]. The Higgs sector of the MSSM is a 2HDM with Type-II Yukawa couplings. At tree level, all Higgs boson masses and couplings can be expressed in terms of two parameters, usually chosen to be $\tan\beta$ and the mass of the pseudoscalar Higgs boson, m_A . Radiative corrections, including contributions from amplitudes mediated by loops of SM particles and their superpartners, lead to modifications of the tree-level relations and can raise the lightest CP-even Higgs boson mass m_h , bounded at tree level by $m_h \leq m_Z |\cos 2\beta|$, to the level of the mass of the Higgs boson discovered by the ATLAS and CMS collaborations. The MSSM predicts a supersymmetric partner for each SM state. In addition to the supersymmetric spin-one partners of the SM fermions (sfermions) and spin-half partners of the gluons (gluinos), there are a total of eight spin-half partners of the electroweak gauge and Higgs bosons: the neutral bino (superpartner of the $U(1)$ gauge field), the winos, which are a charged pair and a neutral particle (superpartners of the $SU(2)_L$ gauge fields) and the higgsinos, which are two neutral particles and a charged pair (superpartners of the Higgs fields). Mixing between the bino, the neutral wino and the neutral higgsinos leads to four neutralinos, while charginos arise from mixing between the charged winos and the charged higgsinos.

In the MSSM, SUSY is explicitly broken by some soft SUSY breaking operators resulting from a separate sector with unspecified dynamics. As a consequence, the MSSM has over a hundred parameters that describe the pattern of sparticle masses and their decays, and the values of the quantum corrections that they induce vary significantly as a function of these parameters. The parameter space is thus too large to be probed exhaustively through a relatively small set of Higgs boson production and decay rate measurements. Therefore, several *benchmark scenarios* have been proposed in which the values of the MSSM parameters other than $\tan\beta$ and m_A are fixed to provide a wide variety of phenomenologies of the MSSM Higgs sector. With only two free parameters left, $\tan\beta$ and m_A , an MSSM interpretation of the Higgs boson production and decay rate measurements is simplified.

4.2.1 MSSM benchmark scenarios

In this study, the measured production and decay rates of the observed Higgs boson with mass close to 125 GeV are compared with the predictions of eight benchmark scenarios [38–40], assuming that the observed boson is the lightest CP-even Higgs boson h of the MSSM. In each benchmark scenario, constraints are set on m_A and $\tan\beta$ after fixing the values of all the other MSSM parameters. The scenarios that are considered can be briefly summarised as follows, with details provided in the references listed in Table 6:

- **M_h^{125} scenario:** All superparticles are assumed to be so heavy that they affect the production and decay rates of the MSSM Higgs bosons very slightly. The loop-induced contributions to the

Table 6: Benchmark scenarios considered for the MSSM interpretation.

Scenario	Reference
M_h^{125}	[39]
$M_h^{125}(\tilde{\tau})$	[39]
$M_h^{125}(\tilde{\chi})$	[39]
$M_h^{125}(\text{alignment})$	[39]
$M_{h_1}^{125}(\text{CPV})$	[39]
$M_{h,\text{EFT}}^{125}$	[40]
$M_{h,\text{EFT}}^{125}(\tilde{\chi})$	[40]
hMSSM	[38]

couplings of the light CP-even scalar are small. The largest ones, enhanced with $\tan\beta$, affect the b -quark Yukawa coupling but cancel out in the decoupling limit $m_A \gg m_Z$.

- **$M_h^{125}(\tilde{\tau})$ scenario:** The parameters related to the stop, sbottom and gluino masses are the same as in the previous case, but the soft-SUSY-breaking masses and trilinear interaction term for the staus are significantly smaller. The resulting light staus and light wino-like charginos can alter the amplitude for the decay of the SM-like Higgs boson into photons, with contributions that scale as $1/\tan\beta$ and $\tan^2\beta$, respectively. As in the previous scenario, at low m_A the hbb coupling is enhanced, leading to a suppression of the branching ratios for all the other decay channels of the SM-like Higgs boson.
- **$M_h^{125}(\tilde{\chi})$ scenario:** The SUSY parameters in this scenario lead to all charginos and neutralinos being relatively light, with significant higgsino–gaugino mixing, and to a compressed electroweakino mass spectrum. The $\tan\beta$ -enhanced corrections to the hbb coupling are suppressed. At low values of $\tan\beta$, the partial width for the SM-like Higgs boson decay into two photons is significantly increased. The corresponding branching ratio for low $\tan\beta$ is larger (smaller) than the SM prediction for large (small) values of m_A .
- **$M_h^{125}(\text{alignment})$ scenario:** In the ‘alignment without decoupling’ scenario, for a given value of $\tan\beta$, one of the two neutral CP-even scalars has SM-like couplings independent of the mass spectrum of the remaining Higgs bosons. In particular, for $\tan\beta \approx 7$ the lightest scalar Higgs boson h is SM-like, for relatively low values of m_A .
- **$M_{h_1}^{125}(\text{CPV})$ scenario:** This scenario is characterised by CP violation in the Higgs sector, where mixing between the neutral CP-even scalar states h and H and the CP-odd scalar state A is induced by a non-zero phase ϕ_{A_t} in the soft-SUSY-breaking Higgs–stop interaction term. This causes significant interference effects in the production and decay of the two heavier neutral states, which leads to weaker constraints from searches for resonances decaying into $\tau^+\tau^-$. The relevant parameters are chosen such that the strongest interference region is located near the exclusion contour of the corresponding scenario with real parameters, the mass of the SM-like scalar is near the observed value, and all electric dipole moments are within the experimentally allowed ranges.
- **$M_{h,\text{EFT}}^{125}$ scenario:** In the previous scenarios, the sfermions have masses close to the TeV scale. As a consequence, values of $\tan\beta < 5$ are automatically excluded as they would lead to values of the Higgs boson mass m_h that are significantly lower than the experimental value of 125 GeV. To circumvent this problem, in this scenario all SUSY particles, including sfermions, are relatively heavy, with masses close to the SUSY scale m_{SUSY} , which is adjusted dynamically from 6 TeV to

10^{13} TeV to yield a Higgs boson mass of 125 GeV for values of $\tan\beta$ between one and ten. As all SUSY particles are so heavy that production and decays of the MSSM Higgs bosons are only mildly affected by their presence, their contribution to the Higgs boson properties is calculated with an effective field theory. As in most of the previous scenarios, for low values of m_A the hbb coupling is enhanced with respect to the SM prediction and reduces to the SM value for large m_A . At very small $\tan\beta$, the model predicts a small suppression of the htt coupling, leading in turn to a slight suppression of the ggF production cross-section.

- **$M_{h,\text{EFT}}^{125}(\tilde{\chi})$ scenario:** Similarly to the previous scenario, the SUSY scale is adjusted dynamically to yield a light Higgs boson mass of $m_h \approx 125$ GeV at each point of the SUSY parameter space. In contrast to the $M_{h,\text{EFT}}^{125}$ scenario, however, this scenario features relatively light neutralinos and charginos. As a consequence, loop-induced amplitudes mediated by these light states can induce modifications of the light Higgs boson properties such as an enhancement of its partial decay width to two photons, in particular for low $\tan\beta$. The model therefore predicts a branching ratio for $h \rightarrow \gamma\gamma$ significantly larger than the SM prediction for large m_A and low $\tan\beta$. At low values of m_A , the enhancement of the hbb coupling leads instead to an increase of the $h \rightarrow b\bar{b}$ branching ratio and a decrease of the $h \rightarrow \gamma\gamma$ branching ratio relative to their SM expectations.
- **hMSSM scenario:** In this scenario, the lightest Higgs boson h is assumed to be the particle discovered by the ATLAS and CMS collaborations in 2012. Its mass m_h is assumed to be equal to the experimental value of 125.09 GeV, and it is used to fix the values of the dominant radiative corrections to the CP-even Higgs mass matrix from the stop–top sector. This hypothesis determines all other parameters of the Higgs sector, including α , m_H , m_{H^\pm} , and the couplings between the Higgs bosons and the other particles, in terms of only the two parameters m_A and $\tan\beta$. Due to its assumptions, the hMSSM is a good approximation of the MSSM only for moderate values of $\tan\beta$, $1 \lesssim \tan\beta \lesssim 10$ [37].

4.2.2 Interpretation procedure

Since Higgs boson production cross-sections in the MSSM are not calculated for the full STXS Stage-1.2 phase space partitioning of production cross-sections, only inclusive production-mode cross-sections are used in this analysis. The signal strength modifiers $\mu^{i,X}$ in the signal yield expression of Eq. (2) are then written as:

$$\mu^{i,X}(m_A, \tan\beta) = \frac{\sigma_{(\text{h})\text{MSSM}}^i(m_A, \tan\beta)}{\sigma_{\text{SM}}^i} \cdot \frac{\mathcal{B}_{(\text{h})\text{MSSM}}^X(m_A, \tan\beta)}{\mathcal{B}_{\text{SM}}^X} \equiv r^i(m_A, \tan\beta) \cdot r^X(m_A, \tan\beta), \quad (17)$$

where the index i enumerates the production processes (ggF, bbH , ttH , VBF, $qq/qg \rightarrow ZH$, $gg \rightarrow ZH$, $qq \rightarrow WH$).

The values of the production cross-section scale factors r^i and branching ratio scale factors r^X are obtained using the procedures defined by the LHC Higgs Cross-section Working Group, described in [49], as well as the calculations detailed in [139]. For all scenarios except the hMSSM, Higgs boson masses and mixing (and effective Yukawa couplings) are calculated with FEYNHIGGS [140–146], and branching ratios are obtained combining the most precise estimates of FEYNHIGGS, HDECAY [147, 148] and PROPHECY4F [149, 150]. In the hMSSM scenario, the Higgs boson mass is fixed at 125 GeV, the mixing is calculated analytically, and the branching ratios are calculated using the κ -framework. Inclusive cross-sections for ggF production are obtained with SUSHi [151, 152], which includes NLO QCD corrections [153], NNLO QCD

corrections for the top-quark contribution in the effective theory of a heavy top quark [154–158] and EW corrections involving light quarks [159, 160]. For the SM-like Higgs boson, SusHi adds N³LO corrections in the effective theory of a heavy top quark in a threshold expansion [61, 161, 162]. Cross-sections for b -quark-initiated Higgs boson production rely on matched predictions [163–166], which are based on the five flavour NNLO QCD calculation [167] and the four flavour NLO QCD calculation [168, 169]. The values of the cross-sections of rare production modes, such as $gg \rightarrow Zh$, $pp \rightarrow thW$ and $pp \rightarrow thq$, which are not calculated in Refs. [139, 170], are determined through their relations, within the κ -framework, to κ_t , κ_b , κ_W , κ_Z , which are determined as either $\kappa_t^2 = \sigma_{\text{MSSM}}^{t\bar{t}h} / \sigma_{\text{SM}}^{t\bar{t}H}$ or $\kappa_x^2 = \Gamma_{\text{MSSM}}^{h \rightarrow xx} / \Gamma_{\text{SM}}^{h \rightarrow xx}$, where $x = b, W, Z$:

$$\begin{aligned} r^{gg \rightarrow Zh} &= 2.456\kappa_Z^2 + 0.456\kappa_t^2 - 1.903\kappa_Z\kappa_t - 0.011\kappa_Z\kappa_b + 0.003\kappa_t\kappa_b, \\ r^{thW} &= 2.909\kappa_t^2 + 2.310\kappa_W^2 - 4.220\kappa_t\kappa_W, \\ r^{thq} &= 2.633\kappa_t^2 + 3.578\kappa_W^2 - 5.211\kappa_t\kappa_W. \end{aligned}$$

As the cross-section and branching ratio scale factors r^i and r^X are only calculated for a discrete set of points in the two-dimensional benchmark parameter space, a likelihood based on Eq. (17) is not differentially expressed in terms of its theory parameters ($m_A, \tan\beta$). Instead, the values of the profile likelihood ratio are calculated at fixed points in the ($m_A, \tan\beta$) plane, profiling only the nuisance parameters, and then numerically interpolated. Regions of the benchmark scenario parameter space for which the mass m_h of the lightest CP-even Higgs boson h predicted by the MSSM differs from the experimental value of 125.09 GeV by more than 3 GeV, which is the estimated bound of allowed values of the calculated Higgs boson mass in FEYNHIGGS [142, 171], are considered as excluded.

4.2.3 Results

Figure 22 shows the observed and expected 95% CL exclusion limits of the MSSM in the two-dimensional plane ($m_A, \tan\beta$) for the M_h^{125} , $M_h^{125}(\tilde{\chi})$, $M_h^{125}(\tilde{\tau})$, and M_h^{125} (alignment) benchmark scenarios. For all four scenarios, the regions excluded by the Higgs boson mass requirement ($|m_h - 125.09 \text{ GeV}| < 3 \text{ GeV}$) are separately indicated with gray shaded areas. For completeness, the constraints from previous searches for $H/A \rightarrow \tau\tau$ [172] and $H^+ \rightarrow t\bar{b}$ [173] are also overlaid.

In the M_h^{125} and $M_h^{125}(\tilde{\chi})$ scenarios, the low m_A region, in which an enhancement of the $h \rightarrow b\bar{b}$ branching ratio and a suppression of the branching ratios of the other decay modes such as $h \rightarrow \gamma\gamma$ are predicted, is disfavoured. Since the data marginally prefer $B(h \rightarrow b\bar{b})/B_{\text{SM}}(h \rightarrow b\bar{b}) < 1$ and $B(h \rightarrow \gamma\gamma)/B_{\text{SM}}(h \rightarrow \gamma\gamma) > 1$, the observed constraints are tighter than those expected under the SM hypothesis. In the $M_h^{125}(\tilde{\tau})$ scenario, the region at low m_A and $\tan\beta < 53$ is excluded for the same reason. The region $\tan\beta > 53$, in which the $\tilde{\tau}$ loop reduces the hbb coupling, leading to an enhanced prediction of $B(h \rightarrow \gamma\gamma)$, is also largely excluded. In the M_h^{125} (alignment) scenario, the limit of alignment without decoupling is only realised for $\tan\beta \approx 7$ and $m_A \approx 200 \text{ GeV}$. For larger values of m_A , MSSM couplings are more similar to SM couplings, causing the allowed $\tan\beta$ region to broaden.

In all four benchmark scenarios, the MSSM analysis generally excludes the low m_A regime for most of the scanned $\tan\beta$ range, while the requirement $|m_h - 125.09 \text{ GeV}| < 3 \text{ GeV}$ excludes the low $\tan\beta$ range for all scanned values of m_A , a range not fully covered by the direct searches for $H/A \rightarrow \tau\tau$ and $H^+ \rightarrow t\bar{b}$ decays. The exclusion range due to the m_h requirement is qualitatively different in the M_h^{125} (alignment) scenario, as this scenario can only realized for a value of $\tan(\beta) \approx 7$.




Figure 22: Observed (solid yellow area) and expected (dashed line) exclusion contours at 95% CL in the $(m_A, \tan\beta)$ plane for the (a) M_h^{125} , (b) $M_h^{125}(\tilde{\tau})$, (c) $M_h^{125}(\tilde{\chi})$ and (d) $M_h^{125}(\text{alignment})$ scenarios. The parameter space excluded by the condition $|m_h - 125.09| < 3$ GeV is marked in gray. For comparison, the parameter space excluded by the searches for $H/A \rightarrow \tau\tau$ [172] and for $H^\pm \rightarrow t\bar{b}$ [173] are overlaid in blue and purple, respectively. For the $M_h^{125}(\text{alignment})$ scenario, no constraints from the search for $H^\pm \rightarrow t\bar{b}$ are available.

Figure 23 shows the observed and expected 95% CL exclusion limits on the MSSM in the two-dimensional plane of m_{H^\pm} versus $\tan\beta$ for the $M_{h_1}^{125}$ (CPV) scenario and of m_A versus $\tan\beta$ for the $M_{h,\text{EFT}}^{125}$, $M_{h,\text{EFT}}^{125}(\tilde{\chi})$ and hMSSM scenarios. The condition $|m_h - 125.09 \text{ GeV}| < 3 \text{ GeV}$ excludes values of $\tan\beta \lesssim 6$ for all values of m_{H^\pm} in the $M_{h_1}^{125}$ (CPV) scenario, while no part of the parameter space is excluded in the $M_{h,\text{EFT}}^{125}$ and $M_{h,\text{EFT}}^{125}(\tilde{\chi})$ benchmark scenarios, in which $m_h = 125.09 \text{ GeV}$ can be achieved through the flexible m_{SUSY} scale. In these four models, the low m_A region, where an enhancement of the hbb coupling is predicted, is excluded, and the excluded region is larger than expected in the SM hypothesis for the same reason as for the previous four models.




Figure 23: Observed (solid yellow area) and expected (dashed line) exclusion contours at 95% CL in (a) the $(m_{H^\pm}, \tan\beta)$ plane for the $M_{h_1}^{125}$ (CPV) scenario and in the $(m_A, \tan\beta)$ plane for the (b) $M_{h,EFT}^{125}$, (c) $M_{h,EFT}^{125}(\tilde{\chi})$ and (d) hMSSM scenarios. In (a), the parameter space excluded by the condition $|m_h - 125.09$ GeV $<$ 3 GeV is marked in gray.

5 Conclusions

Novel interpretations of the recent combined ATLAS measurements of Higgs boson production and decay rates and simplified template cross-sections in several final states are presented. Constraints on linear combinations of Wilson coefficients corresponding to SM effective field theory operators in the Warsaw basis are reported. In this model-independent parameterisation of BSM physics effects, no significant deviations from the SM are observed. A comparison of results interpreted with a linearised SMEFT model that only considers terms suppressed by up to a factor Λ^{-2} , and a variant including quadratic terms that considers all available terms including those with suppression factor Λ^{-4} , shows that the effect of operators suppressed by Λ^{-4} can significantly affect constraints on Wilson coefficients for a mass scale of $\Lambda = 1$ TeV.

Constraints on a subset of these operators have also been set through a joint interpretation of the fiducial cross-section measurements of Higgs boson production as a function of transverse momentum in the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ final states. Three linear combinations of the Wilson coefficients of operators affecting these spectra can be simultaneously constrained. For two of these combinations, the constraints are similar to those that can be derived from the interpretation of Higgs boson simplified template cross-section measurements in the same channels. The constraint on the top–gluon coupling is significantly stronger in the interpretation of simplified template cross-sections.

Finally, the Higgs boson production and decay rate measurements are interpreted in the context of two-Higgs-doublet models and of eight benchmark scenarios of the minimal supersymmetric model. Constraints are set on the $(\cos(\beta - \alpha), \tan\beta)$ plane in 2HDM Type-I, Type-II, Lepton-specific and Flipped models, and on the parameters $(m_A, \tan\beta)$ of the MSSM in eight benchmark scenarios. These results are complementary to limits from direct searches for additional Higgs bosons.

Acknowledgements

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ISF and Benoziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MEiN, Poland; FCT, Portugal; MNE/IFA, Romania; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DSI/NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taipei; TENMAK, Türkiye; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, CANARIE, CRC and DRAC, Canada; PRIMUS 21/SCI/017 and UNCE SCI/013, Czech Republic; COST, ERC, ERDF, Horizon 2020, ICSC-NextGenerationEU and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex, Investissements d’Avenir IDEX and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; Norwegian Financial Mechanism 2014-2021,

Norway; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF/SFU (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [174].

Appendix: Validity of Gaussian approximation of STXS measurements in the SMEFT interpretation

The STXS SMEFT interpretation model constructs the likelihood function in terms of the fit basis parameters c' from three ingredients:

1. The likelihood function of the STXS measurements, measuring 78 cross-section parameters, and featuring over a thousand nuisance parameters expressing systematic uncertainties.
2. A specification of the impact of all relevant CP-even Warsaw basis SMEFT operators on the SM predictions of these cross-sections, including acceptance corrections where necessary, i.e., Eqs. (12) and (13), using the coefficients $A_j^{\sigma_{i,k'}}$, $A_j^{\Gamma_{H \rightarrow X}}$, $B_{jl}^{\sigma_{i,k'}}$ and $B_{jl}^{\Gamma_{H \rightarrow X}}$ as defined in Eqs. (8)-(11), and visualised in Figure 5.
3. A rotation matrix from the 50 Warsaw basis parameters c , which cannot be unambiguously constrained from the STXS data, to a fit basis c' for which all 19 parameters can be simultaneously constrained from the data, as shown in Figure 8.

An important upside of a likelihood model built this way is that all non-Gaussian effects in the STXS measurements are accounted for, and that the correlated effects of all experimental and theoretical systematic uncertainties are accounted at the level of detail implemented in the original underlying measurements. A downside of this likelihood model is that the required complexity of its implementation prevents it from easily being made available.

An alternative likelihood function, based on a multivariate Gaussian approximation of the STXS measurements instead of the full measurement, with an identical SMEFT re-parameterisation, can be constructed in a relatively straightforward way from the information provided and represents a reasonably good approximation of the full likelihood. A comparison of the results obtained with this simplified Gaussian variant of the likelihood model and those obtained with the full likelihood function is shown in Figure 24 for the linear SMEFT model and in Figure 25 for the SMEFT model including the quadratic terms. Both comparisons indicate generally good agreement between expected and observed parameter estimates and their 68% and 95% uncertainty intervals.




Figure 24: Comparison of the observed result from the full likelihood fit in the SMEFT fit basis with only linear terms (blue) to a fit based on a simplified likelihood constructed from the covariance matrix of the STXS measurements rotated to the same SMEFT fit basis (purple).




Figure 25: Comparison of the observed result from the full likelihood fit in the SMEFT fit basis with linear and quadratic terms (orange) to a fit based on a simplified likelihood constructed from the covariance matrix of the STXS measurements rotated to the same SMEFT fit basis (dark red).

References

- [1] F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, [Phys. Rev. Lett. **13** \(1964\) 321](#).
- [2] P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, [Phys. Lett. **12** \(1964\) 132](#).
- [3] P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, [Phys. Rev. Lett. **13** \(1964\) 508](#).
- [4] G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, *Global Conservation Laws and Massless Particles*, [Phys. Rev. Lett. **13** \(1964\) 585](#).
- [5] P. W. Higgs, *Spontaneous Symmetry Breakdown without Massless Bosons*, [Phys. Rev. **145** \(1966\) 1156](#).
- [6] T. W. B. Kibble, *Symmetry Breaking in Non-Abelian Gauge Theories*, [Phys. Rev. **155** \(1967\) 1554](#).
- [7] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, [Phys. Lett. B **716** \(2012\) 1](#), arXiv: [1207.7214 \[hep-ex\]](#).
- [8] CMS Collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, [Phys. Lett. B **716** \(2012\) 30](#), arXiv: [1207.7235 \[hep-ex\]](#).
- [9] ATLAS Collaboration, *A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery*, [Nature **607** \(2022\) 52](#), arXiv: [2207.00092 \[hep-ex\]](#).
- [10] CMS Collaboration, *A portrait of the Higgs boson by the CMS experiment ten years after the discovery*, [Nature **607** \(2022\) 60](#), arXiv: [2207.00043 \[hep-ex\]](#).
- [11] T. D. Lee, *A Theory of Spontaneous T Violation*, [Phys. Rev. D **8** \(1973\) 1226](#).
- [12] J. F. Gunion and H. E. Haber, *CP-conserving two-Higgs-doublet model: The approach to the decoupling limit*, [Phys. Rev. D **67** \(2003\) 075019](#), arXiv: [hep-ph/0207010 \[hep-ph\]](#).
- [13] G. Branco et al., *Theory and phenomenology of two-Higgs-doublet models*, [Phys. Rept. **516** \(2012\) 1](#), arXiv: [1106.0034 \[hep-ph\]](#).
- [14] LHC Higgs Cross Section Working Group, *Handbook of LHC Higgs Cross Sections: 3. Higgs Properties*, [CERN-2013-004 \(2013\)](#), arXiv: [1307.1347 \[hep-ph\]](#).
- [15] P. Fayet, *Supersymmetry and weak, electromagnetic and strong interactions*, [Phys. Lett. B **64** \(1976\) 159](#).
- [16] P. Fayet, *Spontaneously broken supersymmetric theories of weak, electromagnetic and strong interactions*, [Phys. Lett. B **69** \(1977\) 489](#).
- [17] G. R. Farrar and P. Fayet, *Phenomenology of the production, decay, and detection of new hadronic states associated with supersymmetry*, [Phys. Lett. B **76** \(1978\) 575](#).
- [18] P. Fayet, *Relations between the masses of the superpartners of leptons and quarks, the goldstino coupling and the neutral currents*, [Phys. Lett. B **84** \(1979\) 416](#).

- [19] S. Dimopoulos and H. Georgi, *Softly broken supersymmetry and SU(5)*, [Nucl. Phys. B **193** \(1981\) 150](#).
- [20] ATLAS Collaboration, *Measurements of the Higgs boson inclusive and differential fiducial cross sections in the 4ℓ decay channel at $\sqrt{s} = 13$ TeV*, [Eur. Phys. J. C **80** \(2020\) 942](#), arXiv: [2004.03969 \[hep-ex\]](#).
- [21] ATLAS Collaboration, *Measurements of the Higgs boson inclusive and differential fiducial cross-sections in the diphoton decay channel with pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [JHEP **08** \(2022\) 027](#), arXiv: [2202.00487 \[hep-ex\]](#).
- [22] ATLAS Collaboration, *Measurements of differential cross sections of Higgs boson production through gluon fusion in the $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ final state at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [Eur. Phys. J. C **83** \(2023\) 774](#), arXiv: [2301.06822 \[hep-ex\]](#).
- [23] ATLAS Collaboration, *Integrated and differential fiducial cross-section measurements for the vector boson fusion production of the Higgs boson in the $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ decay channel at 13 TeV with the ATLAS detector*, [Phys. Rev. D **108** \(2023\) 072003](#), arXiv: [2304.03053 \[hep-ex\]](#).
- [24] ATLAS Collaboration, *Higgs boson production cross-section measurements and their EFT interpretation in the 4ℓ decay channel at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [Eur. Phys. J. C **80** \(2020\) 957](#), arXiv: [2004.03447 \[hep-ex\]](#), Erratum: [Eur. Phys. J. C **81** \(2021\) 29](#), Erratum: [Eur. Phys. J. C **81** \(2021\) 398](#).
- [25] ATLAS Collaboration, *Measurement of the properties of Higgs boson production at $\sqrt{s} = 13$ TeV in the $H \rightarrow \gamma\gamma$ channel using 139fb^{-1} of pp collision data with the ATLAS experiment*, [JHEP **07** \(2023\) 088](#), arXiv: [2207.00348 \[hep-ex\]](#).
- [26] ATLAS Collaboration, *Measurements of WH and ZH production in the $H \rightarrow b\bar{b}$ decay channel in pp collisions at 13 TeV with the ATLAS detector*, [Eur. Phys. J. C **81** \(2021\) 178](#), arXiv: [2007.02873 \[hep-ex\]](#).
- [27] ATLAS Collaboration, *Measurement of the associated production of a Higgs boson decaying into b -quarks with a vector boson at high transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [Phys. Lett. B **816** \(2021\) 136204](#), arXiv: [2008.02508 \[hep-ex\]](#).
- [28] ATLAS Collaboration, *Measurements of Higgs bosons decaying to bottom quarks from vector boson fusion production with the ATLAS experiment at $\sqrt{s} = 13$ TeV*, [Eur. Phys. J. C **81** \(2021\) 537](#), arXiv: [2011.08280 \[hep-ex\]](#).
- [29] ATLAS Collaboration, *Constraints on Higgs boson production with large transverse momentum using $H \rightarrow b\bar{b}$ decays in the ATLAS detector*, [Phys. Rev. D **105** \(2022\) 092003](#), arXiv: [2111.08340 \[hep-ex\]](#).
- [30] ATLAS Collaboration, *Measurement of Higgs boson decay into b -quarks in associated production with a top-quark pair in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [JHEP **06** \(2022\) 097](#), arXiv: [2111.06712 \[hep-ex\]](#).
- [31] ATLAS Collaboration, *Measurements of Higgs boson production cross-sections in the $H \rightarrow \tau^+\tau^-$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [JHEP **08** \(2022\) 175](#), arXiv: [2201.08269 \[hep-ex\]](#).

- [32] ATLAS Collaboration, *Measurements of Higgs boson production by gluon–gluon fusion and vector-boson fusion using $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ decays in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **108** (2023) 032005, arXiv: [2207.00338 \[hep-ex\]](#).
- [33] ATLAS Collaboration, *A search for the $Z\gamma$ decay mode of the Higgs boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **809** (2020) 135754, arXiv: [2005.05382 \[hep-ex\]](#).
- [34] ATLAS Collaboration, *A search for the dimuon decay of the Standard Model Higgs boson with the ATLAS detector*, *Phys. Lett. B* **812** (2021) 135980, arXiv: [2007.07830 \[hep-ex\]](#).
- [35] ATLAS Collaboration, *Measurements of gluon-gluon fusion and vector-boson fusion Higgs boson production cross-sections in the $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **789** (2019) 508, arXiv: [1808.09054 \[hep-ex\]](#).
- [36] ATLAS Collaboration, *Evidence for the associated production of the Higgs boson and a top quark pair with the ATLAS detector*, *Phys. Rev. D* **97** (2018) 072003, arXiv: [1712.08891 \[hep-ex\]](#).
- [37] ATLAS Collaboration, *Combined measurements of Higgs boson production and decay using up to 80fb^{-1} of proton–proton collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS experiment*, *Phys. Rev. D* **101** (2020) 012002, arXiv: [1909.02845 \[hep-ex\]](#).
- [38] A. Djouadi et al., *The post-Higgs MSSM scenario: habemus MSSM?*, *Eur. Phys. J. C* **73** (2013) 2650, arXiv: [1307.5205 \[hep-ph\]](#).
- [39] E. Bagnaschi et al., *MSSM Higgs boson searches at the LHC: benchmark scenarios for Run 2 and beyond*, *Eur. Phys. J. C* **79** (2019) 617, arXiv: [1808.07542 \[hep-ph\]](#).
- [40] H. Bahl, S. Liebler and T. Stefaniak, *MSSM Higgs benchmark scenarios for Run 2 and beyond: the low $\tan\beta$ region*, *Eur. Phys. J. C* **79** (2019) 279, arXiv: [1901.05933 \[hep-ph\]](#).
- [41] ATLAS and CMS Collaborations, *Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments*, *Phys. Rev. Lett.* **114** (2015) 191803, arXiv: [1503.07589 \[hep-ex\]](#).
- [42] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, *JINST* **3** (2008) S08003.
- [43] ATLAS Collaboration, *ATLAS Insertable B-Layer: Technical Design Report*, ATLAS-TDR-19; CERN-LHCC-2010-013, 2010, URL: <https://cds.cern.ch/record/1291633>, Addendum: ATLAS-TDR-19-ADD-1; CERN-LHCC-2012-009, 2012, URL: <https://cds.cern.ch/record/1451888>.
- [44] B. Abbott et al., *Production and integration of the ATLAS Insertable B-Layer*, *JINST* **13** (2018) T05008, arXiv: [1803.00844 \[physics.ins-det\]](#).
- [45] ATLAS Collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector at the LHC*, *Eur. Phys. J. C* **76** (2016) 653, arXiv: [1608.03953 \[hep-ex\]](#).
- [46] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*, *JINST* **13** (2018) P07017.

- [47] ATLAS Collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC*, *Eur. Phys. J. C* **83** (2023) 982, arXiv: 2212.09379 [hep-ex].
- [48] ATLAS Collaboration, *Measurement of the production cross section for a Higgs boson in association with a vector boson in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **798** (2019) 134949, arXiv: 1903.10052 [hep-ex].
- [49] LHC Higgs Cross Section Working Group, *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, CERN-2017-002-M (2017), arXiv: 1610.07922 [hep-ph].
- [50] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040, arXiv: hep-ph/0409146.
- [51] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070, arXiv: 0709.2092 [hep-ph].
- [52] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043, arXiv: 1002.2581 [hep-ph].
- [53] S. Alioli, P. Nason, C. Oleari and E. Re, *NLO Higgs boson production via gluon fusion matched with shower in POWHEG*, *JHEP* **04** (2009) 002, arXiv: 0812.0578 [hep-ph].
- [54] K. Hamilton, P. Nason, E. Re and G. Zanderighi, *NNLOPS simulation of Higgs boson production*, *JHEP* **10** (2013) 222, arXiv: 1309.0017 [hep-ph].
- [55] K. Hamilton, P. Nason and G. Zanderighi, *Finite quark-mass effects in the NNLOPS POWHEG+MiNLO Higgs generator*, *JHEP* **05** (2015) 140, arXiv: 1501.04637 [hep-ph].
- [56] S. Catani and M. Grazzini, *Next-to-Next-to-Leading-Order Subtraction Formalism in Hadron Collisions and its Application to Higgs-boson Production at the Large Hadron Collider*, *Phys. Rev. Lett.* **98** (2007) 222002, arXiv: hep-ph/0703012 [hep-ph].
- [57] K. Hamilton, P. Nason, C. Oleari and G. Zanderighi, *Merging H/W/Z + 0 and 1 jet at NLO with no merging scale: a path to parton shower + NNLO matching*, *JHEP* **05** (2013) 082, arXiv: 1212.4504 [hep-ph].
- [58] J. Butterworth et al., *PDF4LHC recommendations for LHC Run II*, *J. Phys. G* **43** (2016) 023001, arXiv: 1510.03865 [hep-ph].
- [59] NNPDF Collaboration, R. D. Ball et al., *Parton distributions for the LHC run II*, *JHEP* **04** (2015) 040, arXiv: 1410.8849 [hep-ph].
- [60] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog and B. Mistlberger, *Higgs Boson Gluon-Fusion Production in QCD at Three Loops*, *Phys. Rev. Lett.* **114** (2015) 212001, arXiv: 1503.06056 [hep-ph].
- [61] C. Anastasiou et al., *High precision determination of the gluon fusion Higgs boson cross-section at the LHC*, *JHEP* **05** (2016) 058, arXiv: 1602.00695 [hep-ph].

- [62] S. Actis, G. Passarino, C. Sturm and S. Uccirati, *NLO electroweak corrections to Higgs boson production at hadron colliders*, [Phys. Lett. B **670** \(2008\) 12](#), arXiv: [0809.1301 \[hep-ph\]](#).
- [63] C. Anastasiou, R. Boughezal and F. Petriello, *Mixed QCD-electroweak corrections to Higgs boson production in gluon fusion*, [JHEP **04** \(2009\) 003](#), arXiv: [0811.3458 \[hep-ph\]](#).
- [64] M. Grazzini and H. Sargsyan, *Heavy-quark mass effects in Higgs boson production at the LHC*, [JHEP **09** \(2013\) 129](#), arXiv: [1306.4581 \[hep-ph\]](#).
- [65] P. Nason and C. Oleari, *NLO Higgs boson production via vector-boson fusion matched with shower in POWHEG*, [JHEP **02** \(2010\) 037](#), arXiv: [0911.5299 \[hep-ph\]](#).
- [66] K. Hamilton, P. Nason and G. Zanderighi, *MINLO: multi-scale improved NLO*, [JHEP **10** \(2012\) 155](#), arXiv: [1206.3572 \[hep-ph\]](#).
- [67] M. Ciccolini, A. Denner and S. Dittmaier, *Strong and Electroweak Corrections to the Production of a Higgs Boson + 2 Jets via Weak Interactions at the Large Hadron Collider*, [Phys. Rev. Lett. **99** \(2007\) 161803](#), arXiv: [0707.0381 \[hep-ph\]](#).
- [68] M. Ciccolini, A. Denner and S. Dittmaier, *Electroweak and QCD corrections to Higgs production via vector-boson fusion at the CERN LHC*, [Phys. Rev. D **77** \(2008\) 013002](#), arXiv: [0710.4749 \[hep-ph\]](#).
- [69] P. Bolzoni, F. Maltoni, S.-O. Moch and M. Zaro, *Higgs Boson Production via Vector-Boson Fusion at Next-to-Next-to-Leading Order in QCD*, [Phys. Rev. Lett. **105** \(2010\) 011801](#), arXiv: [1003.4451 \[hep-ph\]](#).
- [70] O. Brein, A. Djouadi and R. Harlander, *NNLO QCD corrections to the Higgs-strahlung processes at hadron colliders*, [Phys. Lett. B **579** \(2004\) 149](#), arXiv: [hep-ph/0307206](#).
- [71] A. Denner, S. Dittmaier, S. Kallweit and A. Mück, *Electroweak corrections to Higgs-strahlung off W/Z bosons at the Tevatron and the LHC with HAWK*, [JHEP **03** \(2012\) 075](#), arXiv: [1112.5142 \[hep-ph\]](#).
- [72] L. Altenkamp, S. Dittmaier, R. V. Harlander, H. Rzehak and T. J. E. Zirke, *Gluon-induced Higgs-strahlung at next-to-leading order QCD*, [JHEP **02** \(2013\) 078](#), arXiv: [1211.5015 \[hep-ph\]](#).
- [73] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, [JHEP **07** \(2014\) 079](#), arXiv: [1405.0301 \[hep-ph\]](#).
- [74] W. Beenakker et al., *NLO QCD corrections to $t\bar{t}H$ production in hadron collisions*, [Nucl. Phys. B **653** \(2003\) 151](#), arXiv: [hep-ph/0211352 \[hep-ph\]](#).
- [75] S. Dawson, C. Jackson, L. Orr, L. Reina and D. Wackerroth, *Associated Higgs boson production with top quarks at the CERN Large Hadron Collider: NLO QCD corrections*, [Phys. Rev. D **68** \(2003\) 034022](#), arXiv: [hep-ph/0305087 \[hep-ph\]](#).
- [76] Y. Zhang, W.-G. Ma, R.-Y. Zhang, C. Chen and L. Guo, *QCD NLO and EW NLO corrections to $t\bar{t}H$ production with top quark decays at hadron collider*, [Phys. Lett. B **738** \(2014\) 1](#), arXiv: [1407.1110 \[hep-ph\]](#).

- [77] S. Frixione, V. Hirschi, D. Pagani, H.-S. Shao and M. Zaro, *Weak corrections to Higgs hadroproduction in association with a top-quark pair*, [JHEP **09** \(2014\) 065](#), arXiv: [1407.0823 \[hep-ph\]](#).
- [78] F. Demartin, F. Maltoni, K. Mawatari and M. Zaro, *Higgs production in association with a single top quark at the LHC*, [Eur. Phys. J. C **75** \(2015\) 267](#), arXiv: [1504.00611 \[hep-ph\]](#).
- [79] T. Sjöstrand, S. Mrenna and P. Skands, *A brief introduction to PYTHIA 8.1*, [Comput. Phys. Commun. **178** \(2008\) 852](#), arXiv: [0710.3820 \[hep-ph\]](#).
- [80] ATLAS Collaboration, *Measurement of the Z/γ^* boson transverse momentum distribution in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, [JHEP **09** \(2014\) 145](#), arXiv: [1406.3660 \[hep-ex\]](#).
- [81] ATLAS Collaboration, *ATLAS Pythia 8 tunes to 7 TeV data*, ATL-PHYS-PUB-2014-021, 2014, URL: <https://cds.cern.ch/record/1966419>.
- [82] S. Agostinelli et al., *GEANT4 – a simulation toolkit*, [Nucl. Instrum. Meth. A **506** \(2003\) 250](#).
- [83] ATLAS Collaboration, *The ATLAS Simulation Infrastructure*, [Eur. Phys. J. C **70** \(2010\) 823](#), arXiv: [1005.4568 \[physics.ins-det\]](#).
- [84] A. D. Martin, W. J. Stirling, R. S. Thorne and G. Watt, *Parton distributions for the LHC*, [Eur. Phys. J. C **63** \(2009\) 189](#), arXiv: [0901.0002 \[hep-ph\]](#).
- [85] ATLAS Collaboration, *Summary of ATLAS Pythia 8 tunes*, ATL-PHYS-PUB-2012-003, 2012, URL: <https://cds.cern.ch/record/1474107>.
- [86] NNPDF Collaboration, R. D. Ball et al., *Parton distributions with LHC data*, [Nucl. Phys. B **867** \(2013\) 244](#), arXiv: [1207.1303 \[hep-ph\]](#).
- [87] ATLAS Collaboration, *The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie–Landshoff diffractive model*, ATL-PHYS-PUB-2016-017, 2016, URL: <https://cds.cern.ch/record/2206965>.
- [88] S. Badger et al., *Les Houches 2015: Physics at TeV Colliders Standard Model Working Group Report*, (2016), arXiv: [1605.04692 \[hep-ph\]](#).
- [89] N. Berger et al., *Simplified Template Cross Sections - Stage 1.1*, (2019), arXiv: [1906.02754 \[hep-ph\]](#).
- [90] S. Amoroso et al., ‘Les Houches 2019: Physics at TeV Colliders: Standard Model Working Group Report’, *11th Les Houches Workshop on Physics at TeV Colliders: PhysTeV Les Houches*, 2020, arXiv: [2003.01700 \[hep-ph\]](#).
- [91] ATLAS Collaboration, *Evaluation of theoretical uncertainties for simplified template cross section measurements of V -associated production of the Higgs boson*, ATL-PHYS-PUB-2018-035, 2018, URL: <https://cds.cern.ch/record/2649241>.
- [92] I. Brivio and M. Trott, *The standard model as an effective field theory*, [Phys. Rept. **793** \(2019\) 1](#), arXiv: [1706.08945 \[hep-ph\]](#).
- [93] ATLAS Collaboration, *Methodology for EFT interpretation of Higgs boson simplified template cross-section results in ATLAS*, ATL-PHYS-PUB-2019-042, 2019, URL: <https://cds.cern.ch/record/2694284>.

- [94] B. Grzadkowski, M. Iskrzyński, M. Misiak and J. Rosiek, *Dimension-six terms in the Standard Model Lagrangian*, **JHEP** **10** (2010) 085, arXiv: [1008.4884 \[hep-ph\]](#).
- [95] J. A. Aguilar-Saavedra et al., *Interpreting top-quark LHC measurements in the standard-model effective field theory*, CERN-LPCC-2018-01 (2018), arXiv: [1802.07237 \[hep-ph\]](#).
- [96] I. Brivio, Y. Jiang and M. Trott, *The SMEFTsim package, theory and tools*, **JHEP** **12** (2017) 070, arXiv: [1709.06492 \[hep-ph\]](#).
- [97] C. Degrande et al., *Standard Model Effective Theory at Next-to-Leading-Order in QCD*, URL: <http://feynrules.irmp.ucl.ac.be/wiki/SMEFTatNLO>.
- [98] S. Dawson and P. P. Giardino, *Electroweak corrections to Higgs boson decays to $\gamma\gamma$ and W^+W^- in standard model EFT*, **Phys. Rev. D** **98** (2018) 095005, arXiv: [1807.11504 \[hep-ph\]](#).
- [99] S. Dawson and P. P. Giardino, *Higgs decays to ZZ and $Z\gamma$ in the standard model effective field theory: An NLO analysis*, **Phys. Rev. D** **97** (2018) 093003, arXiv: [1801.01136 \[hep-ph\]](#).
- [100] M. L. Mangano, M. Moretti, F. Piccinini and M. Treccani, *Matching matrix elements and shower evolution for top-pair production in hadronic collisions*, **JHEP** **01** (2007) 013, arXiv: [hep-ph/0611129](#).
- [101] S. Mrenna and P. Richardson, *Matching matrix elements and parton showers with HERWIG and PYTHIA*, **JHEP** **05** (2004) 040, arXiv: [hep-ph/0312274](#).
- [102] A. Buckley et al., *Rivet user manual*, **Comput. Phys. Commun.** **184** (2013) 2803, arXiv: [1003.0694 \[hep-ph\]](#).
- [103] T. Corbett, A. Martin and M. Trott, *Consistent higher order $\sigma(\mathcal{G}\mathcal{G} \rightarrow h)$, $\Gamma(h \rightarrow \mathcal{G}\mathcal{G})$ and $\Gamma(h \rightarrow \gamma\gamma)$ in geoSMEFT*, **JHEP** **12** (2021) 147, arXiv: [2107.07470 \[hep-ph\]](#).
- [104] T. Corbett and A. Martin, *Higgs associated production with a vector decaying to two fermions in the geoSMEFT*, (2023), arXiv: [2306.00053 \[hep-ph\]](#).
- [105] A. Martin and M. Trott, *More accurate $\sigma(\mathcal{G}\mathcal{G} \rightarrow h)$, $\Gamma(h \rightarrow \mathcal{G}\mathcal{G}, \mathcal{A}\mathcal{A}, \bar{\Psi}\Psi)$ and Higgs width results via the geoSMEFT*, (2023), arXiv: [2305.05879 \[hep-ph\]](#).
- [106] C. Degrande, B. Fuks, K. Mawatari, K. Mimasu and V. Sanz, *Electroweak Higgs boson production in the standard model effective field theory beyond leading order in QCD*, **Eur. Phys. J. C** **77** (2017) 262, arXiv: [1609.04833 \[hep-ph\]](#).
- [107] S. L. Glashow and S. Weinberg, *Natural conservation laws for neutral currents*, **Phys. Rev. D** **15** (1977) 1958.
- [108] E. A. Paschos, *Diagonal neutral currents*, **Phys. Rev. D** **15** (1977) 1966.
- [109] F. Monti et al., *Modelling of the single-Higgs simplified template cross-sections (STXS 1.2) for the determination of the Higgs boson trilinear self-coupling*, tech. rep., 2022, URL: <https://cds.cern.ch/record/2803606>.

- [110] F. Arco, S. Heinemeyer and M. J. Herrero, *Exploring sizable triple Higgs couplings in the 2HDM*, *Eur. Phys. J. C* **80** (2020) 884, arXiv: 2005.10576 [hep-ph].
- [111] ATLAS Collaboration, *Constraining the Higgs boson self-coupling from single- and double-Higgs production with the ATLAS detector using pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **843** (2023) 137745, arXiv: 2211.01216 [hep-ex].
- [112] S. Dawson, S. Homiller and S. D. Lane, *Putting standard model EFT fits to work*, *Phys. Rev. D* **102** (2020) 055012, arXiv: 2007.01296 [hep-ph].
- [113] S. Dawson, D. Fontes, S. Homiller and M. Sullivan, *Role of dimension-eight operators in an EFT for the 2HDM*, *Phys. Rev. D* **106** (2022) 055012, arXiv: 2205.01561 [hep-ph].
- [114] H. Bélusca-Maïto, A. Falkowski, D. Fontes, J. C. Romão and J. P. Silva, *Higgs EFT for 2HDM and beyond*, *Eur. Phys. J. C* **77** (2017) 176, arXiv: 1611.01112 [hep-ph].
- [115] H. Miyazawa, *Baryon Number Changing Currents*, *Progress of Theoretical Physics* **36** (1966) 1266.
- [116] P. Ramond, *Dual Theory for Free Fermions*, *Phys. Rev. D* **3** (1971) 2415.
- [117] Y. A. Gol'fand and E. P. Likhtman, *Extension of the algebra of Poincare group generators and violation of P invariance*, *JETP Lett.* **13** (1971) 323, reprinted in *Supergravities in Diverse Dimensions* (1989) 20.
- [118] A. Neveu and J. H. Schwarz, *Factorizable dual model of pions*, *Nucl. Phys. B* **31** (1971) 86.
- [119] A. Neveu and J. H. Schwarz, *Quark Model of Dual Pions*, *Phys. Rev. D* **4** (1971) 1109.
- [120] J.-L. Gervais and B. Sakita, *Field theory interpretation of supergauges in dual models*, *Nucl. Phys. B* **34** (1971) 632.
- [121] D. V. Volkov and V. P. Akulov, *Is the neutrino a goldstone particle?*, *Phys. Lett. B* **46** (1973) 109.
- [122] J. Wess and B. Zumino, *A lagrangian model invariant under supergauge transformations*, *Phys. Lett. B* **49** (1974) 52.
- [123] J. Wess and B. Zumino, *Supergauge transformations in four dimensions*, *Nucl. Phys. B* **70** (1974) 39.
- [124] S. Weinberg, *Implications of dynamical symmetry breaking*, *Phys. Rev. D* **13** (1976) 974.
- [125] E. Gildener, *Gauge-symmetry hierarchies*, *Phys. Rev. D* **14** (1976) 1667.
- [126] S. Weinberg, *Implications of dynamical symmetry breaking: An addendum*, *Phys. Rev. D* **19** (1979) 1277.
- [127] L. Susskind, *Dynamics of spontaneous symmetry breaking in the Weinberg-Salam theory*, *Phys. Rev. D* **20** (1979) 2619.
- [128] H. Goldberg, *Constraint on the Photino Mass from Cosmology*, *Phys. Rev. Lett.* **50** (1983) 1419.
- [129] J. Ellis, J. Hagelin, D. Nanopoulos, K. Olive and M. Srednicki, *Supersymmetric relics from the big bang*, *Nucl. Phys. B* **238** (1984) 453.
- [130] S. Dimopoulos, S. Raby and F. Wilczek, *Supersymmetry and the scale of unification*, *Phys. Rev. D* **24** (1981) 1681.
- [131] N. Sakai, *Naturalnes in supersymmetric GUTS*, *Z. Phys. C* **11** (1981) 153.

- [132] L. Ibáñez and G. Ross, *Low-energy predictions in supersymmetric grand unified theories*, *Phys. Lett. B* **105** (1981) 439.
- [133] M. Einhorn and D. Jones, *The weak mixing angle and unification mass in supersymmetric SU(5)*, *Nucl. Phys. B* **196** (1982) 475.
- [134] W. J. Marciano and G. Senjanović, *Predictions of supersymmetric grand unified theories*, *Phys. Rev. D* **25** (1982) 3092.
- [135] C. Giunti, C. W. Kim and U. W. Lee, *Running coupling constants and grand unification models*, *Mod. Phys. Lett. A* **06** (1991) 1745.
- [136] J. Ellis, S. Kelley and D. Nanopoulos, *Probing the desert using gauge coupling unification*, *Phys. Lett. B* **260** (1991) 131.
- [137] U. Amaldi, W. de Boer and H. Fürstenau, *Comparison of grand unified theories with electroweak and strong coupling constants measured at LEP*, *Phys. Lett. B* **260** (1991) 447.
- [138] P. Langacker and M. Luo, *Implications of precision electroweak experiments for m_t , ρ_0 , $\sin^2\theta_W$, and grand unification*, *Phys. Rev. D* **44** (1991) 817.
- [139] E. A. Bagnaschi, S. Heinemeyer, S. Liebler, P. Slavich and M. Spira, *Benchmark Scenarios for MSSM Higgs Boson Searches at the LHC*, LHCHWG-2021-001, 2021, URL: <https://cds.cern.ch/record/2791954>.
- [140] S. Heinemeyer, W. Hollik and G. Weiglein, *FeynHiggs: a program for the calculation of the masses of the neutral CP-even Higgs bosons in the MSSM*, *Comput. Phys. Commun.* **124** (2000) 76, arXiv: [hep-ph/9812320](https://arxiv.org/abs/hep-ph/9812320) [[hep-ph](https://arxiv.org/abs/hep-ph/9812320)].
- [141] S. Heinemeyer, W. Hollik and G. Weiglein, *The masses of the neutral CP-even Higgs bosons in the MSSM: Accurate analysis at the two-loop level*, *Eur. Phys. J. C* **9** (1999) 343, arXiv: [hep-ph/9812472](https://arxiv.org/abs/hep-ph/9812472) [[hep-ph](https://arxiv.org/abs/hep-ph/9812472)].
- [142] G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich and G. Weiglein, *Towards high-precision predictions for the MSSM Higgs sector*, *Eur. Phys. J. C* **28** (2003) 133, arXiv: [hep-ph/0212020](https://arxiv.org/abs/hep-ph/0212020) [[hep-ph](https://arxiv.org/abs/hep-ph/0212020)].
- [143] M. Frank et al., *The Higgs boson masses and mixings of the complex MSSM in the Feynman-diagrammatic approach*, *JHEP* **02** (2007) 047.
- [144] T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak and G. Weiglein, *High-Precision Predictions for the Light CP-Even Higgs Boson Mass of the Minimal Supersymmetric Standard Model*, *Phys. Rev. Lett.* **112** (2014).
- [145] H. Bahl and W. Hollik, *Precise prediction for the light MSSM Higgs-boson mass combining effective field theory and fixed-order calculations*, *Eur. Phys. J. C* **76** (2016) 499, arXiv: [1608.01880](https://arxiv.org/abs/1608.01880) [[hep-ph](https://arxiv.org/abs/1608.01880)].
- [146] H. Bahl, S. Heinemeyer, W. Hollik and G. Weiglein, *Reconciling EFT and hybrid calculations of the light MSSM Higgs-boson mass*, *Eur. Phys. J. C* **78** (2018) 57, arXiv: [1706.00346](https://arxiv.org/abs/1706.00346) [[hep-ph](https://arxiv.org/abs/1706.00346)].
- [147] A. Djouadi, J. Kalinowski and M. Spira, *HDECAY: A program for Higgs boson decays in the Standard Model and its supersymmetric extension*, *Comput. Phys. Commun.* **108** (1998) 56, arXiv: [hep-ph/9704448](https://arxiv.org/abs/hep-ph/9704448).

- [148] A. Djouadi, J. Kalinowski, M. Mühlleitner and M. Spira, *HDECAY: Twenty++ years after*, *Comput. Phys. Commun.* **238** (2019) 214, arXiv: [1801.09506 \[hep-ph\]](#).
- [149] A. Bredenstein, A. Denner, S. Dittmaier and M. M. Weber, *Precise predictions for the Higgs-boson decay $H \rightarrow WW/ZZ \rightarrow 4$ leptons*, *Phys. Rev. D* **74** (2006) 013004, arXiv: [hep-ph/0604011 \[hep-ph\]](#).
- [150] A. Bredenstein, A. Denner, S. Dittmaier and M. M. Weber, *Radiative corrections to the semileptonic and hadronic Higgs-boson decays $H \rightarrow WW/ZZ \rightarrow 4$ fermions*, *JHEP* **02** (2007) 080, arXiv: [hep-ph/0611234](#).
- [151] R. V. Harlander, S. Liebler and H. Mantler, *SusHi: A program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model and the MSSM*, *Comput. Phys. Commun.* **184** (2013) 1605, arXiv: [1212.3249 \[hep-ph\]](#).
- [152] R. V. Harlander, S. Liebler and H. Mantler, *SusHi Bento: Beyond NNLO and the heavy-top limit*, *Comput. Phys. Commun.* **212** (2017) 239, arXiv: [1605.03190 \[hep-ph\]](#).
- [153] M. Spira, A. Djouadi, D. Graudenz and P. Zerwas, *Higgs boson production at the LHC*, *Nucl. Phys. B* **453** (1995) 17, arXiv: [hep-ph/9504378](#).
- [154] R. V. Harlander and W. B. Kilgore, *Next-to-Next-to-Leading Order Higgs Production at Hadron Colliders*, *Phys. Rev. Lett.* **88** (2002) 201801, arXiv: [hep-ph/0201206](#).
- [155] C. Anastasiou and K. Melnikov, *Higgs boson production at hadron colliders in NNLO QCD*, *Nucl. Phys. B* **646** (2002) 220, arXiv: [hep-ph/0207004](#).
- [156] V. Ravindran, J. Smith and W. L. van Neerven, *NNLO corrections to the total cross section for Higgs boson production in hadron-hadron collisions*, *Nucl. Phys. B* **665** (2003) 325, arXiv: [hep-ph/0302135](#).
- [157] R. V. Harlander and W. B. Kilgore, *Production of a pseudo-scalar Higgs boson at hadron colliders at next-to-next-to leading order*, *JHEP* **10** (2002) 017, arXiv: [hep-ph/0208096 \[hep-ph\]](#).
- [158] C. Anastasiou and K. Melnikov, *Pseudoscalar Higgs boson production at hadron colliders in next-to-next-to-leading order QCD*, *Phys. Rev. D* **67** (2003) 037501, arXiv: [hep-ph/0208115 \[hep-ph\]](#).
- [159] U. Aglietti, R. Bonciani, G. Degrossi and A. Vicini, *Two-loop light fermion contribution to Higgs production and decays*, *Phys. Lett. B* **595** (2004) 432, arXiv: [hep-ph/0404071](#).
- [160] R. Bonciani, G. Degrossi and A. Vicini, *On the generalized harmonic polylogarithms of one complex variable*, *Comput. Phys. Commun.* **182** (2011) 1253, arXiv: [1007.1891 \[hep-ph\]](#).
- [161] C. Anastasiou et al., *Higgs boson gluon-fusion production beyond threshold in N^3 LO QCD*, *JHEP* **03** (2015) 091, arXiv: [1411.3584 \[hep-ph\]](#).
- [162] C. Anastasiou et al., *Soft expansion of double-real-virtual corrections to Higgs production at N^3 LO*, *JHEP* **08** (2015) 051, arXiv: [1505.04110 \[hep-ph\]](#).
- [163] M. Bonvini, A. S. Papanastasiou and F. J. Tackmann, *Resummation and matching of b -quark mass effects in $b\bar{b}H$ production*, *JHEP* **11** (2015) 196, arXiv: [1508.03288 \[hep-ph\]](#).

- [164] M. Bonvini, A. S. Papanastasiou and F. J. Tackmann,
Matched predictions for the $b\bar{b}H$ cross section at the 13 TeV LHC, *JHEP* **10** (2016) 053,
arXiv: [1605.01733 \[hep-ph\]](#).
- [165] S. Forte, D. Napoletano and M. Ubiali,
Higgs production in bottom-quark fusion in a matched scheme, *Phys. Lett. B* **751** (2015) 331,
arXiv: [1508.01529 \[hep-ph\]](#).
- [166] S. Forte, D. Napoletano and M. Ubiali,
Higgs production in bottom-quark fusion: matching beyond leading order,
Phys. Lett. B **763** (2016) 190, arXiv: [1607.00389 \[hep-ph\]](#).
- [167] R. V. Harlander and W. B. Kilgore,
Higgs boson production in bottom quark fusion at next-to-next-to-leading order,
Phys. Rev. D **68** (2003) 013001, arXiv: [hep-ph/0304035 \[hep-ph\]](#).
- [168] S. Dittmaier, M. Krämer and M. Spira,
Higgs radiation off bottom quarks at the Fermilab Tevatron and the CERN LHC,
Phys. Rev. D **70** (2004) 074010, arXiv: [hep-ph/0309204 \[hep-ph\]](#).
- [169] S. Dawson, C. Jackson, L. Reina and D. Wackerroth,
Exclusive Higgs boson production with bottom quarks at hadron colliders,
Phys. Rev. D **69** (2004) 074027, arXiv: [hep-ph/0311067 \[hep-ph\]](#).
- [170] LHC Higgs WG, *MSSM Neutral Higgs LHCXSWG3 Twiki page*,
URL: <https://twiki.cern.ch/twiki/bin/view/LHCPhysics/LHCHXSWGSSMNeutral>.
- [171] B. C. Allanach, A. Djouadi, J.-L. Kneur, W. Porod and P. Slavich,
Precise determination of the neutral Higgs boson masses in the MSSM, *JHEP* **09** (2004) 044,
arXiv: [hep-ph/0406166](#).
- [172] ATLAS Collaboration, *Search for Heavy Higgs Bosons Decaying into Two Tau Leptons with the ATLAS Detector Using pp Collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **125** (2020) 051801,
arXiv: [2002.12223 \[hep-ex\]](#).
- [173] ATLAS Collaboration, *Search for charged Higgs bosons decaying into a top quark and a bottom quark at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **06** (2021) 145,
arXiv: [2102.10076 \[hep-ex\]](#).
- [174] ATLAS Collaboration, *ATLAS Computing Acknowledgements*, ATL-SOFT-PUB-2023-001, 2023,
URL: <https://cds.cern.ch/record/2869272>.

The ATLAS Collaboration

G. Aad ¹⁰², B. Abbott ¹²⁰, K. Abeling ⁵⁵, N.J. Abicht ⁴⁹, S.H. Abidi ²⁹, A. Aboulhorma ^{35e}, H. Abramowicz ¹⁵¹, H. Abreu ¹⁵⁰, Y. Abulaiti ¹¹⁷, B.S. Acharya ^{69a,69b,m}, C. Adam Bourdarios ⁴, L. Adamczyk ^{86a}, S.V. Addepalli ²⁶, M.J. Addison ¹⁰¹, J. Adelman ¹¹⁵, A. Adiguzel ^{21c}, T. Adye ¹³⁴, A.A. Affolder ¹³⁶, Y. Afik ³⁶, M.N. Agaras ¹³, J. Agarwala ^{73a,73b}, A. Aggarwal ¹⁰⁰, C. Agheorghiesei ^{27c}, A. Ahmad ³⁶, F. Ahmadov ^{38,y}, W.S. Ahmed ¹⁰⁴, S. Ahuja ⁹⁵, X. Ai ^{62a}, G. Aielli ^{76a,76b}, A. Aikot ¹⁶³, M. Ait Tamlihat ^{35e}, B. Aitbenchikh ^{35a}, I. Aizenberg ¹⁶⁹, M. Akbiyik ¹⁰⁰, T.P.A. Åkesson ⁹⁸, A.V. Akimov ³⁷, D. Akiyama ¹⁶⁸, N.N. Akolkar ²⁴, S. Aktas ^{21a}, K. Al Houry ⁴¹, G.L. Alberghi ^{23b}, J. Albert ¹⁶⁵, P. Albicocco ⁵³, G.L. Albouy ⁶⁰, S. Alderweireldt ⁵², Z.L. Alegria ¹²¹, M. Aleksa ³⁶, I.N. Aleksandrov ³⁸, C. Alexa ^{27b}, T. Alexopoulos ¹⁰, F. Alfonsi ^{23b}, M. Algren ⁵⁶, M. Alhroob ¹²⁰, B. Ali ¹³², H.M.J. Ali ⁹¹, S. Ali ¹⁴⁸, S.W. Alibocus ⁹², M. Aliev ¹⁴⁵, G. Alimonti ^{71a}, W. Alkakhki ⁵⁵, C. Allaire ⁶⁶, B.M.M. Allbrooke ¹⁴⁶, J.F. Allen ⁵², C.A. Allendes Flores ^{137f}, P.P. Allport ²⁰, A. Aloisio ^{72a,72b}, F. Alonso ⁹⁰, C. Alpigiani ¹³⁸, M. Alvarez Estevez ⁹⁹, A. Alvarez Fernandez ¹⁰⁰, M. Alves Cardoso ⁵⁶, M.G. Alviggi ^{72a,72b}, M. Aly ¹⁰¹, Y. Amaral Coutinho ^{83b}, A. Ambler ¹⁰⁴, C. Amelung ³⁶, M. Amerl ¹⁰¹, C.G. Ames ¹⁰⁹, D. Amidei ¹⁰⁶, S.P. Amor Dos Santos ^{130a}, K.R. Amos ¹⁶³, V. Ananiev ¹²⁵, C. Anastopoulos ¹³⁹, T. Andeen ¹¹, J.K. Anders ³⁶, S.Y. Andrean ^{47a,47b}, A. Andreatta ^{71a,71b}, S. Angelidakis ⁹, A. Angerami ^{41,ab}, A.V. Anisenkov ³⁷, A. Annovi ^{74a}, C. Antel ⁵⁶, M.T. Anthony ¹³⁹, E. Antipov ¹⁴⁵, M. Antonelli ⁵³, F. Anulli ^{75a}, M. Aoki ⁸⁴, T. Aoki ¹⁵³, J.A. Aparisi Pozo ¹⁶³, M.A. Aparo ¹⁴⁶, L. Aperio Bella ⁴⁸, C. Appelt ¹⁸, A. Apyan ²⁶, N. Aranzabal ³⁶, S.J. Arbiol Val ⁸⁷, C. Arcangeletti ⁵³, A.T.H. Arce ⁵¹, E. Arena ⁹², J-F. Arguin ¹⁰⁸, S. Argyropoulos ⁵⁴, J.-H. Arling ⁴⁸, O. Arnaez ⁴, H. Arnold ¹¹⁴, G. Artoni ^{75a,75b}, H. Asada ¹¹¹, K. Asai ¹¹⁸, S. Asai ¹⁵³, N.A. Asbah ⁶¹, K. Assamagan ²⁹, R. Astalos ^{28a}, S. Atashi ¹⁶⁰, R.J. Atkin ^{33a}, M. Atkinson ¹⁶², H. Atmani ^{35f}, P.A. Atlasiddha ¹²⁸, K. Augsten ¹³², S. Auricchio ^{72a,72b}, A.D. Auriol ²⁰, V.A. Austrup ¹⁰¹, G. Avolio ³⁶, K. Axiotis ⁵⁶, G. Azuelos ^{108,af}, D. Babal ^{28b}, H. Bachacou ¹³⁵, K. Bachas ^{152,p}, A. Bachi ³⁴, F. Backman ^{47a,47b}, A. Badea ⁶¹, T.M. Baer ¹⁰⁶, P. Bagnaia ^{75a,75b}, M. Bahmani ¹⁸, D. Bahner ⁵⁴, A.J. Bailey ¹⁶³, V.R. Bailey ¹⁶², J.T. Baines ¹³⁴, L. Baines ⁹⁴, O.K. Baker ¹⁷², E. Bakos ¹⁵, D. Bakshi Gupta ⁸, V. Balakrishnan ¹²⁰, R. Balasubramanian ¹¹⁴, E.M. Baldin ³⁷, P. Balek ^{86a}, E. Ballabene ^{23b,23a}, F. Balli ¹³⁵, L.M. Baltes ^{63a}, W.K. Balunas ³², J. Balz ¹⁰⁰, E. Banas ⁸⁷, M. Bandieramonte ¹²⁹, A. Bandyopadhyay ²⁴, S. Bansal ²⁴, L. Barak ¹⁵¹, M. Barakat ⁴⁸, E.L. Barberio ¹⁰⁵, D. Barberis ^{57b,57a}, M. Barbero ¹⁰², M.Z. Barel ¹¹⁴, K.N. Barends ^{33a}, T. Barillari ¹¹⁰, M-S. Barisits ³⁶, T. Barklow ¹⁴³, P. Baron ¹²², D.A. Baron Moreno ¹⁰¹, A. Baroncelli ^{62a}, G. Barone ²⁹, A.J. Barr ¹²⁶, J.D. Barr ⁹⁶, L. Barranco Navarro ^{47a,47b}, F. Barreiro ⁹⁹, J. Barreiro Guimarães da Costa ^{14a}, U. Barron ¹⁵¹, M.G. Barros Teixeira ^{130a}, S. Barsov ³⁷, F. Bartels ^{63a}, R. Bartoldus ¹⁴³, A.E. Barton ⁹¹, P. Bartos ^{28a}, A. Basan ¹⁰⁰, M. Baselga ⁴⁹, A. Bassalat ^{66,b}, M.J. Basso ^{156a}, C.R. Basson ¹⁰¹, R.L. Bates ⁵⁹, S. Batlamous ^{35e}, J.R. Batley ³², B. Batool ¹⁴¹, M. Battaglia ¹³⁶, D. Battulga ¹⁸, M. Baunce ^{75a,75b}, M. Bauer ³⁶, P. Bauer ²⁴, L.T. Bazzano Hurrell ³⁰, J.B. Beacham ⁵¹, T. Beau ¹²⁷, J.Y. Beauchamp ⁹⁰, P.H. Beauchemin ¹⁵⁸, F. Becherer ⁵⁴, P. Bechtel ²⁴, H.P. Beck ^{19,o}, K. Becker ¹⁶⁷, A.J. Beddall ⁸², V.A. Bednyakov ³⁸, C.P. Bee ¹⁴⁵, L.J. Beamster ¹⁵, T.A. Beermann ³⁶, M. Begalli ^{83d}, M. Begel ²⁹, A. Behera ¹⁴⁵, J.K. Behr ⁴⁸, J.F. Beirer ³⁶, F. Beisiegel ²⁴, M. Belfkir ¹⁵⁹, G. Bella ¹⁵¹, L. Bellagamba ^{23b}, A. Bellerive ³⁴, P. Bellos ²⁰, K. Beloborodov ³⁷, D. Bencheekroun ^{35a}, F. Bendebba ^{35a}, Y. Benhammou ¹⁵¹, M. Benoit ²⁹, J.R. Bensinger ²⁶, S. Bentvelsen ¹¹⁴, L. Beresford ⁴⁸,

M. Beretta ⁵³, E. Bergeaas Kuutmann ¹⁶¹, N. Berger ⁴, B. Bergmann ¹³², J. Beringer ^{17a},
G. Bernardi ⁵, C. Bernius ¹⁴³, F.U. Bernlochner ²⁴, F. Bernon ^{36,102}, A. Berrocal Guardia ¹³,
T. Berry ⁹⁵, P. Berta ¹³³, A. Berthold ⁵⁰, I.A. Bertram ⁹¹, S. Bethke ¹¹⁰, A. Betti ^{75a,75b},
A.J. Bevan ⁹⁴, N.K. Bhalla ⁵⁴, M. Bhamjee ^{33c}, S. Bhatta ¹⁴⁵, D.S. Bhattacharya ¹⁶⁶,
P. Bhattarai ¹⁴³, V.S. Bhopatkar ¹²¹, R. Bi ^{29,ai}, R.M. Bianchi ¹²⁹, G. Bianco ^{23b,23a}, O. Biebel ¹⁰⁹,
R. Bielski ¹²³, M. Biglietti ^{77a}, M. Bindi ⁵⁵, A. Bingul ^{21b}, C. Bini ^{75a,75b}, A. Biondini ⁹²,
C.J. Birch-sykes ¹⁰¹, G.A. Bird ^{20,134}, M. Birman ¹⁶⁹, M. Biros ¹³³, S. Biryukov ¹⁴⁶,
T. Bisanz ⁴⁹, E. Bisceglie ^{43b,43a}, J.P. Biswal ¹³⁴, D. Biswas ¹⁴¹, A. Bitadze ¹⁰¹, K. Bjørke ¹²⁵,
I. Bloch ⁴⁸, A. Blue ⁵⁹, U. Blumenschein ⁹⁴, J. Blumenthal ¹⁰⁰, G.J. Bobbink ¹¹⁴,
V.S. Bobrovnikov ³⁷, M. Boehler ⁵⁴, B. Boehm ¹⁶⁶, D. Bogavac ³⁶, A.G. Bogdanchikov ³⁷,
C. Bohm ^{47a}, V. Boisvert ⁹⁵, P. Bokan ⁴⁸, T. Bold ^{86a}, M. Bomben ⁵, M. Bona ⁹⁴,
M. Boonekamp ¹³⁵, C.D. Booth ⁹⁵, A.G. Borbély ⁵⁹, I.S. Bordulev ³⁷, H.M. Borecka-Bielska ¹⁰⁸,
G. Borissov ⁹¹, D. Bortoletto ¹²⁶, D. Boscherini ^{23b}, M. Bosman ¹³, J.D. Bossio Sola ³⁶,
K. Bouaouda ^{35a}, N. Bouchhar ¹⁶³, J. Boudreau ¹²⁹, E.V. Bouhova-Thacker ⁹¹, D. Boumediene ⁴⁰,
R. Bouquet ¹⁶⁵, A. Boveia ¹¹⁹, J. Boyd ³⁶, D. Boye ²⁹, I.R. Boyko ³⁸, J. Bracinik ²⁰,
N. Brahimy ^{62d}, G. Brandt ¹⁷¹, O. Brandt ³², F. Braren ⁴⁸, B. Brau ¹⁰³, J.E. Brau ¹²³,
R. Brenner ¹⁶⁹, L. Brenner ¹¹⁴, R. Brenner ¹⁶¹, S. Bressler ¹⁶⁹, D. Britton ⁵⁹, D. Britzger ¹¹⁰,
I. Brock ²⁴, G. Brooijmans ⁴¹, W.K. Brooks ^{137f}, E. Brost ²⁹, L.M. Brown ¹⁶⁵, L.E. Bruce ⁶¹,
T.L. Bruckler ¹²⁶, P.A. Bruckman de Renstrom ⁸⁷, B. Brüers ⁴⁸, A. Bruni ^{23b}, G. Bruni ^{23b},
M. Bruschi ^{23b}, N. Bruscinò ^{75a,75b}, T. Buanes ¹⁶, Q. Buat ¹³⁸, D. Buchin ¹¹⁰, A.G. Buckley ⁵⁹,
O. Bulekov ³⁷, B.A. Bullard ¹⁴³, S. Burdin ⁹², C.D. Burgard ⁴⁹, A.M. Burger ⁴⁰,
B. Burghgrave ⁸, O. Burlayenko ⁵⁴, J.T.P. Burr ³², C.D. Burton ¹¹, J.C. Burzynski ¹⁴²,
E.L. Busch ⁴¹, V. Büscher ¹⁰⁰, P.J. Bussey ⁵⁹, J.M. Butler ²⁵, C.M. Buttar ⁵⁹,
J.M. Butterworth ⁹⁶, W. Buttinger ¹³⁴, C.J. Buxo Vazquez ¹⁰⁷, A.R. Buzykaev ³⁷,
S. Cabrera Urbán ¹⁶³, L. Cadamuro ⁶⁶, D. Caforio ⁵⁸, H. Cai ¹²⁹, Y. Cai ^{14a,14e}, Y. Cai ^{14c},
V.M.M. Cairo ³⁶, O. Cakir ^{3a}, N. Calace ³⁶, P. Calafiura ^{17a}, G. Calderini ¹²⁷, P. Calfayan ⁶⁸,
G. Callea ⁵⁹, L.P. Caloba ^{83b}, D. Calvet ⁴⁰, S. Calvet ⁴⁰, T.P. Calvet ¹⁰², M. Calvetti ^{74a,74b},
R. Camacho Toro ¹²⁷, S. Camarda ³⁶, D. Camarero Munoz ²⁶, P. Camarri ^{76a,76b},
M.T. Camerlingo ^{72a,72b}, D. Cameron ³⁶, C. Camincher ¹⁶⁵, M. Campanelli ⁹⁶, A. Camplani ⁴²,
V. Canale ^{72a,72b}, A. Canesse ¹⁰⁴, J. Cantero ¹⁶³, Y. Cao ¹⁶², F. Capocasa ²⁶, M. Capua ^{43b,43a},
A. Carbone ^{71a,71b}, R. Cardarelli ^{76a}, J.C.J. Cardenas ⁸, F. Cardillo ¹⁶³, G. Carducci ^{43b,43a},
T. Carli ³⁶, G. Carlino ^{72a}, J.I. Carlotto ¹³, B.T. Carlson ^{129,q}, E.M. Carlson ^{165,156a},
L. Carminati ^{71a,71b}, A. Carnelli ¹³⁵, M. Carnesale ^{75a,75b}, S. Caron ¹¹³, E. Carquin ^{137f},
S. Carrá ^{71a}, G. Carratta ^{23b,23a}, F. Carrio Argos ^{33g}, J.W.S. Carter ¹⁵⁵, T.M. Carter ⁵²,
M.P. Casado ^{13,i}, M. Caspar ⁴⁸, F.L. Castillo ⁴, L. Castillo Garcia ¹³, V. Castillo Gimenez ¹⁶³,
N.F. Castro ^{130a,130e}, A. Catinaccio ³⁶, J.R. Catmore ¹²⁵, V. Cavaliere ²⁹, N. Cavalli ^{23b,23a},
V. Cavasinni ^{74a,74b}, Y.C. Cekmecelioglu ⁴⁸, E. Celebi ^{21a}, F. Celli ¹²⁶, M.S. Centonze ^{70a,70b},
V. Cepaitis ⁵⁶, K. Cerny ¹²², A.S. Cerqueira ^{83a}, A. Cerri ¹⁴⁶, L. Cerrito ^{76a,76b}, F. Cerutti ^{17a},
B. Cervato ¹⁴¹, A. Cervelli ^{23b}, G. Cesarini ⁵³, S.A. Cetin ⁸², D. Chakraborty ¹¹⁵, J. Chan ¹⁷⁰,
W.Y. Chan ¹⁵³, J.D. Chapman ³², E. Chapon ¹³⁵, B. Chargeishvili ^{149b}, D.G. Charlton ²⁰,
M. Chatterjee ¹⁹, C. Chauhan ¹³³, S. Chekanov ⁶, S.V. Chekulaev ^{156a}, G.A. Chelkov ^{38,a},
A. Chen ¹⁰⁶, B. Chen ¹⁵¹, B. Chen ¹⁶⁵, H. Chen ^{14c}, H. Chen ²⁹, J. Chen ^{62c}, J. Chen ¹⁴²,
M. Chen ¹²⁶, S. Chen ¹⁵³, S.J. Chen ^{14c}, X. Chen ^{62c,135}, X. Chen ^{14b,ae}, Y. Chen ^{62a},
C.L. Cheng ¹⁷⁰, H.C. Cheng ^{64a}, S. Cheong ¹⁴³, A. Cheplakov ³⁸, E. Cheremushkina ⁴⁸,
E. Cherepanova ¹¹⁴, R. Cherkaoui El Moursli ^{35e}, E. Cheu ⁷, K. Cheung ⁶⁵, L. Chevalier ¹³⁵,
V. Chiarella ⁵³, G. Chiarelli ^{74a}, N. Chiedde ¹⁰², G. Chiodini ^{70a}, A.S. Chisholm ²⁰,
A. Chitan ^{27b}, M. Chitishvili ¹⁶³, M.V. Chizhov ³⁸, K. Choi ¹¹, A.R. Chomont ^{75a,75b},

Y. Chou [id¹⁰³](#), E.Y.S. Chow [id¹¹³](#), T. Chowdhury [id^{33g}](#), K.L. Chu [id¹⁶⁹](#), M.C. Chu [id^{64a}](#), X. Chu [id^{14a,14e}](#),
 J. Chudoba [id¹³¹](#), J.J. Chwastowski [id⁸⁷](#), D. Cieri [id¹¹⁰](#), K.M. Ciesla [id^{86a}](#), V. Cindro [id⁹³](#), A. Ciocio [id^{17a}](#),
 F. Cirotto [id^{72a,72b}](#), Z.H. Citron [id^{169,k}](#), M. Citterio [id^{71a}](#), D.A. Ciubotaru^{27b}, A. Clark [id⁵⁶](#), P.J. Clark [id⁵²](#),
 C. Clarry [id¹⁵⁵](#), J.M. Clavijo Columbie [id⁴⁸](#), S.E. Clawson [id⁴⁸](#), C. Clement [id^{47a,47b}](#), J. Clercx [id⁴⁸](#),
 Y. Coadou [id¹⁰²](#), M. Cobal [id^{69a,69c}](#), A. Coccaro [id^{57b}](#), R.F. Coelho Barrue [id^{130a}](#),
 R. Coelho Lopes De Sa [id¹⁰³](#), S. Coelli [id^{71a}](#), A.E.C. Coimbra [id^{71a,71b}](#), B. Cole [id⁴¹](#), J. Collot [id⁶⁰](#),
 P. Conde Muiño [id^{130a,130g}](#), M.P. Connell [id^{33c}](#), S.H. Connell [id^{33c}](#), I.A. Connelly [id⁵⁹](#), E.I. Conroy [id¹²⁶](#),
 F. Conventi [id^{72a,ag}](#), H.G. Cooke [id²⁰](#), A.M. Cooper-Sarkar [id¹²⁶](#), A. Cordeiro Oudot Choi [id¹²⁷](#),
 L.D. Corpe [id⁴⁰](#), M. Corradi [id^{75a,75b}](#), F. Corriveau [id^{104,w}](#), A. Cortes-Gonzalez [id¹⁸](#), M.J. Costa [id¹⁶³](#),
 F. Costanza [id⁴](#), D. Costanzo [id¹³⁹](#), B.M. Cote [id¹¹⁹](#), G. Cowan [id⁹⁵](#), K. Cranmer [id¹⁷⁰](#),
 D. Cremonini [id^{23b,23a}](#), S. Crépe-Renaudin [id⁶⁰](#), F. Crescioli [id¹²⁷](#), M. Cristinziani [id¹⁴¹](#),
 M. Cristoforetti [id^{78a,78b}](#), V. Croft [id¹¹⁴](#), J.E. Crosby [id¹²¹](#), G. Crosetti [id^{43b,43a}](#), A. Cueto [id⁹⁹](#),
 T. Cuhadar Donszelmann [id¹⁶⁰](#), H. Cui [id^{14a,14e}](#), Z. Cui [id⁷](#), W.R. Cunningham [id⁵⁹](#), F. Curcio [id^{43b,43a}](#),
 P. Czodrowski [id³⁶](#), M.M. Czurylo [id^{63b}](#), M.J. Da Cunha Sargedas De Sousa [id^{57b,57a}](#),
 J.V. Da Fonseca Pinto [id^{83b}](#), C. Da Via [id¹⁰¹](#), W. Dabrowski [id^{86a}](#), T. Dado [id⁴⁹](#), S. Dahbi [id^{33g}](#),
 T. Dai [id¹⁰⁶](#), D. Dal Santo [id¹⁹](#), C. Dallapiccola [id¹⁰³](#), M. Dam [id⁴²](#), G. D'amen [id²⁹](#), V. D'Amico [id¹⁰⁹](#),
 J. Damp [id¹⁰⁰](#), J.R. Dandoy [id³⁴](#), M.F. Daneri [id³⁰](#), M. Danninger [id¹⁴²](#), V. Dao [id³⁶](#), G. Darbo [id^{57b}](#),
 S. Darmora [id⁶](#), S.J. Das [id^{29,ai}](#), S. D'Auria [id^{71a,71b}](#), C. David [id^{156b}](#), T. Davidek [id¹³³](#),
 B. Davis-Purcell [id³⁴](#), I. Dawson [id⁹⁴](#), H.A. Day-hall [id¹³²](#), K. De [id⁸](#), R. De Asmundis [id^{72a}](#),
 N. De Biase [id⁴⁸](#), S. De Castro [id^{23b,23a}](#), N. De Groot [id¹¹³](#), P. de Jong [id¹¹⁴](#), H. De la Torre [id¹¹⁵](#),
 A. De Maria [id^{14c}](#), A. De Salvo [id^{75a}](#), U. De Sanctis [id^{76a,76b}](#), F. De Santis [id^{70a,70b}](#), A. De Santo [id¹⁴⁶](#),
 J.B. De Vivie De Regie [id⁶⁰](#), D.V. Dedovich³⁸, J. Degens [id¹¹⁴](#), A.M. Deiana [id⁴⁴](#), F. Del Corso [id^{23b,23a}](#),
 J. Del Peso [id⁹⁹](#), F. Del Rio [id^{63a}](#), L. Delagrangé [id¹²⁷](#), F. Deliot [id¹³⁵](#), C.M. Delitzsch [id⁴⁹](#),
 M. Della Pietra [id^{72a,72b}](#), D. Della Volpe [id⁵⁶](#), A. Dell'Acqua [id³⁶](#), L. Dell'Asta [id^{71a,71b}](#), M. Delmastro [id⁴](#),
 P.A. Delsart [id⁶⁰](#), S. Demers [id¹⁷²](#), M. Demichev [id³⁸](#), S.P. Denisov [id³⁷](#), L. D'Eramo [id⁴⁰](#),
 D. Derendarz [id⁸⁷](#), F. Derue [id¹²⁷](#), P. Dervan [id⁹²](#), K. Desch [id²⁴](#), C. Deutsch [id²⁴](#), F.A. Di Bello [id^{57b,57a}](#),
 A. Di Ciaccio [id^{76a,76b}](#), L. Di Ciaccio [id⁴](#), A. Di Domenico [id^{75a,75b}](#), C. Di Donato [id^{72a,72b}](#),
 A. Di Girolamo [id³⁶](#), G. Di Gregorio [id³⁶](#), A. Di Luca [id^{78a,78b}](#), B. Di Micco [id^{77a,77b}](#), R. Di Nardo [id^{77a,77b}](#),
 C. Diaconu [id¹⁰²](#), M. Diamantopoulou [id³⁴](#), F.A. Dias [id¹¹⁴](#), T. Dias Do Vale [id¹⁴²](#), M.A. Diaz [id^{137a,137b}](#),
 F.G. Diaz Capriles [id²⁴](#), M. Didenko [id¹⁶³](#), E.B. Diehl [id¹⁰⁶](#), L. Diehl [id⁵⁴](#), S. Díez Cornell [id⁴⁸](#),
 C. Díez Pardos [id¹⁴¹](#), C. Dimitriadi [id^{161,24}](#), A. Dimitrievska [id^{17a}](#), J. Dingfelder [id²⁴](#), I-M. Dinu [id^{27b}](#),
 S.J. Dittmeier [id^{63b}](#), F. Dittus [id³⁶](#), F. Djama [id¹⁰²](#), T. Djobava [id^{149b}](#), J.I. Djuvslund [id¹⁶](#),
 C. Doglioni [id^{101,98}](#), A. Dohnalova [id^{28a}](#), J. Dolejsi [id¹³³](#), Z. Dolezal [id¹³³](#), K.M. Dona [id³⁹](#),
 M. Donadelli [id^{83c}](#), B. Dong [id¹⁰⁷](#), J. Donini [id⁴⁰](#), A. D'Onofrio [id^{72a,72b}](#), M. D'Onofrio [id⁹²](#),
 J. Dopke [id¹³⁴](#), A. Doria [id^{72a}](#), N. Dos Santos Fernandes [id^{130a}](#), P. Dougan [id¹⁰¹](#), M.T. Dova [id⁹⁰](#),
 A.T. Doyle [id⁵⁹](#), M.A. Draguet [id¹²⁶](#), E. Dreyer [id¹⁶⁹](#), I. Drivas-koulouris [id¹⁰](#), M. Drnevich [id¹¹⁷](#),
 A.S. Drobac [id¹⁵⁸](#), M. Drozdova [id⁵⁶](#), D. Du [id^{62a}](#), T.A. du Pree [id¹¹⁴](#), F. Dubinin [id³⁷](#), M. Dubovsky [id^{28a}](#),
 E. Duchovni [id¹⁶⁹](#), G. Duckeck [id¹⁰⁹](#), O.A. Ducu [id^{27b}](#), D. Duda [id⁵²](#), A. Dudarev [id³⁶](#), E.R. Duden [id²⁶](#),
 M. D'uffizi [id¹⁰¹](#), L. Dufлот [id⁶⁶](#), M. Dührssen [id³⁶](#), C. Dülsen [id¹⁷¹](#), A.E. Dumitriu [id^{27b}](#), M. Dunford [id^{63a}](#),
 S. Dungs [id⁴⁹](#), K. Dunne [id^{47a,47b}](#), A. Duperrin [id¹⁰²](#), H. Duran Yildiz [id^{3a}](#), M. Düren [id⁵⁸](#),
 A. Durglishvili [id^{149b}](#), B.L. Dwyer [id¹¹⁵](#), G.I. Dyckes [id^{17a}](#), M. Dyndal [id^{86a}](#), B.S. Dziedzic [id⁸⁷](#),
 Z.O. Earnshaw [id¹⁴⁶](#), G.H. Eberwein [id¹²⁶](#), B. Eckerova [id^{28a}](#), S. Eggebrecht [id⁵⁵](#),
 E. Egidio Purcino De Souza [id¹²⁷](#), L.F. Ehrke [id⁵⁶](#), G. Eigen [id¹⁶](#), K. Einsweiler [id^{17a}](#), T. Ekelof [id¹⁶¹](#),
 P.A. Ekman [id⁹⁸](#), S. El Farkh [id^{35b}](#), Y. El Ghazali [id^{35b}](#), H. El Jarrari [id³⁶](#), A. El Moussaouy [id¹⁰⁸](#),
 V. Ellajosyula [id¹⁶¹](#), M. Ellert [id¹⁶¹](#), F. Ellinghaus [id¹⁷¹](#), N. Ellis [id³⁶](#), J. Elmsheuser [id²⁹](#), M. Elsing [id³⁶](#),
 D. Emelianov [id¹³⁴](#), Y. Enari [id¹⁵³](#), I. Ene [id^{17a}](#), S. Epari [id¹³](#), J. Erdmann [id⁴⁹](#), P.A. Erland [id⁸⁷](#),
 M. Errenst [id¹⁷¹](#), M. Escalier [id⁶⁶](#), C. Escobar [id¹⁶³](#), E. Etzion [id¹⁵¹](#), G. Evans [id^{130a}](#), H. Evans [id⁶⁸](#),

L.S. Evans [id](#)⁹⁵, M.O. Evans [id](#)¹⁴⁶, A. Ezhilov [id](#)³⁷, S. Ezzarqtouni [id](#)^{35a}, F. Fabbri [id](#)⁵⁹, L. Fabbri [id](#)^{23b,23a},
 G. Facini [id](#)⁹⁶, V. Fadeyev [id](#)¹³⁶, R.M. Fakhrutdinov [id](#)³⁷, D. Fakoudis [id](#)¹⁰⁰, S. Falciano [id](#)^{75a},
 L.F. Falda Ulhoa Coelho [id](#)³⁶, P.J. Falke [id](#)²⁴, J. Faltova [id](#)¹³³, C. Fan [id](#)¹⁶², Y. Fan [id](#)^{14a}, Y. Fang [id](#)^{14a,14e},
 M. Fanti [id](#)^{71a,71b}, M. Faraj [id](#)^{69a,69b}, Z. Farazpay [id](#)⁹⁷, A. Farbin [id](#)⁸, A. Farilla [id](#)^{77a}, T. Farooque [id](#)¹⁰⁷,
 S.M. Farrington [id](#)⁵², F. Fassi [id](#)^{35e}, D. Fassouliotis [id](#)⁹, M. Faucci Giannelli [id](#)^{76a,76b}, W.J. Fawcett [id](#)³²,
 L. Fayard [id](#)⁶⁶, P. Federic [id](#)¹³³, P. Federicova [id](#)¹³¹, O.L. Fedin [id](#)^{37,a}, G. Fedotov [id](#)³⁷, M. Feickert [id](#)¹⁷⁰,
 L. Feligioni [id](#)¹⁰², D.E. Fellers [id](#)¹²³, C. Feng [id](#)^{62b}, M. Feng [id](#)^{14b}, Z. Feng [id](#)¹¹⁴, M.J. Fenton [id](#)¹⁶⁰,
 A.B. Fenyuk³⁷, L. Ferencz [id](#)⁴⁸, R.A.M. Ferguson [id](#)⁹¹, S.I. Fernandez Luengo [id](#)^{137f},
 P. Fernandez Martinez [id](#)¹³, M.J.V. Fernoux [id](#)¹⁰², J. Ferrando [id](#)⁴⁸, A. Ferrari [id](#)¹⁶¹, P. Ferrari [id](#)^{114,113},
 R. Ferrari [id](#)^{73a}, D. Ferrere [id](#)⁵⁶, C. Ferretti [id](#)¹⁰⁶, F. Fiedler [id](#)¹⁰⁰, P. Fiedler [id](#)¹³², A. Filipčič [id](#)⁹³,
 E.K. Filmer [id](#)¹, F. Filthaut [id](#)¹¹³, M.C.N. Fiolhais [id](#)^{130a,130c,c}, L. Fiorini [id](#)¹⁶³, W.C. Fisher [id](#)¹⁰⁷,
 T. Fitschen [id](#)¹⁰¹, P.M. Fitzhugh¹³⁵, I. Fleck [id](#)¹⁴¹, P. Fleischmann [id](#)¹⁰⁶, T. Flick [id](#)¹⁷¹, M. Flores [id](#)^{33d,ac},
 L.R. Flores Castillo [id](#)^{64a}, L. Flores Sanz De Acedo [id](#)³⁶, F.M. Follega [id](#)^{78a,78b}, N. Fomin [id](#)¹⁶,
 J.H. Foo [id](#)¹⁵⁵, B.C. Forland⁶⁸, A. Formica [id](#)¹³⁵, A.C. Forti [id](#)¹⁰¹, E. Fortin [id](#)³⁶, A.W. Fortman [id](#)⁶¹,
 M.G. Foti [id](#)^{17a}, L. Fountas [id](#)^{9,j}, D. Fournier [id](#)⁶⁶, H. Fox [id](#)⁹¹, P. Francavilla [id](#)^{74a,74b}, S. Francescato [id](#)⁶¹,
 S. Franchellucci [id](#)⁵⁶, M. Franchini [id](#)^{23b,23a}, S. Franchino [id](#)^{63a}, D. Francis³⁶, L. Franco [id](#)¹¹³,
 V. Franco Lima [id](#)³⁶, L. Franconi [id](#)⁴⁸, M. Franklin [id](#)⁶¹, G. Frattari [id](#)²⁶, A.C. Freegard [id](#)⁹⁴,
 W.S. Freund [id](#)^{83b}, Y.Y. Frid [id](#)¹⁵¹, J. Friend [id](#)⁵⁹, N. Fritzsche [id](#)⁵⁰, A. Froch [id](#)⁵⁴, D. Froidevaux [id](#)³⁶,
 J.A. Frost [id](#)¹²⁶, Y. Fu [id](#)^{62a}, S. Fuenzalida Garrido [id](#)^{137f}, M. Fujimoto [id](#)¹⁰², K.Y. Fung [id](#)^{64a},
 E. Furtado De Simas Filho [id](#)^{83b}, M. Furukawa [id](#)¹⁵³, J. Fuster [id](#)¹⁶³, A. Gabrielli [id](#)^{23b,23a},
 A. Gabrielli [id](#)¹⁵⁵, P. Gadow [id](#)³⁶, G. Gagliardi [id](#)^{57b,57a}, L.G. Gagnon [id](#)^{17a}, E.J. Gallas [id](#)¹²⁶,
 B.J. Gallop [id](#)¹³⁴, K.K. Gan [id](#)¹¹⁹, S. Ganguly [id](#)¹⁵³, Y. Gao [id](#)⁵², F.M. Garay Walls [id](#)^{137a,137b}, B. Garcia²⁹,
 C. García [id](#)¹⁶³, A. Garcia Alonso [id](#)¹¹⁴, A.G. Garcia Caffaro [id](#)¹⁷², J.E. García Navarro [id](#)¹⁶³,
 M. Garcia-Sciveres [id](#)^{17a}, G.L. Gardner [id](#)¹²⁸, R.W. Gardner [id](#)³⁹, N. Garelli [id](#)¹⁵⁸, D. Garg [id](#)⁸⁰,
 R.B. Garg [id](#)^{143,n}, J.M. Gargan⁵², C.A. Garner¹⁵⁵, C.M. Garvey [id](#)^{33a}, P. Gaspar [id](#)^{83b}, V.K. Gassmann¹⁵⁸,
 G. Gaudio [id](#)^{73a}, V. Gautam¹³, P. Gauzzi [id](#)^{75a,75b}, I.L. Gavrilenko [id](#)³⁷, A. Gavrilyuk [id](#)³⁷, C. Gay [id](#)¹⁶⁴,
 G. Gaycken [id](#)⁴⁸, E.N. Gazis [id](#)¹⁰, A.A. Geanta [id](#)^{27b}, C.M. Gee [id](#)¹³⁶, A. Gekow¹¹⁹, C. Gemme [id](#)^{57b},
 M.H. Genest [id](#)⁶⁰, S. Gentile [id](#)^{75a,75b}, A.D. Gentry [id](#)¹¹², S. George [id](#)⁹⁵, W.F. George [id](#)²⁰, T. Geralis [id](#)⁴⁶,
 P. Gessinger-Befurt [id](#)³⁶, M.E. Geyik [id](#)¹⁷¹, M. Ghani [id](#)¹⁶⁷, M. Ghneimat [id](#)¹⁴¹, K. Ghorbanian [id](#)⁹⁴,
 A. Ghosal [id](#)¹⁴¹, A. Ghosh [id](#)¹⁶⁰, A. Ghosh [id](#)⁷, B. Giacobbe [id](#)^{23b}, S. Giagu [id](#)^{75a,75b}, T. Giani [id](#)¹¹⁴,
 P. Giannetti [id](#)^{74a}, A. Giannini [id](#)^{62a}, S.M. Gibson [id](#)⁹⁵, M. Gignac [id](#)¹³⁶, D.T. Gil [id](#)^{86b}, A.K. Gilbert [id](#)^{86a},
 B.J. Gilbert [id](#)⁴¹, D. Gillberg [id](#)³⁴, G. Gilles [id](#)¹¹⁴, N.E.K. Gillwald [id](#)⁴⁸, L. Ginabat [id](#)¹²⁷,
 D.M. Gingrich [id](#)^{2,af}, M.P. Giordani [id](#)^{69a,69c}, P.F. Giraud [id](#)¹³⁵, G. Giugliarelli [id](#)^{69a,69c}, D. Giugni [id](#)^{71a},
 F. Giuli [id](#)³⁶, I. Gkialas [id](#)^{9,j}, L.K. Gladilin [id](#)³⁷, C. Glasman [id](#)⁹⁹, G.R. Gledhill [id](#)¹²³, G. Glemža [id](#)⁴⁸,
 M. Glisic¹²³, I. Gnesi [id](#)^{43b,f}, Y. Go [id](#)^{29,ai}, M. Goblirsch-Kolb [id](#)³⁶, B. Gocke [id](#)⁴⁹, D. Godin¹⁰⁸,
 B. Gokturk [id](#)^{21a}, S. Goldfarb [id](#)¹⁰⁵, T. Golling [id](#)⁵⁶, M.G.D. Gololo^{33g}, D. Golubkov [id](#)³⁷,
 J.P. Gombas [id](#)¹⁰⁷, A. Gomes [id](#)^{130a,130b}, G. Gomes Da Silva [id](#)¹⁴¹, A.J. Gomez Delegido [id](#)¹⁶³,
 R. Gonçalves [id](#)^{130a,130c}, G. Gonella [id](#)¹²³, L. Gonella [id](#)²⁰, A. Gongadze [id](#)^{149c}, F. Gonnella [id](#)²⁰,
 J.L. Gonski [id](#)⁴¹, R.Y. González Andana [id](#)⁵², S. González de la Hoz [id](#)¹⁶³, S. Gonzalez Fernandez [id](#)¹³,
 R. Gonzalez Lopez [id](#)⁹², C. Gonzalez Renteria [id](#)^{17a}, M.V. Gonzalez Rodrigues [id](#)⁴⁸,
 R. Gonzalez Suarez [id](#)¹⁶¹, S. Gonzalez-Sevilla [id](#)⁵⁶, G.R. Gonzalvo Rodriguez [id](#)¹⁶³, L. Goossens [id](#)³⁶,
 B. Gorini [id](#)³⁶, E. Gorini [id](#)^{70a,70b}, A. Gorišek [id](#)⁹³, T.C. Gosart [id](#)¹²⁸, A.T. Goshaw [id](#)⁵¹, M.I. Gostkin [id](#)³⁸,
 S. Goswami [id](#)¹²¹, C.A. Gottardo [id](#)³⁶, S.A. Gotz [id](#)¹⁰⁹, M. Gouighri [id](#)^{35b}, V. Goumarre [id](#)⁴⁸,
 A.G. Goussiou [id](#)¹³⁸, N. Govender [id](#)^{33c}, I. Grabowska-Bold [id](#)^{86a}, K. Graham [id](#)³⁴, E. Gramstad [id](#)¹²⁵,
 S. Grancagnolo [id](#)^{70a,70b}, M. Grandi [id](#)¹⁴⁶, C.M. Grant^{1,135}, P.M. Gravila [id](#)^{27f}, F.G. Gravili [id](#)^{70a,70b},
 H.M. Gray [id](#)^{17a}, M. Greco [id](#)^{70a,70b}, C. Grefe [id](#)²⁴, I.M. Gregor [id](#)⁴⁸, P. Grenier [id](#)¹⁴³, S.G. Grewe¹¹⁰,
 C. Grieco [id](#)¹³, A.A. Grillo [id](#)¹³⁶, K. Grimm [id](#)³¹, S. Grinstein [id](#)^{13,s}, J.-F. Grivaz [id](#)⁶⁶, E. Gross [id](#)¹⁶⁹,

J. Grosse-Knetter ⁵⁵, C. Grud ¹⁰⁶, J.C. Grundy ¹²⁶, L. Guan ¹⁰⁶, W. Guan ²⁹, C. Gubbels ¹⁶⁴, J.G.R. Guerrero Rojas ¹⁶³, G. Guerrieri ^{69a,69c}, F. Guescini ¹¹⁰, R. Gugel ¹⁰⁰, J.A.M. Guhit ¹⁰⁶, A. Guida ¹⁸, E. Guilloton ^{167,134}, S. Guindon ³⁶, F. Guo ^{14a,14e}, J. Guo ^{62c}, L. Guo ⁴⁸, Y. Guo ¹⁰⁶, R. Gupta ⁴⁸, R. Gupta ¹²⁹, S. Gurbuz ²⁴, S.S. Gurdasani ⁵⁴, G. Gustavino ³⁶, M. Guth ⁵⁶, P. Gutierrez ¹²⁰, L.F. Gutierrez Zagazeta ¹²⁸, M. Gutsche ⁵⁰, C. Gutschow ⁹⁶, C. Gwenlan ¹²⁶, C.B. Gwilliam ⁹², E.S. Haaland ¹²⁵, A. Haas ¹¹⁷, M. Habedank ⁴⁸, C. Haber ^{17a}, H.K. Hadavand ⁸, A. Hadeef ⁵⁰, S. Hadzic ¹¹⁰, A.I. Hagan ⁹¹, J.J. Hahn ¹⁴¹, E.H. Haines ⁹⁶, M. Haleem ¹⁶⁶, J. Haley ¹²¹, J.J. Hall ¹³⁹, G.D. Hallewell ¹⁰², L. Halser ¹⁹, K. Hamano ¹⁶⁵, M. Hamer ²⁴, G.N. Hamity ⁵², E.J. Hampshire ⁹⁵, J. Han ^{62b}, K. Han ^{62a}, L. Han ^{14c}, L. Han ^{62a}, S. Han ^{17a}, Y.F. Han ¹⁵⁵, K. Hanagaki ⁸⁴, M. Hance ¹³⁶, D.A. Hangal ^{41,ab}, H. Hanif ¹⁴², M.D. Hank ¹²⁸, R. Hankache ¹⁰¹, J.B. Hansen ⁴², J.D. Hansen ⁴², P.H. Hansen ⁴², K. Hara ¹⁵⁷, D. Harada ⁵⁶, T. Harenberg ¹⁷¹, S. Harkusha ³⁷, M.L. Harris ¹⁰³, Y.T. Harris ¹²⁶, J. Harrison ¹³, N.M. Harrison ¹¹⁹, P.F. Harrison ¹⁶⁷, N.M. Hartman ¹¹⁰, N.M. Hartmann ¹⁰⁹, Y. Hasegawa ¹⁴⁰, R. Hauser ¹⁰⁷, C.M. Hawkes ²⁰, R.J. Hawkings ³⁶, Y. Hayashi ¹⁵³, S. Hayashida ¹¹¹, D. Hayden ¹⁰⁷, C. Hayes ¹⁰⁶, R.L. Hayes ¹¹⁴, C.P. Hays ¹²⁶, J.M. Hays ⁹⁴, H.S. Hayward ⁹², F. He ^{62a}, M. He ^{14a,14e}, Y. He ¹⁵⁴, Y. He ⁴⁸, N.B. Heatley ⁹⁴, V. Hedberg ⁹⁸, A.L. Heggelund ¹²⁵, N.D. Hehir ^{94,*}, C. Heidegger ⁵⁴, K.K. Heidegger ⁵⁴, W.D. Heidorn ⁸¹, J. Heilman ³⁴, S. Heim ⁴⁸, T. Heim ^{17a}, J.G. Heinlein ¹²⁸, J.J. Heinrich ¹²³, L. Heinrich ^{110,ad}, J. Hejbal ¹³¹, L. Helary ⁴⁸, A. Held ¹⁷⁰, S. Hellesund ¹⁶, C.M. Helling ¹⁶⁴, S. Hellman ^{47a,47b}, R.C.W. Henderson ⁹¹, L. Henkelmann ³², A.M. Henriques Correia ³⁶, H. Herde ⁹⁸, Y. Hernández Jiménez ¹⁴⁵, L.M. Herrmann ²⁴, T. Herrmann ⁵⁰, G. Herten ⁵⁴, R. Hertenberger ¹⁰⁹, L. Hervas ³⁶, M.E. Hesping ¹⁰⁰, N.P. Hessey ^{156a}, H. Hibi ⁸⁵, E. Hill ¹⁵⁵, S.J. Hillier ²⁰, J.R. Hinds ¹⁰⁷, F. Hinterkeuser ²⁴, M. Hirose ¹²⁴, S. Hirose ¹⁵⁷, D. Hirschbuehl ¹⁷¹, T.G. Hitchings ¹⁰¹, B. Hiti ⁹³, J. Hobbs ¹⁴⁵, R. Hobincu ^{27e}, N. Hod ¹⁶⁹, M.C. Hodgkinson ¹³⁹, B.H. Hodgkinson ³², A. Hoecker ³⁶, D.D. Hofer ¹⁰⁶, J. Hofer ⁴⁸, T. Holm ²⁴, M. Holzbock ¹¹⁰, L.B.A.H. Hommels ³², B.P. Honan ¹⁰¹, J. Hong ^{62c}, T.M. Hong ¹²⁹, B.H. Hooberman ¹⁶², W.H. Hopkins ⁶, Y. Horii ¹¹¹, S. Hou ¹⁴⁸, A.S. Howard ⁹³, J. Howarth ⁵⁹, J. Hoya ⁶, M. Hrabovsky ¹²², A. Hrynevich ⁴⁸, T. Hryn'ova ⁴, P.J. Hsu ⁶⁵, S.-C. Hsu ¹³⁸, Q. Hu ^{62a}, Y.F. Hu ^{14a,14e}, S. Huang ^{64b}, X. Huang ^{14c}, X. Huang ^{14a,14e}, Y. Huang ¹³⁹, Y. Huang ^{14a}, Z. Huang ¹⁰¹, Z. Hubacek ¹³², M. Huebner ²⁴, F. Huegging ²⁴, T.B. Huffman ¹²⁶, C.A. Hugli ⁴⁸, M. Huhtinen ³⁶, S.K. Huiberts ¹⁶, R. Hulsken ¹⁰⁴, N. Huseynov ¹², J. Huston ¹⁰⁷, J. Huth ⁶¹, R. Hyneman ¹⁴³, G. Iacobucci ⁵⁶, G. Iakovidis ²⁹, I. Ibragimov ¹⁴¹, L. Iconomidou-Fayard ⁶⁶, P. Iengo ^{72a,72b}, R. Iguchi ¹⁵³, T. Iizawa ¹²⁶, Y. Ikegami ⁸⁴, N. Ilic ¹⁵⁵, H. Imam ^{35a}, M. Ince Lezki ⁵⁶, T. Ingebretsen Carlson ^{47a,47b}, G. Introzzi ^{73a,73b}, M. Iodice ^{77a}, V. Ippolito ^{75a,75b}, R.K. Irwin ⁹², M. Ishino ¹⁵³, W. Islam ¹⁷⁰, C. Issever ^{18,48}, S. Istin ^{21a,al}, H. Ito ¹⁶⁸, J.M. Iturbe Ponce ^{64a}, R. Iuppa ^{78a,78b}, A. Ivina ¹⁶⁹, J.M. Izen ⁴⁵, V. Izzo ^{72a}, P. Jacka ^{131,132}, P. Jackson ¹, R.M. Jacobs ⁴⁸, B.P. Jaeger ¹⁴², C.S. Jagfeld ¹⁰⁹, G. Jain ^{156a}, P. Jain ⁵⁴, K. Jakobs ⁵⁴, T. Jakoubek ¹⁶⁹, J. Jamieson ⁵⁹, K.W. Janas ^{86a}, M. Javurkova ¹⁰³, F. Jeanneau ¹³⁵, L. Jeanty ¹²³, J. Jejelava ^{149a,z}, P. Jenni ^{54,g}, C.E. Jessiman ³⁴, S. Jézéquel ⁴, C. Jia ^{62b}, J. Jia ¹⁴⁵, X. Jia ⁶¹, X. Jia ^{14a,14e}, Z. Jia ^{14c}, S. Jiggins ⁴⁸, J. Jimenez Pena ¹³, S. Jin ^{14c}, A. Jinaru ^{27b}, O. Jinnouchi ¹⁵⁴, P. Johansson ¹³⁹, K.A. Johns ⁷, J.W. Johnson ¹³⁶, D.M. Jones ³², E. Jones ⁴⁸, P. Jones ³², R.W.L. Jones ⁹¹, T.J. Jones ⁹², H.L. Joos ^{55,36}, R. Joshi ¹¹⁹, J. Jovicevic ¹⁵, X. Ju ^{17a}, J.J. Junggeburth ¹⁰³, T. Junkermann ^{63a}, A. Juste Rozas ^{13,s}, M.K. Juzek ⁸⁷, S. Kabana ^{137e}, A. Kaczmarska ⁸⁷, M. Kado ¹¹⁰, H. Kagan ¹¹⁹, M. Kagan ¹⁴³, A. Kahn ⁴¹, A. Kahn ¹²⁸, C. Kahra ¹⁰⁰, T. Kaji ¹⁵³, E. Kajomovitz ¹⁵⁰, N. Kakati ¹⁶⁹, I. Kalaitzidou ⁵⁴, C.W. Kalderon ²⁹, A. Kamenshchikov ¹⁵⁵, N.J. Kang ¹³⁶, D. Kar ^{33g}, K. Karava ¹²⁶,

M.J. Kareem ^{156b}, E. Karentzos ⁵⁴, I. Karkanias ¹⁵², O. Karkout ¹¹⁴, S.N. Karpov ³⁸,
Z.M. Karpova ³⁸, V. Kartvelishvili ⁹¹, A.N. Karyukhin ³⁷, E. Kasimi ¹⁵², J. Katzy ⁴⁸,
S. Kaur ³⁴, K. Kawade ¹⁴⁰, M.P. Kawale ¹²⁰, C. Kawamoto ⁸⁸, T. Kawamoto ^{62a}, E.F. Kay ³⁶,
F.I. Kaya ¹⁵⁸, S. Kazakos ¹⁰⁷, V.F. Kazanin ³⁷, Y. Ke ¹⁴⁵, J.M. Keaveney ^{33a}, R. Keeler ¹⁶⁵,
G.V. Kehris ⁶¹, J.S. Keller ³⁴, A.S. Kelly ⁹⁶, J.J. Kempster ¹⁴⁶, K.E. Kennedy ⁴¹,
P.D. Kennedy ¹⁰⁰, O. Kepka ¹³¹, B.P. Kerridge ¹⁶⁷, S. Kersten ¹⁷¹, B.P. Kerševan ⁹³,
S. Keshri ⁶⁶, L. Keszeghova ^{28a}, S. Ketabchi Haghighat ¹⁵⁵, R.A. Khan ¹²⁹, M. Khandoga ¹²⁷,
A. Khanov ¹²¹, A.G. Kharlamov ³⁷, T. Kharlamova ³⁷, E.E. Khoda ¹³⁸, M. Kholodenko ³⁷,
T.J. Khoo ¹⁸, G. Khoraiuli ¹⁶⁶, J. Khubua ^{149b}, Y.A.R. Khwaira ⁶⁶, A. Kilgallon ¹²³,
D.W. Kim ^{47a,47b}, Y.K. Kim ³⁹, N. Kimura ⁹⁶, M.K. Kingston ⁵⁵, A. Kirchhoff ⁵⁵, C. Kirfel ²⁴,
F. Kirfel ²⁴, J. Kirk ¹³⁴, A.E. Kiryunin ¹¹⁰, C. Kitsaki ¹⁰, O. Kivernyk ²⁴, M. Klassen ^{63a},
C. Klein ³⁴, L. Klein ¹⁶⁶, M.H. Klein ¹⁰⁶, M. Klein ⁹², S.B. Klein ⁵⁶, U. Klein ⁹²,
P. Klimek ³⁶, A. Klimentov ²⁹, T. Klioutchnikova ³⁶, P. Kluit ¹¹⁴, S. Kluth ¹¹⁰, E. Kneringer ⁷⁹,
T.M. Knight ¹⁵⁵, A. Knue ⁴⁹, R. Kobayashi ⁸⁸, D. Kobylanski ¹⁶⁹, S.F. Koch ¹²⁶,
M. Kocian ¹⁴³, P. Kodyš ¹³³, D.M. Koeck ¹²³, P.T. Koenig ²⁴, T. Koffas ³⁴, O. Kolay ⁵⁰,
I. Koletsou ⁴, T. Komarek ¹²², K. Köneke ⁵⁴, A.X.Y. Kong ¹, T. Kono ¹¹⁸, N. Konstantinidis ⁹⁶,
P. Kontaxakis ⁵⁶, B. Konya ⁹⁸, R. Kopeliansky ⁶⁸, S. Koperny ^{86a}, K. Korcyl ⁸⁷, K. Kordas ^{152,e},
G. Koren ¹⁵¹, A. Korn ⁹⁶, S. Korn ⁵⁵, I. Korolkov ¹³, N. Korotkova ³⁷, B. Kortman ¹¹⁴,
O. Kortner ¹¹⁰, S. Kortner ¹¹⁰, W.H. Kostecka ¹¹⁵, V.V. Kostyukhin ¹⁴¹, A. Kotskechagia ¹³⁵,
A. Kotwal ⁵¹, A. Koulouris ³⁶, A. Kourkoumeli-Charalampidi ^{73a,73b}, C. Kourkoumelis ⁹,
E. Kourlitis ^{110,ad}, O. Kovanda ¹⁴⁶, R. Kowalewski ¹⁶⁵, W. Kozanecki ¹³⁵, A.S. Kozhin ³⁷,
V.A. Kramarenko ³⁷, G. Kramberger ⁹³, P. Kramer ¹⁰⁰, M.W. Krasny ¹²⁷, A. Krasznahorkay ³⁶,
J.W. Kraus ¹⁷¹, J.A. Kremer ⁴⁸, T. Kresse ⁵⁰, J. Kretschmar ⁹², K. Kreul ¹⁸, P. Krieger ¹⁵⁵,
S. Krishnamurthy ¹⁰³, M. Krivos ¹³³, K. Krizka ²⁰, K. Kroeninger ⁴⁹, H. Kroha ¹¹⁰, J. Kroll ¹³¹,
J. Kroll ¹²⁸, K.S. Krowpman ¹⁰⁷, U. Kruchonak ³⁸, H. Krüger ²⁴, N. Krumnack ⁸¹, M.C. Kruse ⁵¹,
O. Kuchinskaia ³⁷, S. Kuday ^{3a}, S. Kuehn ³⁶, R. Kuesters ⁵⁴, T. Kuhl ⁴⁸, V. Kukhtin ³⁸,
Y. Kulchitsky ^{37,a}, S. Kuleshov ^{137d,137b}, M. Kumar ^{33g}, N. Kumari ⁴⁸, P. Kumari ^{156b},
A. Kupco ¹³¹, T. Kupfer ⁴⁹, A. Kupich ³⁷, O. Kuprash ⁵⁴, H. Kurashige ⁸⁵, L.L. Kurchaninov ^{156a},
O. Kurdysh ⁶⁶, Y.A. Kurochkin ³⁷, A. Kurova ³⁷, M. Kuze ¹⁵⁴, A.K. Kvam ¹⁰³, J. Kvitá ¹²²,
T. Kwan ¹⁰⁴, N.G. Kyriacou ¹⁰⁶, L.A.O. Laatu ¹⁰², C. Lacasta ¹⁶³, F. Lacava ^{75a,75b},
H. Lacker ¹⁸, D. Lacour ¹²⁷, N.N. Lad ⁹⁶, E. Ladygin ³⁸, B. Laforge ¹²⁷, T. Lagouri ^{137e},
F.Z. Lahbabi ^{35a}, S. Lai ⁵⁵, I.K. Lakomic ^{86a}, N. Lalloue ⁶⁰, J.E. Lambert ¹⁶⁵, S. Lammers ⁶⁸,
W. Lampl ⁷, C. Lampoudis ^{152,e}, A.N. Lancaster ¹¹⁵, E. Lançon ²⁹, U. Landgraf ⁵⁴,
M.P.J. Landon ⁹⁴, V.S. Lang ⁵⁴, R.J. Langenberg ¹⁰³, O.K.B. Langrekken ¹²⁵, A.J. Lankford ¹⁶⁰,
F. Lanni ³⁶, K. Lantzs ²⁴, A. Lanza ^{73a}, A. Lapertosa ^{57b,57a}, J.F. Laporte ¹³⁵, T. Lari ^{71a},
F. Lasagni Manghi ^{23b}, M. Lassnig ³⁶, V. Latonova ¹³¹, A. Laudrain ¹⁰⁰, A. Laurier ¹⁵⁰,
S.D. Lawlor ¹³⁹, Z. Lawrence ¹⁰¹, R. Lazaridou ¹⁶⁷, M. Lazzaroni ^{71a,71b}, B. Le ¹⁰¹,
E.M. Le Boulicaut ⁵¹, B. Leban ⁹³, A. Lebedev ⁸¹, M. LeBlanc ¹⁰¹, F. Ledroit-Guillon ⁶⁰,
A.C.A. Lee ⁹⁶, S.C. Lee ¹⁴⁸, S. Lee ^{47a,47b}, T.F. Lee ⁹², L.L. Leeuw ^{33c}, H.P. Lefebvre ⁹⁵,
M. Lefebvre ¹⁶⁵, C. Leggett ^{17a}, G. Lehmann Miotto ³⁶, M. Leigh ⁵⁶, W.A. Leight ¹⁰³,
W. Leinonen ¹¹³, A. Leisos ^{152,r}, M.A.L. Leite ^{83c}, C.E. Leitgeb ⁴⁸, R. Leitner ¹³³,
K.J.C. Leney ⁴⁴, T. Lenz ²⁴, S. Leone ^{74a}, C. Leonidopoulos ⁵², A. Leopold ¹⁴⁴, C. Leroy ¹⁰⁸,
R. Les ¹⁰⁷, C.G. Lester ³², M. Levchenko ³⁷, J. Levêque ⁴, D. Levin ¹⁰⁶, L.J. Levinson ¹⁶⁹,
M.P. Lewicki ⁸⁷, D.J. Lewis ⁴, A. Li ⁵, B. Li ^{62b}, C. Li ^{62a}, C-Q. Li ¹¹⁰, H. Li ^{62a}, H. Li ^{62b},
H. Li ^{14c}, H. Li ^{14b}, H. Li ^{62b}, J. Li ^{62c}, K. Li ¹³⁸, L. Li ^{62c}, M. Li ^{14a,14e}, Q.Y. Li ^{62a},
S. Li ^{14a,14e}, S. Li ^{62d,62c,d}, T. Li ⁵, X. Li ¹⁰⁴, Z. Li ¹²⁶, Z. Li ¹⁰⁴, Z. Li ⁹², Z. Li ^{14a,14e},
S. Liang ^{14a,14e}, Z. Liang ^{14a}, M. Liberatore ¹³⁵, B. Liberti ^{76a}, K. Lie ^{64c}, J. Lieber Marin ^{83b},

H. Lien ⁶⁸, K. Lin ¹⁰⁷, R.E. Lindley ⁷, J.H. Lindon ², E. Lipeles ¹²⁸, A. Lipniacka ¹⁶,
 A. Lister ¹⁶⁴, J.D. Little ⁴, B. Liu ^{14a}, B.X. Liu ¹⁴², D. Liu ^{62d,62c}, J.B. Liu ^{62a}, J.K.K. Liu ³²,
 K. Liu ^{62d,62c}, M. Liu ^{62a}, M.Y. Liu ^{62a}, P. Liu ^{14a}, Q. Liu ^{62d,138,62c}, X. Liu ^{62a}, Y. Liu ^{14d,14e},
 Y.L. Liu ^{62b}, Y.W. Liu ^{62a}, J. Llorente Merino ¹⁴², S.L. Lloyd ⁹⁴, E.M. Lobodzinska ⁴⁸,
 P. Loch ⁷, T. Lohse ¹⁸, K. Lohwasser ¹³⁹, E. Loiacono ⁴⁸, M. Lokajicek ^{131,*}, J.D. Lomas ²⁰,
 J.D. Long ¹⁶², I. Longarini ¹⁶⁰, L. Longo ^{70a,70b}, R. Longo ¹⁶², I. Lopez Paz ⁶⁷,
 A. Lopez Solis ⁴⁸, N. Lorenzo Martinez ⁴, A.M. Lory ¹⁰⁹, G. Löschcke Centeno ¹⁴⁶,
 O. Loseva ³⁷, X. Lou ^{47a,47b}, X. Lou ^{14a,14e}, A. Lounis ⁶⁶, J. Love ⁶, P.A. Love ⁹¹,
 G. Lu ^{14a,14e}, M. Lu ⁸⁰, S. Lu ¹²⁸, Y.J. Lu ⁶⁵, H.J. Lubatti ¹³⁸, C. Luci ^{75a,75b},
 F.L. Lucio Alves ^{14c}, A. Lucotte ⁶⁰, F. Luehring ⁶⁸, I. Luise ¹⁴⁵, O. Lukianchuk ⁶⁶,
 O. Lundberg ¹⁴⁴, B. Lund-Jensen ¹⁴⁴, N.A. Luongo ⁶, M.S. Lutz ¹⁵¹, A.B. Lux ²⁵, D. Lynn ²⁹,
 H. Lyons ⁹², R. Lysak ¹³¹, E. Lytken ⁹⁸, V. Lyubushkin ³⁸, T. Lyubushkina ³⁸, M.M. Lyukova ¹⁴⁵,
 H. Ma ²⁹, K. Ma ^{62a}, L.L. Ma ^{62b}, W. Ma ^{62a}, Y. Ma ¹²¹, D.M. Mac Donell ¹⁶⁵,
 G. Maccarrone ⁵³, J.C. MacDonald ¹⁰⁰, P.C. Machado De Abreu Farias ^{83b}, R. Madar ⁴⁰,
 W.F. Mader ⁵⁰, T. Madula ⁹⁶, J. Maeda ⁸⁵, T. Maeno ²⁹, H. Maguire ¹³⁹, V. Maiboroda ¹³⁵,
 A. Maio ^{130a,130b,130d}, K. Maj ^{86a}, O. Majersky ⁴⁸, S. Majewski ¹²³, N. Makovec ⁶⁶,
 V. Maksimovic ¹⁵, B. Malaescu ¹²⁷, Pa. Malecki ⁸⁷, V.P. Maleev ³⁷, F. Malek ⁶⁰, M. Mali ⁹³,
 D. Malito ⁹⁵, U. Mallik ⁸⁰, S. Maltezos ¹⁰, S. Malyukov ³⁸, J. Mamuzic ¹³, G. Mancini ⁵³,
 G. Manco ^{73a,73b}, J.P. Mandalia ⁹⁴, I. Mandić ⁹³, L. Manhaes de Andrade Filho ^{83a},
 I.M. Maniatis ¹⁶⁹, J. Manjarres Ramos ^{102,aa}, D.C. Mankad ¹⁶⁹, A. Mann ¹⁰⁹, B. Mansoulie ¹³⁵,
 S. Manzoni ³⁶, L. Mao ^{62c}, X. Mapekula ^{33c}, A. Marantis ^{152,r}, G. Marchiori ⁵,
 M. Marcisovsky ¹³¹, C. Marcon ^{71a}, M. Marinescu ²⁰, S. Marium ⁴⁸, M. Marjanovic ¹²⁰,
 E.J. Marshall ⁹¹, Z. Marshall ^{17a}, S. Marti-Garcia ¹⁶³, T.A. Martin ¹⁶⁷, V.J. Martin ⁵²,
 B. Martin dit Latour ¹⁶, L. Martinelli ^{75a,75b}, M. Martinez ^{13,s}, P. Martinez Agullo ¹⁶³,
 V.I. Martinez Outschoorn ¹⁰³, P. Martinez Suarez ¹³, S. Martin-Haugh ¹³⁴, V.S. Martoiu ^{27b},
 A.C. Martyniuk ⁹⁶, A. Marzin ³⁶, D. Mascione ^{78a,78b}, L. Masetti ¹⁰⁰, T. Mashimo ¹⁵³,
 J. Masik ¹⁰¹, A.L. Maslennikov ³⁷, L. Massa ^{23b}, P. Massarotti ^{72a,72b}, P. Mastrandrea ^{74a,74b},
 A. Mastroberardino ^{43b,43a}, T. Masubuchi ¹⁵³, T. Mathisen ¹⁶¹, J. Matousek ¹³³, N. Matsuzawa ¹⁵³,
 J. Maurer ^{27b}, B. Maček ⁹³, D.A. Maximov ³⁷, R. Mazini ¹⁴⁸, I. Maznas ¹⁵², M. Mazza ¹⁰⁷,
 S.M. Mazza ¹³⁶, E. Mazzeo ^{71a,71b}, C. Mc Ginn ²⁹, J.P. Mc Gowan ¹⁰⁴, S.P. Mc Kee ¹⁰⁶,
 E.F. McDonald ¹⁰⁵, A.E. McDougall ¹¹⁴, J.A. Mcfayden ¹⁴⁶, R.P. McGovern ¹²⁸,
 G. Mchedlidze ^{149b}, R.P. Mckenzie ^{33g}, T.C. Mclachlan ⁴⁸, D.J. Mclaughlin ⁹⁶,
 S.J. McMahon ¹³⁴, C.M. Mcpartland ⁹², R.A. McPherson ^{165,w}, S. Mehlhase ¹⁰⁹, A. Mehta ⁹²,
 D. Melini ¹⁵⁰, B.R. Mellado Garcia ^{33g}, A.H. Melo ⁵⁵, F. Meloni ⁴⁸,
 A.M. Mendes Jacques Da Costa ¹⁰¹, H.Y. Meng ¹⁵⁵, L. Meng ⁹¹, S. Menke ¹¹⁰, M. Mentink ³⁶,
 E. Meoni ^{43b,43a}, G. Mercado ¹¹⁵, C. Merlassino ^{69a,69c}, L. Merola ^{72a,72b}, C. Meroni ^{71a,71b},
 G. Merz ¹⁰⁶, J. Metcalfe ⁶, A.S. Mete ⁶, C. Meyer ⁶⁸, J-P. Meyer ¹³⁵, R.P. Middleton ¹³⁴,
 L. Mijović ⁵², G. Mikenberg ¹⁶⁹, M. Mikestikova ¹³¹, M. Mikuž ⁹³, H. Mildner ¹⁰⁰, A. Milic ³⁶,
 C.D. Milke ⁴⁴, D.W. Miller ³⁹, L.S. Miller ³⁴, A. Milov ¹⁶⁹, D.A. Milstead ^{47a,47b}, T. Min ^{14c},
 A.A. Minaenko ³⁷, I.A. Minashvili ^{149b}, L. Mince ⁵⁹, A.I. Mincer ¹¹⁷, B. Mindur ^{86a},
 M. Mineev ³⁸, Y. Mino ⁸⁸, L.M. Mir ¹³, M. Miralles Lopez ¹⁶³, M. Mironova ^{17a}, A. Mishima ¹⁵³,
 M.C. Missio ¹¹³, A. Mitra ¹⁶⁷, V.A. Mitsou ¹⁶³, Y. Mitsumori ¹¹¹, O. Miu ¹⁵⁵,
 P.S. Miyagawa ⁹⁴, T. Mkrtychyan ^{63a}, M. Mlinarevic ⁹⁶, T. Mlinarevic ⁹⁶, M. Mlynarikova ³⁶,
 S. Mobius ¹⁹, P. Moder ⁴⁸, P. Mogg ¹⁰⁹, M.H. Mohamed Farook ¹¹², A.F. Mohammed ^{14a,14e},
 S. Mohapatra ⁴¹, G. Mokgatitwane ^{33g}, L. Moleri ¹⁶⁹, B. Mondal ¹⁴¹, S. Mondal ¹³²,
 K. Mönig ⁴⁸, E. Monnier ¹⁰², L. Monsonis Romero ¹⁶³, J. Montejo Berlingen ¹³, M. Montella ¹¹⁹,
 F. Montekali ^{77a,77b}, F. Monticelli ⁹⁰, S. Monzani ^{69a,69c}, N. Morange ⁶⁶,

A.L. Moreira De Carvalho [ID130a](#), M. Moreno Llácer [ID163](#), C. Moreno Martinez [ID56](#), P. Morettini [ID57b](#),
 S. Morgenstern [ID36](#), M. Morii [ID61](#), M. Morinaga [ID153](#), A.K. Morley [ID36](#), F. Morodei [ID75a,75b](#),
 L. Morvaj [ID36](#), P. Moschovakos [ID36](#), B. Moser [ID36](#), M. Mosidze [ID149b](#), T. Moskalets [ID54](#),
 P. Moskvitina [ID113](#), J. Moss [ID31.1](#), E.J.W. Moyse [ID103](#), O. Mtintsilana [ID33g](#), S. Muanza [ID102](#),
 J. Mueller [ID129](#), D. Muenstermann [ID91](#), R. Müller [ID19](#), G.A. Mullier [ID161](#), A.J. Mullin³², J.J. Mullin¹²⁸,
 D.P. Mungo [ID155](#), D. Munoz Perez [ID163](#), F.J. Munoz Sanchez [ID101](#), M. Murin [ID101](#), W.J. Murray [ID167,134](#),
 A. Murrone [ID71a,71b](#), M. Muškinja [ID17a](#), C. Mwewa [ID29](#), A.G. Myagkov [ID37,a](#), A.J. Myers [ID8](#),
 G. Myers [ID68](#), M. Myska [ID132](#), B.P. Nachman [ID17a](#), O. Nackenhorst [ID49](#), A. Nag [ID50](#), K. Nagai [ID126](#),
 K. Nagano [ID84](#), J.L. Nagle [ID29,ai](#), E. Nagy [ID102](#), A.M. Nairz [ID36](#), Y. Nakahama [ID84](#), K. Nakamura [ID84](#),
 K. Nakkalil [ID5](#), H. Nanjo [ID124](#), R. Narayan [ID44](#), E.A. Narayanan [ID112](#), I. Naryshkin [ID37](#), M. Naseri [ID34](#),
 S. Nasri [ID159](#), C. Nass [ID24](#), G. Navarro [ID22a](#), J. Navarro-Gonzalez [ID163](#), R. Nayak [ID151](#), A. Nayaz [ID18](#),
 P.Y. Nechaeva [ID37](#), F. Nechansky [ID48](#), L. Nedic [ID126](#), T.J. Neep [ID20](#), A. Negri [ID73a,73b](#), M. Negrini [ID23b](#),
 C. Nellist [ID114](#), C. Nelson [ID104](#), K. Nelson [ID106](#), S. Nemecek [ID131](#), M. Nessi [ID36,h](#), M.S. Neubauer [ID162](#),
 F. Neuhaus [ID100](#), J. Neundorf [ID48](#), R. Newhouse [ID164](#), P.R. Newman [ID20](#), C.W. Ng [ID129](#), Y.W.Y. Ng [ID48](#),
 B. Ngair [ID35e](#), H.D.N. Nguyen [ID108](#), R.B. Nickerson [ID126](#), R. Nicolaidou [ID135](#), J. Nielsen [ID136](#),
 M. Niemeyer [ID55](#), J. Niermann [ID55,36](#), N. Nikiforou [ID36](#), V. Nikolaenko [ID37,a](#), I. Nikolic-Audit [ID127](#),
 K. Nikolopoulos [ID20](#), P. Nilsson [ID29](#), I. Ninca [ID48](#), H.R. Nindhito [ID56](#), G. Ninio [ID151](#), A. Nisati [ID75a](#),
 N. Nishu [ID2](#), R. Nisius [ID110](#), J-E. Nitschke [ID50](#), E.K. Nkadimeng [ID33g](#), T. Nobe [ID153](#), D.L. Noel [ID32](#),
 T. Nommensen [ID147](#), M.B. Norfolk [ID139](#), R.R.B. Norisam [ID96](#), B.J. Norman [ID34](#), M. Noury [ID35a](#),
 J. Novak [ID93](#), T. Novak [ID48](#), L. Novotny [ID132](#), R. Novotny [ID112](#), L. Nozka [ID122](#), K. Ntekas [ID160](#),
 N.M.J. Nunes De Moura Junior [ID83b](#), E. Nurse⁹⁶, J. Ocariz [ID127](#), A. Ochi [ID85](#), I. Ochoa [ID130a](#),
 S. Oerdek [ID48,t](#), J.T. Offermann [ID39](#), A. Ogrodnik [ID133](#), A. Oh [ID101](#), C.C. Ohm [ID144](#), H. Oide [ID84](#),
 R. Oishi [ID153](#), M.L. Ojeda [ID48](#), M.W. O'Keefe⁹², Y. Okumura [ID153](#), L.F. Oleiro Seabra [ID130a](#),
 S.A. Olivares Pino [ID137d](#), D. Oliveira Damazio [ID29](#), D. Oliveira Goncalves [ID83a](#), J.L. Oliver [ID160](#),
 Ö.O. Öncel [ID54](#), A.P. O'Neill [ID19](#), A. Onofre [ID130a,130e](#), P.U.E. Onyisi [ID11](#), M.J. Oreglia [ID39](#),
 G.E. Orellana [ID90](#), D. Orestano [ID77a,77b](#), N. Orlando [ID13](#), R.S. Orr [ID155](#), V. O'Shea [ID59](#),
 L.M. Osojnak [ID128](#), R. Ospanov [ID62a](#), G. Otero y Garzon [ID30](#), H. Otono [ID89](#), P.S. Ott [ID63a](#),
 G.J. Ottino [ID17a](#), M. Ouchrif [ID35d](#), J. Ouellette [ID29](#), F. Ould-Saada [ID125](#), M. Owen [ID59](#), R.E. Owen [ID134](#),
 K.Y. Oyulmaz [ID21a](#), V.E. Ozcan [ID21a](#), F. Ozturk [ID87](#), N. Ozturk [ID8](#), S. Ozturk [ID82](#), H.A. Pacey [ID126](#),
 A. Pacheco Pages [ID13](#), C. Padilla Aranda [ID13](#), G. Padovano [ID75a,75b](#), S. Pagan Griso [ID17a](#),
 G. Palacino [ID68](#), A. Palazzo [ID70a,70b](#), S. Palestini [ID36](#), J. Pan [ID172](#), T. Pan [ID64a](#), D.K. Panchal [ID11](#),
 C.E. Pandini [ID114](#), J.G. Panduro Vazquez [ID95](#), H.D. Pandya [ID1](#), H. Pang [ID14b](#), P. Pani [ID48](#),
 G. Panizzo [ID69a,69c](#), L. Paolozzi [ID56](#), C. Papadatos [ID108](#), S. Parajuli [ID44](#), A. Paramonov [ID6](#),
 C. Paraskevopoulos [ID10](#), D. Paredes Hernandez [ID64b](#), K.R. Park [ID41](#), T.H. Park [ID155](#), M.A. Parker [ID32](#),
 F. Parodi [ID57b,57a](#), E.W. Parrish [ID115](#), V.A. Parrish [ID52](#), J.A. Parsons [ID41](#), U. Parzefall [ID54](#),
 B. Pascual Dias [ID108](#), L. Pascual Dominguez [ID151](#), E. Pasqualucci [ID75a](#), S. Passaggio [ID57b](#), F. Pastore [ID95](#),
 P. Pasuwan [ID47a,47b](#), P. Patel [ID87](#), U.M. Patel [ID51](#), J.R. Pater [ID101](#), T. Pauly [ID36](#), J. Pearkes [ID143](#),
 M. Pedersen [ID125](#), R. Pedro [ID130a](#), S.V. Peleganchuk [ID37](#), O. Penc [ID36](#), E.A. Pender [ID52](#),
 K.E. Penski [ID109](#), M. Penzin [ID37](#), B.S. Peralva [ID83d](#), A.P. Pereira Peixoto [ID60](#), L. Pereira Sanchez [ID47a,47b](#),
 D.V. Perepelitsa [ID29,ai](#), E. Perez Codina [ID156a](#), M. Perganti [ID10](#), L. Perini [ID71a,71b,*](#), H. Pernegger [ID36](#),
 O. Perrin [ID40](#), K. Peters [ID48](#), R.F.Y. Peters [ID101](#), B.A. Petersen [ID36](#), T.C. Petersen [ID42](#), E. Petit [ID102](#),
 V. Petousis [ID132](#), C. Petridou [ID152,e](#), A. Petrukhin [ID141](#), M. Pettee [ID17a](#), N.E. Pettersson [ID36](#),
 A. Petukhov [ID37](#), K. Petukhova [ID133](#), R. Pezoa [ID137f](#), L. Pezzotti [ID36](#), G. Pezzullo [ID172](#), T.M. Pham [ID170](#),
 T. Pham [ID105](#), P.W. Phillips [ID134](#), G. Piacquadio [ID145](#), E. Pianori [ID17a](#), F. Piazza [ID123](#), R. Piegai [ID30](#),
 D. Pietreanu [ID27b](#), A.D. Pilkington [ID101](#), M. Pinamonti [ID69a,69c](#), J.L. Pinfeld [ID2](#),
 B.C. Pinheiro Pereira [ID130a](#), A.E. Pinto Pinoargote [ID100,135](#), L. Pintucci [ID69a,69c](#), K.M. Piper [ID146](#),
 A. Pirttikoski [ID56](#), D.A. Pizzi [ID34](#), L. Pizzimento [ID64b](#), A. Pizzini [ID114](#), M.-A. Pleier [ID29](#), V. Plesanovs⁵⁴,

V. Pleskot ¹³³, E. Plotnikova³⁸, G. Poddar ⁴, R. Poettgen ⁹⁸, L. Poggioli ¹²⁷, I. Pokharel ⁵⁵, S. Polacek ¹³³, G. Polesello ^{73a}, A. Poley ^{142,156a}, R. Polifka ¹³², A. Polini ^{23b}, C.S. Pollard ¹⁶⁷, Z.B. Pollock ¹¹⁹, V. Polychronakos ²⁹, E. Pompa Pacchi ^{75a,75b}, D. Ponomarenko ¹¹³, L. Pontecorvo ³⁶, S. Popa ^{27a}, G.A. Popeneciu ^{27d}, A. Poreba ³⁶, D.M. Portillo Quintero ^{156a}, S. Pospisil ¹³², M.A. Postill ¹³⁹, P. Postolache ^{27c}, K. Potamianos ¹⁶⁷, P.A. Potepa ^{86a}, I.N. Potrap ³⁸, C.J. Potter ³², H. Potti ¹, T. Poulsen ⁴⁸, J. Poveda ¹⁶³, M.E. Pozo Astigarraga ³⁶, A. Prades Ibanez ¹⁶³, J. Pretel ⁵⁴, D. Price ¹⁰¹, M. Primavera ^{70a}, M.A. Principe Martin ⁹⁹, R. Privara ¹²², T. Procter ⁵⁹, M.L. Proffitt ¹³⁸, N. Proklova ¹²⁸, K. Prokofiev ^{64c}, G. Proto ¹¹⁰, S. Protopopescu ²⁹, J. Proudfoot ⁶, M. Przybycien ^{86a}, W.W. Przygoda ^{86b}, J.E. Puddefoot ¹³⁹, D. Pudzha ³⁷, D. Pyatiizbyantseva ³⁷, J. Qian ¹⁰⁶, D. Qichen ¹⁰¹, Y. Qin ¹⁰¹, T. Qiu ⁵², A. Quadt ⁵⁵, M. Queitsch-Maitland ¹⁰¹, G. Quetant ⁵⁶, R.P. Quinn ¹⁶⁴, G. Rabanal Bolanos ⁶¹, D. Rafanoharana ⁵⁴, F. Ragusa ^{71a,71b}, J.L. Rainbolt ³⁹, J.A. Raine ⁵⁶, S. Rajagopalan ²⁹, E. Ramakoti ³⁷, I.A. Ramirez-Berend ³⁴, K. Ran ^{48,14e}, N.P. Rapheeha ^{33g}, H. Rasheed ^{27b}, V. Raskina ¹²⁷, D.F. Rassloff ^{63a}, S. Rave ¹⁰⁰, B. Ravina ⁵⁵, I. Ravinovich ¹⁶⁹, M. Raymond ³⁶, A.L. Read ¹²⁵, N.P. Readioff ¹³⁹, D.M. Rebutzi ^{73a,73b}, G. Redlinger ²⁹, A.S. Reed ¹¹⁰, K. Reeves ²⁶, J.A. Reidelsturz ¹⁷¹, D. Reikher ¹⁵¹, A. Rej ⁴⁹, C. Rembser ³⁶, A. Renardi ⁴⁸, M. Renda ^{27b}, M.B. Rendel¹¹⁰, F. Renner ⁴⁸, A.G. Rennie ¹⁶⁰, A.L. Rescia ⁴⁸, S. Resconi ^{71a}, M. Ressegotti ^{57b,57a}, S. Rettie ³⁶, J.G. Reyes Rivera ¹⁰⁷, E. Reynolds ^{17a}, O.L. Rezanova ³⁷, P. Reznicek ¹³³, N. Ribaric ⁹¹, E. Ricci ^{78a,78b}, R. Richter ¹¹⁰, S. Richter ^{47a,47b}, E. Richter-Was ^{86b}, M. Ridel ¹²⁷, S. Ridouani ^{35d}, P. Rieck ¹¹⁷, P. Riedler ³⁶, E.M. Riefel ^{47a,47b}, J.O. Rieger ¹¹⁴, M. Rijssenbeek ¹⁴⁵, A. Rimoldi ^{73a,73b}, M. Rimoldi ³⁶, L. Rinaldi ^{23b,23a}, T.T. Rinn ²⁹, M.P. Rinnagel ¹⁰⁹, G. Ripellino ¹⁶¹, I. Riu ¹³, P. Rivadeneira ⁴⁸, J.C. Rivera Vergara ¹⁶⁵, F. Rizatdinova ¹²¹, E. Rizvi ⁹⁴, B.A. Roberts ¹⁶⁷, B.R. Roberts ^{17a}, S.H. Robertson ^{104,w}, D. Robinson ³², C.M. Robles Gajardo^{137f}, M. Robles Manzano ¹⁰⁰, A. Robson ⁵⁹, A. Rocchi ^{76a,76b}, C. Roda ^{74a,74b}, S. Rodriguez Bosca ^{63a}, Y. Rodriguez Garcia ^{22a}, A. Rodriguez Rodriguez ⁵⁴, A.M. Rodríguez Vera ^{156b}, S. Roe³⁶, J.T. Roemer ¹⁶⁰, A.R. Roepe-Gier ¹³⁶, J. Roggel ¹⁷¹, O. Røhne ¹²⁵, R.A. Rojas ¹⁰³, C.P.A. Roland ¹²⁷, J. Roloff ²⁹, A. Romaniouk ³⁷, E. Romano ^{73a,73b}, M. Romano ^{23b}, A.C. Romero Hernandez ¹⁶², N. Rompotis ⁹², L. Roos ¹²⁷, S. Rosati ^{75a}, B.J. Rosser ³⁹, E. Rossi ¹²⁶, E. Rossi ^{72a,72b}, L.P. Rossi ^{57b}, L. Rossini ⁵⁴, R. Rosten ¹¹⁹, M. Rotaru ^{27b}, B. Rottler ⁵⁴, C. Rougier ^{102,aa}, D. Rousseau ⁶⁶, D. Rousso ³², A. Roy ¹⁶², S. Roy-Garand ¹⁵⁵, A. Rozanov ¹⁰², Z.M.A. Rozario ⁵⁹, Y. Rozen ¹⁵⁰, X. Ruan ^{33g}, A. Rubio Jimenez ¹⁶³, A.J. Ruby ⁹², V.H. Ruelas Rivera ¹⁸, T.A. Ruggeri ¹, A. Ruggiero ¹²⁶, A. Ruiz-Martinez ¹⁶³, A. Rummeler ³⁶, Z. Rurikova ⁵⁴, N.A. Rusakovich ³⁸, H.L. Russell ¹⁶⁵, G. Russo ^{75a,75b}, J.P. Rutherford ⁷, S. Rutherford Colmenares ³², K. Rybacki⁹¹, M. Rybar ¹³³, E.B. Rye ¹²⁵, A. Ryzhov ⁴⁴, J.A. Sabater Iglesias ⁵⁶, P. Sabatini ¹⁶³, H.F-W. Sadrozinski ¹³⁶, F. Safai Tehrani ^{75a}, B. Safarzadeh Samani ¹³⁴, M. Safdari ¹⁴³, S. Saha ¹⁶⁵, M. Sahinsoy ¹¹⁰, A. Saibel ¹⁶³, M. Saimpert ¹³⁵, M. Saito ¹⁵³, T. Saito ¹⁵³, D. Salamani ³⁶, A. Salnikov ¹⁴³, J. Salt ¹⁶³, A. Salvador Salas ¹⁵¹, D. Salvatore ^{43b,43a}, F. Salvatore ¹⁴⁶, A. Salzburger ³⁶, D. Sammel ⁵⁴, D. Sampsonidis ^{152,e}, D. Sampsonidou ¹²³, J. Sánchez ¹⁶³, A. Sanchez Pineda ⁴, V. Sanchez Sebastian ¹⁶³, H. Sandaker ¹²⁵, C.O. Sander ⁴⁸, J.A. Sandesara ¹⁰³, M. Sandhoff ¹⁷¹, C. Sandoval ^{22b}, D.P.C. Sankey ¹³⁴, T. Sano ⁸⁸, A. Sansoni ⁵³, L. Santi ^{75a,75b}, C. Santoni ⁴⁰, H. Santos ^{130a,130b}, S.N. Santpur ^{17a}, A. Santra ¹⁶⁹, K.A. Saoucha ^{116b}, J.G. Saraiva ^{130a,130d}, J. Sardain ⁷, O. Sasaki ⁸⁴, K. Sato ¹⁵⁷, C. Sauer^{63b}, F. Sauerburger ⁵⁴, E. Sauvan ⁴, P. Savard ^{155,af}, R. Sawada ¹⁵³, C. Sawyer ¹³⁴, L. Sawyer ⁹⁷, I. Sayago Galvan¹⁶³, C. Sbarra ^{23b}, A. Sbrizzi ^{23b,23a}, T. Scanlon ⁹⁶, J. Schaarschmidt ¹³⁸, P. Schacht ¹¹⁰, U. Schäfer ¹⁰⁰, A.C. Schaffer ^{66,44}, D. Schaile ¹⁰⁹, R.D. Schamberger ¹⁴⁵, C. Scharf ¹⁸, M.M. Schefer ¹⁹,

V.A. Schegelsky [ID37](#), D. Scheirich [ID133](#), F. Schenck [ID18](#), M. Schernau [ID160](#), C. Scheulen [ID55](#), C. Schiavi [ID57b,57a](#), E.J. Schioppa [ID70a,70b](#), M. Schioppa [ID43b,43a](#), B. Schlag [ID143,n](#), K.E. Schleicher [ID54](#), S. Schlenker [ID36](#), J. Schmeing [ID171](#), M.A. Schmidt [ID171](#), K. Schmieden [ID100](#), C. Schmitt [ID100](#), N. Schmitt [ID100](#), S. Schmitt [ID48](#), L. Schoeffel [ID135](#), A. Schoening [ID63b](#), P.G. Scholer [ID54](#), E. Schopf [ID126](#), M. Schott [ID100](#), J. Schovancova [ID36](#), S. Schramm [ID56](#), F. Schroeder [ID171](#), T. Schroer [ID56](#), H-C. Schultz-Coulon [ID63a](#), M. Schumacher [ID54](#), B.A. Schumm [ID136](#), Ph. Schune [ID135](#), A.J. Schuy [ID138](#), H.R. Schwartz [ID136](#), A. Schwartzman [ID143](#), T.A. Schwarz [ID106](#), Ph. Schwemling [ID135](#), R. Schwienhorst [ID107](#), A. Sciandra [ID136](#), G. Sciolla [ID26](#), F. Scuri [ID74a](#), C.D. Sebastiani [ID92](#), K. Sedlaczek [ID115](#), P. Seema [ID18](#), S.C. Seidel [ID112](#), A. Seiden [ID136](#), B.D. Seidlitz [ID41](#), C. Seitz [ID48](#), J.M. Seixas [ID83b](#), G. Sekhniaidze [ID72a](#), S.J. Sekula [ID44](#), L. Selem [ID60](#), N. Semprini-Cesari [ID23b,23a](#), D. Sengupta [ID56](#), V. Senthilkumar [ID163](#), L. Serin [ID66](#), L. Serkin [ID69a,69b](#), M. Sessa [ID76a,76b](#), H. Severini [ID120](#), F. Sforza [ID57b,57a](#), A. Sfyrta [ID56](#), E. Shabalina [ID55](#), R. Shaheen [ID144](#), J.D. Shahinian [ID128](#), D. Shaked Renous [ID169](#), L.Y. Shan [ID14a](#), M. Shapiro [ID17a](#), A. Sharma [ID36](#), A.S. Sharma [ID164](#), P. Sharma [ID80](#), S. Sharma [ID48](#), P.B. Shatalov [ID37](#), K. Shaw [ID146](#), S.M. Shaw [ID101](#), A. Shcherbakova [ID37](#), Q. Shen [ID62c,5](#), D.J. Sheppard [ID142](#), P. Sherwood [ID96](#), L. Shi [ID96](#), X. Shi [ID14a](#), C.O. Shimmin [ID172](#), J.D. Shinner [ID95](#), I.P.J. Shipsey [ID126](#), S. Shirabe [ID56,h](#), M. Shiyakova [ID38,u](#), J. Shlomi [ID169](#), M.J. Shochet [ID39](#), J. Shojaii [ID105](#), D.R. Shope [ID125](#), B. Shrestha [ID120](#), S. Shrestha [ID119,ak](#), E.M. Shrif [ID33g](#), M.J. Shroff [ID165](#), P. Sicho [ID131](#), A.M. Sickles [ID162](#), E. Sideras Haddad [ID33g](#), A. Sidoti [ID23b](#), F. Siegert [ID50](#), Dj. Sijacki [ID15](#), F. Sili [ID90](#), J.M. Silva [ID20](#), M.V. Silva Oliveira [ID29](#), S.B. Silverstein [ID47a](#), S. Simion [ID66](#), R. Simoniello [ID36](#), E.L. Simpson [ID59](#), H. Simpson [ID146](#), L.R. Simpson [ID106](#), N.D. Simpson [ID98](#), S. Simsek [ID82](#), S. Sindhu [ID55](#), P. Sinervo [ID155](#), S. Singh [ID155](#), S. Sinha [ID48](#), S. Sinha [ID101](#), M. Sioli [ID23b,23a](#), I. Siral [ID36](#), E. Sitnikova [ID48](#), S.Yu. Sivoklov [ID37,*](#), J. Sjölin [ID47a,47b](#), A. Skaf [ID55](#), E. Skorda [ID20](#), P. Skubic [ID120](#), M. Slawinska [ID87](#), V. Smakhtin [ID169](#), B.H. Smart [ID134](#), J. Smiesko [ID36](#), S.Yu. Smirnov [ID37](#), Y. Smirnov [ID37](#), L.N. Smirnova [ID37,a](#), O. Smirnova [ID98](#), A.C. Smith [ID41](#), E.A. Smith [ID39](#), H.A. Smith [ID126](#), J.L. Smith [ID92](#), R. Smith [ID143](#), M. Smizanska [ID91](#), K. Smolek [ID132](#), A.A. Snesarev [ID37](#), S.R. Snider [ID155](#), H.L. Snoek [ID114](#), S. Snyder [ID29](#), R. Sobie [ID165,w](#), A. Soffer [ID151](#), C.A. Solans Sanchez [ID36](#), E.Yu. Soldatov [ID37](#), U. Soldevila [ID163](#), A.A. Solodkov [ID37](#), S. Solomon [ID26](#), A. Soloshenko [ID38](#), K. Solovieva [ID54](#), O.V. Solovyanov [ID40](#), V. Solovyev [ID37](#), P. Sommer [ID36](#), A. Sonay [ID13](#), W.Y. Song [ID156b](#), J.M. Sonneveld [ID114](#), A. Sopczak [ID132](#), A.L. Soppio [ID96](#), F. Sopkova [ID28b](#), I.R. Sotarriva Alvarez [ID154](#), V. Sothilingam [ID63a](#), O.J. Soto Sandoval [ID137c,137b](#), S. Sottocornola [ID68](#), R. Soualah [ID116b](#), Z. Soumami [ID35e](#), D. South [ID48](#), N. Soybelman [ID169](#), S. Spagnolo [ID70a,70b](#), M. Spalla [ID110](#), D. Sperlich [ID54](#), G. Spigo [ID36](#), S. Spinali [ID91](#), D.P. Spiteri [ID59](#), M. Spousta [ID133](#), E.J. Staats [ID34](#), A. Stabile [ID71a,71b](#), R. Stamen [ID63a](#), A. Stampekis [ID20](#), M. Standke [ID24](#), E. Stanecka [ID87](#), M.V. Stange [ID50](#), B. Stanislaus [ID17a](#), M.M. Stanitzki [ID48](#), B. Stapf [ID48](#), E.A. Starchenko [ID37](#), G.H. Stark [ID136](#), J. Stark [ID102,aa](#), D.M. Starke [ID156b](#), P. Staroba [ID131](#), P. Starovoitov [ID63a](#), S. Stärz [ID104](#), R. Staszewski [ID87](#), G. Stavropoulos [ID46](#), J. Steentoft [ID161](#), P. Steinberg [ID29](#), B. Stelzer [ID142,156a](#), H.J. Stelzer [ID129](#), O. Stelzer-Chilton [ID156a](#), H. Stenzel [ID58](#), T.J. Stevenson [ID146](#), G.A. Stewart [ID36](#), J.R. Stewart [ID121](#), M.C. Stockton [ID36](#), G. Stoicea [ID27b](#), M. Stolarski [ID130a](#), S. Stonjek [ID110](#), A. Straessner [ID50](#), J. Strandberg [ID144](#), S. Strandberg [ID47a,47b](#), M. Stratmann [ID171](#), M. Strauss [ID120](#), T. Streblner [ID102](#), P. Strizenc [ID28b](#), R. Ströhmer [ID166](#), D.M. Strom [ID123](#), R. Stroynowski [ID44](#), A. Strubig [ID47a,47b](#), S.A. Stucci [ID29](#), B. Stugu [ID16](#), J. Stupak [ID120](#), N.A. Styles [ID48](#), D. Su [ID143](#), S. Su [ID62a](#), W. Su [ID62d](#), X. Su [ID62a,66](#), K. Sugizaki [ID153](#), V.V. Sulin [ID37](#), M.J. Sullivan [ID92](#), D.M.S. Sultan [ID78a,78b](#), L. Sultanaliev [ID37](#), S. Sultansoy [ID3b](#), T. Sumida [ID88](#), S. Sun [ID106](#), S. Sun [ID170](#), O. Sunneborn Gudnadottir [ID161](#), N. Sur [ID102](#), M.R. Sutton [ID146](#), H. Suzuki [ID157](#), M. Svatos [ID131](#), M. Swiatlowski [ID156a](#), T. Swirski [ID166](#), I. Sykora [ID28a](#), M. Sykora [ID133](#), T. Sykora [ID133](#), D. Ta [ID100](#), K. Tackmann [ID48,t](#), A. Taffard [ID160](#), R. Tafirout [ID156a](#), J.S. Tafuya Vargas [ID66](#), E.P. Takeva [ID52](#),

Y. Takubo ⁸⁴, M. Talby ¹⁰², A.A. Talyshev ³⁷, K.C. Tam ^{64b}, N.M. Tamir ¹⁵¹, A. Tanaka ¹⁵³,
 J. Tanaka ¹⁵³, R. Tanaka ⁶⁶, M. Tanasini ^{57b,57a}, Z. Tao ¹⁶⁴, S. Tapia Araya ^{137f},
 S. Tapprogge ¹⁰⁰, A. Tarek Abouelfadl Mohamed ¹⁰⁷, S. Tarem ¹⁵⁰, K. Tariq ^{14a}, G. Tarna ^{102,27b},
 G.F. Tartarelli ^{71a}, P. Tas ¹³³, M. Tasevsky ¹³¹, E. Tassi ^{43b,43a}, A.C. Tate ¹⁶², G. Tateno ¹⁵³,
 Y. Tayalati ^{35e,v}, G.N. Taylor ¹⁰⁵, W. Taylor ^{156b}, A.S. Tee ¹⁷⁰, R. Teixeira De Lima ¹⁴³,
 P. Teixeira-Dias ⁹⁵, J.J. Teoh ¹⁵⁵, K. Terashi ¹⁵³, J. Terron ⁹⁹, S. Terzo ¹³, M. Testa ⁵³,
 R.J. Teuscher ^{155,w}, A. Thaler ⁷⁹, O. Theiner ⁵⁶, N. Themistokleous ⁵², T. Theveneaux-Pelzer ¹⁰²,
 O. Thielmann ¹⁷¹, D.W. Thomas ⁹⁵, J.P. Thomas ²⁰, E.A. Thompson ^{17a}, P.D. Thompson ²⁰,
 E. Thomson ¹²⁸, Y. Tian ⁵⁵, V. Tikhomirov ^{37,a}, Yu.A. Tikhonov ³⁷, S. Timoshenko ³⁷,
 D. Timoshyn ¹³³, E.X.L. Ting ¹, P. Tipton ¹⁷², S.H. Tlou ^{33g}, A. Tnourji ⁴⁰, K. Todome ¹⁵⁴,
 S. Todorova-Nova ¹³³, S. Todt ⁵⁰, M. Togawa ⁸⁴, J. Tojo ⁸⁹, S. Tokár ^{28a}, K. Tokushuku ⁸⁴,
 O. Toldaiev ⁶⁸, R. Tombs ³², M. Tomoto ^{84,111}, L. Tompkins ^{143,n}, K.W. Topolnicki ^{86b},
 E. Torrence ¹²³, H. Torres ^{102,aa}, E. Torró Pastor ¹⁶³, M. Toscani ³⁰, C. Tosciri ³⁹, M. Tost ¹¹,
 D.R. Tovey ¹³⁹, A. Traeet ¹⁶, I.S. Trandafir ^{27b}, T. Trefzger ¹⁶⁶, A. Tricoli ²⁹, I.M. Trigger ^{156a},
 S. Trincaz-Duvoid ¹²⁷, D.A. Trischuk ²⁶, B. Trocmé ⁶⁰, C. Troncon ^{71a}, M.R. Trotta ^{aj},
 L. Truong ^{33c}, M. Trzebinski ⁸⁷, A. Trzupiek ⁸⁷, F. Tsai ¹⁴⁵, M. Tsai ¹⁰⁶, A. Tsiamis ^{152,e},
 P.V. Tsiareshka ³⁷, S. Tsigaridas ^{156a}, A. Tsirigotis ^{152,r}, V. Tsiskaridze ¹⁵⁵, E.G. Tskhadadze ^{149a},
 M. Tsopoulou ^{152,e}, Y. Tsujikawa ⁸⁸, I.I. Tsukerman ³⁷, V. Tsulaia ^{17a}, S. Tsuno ⁸⁴, K. Tsuru ¹¹⁸,
 D. Tsybychev ¹⁴⁵, Y. Tu ^{64b}, A. Tudorache ^{27b}, V. Tudorache ^{27b}, A.N. Tuna ⁶¹,
 S. Turchikhin ^{57b,57a}, I. Turk Cakir ^{3a}, R. Turra ^{71a}, T. Turtuvshin ^{38,x}, P.M. Tuts ⁴¹,
 S. Tzamarias ^{152,e}, P. Tzanis ¹⁰, E. Tzovara ¹⁰⁰, F. Ukegawa ¹⁵⁷, P.A. Ulloa Poblete ^{137c,137b},
 E.N. Umaka ²⁹, G. Unal ³⁶, M. Unal ¹¹, A. Undrus ²⁹, G. Unel ¹⁶⁰, J. Urban ^{28b},
 P. Urquijo ¹⁰⁵, P. Urrejola ^{137a}, G. Usai ⁸, R. Ushioda ¹⁵⁴, M. Usman ¹⁰⁸, Z. Uysal ^{21b},
 V. Vacek ¹³², B. Vachon ¹⁰⁴, K.O.H. Vadla ¹²⁵, T. Vafeiadis ³⁶, A. Vaitkus ⁹⁶, C. Valderanis ¹⁰⁹,
 E. Valdes Santurio ^{47a,47b}, M. Valente ^{156a}, S. Valentinetti ^{23b,23a}, A. Valero ¹⁶³,
 E. Valiente Moreno ¹⁶³, A. Vallier ^{102,aa}, J.A. Valls Ferrer ¹⁶³, D.R. Van Arneman ¹¹⁴,
 T.R. Van Daalen ¹³⁸, A. Van Der Graaf ⁴⁹, P. Van Gemmeren ⁶, M. Van Rijnbach ^{125,36},
 S. Van Stroud ⁹⁶, I. Van Vulpen ¹¹⁴, M. Vanadia ^{76a,76b}, W. Vandelli ³⁶, M. Vandenbroucke ¹³⁵,
 E.R. Vandewall ¹²¹, D. Vannicola ¹⁵¹, L. Vannoli ^{57b,57a}, R. Vari ^{75a}, E.W. Varnes ⁷,
 C. Varni ^{17b}, T. Varol ¹⁴⁸, D. Varouchas ⁶⁶, L. Varriale ¹⁶³, K.E. Varvell ¹⁴⁷, M.E. Vasile ^{27b},
 L. Vaslin ⁸⁴, G.A. Vasquez ¹⁶⁵, A. Vasyukov ³⁸, F. Vazeille ⁴⁰, T. Vazquez Schroeder ³⁶,
 J. Veatch ³¹, V. Vecchio ¹⁰¹, M.J. Veen ¹⁰³, I. Veliscek ¹²⁶, L.M. Veloce ¹⁵⁵, F. Veloso ^{130a,130c},
 S. Veneziano ^{75a}, A. Ventura ^{70a,70b}, S. Ventura Gonzalez ¹³⁵, A. Verbytskyi ¹¹⁰,
 M. Verducci ^{74a,74b}, C. Vergis ²⁴, M. Verissimo De Araujo ^{83b}, W. Verkerke ¹¹⁴,
 J.C. Vermeulen ¹¹⁴, C. Vernieri ¹⁴³, M. Vessella ¹⁰³, M.C. Vetterli ^{142,af}, A. Vgenopoulos ^{152,e},
 N. Viaux Maira ^{137f}, T. Vickey ¹³⁹, O.E. Vickey Boeriu ¹³⁹, G.H.A. Viehhauser ¹²⁶, L. Vigani ^{63b},
 M. Villa ^{23b,23a}, M. Villaplana Perez ¹⁶³, E.M. Villhauer ⁵², E. Vilucchi ⁵³, M.G. Vinciter ³⁴,
 G.S. Virdee ²⁰, A. Vishwakarma ⁵², A. Visibile ¹¹⁴, C. Vittori ³⁶, I. Vivarelli ¹⁴⁶,
 E. Voevodina ¹¹⁰, F. Vogel ¹⁰⁹, J.C. Voigt ⁵⁰, P. Vokac ¹³², Yu. Volkotrub ^{86a}, J. Von Ahnen ⁴⁸,
 E. Von Toerne ²⁴, B. Vormwald ³⁶, V. Vorobel ¹³³, K. Vorobev ³⁷, M. Vos ¹⁶³, K. Voss ¹⁴¹,
 J.H. Vossebeld ⁹², M. Vozak ¹¹⁴, L. Vozdecky ⁹⁴, N. Vranjes ¹⁵, M. Vranjes Milosavljevic ¹⁵,
 M. Vreeswijk ¹¹⁴, R. Vuillermet ³⁶, O. Vujanovic ¹⁰⁰, I. Vukotic ³⁹, S. Wada ¹⁵⁷, C. Wagner ¹⁰³,
 J.M. Wagner ^{17a}, W. Wagner ¹⁷¹, S. Wahdan ¹⁷¹, H. Wahlberg ⁹⁰, M. Wakida ¹¹¹, J. Walder ¹³⁴,
 R. Walker ¹⁰⁹, W. Walkowiak ¹⁴¹, A. Wall ¹²⁸, T. Wamorkar ⁶, A.Z. Wang ¹³⁶, C. Wang ¹⁰⁰,
 C. Wang ^{62c}, H. Wang ^{17a}, J. Wang ^{64a}, R.-J. Wang ¹⁰⁰, R. Wang ⁶¹, R. Wang ⁶,
 S.M. Wang ¹⁴⁸, S. Wang ^{62b}, T. Wang ^{62a}, W.T. Wang ⁸⁰, W. Wang ^{14a}, X. Wang ^{14c},
 X. Wang ¹⁶², X. Wang ^{62c}, Y. Wang ^{62d}, Y. Wang ^{14c}, Z. Wang ¹⁰⁶, Z. Wang ^{62d,51,62c},

Z. Wang ¹⁰⁶, A. Warburton ¹⁰⁴, R.J. Ward ²⁰, N. Warrack ⁵⁹, A.T. Watson ²⁰, H. Watson ⁵⁹, M.F. Watson ²⁰, E. Watton ^{59,134}, G. Watts ¹³⁸, B.M. Waugh ⁹⁶, C. Weber ²⁹, H.A. Weber ¹⁸, M.S. Weber ¹⁹, S.M. Weber ^{63a}, C. Wei ^{62a}, Y. Wei ¹²⁶, A.R. Weidberg ¹²⁶, E.J. Weik ¹¹⁷, J. Weingarten ⁴⁹, M. Weirich ¹⁰⁰, C. Weiser ⁵⁴, C.J. Wells ⁴⁸, T. Wenaus ²⁹, B. Wendland ⁴⁹, T. Wengler ³⁶, N.S. Wenke ¹¹⁰, N. Wermes ²⁴, M. Wessels ^{63a}, A.M. Wharton ⁹¹, A.S. White ⁶¹, A. White ⁸, M.J. White ¹, D. Whiteson ¹⁶⁰, L. Wickremasinghe ¹²⁴, W. Wiedenmann ¹⁷⁰, C. Wiel ⁵⁰, M. Wielers ¹³⁴, C. Wiglesworth ⁴², D.J. Wilbern ¹²⁰, H.G. Wilkens ³⁶, D.M. Williams ⁴¹, H.H. Williams ¹²⁸, S. Williams ³², S. Willocq ¹⁰³, B.J. Wilson ¹⁰¹, P.J. Windischhofer ³⁹, F.I. Winkel ³⁰, F. Winklmeier ¹²³, B.T. Winter ⁵⁴, J.K. Winter ¹⁰¹, M. Wittgen ¹⁴³, M. Wobisch ⁹⁷, Z. Wolffs ¹¹⁴, J. Wollrath ¹⁶⁰, M.W. Wolter ⁸⁷, H. Wolters ^{130a,130c}, A.F. Wongel ⁴⁸, E.L. Woodward ⁴¹, S.D. Worm ⁴⁸, B.K. Wosiek ⁸⁷, K.W. Woźniak ⁸⁷, S. Wozniowski ⁵⁵, K. Wraight ⁵⁹, C. Wu ²⁰, J. Wu ^{14a,14e}, M. Wu ^{64a}, M. Wu ¹¹³, S.L. Wu ¹⁷⁰, X. Wu ⁵⁶, Y. Wu ^{62a}, Z. Wu ¹³⁵, J. Wuerzinger ^{110,ad}, T.R. Wyatt ¹⁰¹, B.M. Wynne ⁵², S. Xella ⁴², L. Xia ^{14c}, M. Xia ^{14b}, J. Xiang ^{64c}, M. Xie ^{62a}, X. Xie ^{62a}, S. Xin ^{14a,14e}, A. Xiong ¹²³, J. Xiong ^{17a}, D. Xu ^{14a}, H. Xu ^{62a}, L. Xu ^{62a}, R. Xu ¹²⁸, T. Xu ¹⁰⁶, Y. Xu ^{14b}, Z. Xu ⁵², Z. Xu ^{14c}, B. Yabsley ¹⁴⁷, S. Yacoob ^{33a}, Y. Yamaguchi ¹⁵⁴, E. Yamashita ¹⁵³, H. Yamauchi ¹⁵⁷, T. Yamazaki ^{17a}, Y. Yamazaki ⁸⁵, J. Yan ^{62c}, S. Yan ¹²⁶, Z. Yan ²⁵, H.J. Yang ^{62c,62d}, H.T. Yang ^{62a}, S. Yang ^{62a}, T. Yang ^{64c}, X. Yang ³⁶, X. Yang ^{14a}, Y. Yang ⁴⁴, Y. Yang ^{62a}, Z. Yang ^{62a}, W-M. Yao ^{17a}, Y.C. Yap ⁴⁸, H. Ye ^{14c}, H. Ye ⁵⁵, J. Ye ^{14a}, S. Ye ²⁹, X. Ye ^{62a}, Y. Yeh ⁹⁶, I. Yeletsikh ³⁸, B.K. Yeo ^{17b}, M.R. Yexley ⁹⁶, P. Yin ⁴¹, K. Yorita ¹⁶⁸, S. Younas ^{27b}, C.J.S. Young ³⁶, C. Young ¹⁴³, C. Yu ^{14a,14e,ah}, Y. Yu ^{62a}, M. Yuan ¹⁰⁶, R. Yuan ^{62b}, L. Yue ⁹⁶, M. Zaazoua ^{62a}, B. Zabinski ⁸⁷, E. Zaid ⁵², Z.K. Zak ⁸⁷, T. Zakareishvili ^{149b}, N. Zakharchuk ³⁴, S. Zambito ⁵⁶, J.A. Zamora Saa ^{137d,137b}, J. Zang ¹⁵³, D. Zanzi ⁵⁴, O. Zaplatilek ¹³², C. Zeitnitz ¹⁷¹, H. Zeng ^{14a}, J.C. Zeng ¹⁶², D.T. Zenger Jr ²⁶, O. Zenin ³⁷, T. Ženiš ^{28a}, S. Zenz ⁹⁴, S. Zerradi ^{35a}, D. Zerwas ⁶⁶, M. Zhai ^{14a,14e}, B. Zhang ^{14c}, D.F. Zhang ¹³⁹, J. Zhang ^{62b}, J. Zhang ⁶, K. Zhang ^{14a,14e}, L. Zhang ^{14c}, P. Zhang ^{14a,14e}, R. Zhang ¹⁷⁰, S. Zhang ¹⁰⁶, S. Zhang ⁴⁴, T. Zhang ¹⁵³, X. Zhang ^{62c}, X. Zhang ^{62b}, Y. Zhang ^{62c,5}, Y. Zhang ⁹⁶, Y. Zhang ^{14c}, Z. Zhang ^{17a}, Z. Zhang ⁶⁶, H. Zhao ¹³⁸, T. Zhao ^{62b}, Y. Zhao ¹³⁶, Z. Zhao ^{62a}, A. Zhemchugov ³⁸, J. Zheng ^{14c}, K. Zheng ¹⁶², X. Zheng ^{62a}, Z. Zheng ¹⁴³, D. Zhong ¹⁶², B. Zhou ¹⁰⁶, H. Zhou ⁷, N. Zhou ^{62c}, Y. Zhou ⁷, C.G. Zhu ^{62b}, J. Zhu ¹⁰⁶, Y. Zhu ^{62c}, Y. Zhu ^{62a}, X. Zhuang ^{14a}, K. Zhukov ³⁷, V. Zhulanov ³⁷, N.I. Zimine ³⁸, J. Zinsser ^{63b}, M. Ziolkowski ¹⁴¹, L. Živković ¹⁵, A. Zoccoli ^{23b,23a}, K. Zoch ⁶¹, T.G. Zorbas ¹³⁹, O. Zormpa ⁴⁶, W. Zou ⁴¹, L. Zwalinski ³⁶.

¹Department of Physics, University of Adelaide, Adelaide; Australia.

²Department of Physics, University of Alberta, Edmonton AB; Canada.

^{3(a)}Department of Physics, Ankara University, Ankara; ^(b)Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye.

⁴LAPP, Université Savoie Mont Blanc, CNRS/IN2P3, Annecy; France.

⁵APC, Université Paris Cité, CNRS/IN2P3, Paris; France.

⁶High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America.

⁷Department of Physics, University of Arizona, Tucson AZ; United States of America.

⁸Department of Physics, University of Texas at Arlington, Arlington TX; United States of America.

⁹Physics Department, National and Kapodistrian University of Athens, Athens; Greece.

¹⁰Physics Department, National Technical University of Athens, Zografou; Greece.

¹¹Department of Physics, University of Texas at Austin, Austin TX; United States of America.

¹²Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.

- ¹³Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain.
- ¹⁴(^a)Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; (^b)Physics Department, Tsinghua University, Beijing; (^c)Department of Physics, Nanjing University, Nanjing; (^d)School of Science, Shenzhen Campus of Sun Yat-sen University; (^e)University of Chinese Academy of Science (UCAS), Beijing; China.
- ¹⁵Institute of Physics, University of Belgrade, Belgrade; Serbia.
- ¹⁶Department for Physics and Technology, University of Bergen, Bergen; Norway.
- ¹⁷(^a)Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA; (^b)University of California, Berkeley CA; United States of America.
- ¹⁸Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany.
- ¹⁹Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland.
- ²⁰School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom.
- ²¹(^a)Department of Physics, Bogazici University, Istanbul; (^b)Department of Physics Engineering, Gaziantep University, Gaziantep; (^c)Department of Physics, Istanbul University, Istanbul; Türkiye.
- ²²(^a)Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá; (^b)Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia.
- ²³(^a)Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna; (^b)INFN Sezione di Bologna; Italy.
- ²⁴Physikalisches Institut, Universität Bonn, Bonn; Germany.
- ²⁵Department of Physics, Boston University, Boston MA; United States of America.
- ²⁶Department of Physics, Brandeis University, Waltham MA; United States of America.
- ²⁷(^a)Transilvania University of Brasov, Brasov; (^b)Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; (^c)Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi; (^d)National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca; (^e)University Politehnica Bucharest, Bucharest; (^f)West University in Timisoara, Timisoara; (^g)Faculty of Physics, University of Bucharest, Bucharest; Romania.
- ²⁸(^a)Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava; (^b)Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic.
- ²⁹Physics Department, Brookhaven National Laboratory, Upton NY; United States of America.
- ³⁰Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina.
- ³¹California State University, CA; United States of America.
- ³²Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom.
- ³³(^a)Department of Physics, University of Cape Town, Cape Town; (^b)iThemba Labs, Western Cape; (^c)Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg; (^d)National Institute of Physics, University of the Philippines Diliman (Philippines); (^e)University of South Africa, Department of Physics, Pretoria; (^f)University of Zululand, KwaDlangezwa; (^g)School of Physics, University of the Witwatersrand, Johannesburg; South Africa.
- ³⁴Department of Physics, Carleton University, Ottawa ON; Canada.
- ³⁵(^a)Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca; (^b)Faculté des Sciences, Université Ibn-Tofail, Kénitra; (^c)Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech; (^d)LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda; (^e)Faculté des sciences, Université Mohammed V, Rabat; (^f)Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco.

- ³⁶CERN, Geneva; Switzerland.
- ³⁷Affiliated with an institute covered by a cooperation agreement with CERN.
- ³⁸Affiliated with an international laboratory covered by a cooperation agreement with CERN.
- ³⁹Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America.
- ⁴⁰LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France.
- ⁴¹Nevis Laboratory, Columbia University, Irvington NY; United States of America.
- ⁴²Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark.
- ⁴³(^a)Dipartimento di Fisica, Università della Calabria, Rende;(^b)INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy.
- ⁴⁴Physics Department, Southern Methodist University, Dallas TX; United States of America.
- ⁴⁵Physics Department, University of Texas at Dallas, Richardson TX; United States of America.
- ⁴⁶National Centre for Scientific Research "Demokritos", Agia Paraskevi; Greece.
- ⁴⁷(^a)Department of Physics, Stockholm University;(^b)Oskar Klein Centre, Stockholm; Sweden.
- ⁴⁸Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany.
- ⁴⁹Fakultät Physik , Technische Universität Dortmund, Dortmund; Germany.
- ⁵⁰Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany.
- ⁵¹Department of Physics, Duke University, Durham NC; United States of America.
- ⁵²SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom.
- ⁵³INFN e Laboratori Nazionali di Frascati, Frascati; Italy.
- ⁵⁴Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany.
- ⁵⁵II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany.
- ⁵⁶Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.
- ⁵⁷(^a)Dipartimento di Fisica, Università di Genova, Genova;(^b)INFN Sezione di Genova; Italy.
- ⁵⁸II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany.
- ⁵⁹SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom.
- ⁶⁰LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France.
- ⁶¹Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America.
- ⁶²(^a)Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei;(^b)Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao;(^c)School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai;(^d)Tsung-Dao Lee Institute, Shanghai;(^e)School of Physics and Microelectronics, Zhengzhou University; China.
- ⁶³(^a)Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg;(^b)Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany.
- ⁶⁴(^a)Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong;(^b)Department of Physics, University of Hong Kong, Hong Kong;(^c)Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China.
- ⁶⁵Department of Physics, National Tsing Hua University, Hsinchu; Taiwan.
- ⁶⁶IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France.
- ⁶⁷Centro Nacional de Microelectrónica (IMB-CNM-CSIC), Barcelona; Spain.
- ⁶⁸Department of Physics, Indiana University, Bloomington IN; United States of America.
- ⁶⁹(^a)INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine;(^b)ICTP, Trieste;(^c)Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy.
- ⁷⁰(^a)INFN Sezione di Lecce;(^b)Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy.
- ⁷¹(^a)INFN Sezione di Milano;(^b)Dipartimento di Fisica, Università di Milano, Milano; Italy.

- 72^(a) INFN Sezione di Napoli; ^(b) Dipartimento di Fisica, Università di Napoli, Napoli; Italy.
- 73^(a) INFN Sezione di Pavia; ^(b) Dipartimento di Fisica, Università di Pavia, Pavia; Italy.
- 74^(a) INFN Sezione di Pisa; ^(b) Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy.
- 75^(a) INFN Sezione di Roma; ^(b) Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy.
- 76^(a) INFN Sezione di Roma Tor Vergata; ^(b) Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy.
- 77^(a) INFN Sezione di Roma Tre; ^(b) Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy.
- 78^(a) INFN-TIFPA; ^(b) Università degli Studi di Trento, Trento; Italy.
- 79 Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck; Austria.
- 80 University of Iowa, Iowa City IA; United States of America.
- 81 Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America.
- 82 Istinye University, Sariyer, Istanbul; Türkiye.
- 83^(a) Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; ^(b) Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; ^(c) Instituto de Física, Universidade de São Paulo, São Paulo; ^(d) Rio de Janeiro State University, Rio de Janeiro; Brazil.
- 84 KEK, High Energy Accelerator Research Organization, Tsukuba; Japan.
- 85 Graduate School of Science, Kobe University, Kobe; Japan.
- 86^(a) AGH University of Krakow, Faculty of Physics and Applied Computer Science, Krakow; ^(b) Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland.
- 87 Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland.
- 88 Faculty of Science, Kyoto University, Kyoto; Japan.
- 89 Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka ; Japan.
- 90 Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina.
- 91 Physics Department, Lancaster University, Lancaster; United Kingdom.
- 92 Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom.
- 93 Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia.
- 94 School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom.
- 95 Department of Physics, Royal Holloway University of London, Egham; United Kingdom.
- 96 Department of Physics and Astronomy, University College London, London; United Kingdom.
- 97 Louisiana Tech University, Ruston LA; United States of America.
- 98 Fysiska institutionen, Lunds universitet, Lund; Sweden.
- 99 Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain.
- 100 Institut für Physik, Universität Mainz, Mainz; Germany.
- 101 School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom.
- 102 CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France.
- 103 Department of Physics, University of Massachusetts, Amherst MA; United States of America.
- 104 Department of Physics, McGill University, Montreal QC; Canada.
- 105 School of Physics, University of Melbourne, Victoria; Australia.
- 106 Department of Physics, University of Michigan, Ann Arbor MI; United States of America.
- 107 Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.
- 108 Group of Particle Physics, University of Montreal, Montreal QC; Canada.
- 109 Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany.
- 110 Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany.

- ¹¹¹Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan.
- ¹¹²Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America.
- ¹¹³Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen; Netherlands.
- ¹¹⁴Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands.
- ¹¹⁵Department of Physics, Northern Illinois University, DeKalb IL; United States of America.
- ¹¹⁶^(a)New York University Abu Dhabi, Abu Dhabi;^(b)University of Sharjah, Sharjah; United Arab Emirates.
- ¹¹⁷Department of Physics, New York University, New York NY; United States of America.
- ¹¹⁸Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan.
- ¹¹⁹Ohio State University, Columbus OH; United States of America.
- ¹²⁰Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America.
- ¹²¹Department of Physics, Oklahoma State University, Stillwater OK; United States of America.
- ¹²²Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic.
- ¹²³Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America.
- ¹²⁴Graduate School of Science, Osaka University, Osaka; Japan.
- ¹²⁵Department of Physics, University of Oslo, Oslo; Norway.
- ¹²⁶Department of Physics, Oxford University, Oxford; United Kingdom.
- ¹²⁷LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France.
- ¹²⁸Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America.
- ¹²⁹Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America.
- ¹³⁰^(a)Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisboa;^(b)Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa;^(c)Departamento de Física, Universidade de Coimbra, Coimbra;^(d)Centro de Física Nuclear da Universidade de Lisboa, Lisboa;^(e)Departamento de Física, Universidade do Minho, Braga;^(f)Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain);^(g)Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal.
- ¹³¹Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic.
- ¹³²Czech Technical University in Prague, Prague; Czech Republic.
- ¹³³Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic.
- ¹³⁴Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom.
- ¹³⁵IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France.
- ¹³⁶Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America.
- ¹³⁷^(a)Departamento de Física, Pontificia Universidad Católica de Chile, Santiago;^(b)Millennium Institute for Subatomic physics at high energy frontier (SAPHIR), Santiago;^(c)Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena;^(d)Universidad Andres Bello, Department of Physics, Santiago;^(e)Instituto de Alta Investigación, Universidad de Tarapacá, Arica;^(f)Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile.
- ¹³⁸Department of Physics, University of Washington, Seattle WA; United States of America.
- ¹³⁹Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom.
- ¹⁴⁰Department of Physics, Shinshu University, Nagano; Japan.

- ¹⁴¹Department Physik, Universität Siegen, Siegen; Germany.
- ¹⁴²Department of Physics, Simon Fraser University, Burnaby BC; Canada.
- ¹⁴³SLAC National Accelerator Laboratory, Stanford CA; United States of America.
- ¹⁴⁴Department of Physics, Royal Institute of Technology, Stockholm; Sweden.
- ¹⁴⁵Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America.
- ¹⁴⁶Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom.
- ¹⁴⁷School of Physics, University of Sydney, Sydney; Australia.
- ¹⁴⁸Institute of Physics, Academia Sinica, Taipei; Taiwan.
- ¹⁴⁹^(a)E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi;^(b)High Energy Physics Institute, Tbilisi State University, Tbilisi;^(c)University of Georgia, Tbilisi; Georgia.
- ¹⁵⁰Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel.
- ¹⁵¹Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel.
- ¹⁵²Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece.
- ¹⁵³International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan.
- ¹⁵⁴Department of Physics, Tokyo Institute of Technology, Tokyo; Japan.
- ¹⁵⁵Department of Physics, University of Toronto, Toronto ON; Canada.
- ¹⁵⁶^(a)TRIUMF, Vancouver BC;^(b)Department of Physics and Astronomy, York University, Toronto ON; Canada.
- ¹⁵⁷Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan.
- ¹⁵⁸Department of Physics and Astronomy, Tufts University, Medford MA; United States of America.
- ¹⁵⁹United Arab Emirates University, Al Ain; United Arab Emirates.
- ¹⁶⁰Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America.
- ¹⁶¹Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden.
- ¹⁶²Department of Physics, University of Illinois, Urbana IL; United States of America.
- ¹⁶³Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia; Spain.
- ¹⁶⁴Department of Physics, University of British Columbia, Vancouver BC; Canada.
- ¹⁶⁵Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada.
- ¹⁶⁶Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany.
- ¹⁶⁷Department of Physics, University of Warwick, Coventry; United Kingdom.
- ¹⁶⁸Waseda University, Tokyo; Japan.
- ¹⁶⁹Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel.
- ¹⁷⁰Department of Physics, University of Wisconsin, Madison WI; United States of America.
- ¹⁷¹Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany.
- ¹⁷²Department of Physics, Yale University, New Haven CT; United States of America.
- ^a Also Affiliated with an institute covered by a cooperation agreement with CERN.
- ^b Also at An-Najah National University, Nablus; Palestine.
- ^c Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America.
- ^d Also at Center for High Energy Physics, Peking University; China.
- ^e Also at Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Thessaloniki; Greece.
- ^f Also at Centro Studi e Ricerche Enrico Fermi; Italy.
- ^g Also at CERN, Geneva; Switzerland.

- h* Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.
- i* Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona; Spain.
- j* Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece.
- k* Also at Department of Physics, Ben Gurion University of the Negev, Beer Sheva; Israel.
- l* Also at Department of Physics, California State University, Sacramento; United States of America.
- m* Also at Department of Physics, King's College London, London; United Kingdom.
- n* Also at Department of Physics, Stanford University, Stanford CA; United States of America.
- o* Also at Department of Physics, University of Fribourg, Fribourg; Switzerland.
- p* Also at Department of Physics, University of Thessaly; Greece.
- q* Also at Department of Physics, Westmont College, Santa Barbara; United States of America.
- r* Also at Hellenic Open University, Patras; Greece.
- s* Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain.
- t* Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany.
- u* Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia; Bulgaria.
- v* Also at Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco.
- w* Also at Institute of Particle Physics (IPP); Canada.
- x* Also at Institute of Physics and Technology, Mongolian Academy of Sciences, Ulaanbaatar; Mongolia.
- y* Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.
- z* Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia.
- aa* Also at L2IT, Université de Toulouse, CNRS/IN2P3, UPS, Toulouse; France.
- ab* Also at Lawrence Livermore National Laboratory, Livermore; United States of America.
- ac* Also at National Institute of Physics, University of the Philippines Diliman (Philippines); Philippines.
- ad* Also at Technical University of Munich, Munich; Germany.
- ae* Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China.
- af* Also at TRIUMF, Vancouver BC; Canada.
- ag* Also at Università di Napoli Parthenope, Napoli; Italy.
- ah* Also at University of Chinese Academy of Sciences (UCAS), Beijing; China.
- ai* Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America.
- aj* Associated at University of Copenhagen, Copenhagen ; Denmark.
- ak* Also at Washington College, Chestertown, MD; United States of America.
- al* Also at Yeditepe University, Physics Department, Istanbul; Türkiye.
- * Deceased