

CERN-EP-2024-019
25 January 2024

Observation of the $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ decay

LHCb collaboration[†]

Abstract

The first observation of the $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ decay is reported with high significance using proton-proton collision data, corresponding to an integrated luminosity of 9 fb^{-1} , collected with the LHCb detector at centre-of-mass energies of 7, 8, and 13 TeV. The ratio of its branching fraction relative to the $B_c^+ \rightarrow J/\psi \pi^+$ channel is measured to be

$$\frac{\mathcal{B}_{B_c^+ \rightarrow J/\psi \pi^+ \pi^0}}{\mathcal{B}_{B_c^+ \rightarrow J/\psi \pi^+}} = 2.80 \pm 0.15 \pm 0.11 \pm 0.16,$$

where the first uncertainty is statistical, the second systematic and the third related to imprecise knowledge of the branching fractions for $B^+ \rightarrow J/\psi K^{*+}$ and $B^+ \rightarrow J/\psi K^+$ decays, which are used to determine the π^0 detection efficiency. The $\pi^+ \pi^0$ mass spectrum is found to be consistent with the dominance of an intermediate ρ^+ contribution in accordance with a model based on QCD factorisation.

Submitted to JHEP

© 2024 CERN for the benefit of the LHCb collaboration. CC BY 4.0 licence.

[†]Authors are listed at the end of this paper.

1 Introduction

The B_c^+ meson, consisting of two heavy quarks of different flavours, was discovered by the CDF collaboration [1,2] at the Tevatron collider, and is the heaviest meson that can decay only through the weak interaction. The large b-quark production cross-section at the Large Hadron Collider (LHC) [3–8] enables the ATLAS, CMS and LHCb experiments to study in detail the production, decays and other properties of the B_c^+ meson [9–39].

The B_c^+ decay may proceed through the weak decay of either of its heavy constituents with the other quark playing the role of spectator, or by a $\bar{b}c$ annihilation into a virtual W^+ boson. The lifetime of the B_c^+ meson is about three times smaller [20,23,32] than that of the B^0 and B^+ mesons [40], confirming the important role played by the charm quark in B_c^+ meson decays. Its decay to charmonium and light hadrons proceeds via the spectator diagram as shown in Fig. 1, which is described by the quantum chromodynamics (QCD) factorisation approach [41,42] using the form factors of the $B_c^+ \rightarrow [c\bar{c}]W^+$ transition [43–47] and the universal spectral function for the transition of the virtual W^+ boson into light hadrons [48–51]. This approach works remarkably well for all known decays of the B_c^+ meson into charmonia with an odd number of light hadrons in the final state [9–17].




Figure 1: Diagram for the decays of the B_c^+ meson into the J/ψ meson and light hadrons.

The $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ mode, which is the simplest decay into a charmonium state and an even number of light hadrons, has not yet been observed. Its final state is expected to be dominated by the $B_c^+ \rightarrow J/\psi\rho^+$ mode with a small admixture of $B_c^+ \rightarrow J/\psi\rho(1450)^+$ decays [52]. There are many predictions for the branching fraction ratio \mathcal{R} of this decay relative to the $B_c^+ \rightarrow J/\psi\pi^+$ mode,

$$\mathcal{R} \equiv \frac{\mathcal{B}_{B_c^+ \rightarrow J/\psi\pi^+\pi^0}}{\mathcal{B}_{B_c^+ \rightarrow J/\psi\pi^+}}, \quad (1)$$

which vary from 2.5 to 5.7 [46,48,53–69]. This decay is an important background source for the study of other decays of B_c^+ mesons [39] as well as rare decays of B^0 mesons [70].

This paper reports on the first observation of $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ decays¹ and the measurement of \mathcal{R} . A sample of the $B_c^+ \rightarrow J/\psi\pi^+$ mode is used for normalisation and a high-yield sample of $B^+ \rightarrow J/\psi(K^{*+} \rightarrow K^+\pi^0)$ decays is used to correct the detector resolution and any potential mass bias. The analysis is performed using proton-proton (pp) collision data,

¹In this paper inclusion of charge-conjugate decays is implied.

corresponding to an integrated luminosity of 9 fb^{-1} , collected with the LHCb detector at centre-of-mass energies of 7, 8, and 13 TeV.

2 Detector and simulation

The LHCb detector [71, 72] is a single-arm forward spectrometer covering the pseudo-rapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region [73], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 T m , and three stations of silicon-strip detectors and straw drift tubes [74, 75] placed downstream of the magnet. The tracking system provides a measurement of the momentum p of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at $200 \text{ GeV}/c$. The minimum distance of a track to a primary pp collision vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T) \mu\text{m}$, where p_T is the component of the momentum transverse to the beam in GeV/c . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov (RICH) detectors [76]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [77].

The online event selection is performed by a trigger [78], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. The hardware trigger selects muon candidates with high transverse momentum or dimuon candidates with a high value of the product of the transverse momenta of the two muons. In the software trigger, two oppositely charged muons are required to form a good-quality vertex that is significantly displaced from any PV, and the mass of the $\mu^+\mu^-$ pair is required to exceed $2.7 \text{ GeV}/c^2$.

Simulated samples of $B_c^+ \rightarrow J/\psi\pi^+\pi^0$, $B^+ \rightarrow J/\psi (K^{*+} \rightarrow K^+\pi^0)$ and $B_c^+ \rightarrow J/\psi\pi^+$ decays are used to model the signal mass shapes, optimise the selection requirements and compute the efficiencies needed to determine the branching fraction ratios. The PYTHIA [79] generator with a specific LHCb configuration [80] is used to simulate B^+ meson production from pp collisions. The BCVEGPy generator [81–84], which is interfaced to the PYTHIA parton shower and hadronisation model, is used to simulate B_c^+ meson production. The generator implements the complete lowest-order (α_s^4) perturbative QCD calculations of the dominant gluon-gluon fusion $gg \rightarrow B_c^+ + \bar{c} + b$ and $gg \rightarrow B_c^{*+} + \bar{c} + b$ processes, neglecting the contribution from the quark-pair annihilation $q\bar{q} \rightarrow B_c^+ + \bar{c} + b$ and $q\bar{q} \rightarrow B_c^{*+} + \bar{c} + b$ channels [85–89].

Decays of unstable particles are described by the EVTGEN package [90], in which final-state radiation is generated using PHOTOS [91]. The “BLL” model for $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ decays [48–51, 92] assumes the factorisation into $B_c^+ \rightarrow J/\psi W^{*+}$ with a subsequent $W^{*+} \rightarrow \rho^+$ transition with a tiny fraction of $W^{*+} \rightarrow \rho(1450)^+$ in accordance with the model by Kühn and Santamaria [93] and the measurement of the hadronic structure of the $\tau^+ \rightarrow \pi^+\pi^0\bar{\nu}_\tau$ decay by the CLEO collaboration [94]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [95] as described in Ref. [96]. The transverse momentum and rapidity spectra of

the B_c^+ mesons in simulated samples are adjusted to match those observed in a high-yield, low-background sample of reconstructed $B_c^+ \rightarrow J/\psi \pi^+$ decays. The detector response used for the identification of pions and kaons is sampled from $D^{*+} \rightarrow (D^0 \rightarrow K^- \pi^+) \pi^+$ and $K_S^0 \rightarrow \pi^+ \pi^-$ calibration channels [76, 97]. To account for imperfections in the simulation of charged-particle reconstruction, the track-reconstruction efficiency determined from simulation is corrected using calibration samples of $J/\psi \rightarrow \mu^+ \mu^-$ decays [98]. Samples of $B^+ \rightarrow J/\psi (K^{*+} \rightarrow K^+ (\pi^0 \rightarrow \gamma\gamma))$ decays are used to correct the photon reconstruction efficiency in the simulation [99–103].

3 Event selection

The signal $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ and control $B^+ \rightarrow J/\psi (K^{*+} \rightarrow K^+ \pi^0)$ candidates are reconstructed using $J/\psi \rightarrow \mu^+ \mu^-$ decays. The same dimuon final state of the J/ψ meson is used to reconstruct the normalisation $B_c^+ \rightarrow J/\psi \pi^+$ candidates. A loose preselection similar to that used in Refs. [14, 39, 104] is applied, followed by a multivariate classifier to select a higher purity subset of candidates.

Muons, pions and kaons are identified by combining information from the RICH, calorimeter and muon detectors, and they are required to have transverse momenta larger than $500 \text{ MeV}/c$. Pions and kaons are further required to have a momentum between 3.2 and $150 \text{ GeV}/c$ to ensure good performance of the particle identification in the RICH detectors [76, 105]. To reduce the combinatorial background due to charged particles produced in pp interactions, only tracks that are inconsistent with originating from any PV are used. Photons are reconstructed from clusters in the electromagnetic calorimeter with transverse energy above 300 MeV . The clusters must not be associated with reconstructed tracks [106, 107]. Photon identification is based on the combined information from the calorimeter and tracking systems.

Pairs of oppositely charged muons consistent with originating from a common vertex are combined to form $J/\psi \rightarrow \mu^+ \mu^-$ candidates. The reconstructed mass of the $\mu^+ \mu^-$ pair is required to be within the range $3.0 < m_{\mu^+ \mu^-} < 3.2 \text{ GeV}/c^2$. The position of the reconstructed dimuon vertex is required to be separated from any reconstructed PV. The π^0 candidates are reconstructed as diphoton pairs with mass within $30 \text{ MeV}/c^2$ of the known mass of the π^0 meson [40].

The selected $J/\psi \pi^+ \pi^0$, $J/\psi \pi^+$ and $J/\psi K^+ \pi^0$ combinations are used to form the B_c^+ and B^+ candidates. The $\pi^+ \pi^0$ mass $m_{\pi^+ \pi^0}$ is required to satisfy $620 < m_{\pi^+ \pi^0} < 920 \text{ MeV}/c^2$ and the $K^+ \pi^0$ mass must be within $50 \text{ MeV}/c^2$ of the known mass of the K^{*+} meson [40]. To improve the mass resolution for the B mesons, a kinematic fit [108] constrains the masses of the J/ψ and π^0 candidates to their known values [40] and requires the B candidates to originate from their associated PV.² The decay time t_B of the B_c^+ (B^+) candidates is required to satisfy $0.1 < t_B < 1.0 \text{ mm}/c$ ($0.2 < t_B < 2.0 \text{ mm}/c$) to suppress the large combinatorial background from tracks produced at the PV and from misreconstructed B candidates. The $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ candidates with a $J/\psi \pi^+$ mass within $45 \text{ MeV}/c^2$ of the known mass of the B_c^+ meson are vetoed to avoid contamination from $B_c^+ \rightarrow J/\psi \pi^+$ decays with a random π^0 added. Similarly, candidates with the $J/\psi \pi^+$ mass between 5.180 and $5.305 \text{ GeV}/c^2$ are

²Each B_c^+ or B^+ candidate is associated with the PV that yields the smallest χ_{IP}^2 , which is defined as the difference in the vertex-fit χ^2 of a given PV reconstructed with and without the particle under consideration.

also vetoed to remove the background from $B^+ \rightarrow J/\psi\pi^+$ and $B^+ \rightarrow J/\psi K^+$ decays with a random π^0 added.

Further selection of the B_c^+ and B^+ candidates is based on a multivariate estimator known as a multi-layer perceptron (MLP) classifier. The MLP classifier is based on an artificial neural network algorithm [109, 110] configured with a cross-entropy cost estimator [111]. It reduces the combinatorial background to a low level while retaining a high signal efficiency. Three MLP classifiers are trained separately for the signal, normalisation and control candidates. The variables used in the classifiers include those related to the reconstruction quality, kinematics and decay time of the B_c^+ and B^+ candidates, kinematics of the final-state hadrons and photons, as well as a variable that characterises the identification of charged pions and kaons. The classifiers are trained using simulated signal samples, while the B_c^+ and B^+ candidates from data with mass outside the regions $6.00 < m_{J/\psi\pi^+\pi^0} < 6.50 \text{ GeV}/c^2$, $6.20 < m_{J/\psi\pi^+} < 6.34 \text{ GeV}/c^2$ and $5.10 < m_{J/\psi K^+\pi^0} < 5.50 \text{ GeV}/c^2$ are used to represent the background for the signal, normalisation and control modes, respectively. The k -fold cross-validation technique [112] with $k = 11$ is used to avoid introducing a bias in the MLP evaluation. The requirement on each of the MLP classifiers is chosen to maximise the figure-of-merit defined as $S/\sqrt{S + \kappa B}$, where S represents the expected signal yield, B is the background yield and κ is the fraction of the total background that affects the signal determination. The background yield B is calculated from fits to data, while $S = \varepsilon S_0$, where ε is the efficiency of the requirement on the response of the MLP classifier determined from simulation and S_0 is the signal yield obtained from the fit to the data with a loose requirement applied. The factor κ is determined from pseudoexperiments. The mass distributions for the selected $B_c^+ \rightarrow J/\psi\pi^+$, $B^+ \rightarrow J/\psi K^{*+}$ and $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ candidates are shown in Figs. 2, 3 and 4, respectively.

4 Determination of signal yields

The yields for the normalisation $B_c^+ \rightarrow J/\psi\pi^+$, control $B^+ \rightarrow J/\psi K^{*+}$ and signal $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ decays are determined using extended unbinned maximum-likelihood fits to the $J/\psi\pi^+$, $J/\psi K^+\pi^0$ and $J/\psi\pi^+\pi^0$ mass spectra shown in Figs. 2, 3 and 4. Each signal component is parameterised as the sum of a modified Gaussian function with power law tails on both sides of the distribution [113, 114] and a standard Gaussian function with common location parameter.³ The background components are parameterised by positive decreasing second-order polynomials for the fits to the $J/\psi\pi^+$ and $J/\psi K^+\pi^0$ mass spectra and with a positive convex decreasing fourth-order polynomial for the fit to the $J/\psi\pi^+\pi^0$ mass spectrum [115]. In addition, the fit model for the $J/\psi\pi^+$ mass spectrum includes a small component corresponding to the contribution from Cabibbo-suppressed $B_c^+ \rightarrow J/\psi K^+$ decays [11, 17] where the charged kaon is reconstructed as a pion. The shape of this component is taken from simulation. The fit model for the $J/\psi K^+\pi^0$ mass spectrum includes an additional component corresponding to contributions from $B^+ \rightarrow J/\psi K^+\pi^0\pi^0$ and $B^0 \rightarrow J/\psi K^+\pi^0\pi^-$ decays with missing π^0 or π^- particles. This component is parameterised using a Gaussian function. The signal shape parameters, except for the location parameters, are determined from simulation and their uncertainties are propagated to

³The resulting signal shapes are parameterised as a location-scale family, $f(\frac{x-\mu}{\sigma})$, using the location parameter μ and the scale parameter σ , referred to as the resolution parameter hereafter.




Figure 2: Mass distribution for selected $B_c^+ \rightarrow J/\psi\pi^+$ candidates with the result of the fit described in the text.

the fits using multivariate Gaussian constraints. For each fit, the location and resolution parameters are expressed as

$$\mu = \delta m_B + m_B, \quad (2a)$$

$$\sigma = s \times \sigma_{MC}, \quad (2b)$$

where m_B is the known mass of the corresponding beauty meson [40], σ_{MC} is the resolution parameter obtained from simulation, the term δm_B corrects for a possible mass bias, and the scale factor s takes into account any difference in the mass resolution between simulation and data. The parameters δm_B and s are allowed to vary in the $J/\psi\pi^+$ and $J/\psi K^+\pi^0$ mass spectra fits, while the $J/\psi\pi^+\pi^0$ fit restricts them to the values obtained for the $B^+ \rightarrow J/\psi K^+\pi^0$ mode with their corresponding uncertainties propagated using Gaussian constraints.

The results of the fits are shown in Figs. 2, 3 and 4 with the parameters summarised in Table 1. The statistical significance for the $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ signal is estimated using Wilks' theorem [116] and is found to exceed 20 standard deviations. To validate the observed $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ signal, several cross-checks are performed. The data are split into data-taking periods with different polarity of the dipole magnet and into B_c^+ and B_c^- samples. Alternative multivariate estimators, such as decision trees with gradient




Figure 3: Mass distribution for selected $B^+ \rightarrow J/\psi K^+ \pi^0$ candidates with the result of the fit described in the text.

boosting [117] and a projective likelihood estimator [118], are used. The results are found to be consistent among all samples and analysis techniques.

To validate the predictions of the BLL model, the $\pi^+ \pi^0$ mass spectrum from $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ decays is studied by extending the $\pi^+ \pi^0$ mass interval to the region $m_{\pi^+ \pi^0} < 1.6 \text{ GeV}/c^2$. The background-subtracted $\pi^+ \pi^0$ mass spectrum is shown in Fig. 5, which uses the *sPlot* technique [119] based on the fit to the $J/\psi \pi^+ \pi^0$ mass

Table 1: Yields N , mass biases δm_B and resolution scale factors s from the fits to the $J/\psi \pi^+$, $J/\psi K^+ \pi^0$ and $J/\psi \pi^+ \pi^0$ mass spectra. The uncertainties for δm_B include those from the known masses of the B_c^+ and B^+ mesons. The mass bias and resolution scale parameters for the $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ channel are constrained by those from the $B^+ \rightarrow J/\psi K^+ \pi^0$ control channel.

Channel	N [10^3]	δm_B [MeV/c^2]	s
$B_c^+ \rightarrow J/\psi \pi^+$	16.11 ± 0.15	-0.59 ± 0.35	1.10 ± 0.01
$B^+ \rightarrow J/\psi K^+ \pi^0$	81.91 ± 0.52	-0.03 ± 0.20	1.11 ± 0.01
$B_c^+ \rightarrow J/\psi \pi^+ \pi^0$	1.08 ± 0.06	-0.11 ± 0.49	1.11 ± 0.01




Figure 4: Mass distribution for selected $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ candidates with the result of the fit, described in the text.

spectrum with the wider $\pi^+\pi^0$ mass interval. There is a clear peak corresponding to the decay $\rho^+ \rightarrow \pi^+\pi^0$, as expected by the BLL model. To quantify the possible deviations from the predictions, a fit to the $\pi^+\pi^0$ mass spectrum is performed using a function consisting of two components. The signal component corresponds to the coherent sum of the $B_c^+ \rightarrow J/\psi\rho^+$ and $B_c^+ \rightarrow J/\psi\rho(1450)^+$ contributions [93], with the shape obtained from the BLL model. The second component describes contributions differing from the BLL model, generically parameterised as

$$f(m_{\pi^+\pi^0}) \propto q^3(m_{\pi^+\pi^0})p(m_{\pi^+\pi^0})\mathcal{P}_1^+(m_{\pi^+\pi^0}), \quad (3)$$

where q is the pion momentum in the dipion rest-frame, p is the $\pi^+\pi^0$ momentum in the B_c^+ rest-frame [120], and \mathcal{P}_1^+ is a positive first-order polynomial function [115]. This parameterisation accounts for the phase-space terms and the suppression of the S-wave contribution. The unknown decay dynamics are absorbed by the polynomial function \mathcal{P}_1^+ . The result of the fit to the background-subtracted $\pi^+\pi^0$ mass spectrum is superimposed in Fig. 5. The contribution of the second component vanishes in the fit, demonstrating good agreement between the observed $\pi^+\pi^0$ mass spectrum and the expectations from the BLL model [48–51, 92].




Figure 5: Background-subtracted $\pi^+\pi^0$ mass distribution from the $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ decays. The orange line shows the result of the two-component fit described in the text. The blue dashed line corresponds to the component parameterised by Eq. (3), which vanishes in the fit.

5 Branching fraction ratio computation

The ratio of branching fractions for the $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ and $B_c^+ \rightarrow J/\psi\pi^+$ decays is calculated as

$$\mathcal{R} = \frac{N_{B_c^+ \rightarrow J/\psi\pi^+\pi^0}}{N_{B_c^+ \rightarrow J/\psi\pi^+}} \bigg/ \frac{\varepsilon_{B_c^+ \rightarrow J/\psi\pi^+\pi^0}}{\varepsilon_{B_c^+ \rightarrow J/\psi\pi^+}}, \quad (4)$$

where the yields $N_{B_c^+ \rightarrow J/\psi\pi^+\pi^0}$ and $N_{B_c^+ \rightarrow J/\psi\pi^+}$ are taken from Table 1, and ε denotes the corresponding efficiency, which is defined as the product of the detector acceptance and the reconstruction, selection and trigger efficiencies, where each subsequent efficiency is defined with respect to the previous one. Each of the partial efficiencies is calculated using the simulation samples described in Sec. 2. The ratio of branching fractions is found to be

$$\mathcal{R} = 2.80 \pm 0.15,$$

where the uncertainty is statistical only.

6 Systematic uncertainties

Many sources of systematic uncertainties cancel in the ratio of branching fractions \mathcal{R} due to the similarity between the signal and normalisation decay channels. The remaining contributions to the systematic uncertainty for \mathcal{R} are summarised in Table 2 and discussed below.

An important source of systematic uncertainty is the imperfect knowledge of the shapes of the signal and background components used in the fits. To estimate this effect, several alternative models are tested. The $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ signal shape is parameterised as a sum of a Gaussian and three alternative functions:

1. a bifurcated generalised Student's t -distribution [121, 122];
2. a Johnson's \mathcal{S}_U -distribution [123, 124];
3. a generalised hyperbolic distribution [125, 126].

Similarly, the $B_c^+ \rightarrow J/\psi\pi^+$ signal component is modelled as the sum of a Gaussian function with three alternative shapes:

1. a bifurcated Student's t -distribution;
2. a generalised hyperbolic distribution;
3. a Pearson's type IV distribution [127].

Five alternative background functions are tested for the fit to the $J/\psi\pi^+\pi^0$ mass spectrum:

Table 2: Relative systematic uncertainties for the ratio \mathcal{R} . The total systematic uncertainty is the quadratic sum of individual contributions. The systematic uncertainty of the π^0 reconstruction efficiency due to imprecise knowledge of the branching fraction of the $B^+ \rightarrow J/\psi K^{*+}$ and $B^+ \rightarrow J/\psi K^+$ decays is treated separately.

Source	Uncertainty [%]
Fit model	
Signal shapes	1.0
Background shapes	2.1
Efficiency determination	
B_c^+ kinematics	< 0.1
B_c^+ decay model	2.2
Track reconstruction efficiency	0.1
π^0 reconstruction efficiency	0.9
Hadron identification	0.6
Trigger efficiency	1.1
Data-simulation difference	2.0
Size of simulated samples	0.5
Total	4.1

1. a positive decreasing convex third-order polynomial function [115];
2. a positive decreasing fourth-order polynomial function;
3. a product of an exponential function and a positive third-order polynomial function;
4. a sum of an exponential function and a positive third-order polynomial function;
5. a sum of a generalised Pareto distribution [128, 129] and a positive first-order polynomial function.

Lastly, for the background component of the $J/\psi\pi^+$ mass spectrum four alternative fit models are tested:

1. a positive decreasing third-order polynomial function;
2. a product of an exponential function and a positive first-order polynomial function;
3. a positive third-order polynomial function;
4. a positive fourth-order polynomial function.

For each alternative signal or background function, a large ensemble of pseudoexperiments is produced and fit with the baseline model. The absolute value of the mean over an ensemble for the relative difference between the fitted $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ or $B_c^+ \rightarrow J/\psi\pi^+$ signal yields and their baseline results is calculated. Its maximal value is found to be 1.0% (2.1%) for the alternative signal (background) shapes and is taken as the systematic uncertainty related to the fit model. For each model test, the signal significance of the newly observed $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ decay is estimated using Wilks' theorem [116] and is always found to exceed 20 standard deviations.

An uncertainty may originate from possible differences in the B_c^+ production kinematics between data and simulation. The transverse momentum and rapidity spectra of the B_c^+ mesons in the simulation are weighted to match those observed in a high-yield, low-background sample of $B_c^+ \rightarrow J/\psi\pi^+$ decays. The systematic uncertainty of the efficiency ratio caused by the finite size of this sample is estimated by varying the B_c^+ -meson production kinematic spectra within their uncertainties in the weighting procedure. The induced variation for the ratio \mathcal{R} is found to be smaller than 0.1%.

Simulated $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ events are corrected for polarisation of the dipion system in the B_c^+ rest frame to reproduce the corresponding angular distribution observed in data. Variations of this correction within statistical uncertainties causes a relative variation of 2.2% for the efficiency for $B_c^+ \rightarrow J/\psi\pi^+\pi^0$ decays, which is taken as the systematic uncertainty related to the B_c^+ decay model.

There are residual differences in the reconstruction efficiency of charged-particle tracks that do not cancel completely in the ratio due to slightly different kinematic distributions of the final-state particles. The track-finding efficiencies obtained from simulated samples are corrected using calibration channels [98]. The uncertainties related to the efficiency correction factors are propagated to the ratios of the total efficiencies using pseudoexperiments and found to be 0.1% for the ratio \mathcal{R} .

The difference of the photon reconstruction efficiency between data and simulation is studied using large samples of $B^+ \rightarrow J/\psi(K^{*+} \rightarrow K^+(\pi^0 \rightarrow \gamma\gamma))$ and $B^+ \rightarrow J/\psi K^+$ decays [99–103]. The uncertainty for the photon efficiency corrections has three components:

statistical, systematic and those related to the imprecise knowledge of the ratio of branching fractions for $B^+ \rightarrow J/\psi K^{*+}$ and $B^+ \rightarrow J/\psi K^+$ decays. The statistical and systematic uncertainties are propagated to the ratio of the total efficiencies using pseudoexperiments and the resulting 0.9% is taken as the systematic uncertainty related to the photon reconstruction. The uncertainty due to imprecise knowledge of external branching fractions of 5.9% [40] is treated separately.

The simulated detector response used for identification of pions is resampled from the control channels [76, 97]. The systematic uncertainty obtained through this procedure arises from the kernel shape used in the estimation of the probability density distributions. An alternative combined response is estimated using a modified kernel shape and the efficiency models are regenerated [130, 131]. The difference between the two efficiency ratios of 0.6% is taken as the systematic uncertainty related to hadron identification.

The systematic uncertainty related to the trigger efficiency has been previously studied by comparing the ratios of the trigger efficiencies in data and simulation using large samples of $B^+ \rightarrow J/\psi K^+$ and $B^+ \rightarrow \psi(2S)K^+$ decays [132]. Based on this comparison, a conservative estimate of 1.1% for the relative difference between data and simulation is taken as the corresponding systematic uncertainty.

The imperfect data description by simulation due to remaining effects not described above is studied by varying the MLP selection criteria for the normalisation decay $B_c^+ \rightarrow J/\psi \pi^+$. The observed maximal difference between the efficiency estimated using data and simulation does not exceed 2.0%. This value is taken as the corresponding systematic uncertainty for the ratio \mathcal{R} . The last systematic uncertainty considered is due to the finite size of the simulated samples, which amounts to 0.5%. The total systematic uncertainty is calculated as the quadratic sum of the individual contributions.

7 Results and discussion

The $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ decay is observed with overwhelming significance using proton-proton collision data, corresponding to an integrated luminosity of 9 fb^{-1} , collected with the LHCb detector at centre-of-mass energies of 7, 8, and 13 TeV. The ratio of the $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ branching fraction relative to the $B_c^+ \rightarrow J/\psi \pi^+$ channel is measured to be:

$$\mathcal{R} = \frac{\mathcal{B}_{B_c^+ \rightarrow J/\psi \pi^+ \pi^0}}{\mathcal{B}_{B_c^+ \rightarrow J/\psi \pi^+}} = 2.80 \pm 0.15 \pm 0.11 \pm 0.16, \quad (5)$$

where the first uncertainty is statistical, the second systematic and the third due to uncertainties in the $B^+ \rightarrow J/\psi K^{*+}$ and $B^+ \rightarrow J/\psi K^+$ branching fractions. The comparison of the measured ratio with theory predictions [46, 48, 53–69] is shown in Fig. 6. While most of the theory predictions agree well with the measurement, calculations from Refs. [65, 66] are only marginally consistent and results from Refs. [63, 64, 67] are inconsistent with the measured value. The study of the $\pi^+ \pi^0$ mass spectrum indicates that the $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ decay is saturated by the $B_c^+ \rightarrow J/\psi \rho^+$ contribution, with a small admixture of the $B_c^+ \rightarrow J/\psi \rho(1450)^+$ decay, in accordance with theoretical expectations [48–52, 92].




Figure 6: The measured ratio of the branching fractions of the $B_c^+ \rightarrow J/\psi \pi^+ \pi^0$ and $B_c^+ \rightarrow J/\psi \pi^+$ decays in comparison with theory predictions. The colored band shows the measured ratio with corresponding uncertainties added in quadrature.

Acknowledgements

This paper is dedicated to the memory of our dear friend and colleague Prof. Yury Zaitsev, who made an important contribution to this analysis. We thank A. K. Likhoded, X. Liu and A. V. Luchinsky for stimulating discussions on the decays of the B_c^+ meson. We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MCID/IFA (Romania); MICINN (Spain);

SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), and Polish WLCG (Poland). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from ARC and ARDC (Australia); Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); Minciencias (Colombia); EPLANET, Marie Skłodowska-Curie Actions, ERC and NextGenerationEU (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); AvH Foundation (Germany); ICSC (Italy); GVA, XuntaGal, GENCAT, Inditex, InTalent and Prog. Atracción Talento, CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society and UKRI (United Kingdom).

References

- [1] CDF collaboration, F. Abe *et al.*, *Observation of the B_c meson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV*, Phys. Rev. Lett. **81** (1998) 2432, [arXiv:hep-ex/9805034](#).
- [2] CDF collaboration, F. Abe *et al.*, *Observation of B_c mesons in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV*, Phys. Rev. **D58** (1998) 112004, [arXiv:hep-ex/9804014](#).
- [3] LHCb collaboration, R. Aaij *et al.*, *Measurement of $\sigma(pp \rightarrow b\bar{b}X)$ at $\sqrt{s} = 7$ TeV in the forward region*, Phys. Lett. **B694** (2010) 209, [arXiv:1009.2731](#).
- [4] LHCb collaboration, R. Aaij *et al.*, *Measurement of J/ψ production in pp collisions at $\sqrt{s} = 7$ TeV*, Eur. Phys. J. **C71** (2011) 1645, [arXiv:1103.0423](#).
- [5] LHCb collaboration, R. Aaij *et al.*, *Measurement of the B^\pm production cross-section in pp collisions at $\sqrt{s} = 7$ TeV*, JHEP **04** (2012) 093, [arXiv:1202.4812](#).
- [6] LHCb collaboration, R. Aaij *et al.*, *Measurement of B meson production cross-sections in proton-proton collisions at $\sqrt{s} = 7$ TeV*, JHEP **08** (2013) 117, [arXiv:1306.3663](#).
- [7] LHCb collaboration, R. Aaij *et al.*, *Production of J/ψ and Υ mesons in pp collisions at $\sqrt{s} = 8$ TeV*, JHEP **06** (2013) 064, [arXiv:1304.6977](#).
- [8] LHCb collaboration, R. Aaij *et al.*, *Measurement of forward J/ψ production cross-sections in pp collisions at $\sqrt{s} = 13$ TeV*, JHEP **10** (2015) 172, Erratum *ibid.* **05** (2017) 063, [arXiv:1509.00771](#).
- [9] LHCb collaboration, R. Aaij *et al.*, *First observation of the decay $B_c^+ \rightarrow J/\psi \pi^+ \pi^- \pi^+$* , Phys. Rev. Lett. **108** (2012) 251802, [arXiv:1204.0079](#).
- [10] LHCb collaboration, R. Aaij *et al.*, *Observation of the decay $B_c^+ \rightarrow \psi(2S) \pi^+$* , Phys. Rev. **D87** (2013) 071103(R), [arXiv:1303.1737](#).

- [11] LHCb collaboration, R. Aaij *et al.*, *First observation of the decay $B_c^+ \rightarrow J/\psi K^+$* , JHEP **09** (2013) 075, arXiv:1306.6723.
- [12] LHCb collaboration, R. Aaij *et al.*, *Observation of the decay $B_c^+ \rightarrow J/\psi K^+ K^- \pi^+$* , JHEP **11** (2013) 094, arXiv:1309.0587.
- [13] LHCb collaboration, R. Aaij *et al.*, *Evidence for the decay $B_c^+ \rightarrow J/\psi 3\pi^+ 2\pi^-$* , JHEP **05** (2014) 148, arXiv:1404.0287.
- [14] LHCb collaboration, R. Aaij *et al.*, *Study of B_c^+ decays to charmonia and three light hadrons*, JHEP **01** (2022) 065, arXiv:2111.03001.
- [15] LHCb collaboration, R. Aaij *et al.*, *Study of B_c^+ meson decays to charmonia plus multihadron final states*, JHEP **07** (2023) 198, arXiv:2208.08660.
- [16] LHCb collaboration, R. Aaij *et al.*, *Measurement of the branching fraction ratio $\mathcal{B}(B_c^+ \rightarrow \psi(2S)\pi^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)$* , Phys. Rev. **D92** (2015) 072007, arXiv:1507.03516.
- [17] LHCb collaboration, R. Aaij *et al.*, *Measurement of the ratio of branching fractions $\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)$* , JHEP **09** (2016) 153, arXiv:1607.06823.
- [18] LHCb collaboration, R. Aaij *et al.*, *Observation of $B_c^+ \rightarrow J/\psi D_s^+$ and $B_c^+ \rightarrow J/\psi D_s^{*+}$ decays*, Phys. Rev. **D87** (2013) 112012, arXiv:1304.4530.
- [19] LHCb collaboration, R. Aaij *et al.*, *Observation of the decay $B_c^+ \rightarrow B_s^0 \pi^+$* , Phys. Rev. Lett. **111** (2013) 181801, arXiv:1308.4544.
- [20] LHCb collaboration, R. Aaij *et al.*, *Measurement of the B_c^+ meson lifetime using $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu X$ decays*, Eur. Phys. J. **C74** (2014) 2839, arXiv:1401.6932.
- [21] LHCb collaboration, R. Aaij *et al.*, *First observation of a baryonic B_c^+ decay*, Phys. Rev. Lett. **113** (2014) 152003, arXiv:1408.0971.
- [22] LHCb collaboration, R. Aaij *et al.*, *Measurement of B_c^+ production in proton-proton collisions at $\sqrt{s} = 8$ TeV*, Phys. Rev. Lett. **114** (2015) 132001, arXiv:1411.2943.
- [23] LHCb collaboration, R. Aaij *et al.*, *Measurement of the lifetime of the B_c^+ meson using the $B_c^+ \rightarrow J/\psi\pi^+$ decay mode*, Phys. Lett. **B742** (2015) 29, arXiv:1411.6899.
- [24] CMS collaboration, V. Khachatryan *et al.*, *Measurement of the ratio of the production cross sections times branching fractions of $B_c^\pm \rightarrow J/\psi\pi^\pm$ and $B^\pm \rightarrow J/\psi K^\pm$ and $\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm\pi^\pm\pi^\mp)/\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm)$ in pp collisions at $\sqrt{s} = 8$ TeV*, JHEP **01** (2015) 063, arXiv:1410.5729.
- [25] ATLAS collaboration, G. Aad *et al.*, *Study of the $B_c^+ \rightarrow J/\psi D_s^+$ and $B_c^+ \rightarrow J/\psi D_s^{*+}$ decays with the ATLAS detector*, Eur. Phys. J. **C76** (2016) 24, arXiv:1507.07099.
- [26] LHCb collaboration, R. Aaij *et al.*, *Search for B_c^+ decays to the $p\bar{p}\pi^+$ final state*, Phys. Lett. **B759** (2016) 313, arXiv:1603.07037.

- [27] LHCb collaboration, R. Aaij *et al.*, *Study of B_c^+ decays to the $K^+K^-\pi^+$ final state and evidence for the decay $B_c^+ \rightarrow \chi_{c0}\pi^+$* , Phys. Rev. **D94** (2016) 091102(R), [arXiv:1607.06134](#).
- [28] LHCb collaboration, R. Aaij *et al.*, *Observation of $B_c^+ \rightarrow J/\psi D^{(*)}K^{(*)}$ decays*, Phys. Rev. **D95** (2017) 032005, [arXiv:1612.07421](#).
- [29] LHCb collaboration, R. Aaij *et al.*, *Observation of $B_c^+ \rightarrow D^0K^+$ decays*, Phys. Rev. Lett. **118** (2017) 111803, [arXiv:1701.01856](#).
- [30] LHCb collaboration, R. Aaij *et al.*, *Measurement of the ratio of branching fractions $\mathcal{B}(B_c^+ \rightarrow J/\psi\tau^+\nu_\tau)/\mathcal{B}(B_c^+ \rightarrow J/\psi\mu^+\nu_\mu)$* , Phys. Rev. Lett. **120** (2018) 121801, [arXiv:1711.05623](#).
- [31] LHCb collaboration, R. Aaij *et al.*, *Search for B_c^+ decays to two charm mesons*, Nucl. Phys. **B930** (2018) 563, [arXiv:1712.04702](#).
- [32] CMS collaboration, A. M. Sirunyan *et al.*, *Measurement of b hadron lifetimes in pp collisions at $\sqrt{s} = 8$ TeV*, Eur. Phys. J. **C78** (2018) 457, Erratum *ibid.* **C78** (2018) 561, [arXiv:1710.08949](#).
- [33] CMS collaboration, A. M. Sirunyan *et al.*, *Observation of two excited B_c^+ states and measurement of the $B_c^+(2S)$ mass in pp collisions at $\sqrt{s} = 13$ TeV*, Phys. Rev. Lett. **122** (2019) 132001, [arXiv:1902.00571](#).
- [34] LHCb collaboration, R. Aaij *et al.*, *Observation of an excited B_c^+ state*, Phys. Rev. Lett. **122** (2019) 232001, [arXiv:1904.00081](#).
- [35] LHCb collaboration, R. Aaij *et al.*, *Measurement of the B_c^- production fraction and asymmetry in 7 and 13 TeV pp collisions*, Phys. Rev. **D100** (2019) 112006, [arXiv:1910.13404](#).
- [36] LHCb collaboration, R. Aaij *et al.*, *Precision measurement of the B_c^+ meson mass*, JHEP **07** (2020) 123, [arXiv:2004.08163](#).
- [37] LHCb collaboration, R. Aaij *et al.*, *Updated search for B_c^+ decays to two charm mesons*, JHEP **12** (2021) 117, [arXiv:2109.00488](#).
- [38] LHCb collaboration, R. Aaij *et al.*, *Search for $B_c^+ \rightarrow \pi^+\mu^+\mu^-$ decays and measurement of the branching fraction ratio $\mathcal{B}(B_c^+ \rightarrow \psi(2S)\pi^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)$* , [arXiv:2312.12228](#), submitted to Eur. Phys. J. C.
- [39] LHCb collaboration, R. Aaij *et al.*, *Study of the $B_c^+ \rightarrow \chi_c\pi^+$ decays*, [arXiv:2312.12987](#), to appear in JHEP.
- [40] Particle Data Group, R. L. Workman *et al.*, *Review of particle physics*, Prog. Theor. Exp. Phys. **2022** (2022) 083C01, and 2023 update.
- [41] M. Bauer, B. Stech, and M. Wirbel, *Exclusive non-leptonic decays of D-, D_s-, and B-mesons*, Z. Phys. **C34** (1987) 103.

- [42] M. Wirbel, *Description of weak decays of D and B mesons*, Prog. Part. Nucl. Phys. **21** (1988) 33.
- [43] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, and A. V. Tkabladze, *Physics of B_c-mesons*, Phys. Usp. **38** (1995) 1, arXiv:hep-ph/9504319.
- [44] S. S. Gershtein *et al.*, *Theoretical status of the B_c meson*, in IV Workshop on Heavy Quark Physics, (Rostock, Germany), 1997, arXiv:hep-ph/9803433.
- [45] V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, *Semileptonic B_c meson decays in sum rules of QCD and NRQCD*, Nucl. Phys. **B569** (2000) 473, arXiv:hep-ph/9905359.
- [46] V. V. Kiselev, A. E. Kovalsky, and A. K. Likhoded, *B_c decays and lifetime in QCD sum rules*, Nucl. Phys. **B585** (2000) 353, arXiv:hep-ph/0002127.
- [47] D. Ebert, R. N. Faustov, and V. O. Galkin, *Properties of heavy quarkonia and B_c mesons in the relativistic quark model*, Phys. Rev. **D67** (2003) 014027, arXiv:hep-ph/0210381.
- [48] A. K. Likhoded and A. V. Luchinsky, *Light hadron production in B_c → J/ψ + X decays*, Phys. Rev. **D81** (2010) 014015, arXiv:0910.3089.
- [49] A. K. Likhoded and A. V. Luchinsky, *Production of a pion system in exclusive B_c → V(P) + nπ decays*, Phys. Atom. Nucl. **76** (2013) 787.
- [50] A. V. Berezhnoy, A. K. Likhoded, and A. V. Luchinsky, *B_c → J/ψ(B_s, B_s^{*}) + nπ decays*, PoS **QFTHEP2011** (2012) 076, arXiv:1111.5952.
- [51] A. V. Luchinsky, *Production of charged π mesons in exclusive B_c → V(P) + nπ decays*, Phys. Rev. **D86** (2012) 074024, arXiv:1208.1398.
- [52] C. Bruch, A. Khodjamirian, and J. H. Kuhn, *Modeling the pion and kaon form factors in the timelike region*, Eur. Phys. J. **C39** (2005) 41, arXiv:hep-ph/0409080.
- [53] C.-H. Chang and Y.-Q. Chen, *The decays of B_c meson*, Phys. Rev. **D49** (1994) 3399.
- [54] J.-F. Liu and K.-T. Chao, *B_c meson weak decays and CP violation*, Phys. Rev. **D56** (1997) 4133.
- [55] P. Colangelo and F. De Fazio, *Using heavy quark spin symmetry in semileptonic B_c decays*, Phys. Rev. **D61** (2000) 034012, arXiv:hep-ph/9909423.
- [56] A. Abd El-Hady, J. H. Munoz, and J. P. Vary, *Semileptonic and nonleptonic B_c decays*, Phys. Rev. **D62** (2000) 014019, arXiv:hep-ph/9909406.
- [57] V. V. Kiselev, A. E. Kovalsky, and A. K. Likhoded, *B_c-meson decays and lifetime within QCD sum rules*, Phys. Atom. Nucl. **64** (2001) 1860.
- [58] D. Ebert, R. N. Faustov, and V. O. Galkin, *Weak decays of the B_c meson to charmonium and D mesons in the relativistic quark model*, Phys. Rev. **D68** (2003) 094020, arXiv:hep-ph/0306306.

- [59] M. A. Ivanov, J. G. Korner, and P. Santorelli, *Exclusive semileptonic and nonleptonic decays of the B_c meson*, Phys. Rev. **D73** (2006) 054024, [arXiv:hep-ph/0602050](#).
- [60] E. Hernandez, J. Nieves, and J. M. Verde-Velasco, *Study of exclusive semileptonic and nonleptonic decays of B_c^- in a nonrelativistic quark model*, Phys. Rev. **D74** (2006) 074008, [arXiv:hep-ph/0607150](#).
- [61] W. Wang, Y.-L. Shen, and C.-D. Lu, *The study of $B_c^- \rightarrow X(3872)\pi^-(K^-)$ decays in the covariant light-front approach*, Eur. Phys. J. **C51** (2007) 841, [arXiv:0704.2493](#).
- [62] C.-F. Qiao, P. Sun, D. Yang, and R.-L. Zhu, *B_c exclusive decays to charmonium and a light meson at next-to-leading order accuracy*, Phys. Rev. **D89** (2014) 034008, [arXiv:1209.5859](#).
- [63] S. Naimuddin *et al.*, *Nonleptonic two-body B_c -meson decays*, Phys. Rev. **D86** (2012) 094028.
- [64] S. Kar *et al.*, *Nonleptonic $B_c^+ \rightarrow VV$ decays*, Phys. Rev. **D88** (2013) 094014.
- [65] Z. Rui and Z.-T. Zou, *S-wave ground state charmonium decays of B_c mesons in the perturbative QCD approach*, Phys. Rev. **D90** (2014) 114030, [arXiv:1407.5550](#).
- [66] A. Issadykov and M. A. Ivanov, *The decays $B_c \rightarrow J/\psi + \bar{\ell}\nu_\ell$ and $B_c \rightarrow J/\psi + \pi(K)$ in covariant confined quark model*, Phys. Lett. **B783** (2018) 178, [arXiv:1804.00472](#).
- [67] W. Cheng *et al.*, *$B_c \rightarrow J/\psi$ helicity form factors and the $B_c^+ \rightarrow J/\psi + (P, V, \ell^+\nu_\ell)$ decays*, Chin. Phys. **C46** (2022) 053103, [arXiv:2107.08405](#).
- [68] Z.-Q. Zhang *et al.*, *Covariant light-front approach for B_c decays into charmonium: Implications on form factors and branching ratios*, Eur. Phys. J. **C83** (2023) 477, [arXiv:2301.11107](#).
- [69] X. Liu, *The B_c -meson decays into J/ψ plus a light meson in the $iPQCD$ formalism*, [arXiv:2305.00713](#).
- [70] LHCb collaboration, R. Aaij *et al.*, *New measurement of the branching fraction of the $B^0 \rightarrow J/\psi\pi^0$ decays*, LHCb-PAPER-2023-041, in preparation.
- [71] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [72] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, [arXiv:1412.6352](#).
- [73] R. Aaij *et al.*, *Performance of the LHCb Vertex Locator*, JINST **9** (2014) P09007, [arXiv:1405.7808](#).
- [74] R. Arink *et al.*, *Performance of the LHCb Outer Tracker*, JINST **9** (2014) P01002, [arXiv:1311.3893](#).
- [75] P. d'Argent *et al.*, *Improved performance of the LHCb Outer Tracker in LHC Run 2*, JINST **12** (2017) P11016, [arXiv:1708.00819](#).

- [76] M. Adinolfi *et al.*, *Performance of the LHCb RICH detector at the LHC*, Eur. Phys. J. **C73** (2013) 2431, [arXiv:1211.6759](#).
- [77] A. A. Alves Jr. *et al.*, *Performance of the LHCb muon system*, JINST **8** (2013) P02022, [arXiv:1211.1346](#).
- [78] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, [arXiv:1211.3055](#).
- [79] T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, [arXiv:0710.3820](#).
- [80] I. Belyaev *et al.*, *Handling of the generation of primary events in GAUSS, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [81] C.-H. Chang, C. Driouichi, P. Eerola, and X.-G. Wu, *BCVEGPy: An event generator for hadronic production of the B_c meson*, Comput. Phys. Commun. **159** (2004) 192, [arXiv:hep-ph/0309120](#).
- [82] C.-H. Chang, J.-X. Wang, and X.-G. Wu, *BCVEGPy 2.0: An upgrade version of the generator BCVEGPy with an addendum about hadroproduction of the P-wave B_c states*, Comput. Phys. Commun. **174** (2006) 241, [arXiv:hep-ph/0504017](#).
- [83] X.-Y. Wang and X.-G. Wu, *A trick to improve the efficiency of generating unweighted B_c events from BCVEGPy*, Comput. Phys. Commun. **183** (2012) 442, [arXiv:1108.2442](#).
- [84] X.-G. Wu, *BCVEGPy and GENXICC for the hadronic production of the doubly heavy mesons and baryons*, J. Phys. Conf. Ser. **523** (2014) 012042, [arXiv:1307.3344](#).
- [85] C.-H. Chang and Y.-Q. Chen, *The hadronic production of the B_c meson at Tevatron, CERN LHC and SSC*, Phys. Rev. **D48** (1993) 4086.
- [86] C.-H. Chang, Y.-Q. Chen, G.-P. Han, and H.-T. Jiang, *On hadronic production of the B_c meson*, Phys. Lett. **B364** (1995) 78, [arXiv:hep-ph/9408242](#).
- [87] K. Kolodziej, A. Leike, and R. Ruckl, *Production of B_c mesons in hadronic collisions*, Phys. Lett. **B355** (1995) 337, [arXiv:hep-ph/9505298](#).
- [88] C.-H. Chang, Y.-Q. Chen, and R. J. Oakes, *Comparative study of the hadronic production of B_c mesons*, Phys. Rev. **D54** (1996) 4344, [arXiv:hep-ph/9602411](#).
- [89] A. V. Berezhnoy, V. V. Kiselev, and A. K. Likhoded, *Hadronic production of S- and P-wave states of $\bar{b}c$ quarkonium*, Z. Phys. **A356** (1996) 79, [arXiv:hep-ph/9602347](#).
- [90] D. J. Lange, *The EVTGEN particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [91] N. Davidson, T. Przedzinski, and Z. Was, *PHOTOS interface in C++: Technical and physics documentation*, Comp. Phys. Comm. **199** (2016) 86, [arXiv:1011.0937](#).

- [92] A. V. Berezhnoy, A. K. Likhoded, and A. V. Luchinsky, *BC_NPI module for the analysis of $B_c \rightarrow J/\psi + n\pi$ and $B_c \rightarrow B_s + n\pi$ decays within the EVTGEN package*, arXiv:1104.0808.
- [93] J. H. Kühn and A. Santamaria, *Tau decays to pions*, Z. Phys. **C48** (1990) 445.
- [94] CLEO collaboration, S. Anderson *et al.*, *Hadronic structure in the decay $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$* , Phys. Rev. **D61** (2000) 112002, arXiv:hep-ex/9910046.
- [95] Geant4 collaboration, J. Allison *et al.*, *GEANT4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *GEANT4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [96] M. Clemencic *et al.*, *The LHCb simulation application, GAUSS: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [97] R. Aaij *et al.*, *Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in Run 2*, Eur. Phys. J. Tech. Instr. **6** (2019) 1, arXiv:1803.00824.
- [98] LHCb collaboration, R. Aaij *et al.*, *Measurement of the track reconstruction efficiency at LHCb*, JINST **10** (2015) P02007, arXiv:1408.1251.
- [99] LHCb collaboration, R. Aaij *et al.*, *Evidence for the decay $B^0 \rightarrow J/\psi \omega$ and measurement of the relative branching fractions of B_s^0 meson decays to $J/\psi \eta$ and $J/\psi \eta'$* , Nucl. Phys. **B867** (2013) 547, arXiv:1210.2631.
- [100] LHCb collaboration, R. Aaij *et al.*, *Observations of $B_s^0 \rightarrow \psi(2S)\eta$ and $B_{(s)}^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays*, Nucl. Phys. **B871** (2013) 403, arXiv:1302.6354.
- [101] E. Govorkova, *Study of π^0/γ efficiency using B meson decays in the LHCb experiment*, Phys. Atom. Nucl. **79** (2016) 1474, arXiv:1505.02960.
- [102] K. Govorkova, *Study of photons and neutral pions reconstruction efficiency in the LHCb experiment*, Master's thesis, Lomonosov Moscow State University, Moscow, Russia, 2015. CERN-THESIS-2015-272.
- [103] I. M. Belyaev, E. M. Govorkova, V. Y. Egorychev, and D. V. Savrina, *Study of π^0/γ reconstruction and selection efficiency in the LHCb experiment*, Moscow Univ. Phys. Bull. **70** (2015) 497.
- [104] LHCb collaboration, R. Aaij *et al.*, *Observation of the $B^+ \rightarrow J/\psi \eta' K^+$ decay*, JHEP **08** (2023) 174, arXiv:2303.09443.
- [105] A. Powell *et al.*, *Particle identification at LHCb*, PoS **ICHEP2010** (2010) 020, LHCb-PROC-2011-008.
- [106] H. Terrier and I. Belyaev, *Particle identification with LHCb calorimeters*, LHCb-2003-092, 2003.
- [107] C. Abellan Beteta *et al.*, *Calibration and performance of the LHCb calorimeters in Run 1 and 2 at the LHC*, arXiv:2008.11556, submitted to JINST.

- [108] W. D. Hulsbergen, *Decay chain fitting with a Kalman filter*, Nucl. Instrum. Meth. **A552** (2005) 566, [arXiv:physics/0503191](#).
- [109] W. S. McCulloch and W. Pitts, *A logical calculus of the ideas immanent in nervous activity*, The bulletin of mathematical biophysics **5** (1943) 115.
- [110] F. Rosenblatt, *The perceptron: A probabilistic model for information storage and organization in the brain*, Psychological Review **65** (1958) 386.
- [111] J.-H. Zhong *et al.*, *A program for the Bayesian neural network in the ROOT framework*, Comput. Phys. Commun. **182** (2011) 2655, [arXiv:1103.2854](#).
- [112] S. Geisser, *Predictive inference: An introduction*, Chapman and Hall/CRC, New York, 1993.
- [113] T. Skwarnicki, *A study of the radiative cascade transitions between the Υ' and Υ resonances*, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [114] LHCb collaboration, R. Aaij *et al.*, *Observation of J/ψ -pair production in pp collisions at $\sqrt{s} = 7$ TeV*, Phys. Lett. **B707** (2012) 52, [arXiv:1109.0963](#).
- [115] S. Karlin and L. S. Shapley, *Geometry of moment spaces*, vol. 12 of *Memoirs of the American Mathematical Society*, American Mathematical Society, Providence, Rhode Island, 1953.
- [116] S. S. Wilks, *The large-sample distribution of the likelihood ratio for testing composite hypotheses*, Ann. Math. Stat. **9** (1938) 60.
- [117] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [118] T. Hastie, R. Tibshirani, and J. Friedman, *The elements of statistical learning: data mining, inference, and prediction*, Springer, New York, NY, USA, 2009.
- [119] M. Pivk and F. R. Le Diberder, *sPlot: A statistical tool to unfold data distributions*, Nucl. Instrum. Meth. **A555** (2005) 356, [arXiv:physics/0402083](#).
- [120] E. Byckling and K. Kajantie, *Particle kinematics*, John Wiley & Sons Inc., New York, 1973.
- [121] Student (W. S. Gosset), *The probable error of a mean*, Biometrika **6** (1908) 1.
- [122] S. Jackman, *Bayesian analysis for the social sciences*, John Wiley & Sons, Inc., Hoboken, New Jersey, USA, 2009.
- [123] N. L. Johnson, *Systems of frequency curves generated by methods of translation*, Biometrika **36** (1949) 149.
- [124] N. L. Johnson, *Bivariate distributions based on simple translation systems*, Biometrika **36** (1949) 297.

- [125] O. E. Barndorff-Nielsen, T. Mikosch, and S. I. Resnick, *Lévy processes: Theory and applications*, Sankhya: The Indian Journal of Statistics **65** (2003) 488.
- [126] O. Barndorff-Nielsen and D. G. Kendall, *Exponentially decreasing distributions for the logarithm of particle size*, Proceedings of the Royal Society of London. A. Mathematical and physical sciences **A353** (1977) 401.
- [127] K. Pearson, *Contributions to the mathematical theory of evolution. II. Skew variation in homogeneous material*, Philosophical Transactions of the Royal Society of London **186** (1895) 343.
- [128] S. Coles, *An introduction to statistical modeling of extreme values*, Springer, London, 2001.
- [129] G. R. Dargahi-Noubary, *On tail estimation: An improved method*, Math. Geol. **21** (1989) 829.
- [130] LHCb collaboration, R. Aaij *et al.*, *Amplitude analysis of the $B^+ \rightarrow D^+D^-K^+$ decay*, Phys. Rev. **D102** (2020) 112003, [arXiv:2009.00026](#).
- [131] A. Poluektov, *Kernel density estimation of a multidimensional efficiency profile*, JINST **10** (2015) P02011, [arXiv:1411.5528](#).
- [132] LHCb collaboration, R. Aaij *et al.*, *Measurement of relative branching fractions of B decays to $\psi(2S)$ and J/ψ mesons*, Eur. Phys. J. **C72** (2012) 2118, [arXiv:1205.0918](#).



















LHCb collaboration

R. Aaij³⁶ , A.S.W. Abdelmotteleb⁵⁵ , C. Abellan Beteta⁴⁹ , F. Abudinén⁵⁵ ,
T. Ackernley⁵⁹ , J. A. Adams⁶⁷ , A. A. Adefisoye⁶⁷ , B. Adeva⁴⁵ , M. Adinolfi⁵³ ,
P. Adlarson⁷⁹ , C. Agapopoulou⁴⁷ , C.A. Aidala⁸⁰ , Z. Ajaltouni¹¹ , S. Akar⁶⁴ ,
K. Akiba³⁶ , P. Albicocco²⁶ , J. Albrecht¹⁸ , F. Alessio⁴⁷ , M. Alexander⁵⁸ ,
Z. Aliouche⁶¹ , P. Alvarez Cartelle⁵⁴ , R. Amalric¹⁵ , S. Amato³ , J.L. Amey⁵³ ,
Y. Amhis^{13,47} , L. An⁶ , L. Anderlini²⁵ , M. Andersson⁴⁹ , A. Andreianov⁴² ,
P. Andreola⁴⁹ , M. Andreotti²⁴ , D. Andreou⁶⁷ , A. Anelli^{29,p} , D. Ao⁷ ,
F. Archilli^{35,v} , M. Argenton²⁴ , S. Arguedas Cuendis⁹ , A. Artamonov⁴² ,
M. Artuso⁶⁷ , E. Aslanides¹² , M. Atzeni⁶³ , B. Audurier¹⁴ , D. Bacher⁶² ,
I. Bachiller Perea¹⁰ , S. Bachmann²⁰ , M. Bachmayer⁴⁸ , J.J. Back⁵⁵ ,
P. Baladron Rodriguez⁴⁵ , V. Balagura¹⁴ , W. Baldini²⁴ , J. Baptista de Souza Leite⁵⁹ ,
M. Barbetti^{25,m} , I. R. Barbosa⁶⁸ , R.J. Barlow⁶¹ , S. Barsuk¹³ , W. Barter⁵⁷ ,
M. Bartolini⁵⁴ , J. Bartz⁶⁷ , F. Baryshnikov⁴² , J.M. Basels¹⁶ , G. Bassi³³ ,
B. Batsukh⁵ , A. Battig¹⁸ , A. Bay⁴⁸ , A. Beck⁵⁵ , M. Becker¹⁸ , F. Bedeschi³³ ,
I.B. Bediaga² , A. Beiter⁶⁷ , S. Belin⁴⁵ , V. Bellee⁴⁹ , K. Belous⁴² , I. Belov²⁷ ,
I. Belyaev³⁴ , G. Benane¹² , G. Bencivenni²⁶ , E. Ben-Haim¹⁵ , A. Berezhnoy⁴² ,
R. Bernet⁴⁹ , S. Bernet Andres⁴³ , C. Bertella⁶¹ , A. Bertolin³¹ , C. Betancourt⁴⁹ ,
F. Betti⁵⁷ , J. Bex⁵⁴ , Ia. Bezshyiko⁴⁹ , J. Bhom³⁹ , M.S. Bieker¹⁸ , N.V. Biesuz²⁴ ,
P. Billoir¹⁵ , A. Biolchini³⁶ , M. Birch⁶⁰ , F.C.R. Bishop¹⁰ , A. Bitadze⁶¹ ,
A. Bizzeti , T. Blake⁵⁵ , F. Blanc⁴⁸ , J.E. Blank¹⁸ , S. Blusk⁶⁷ , V. Bocharnikov⁴² ,
J.A. Boelhauve¹⁸ , O. Boente Garcia¹⁴ , T. Boettcher⁶⁴ , A. Bohare⁵⁷ ,
A. Boldyrev⁴² , C.S. Bolognani⁷⁶ , R. Bolzonella^{24,l} , N. Bondar⁴² , F. Borgato^{31,47} ,
S. Borghi⁶¹ , M. Borsato^{29,p} , J.T. Borsuk³⁹ , S.A. Bouchiba⁴⁸ , T.J.V. Bowcock⁵⁹ ,
A. Boyer⁴⁷ , C. Bozzi²⁴ , M.J. Bradley⁶⁰ , A. Brea Rodriguez⁴⁵ , N. Breer¹⁸ ,
J. Brodzicka³⁹ , A. Brossa Gonzalo⁴⁵ , J. Brown⁵⁹ , D. Brundu³⁰ , E. Buchanan⁵⁷ ,
A. Buonauro⁴⁹ , L. Buonincontri³¹ , A.T. Burke⁶¹ , C. Burr⁴⁷ , A. Bursche⁷⁰ ,
A. Butkevich⁴² , J.S. Butter⁵⁴ , J. Buytaert⁴⁷ , W. Byczynski⁴⁷ , S. Cadeddu³⁰ ,
H. Cai⁷² , R. Calabrese^{24,l} , L. Calefice⁴⁴ , S. Cali²⁶ , M. Calvi^{29,p} , M. Calvo Gomez⁴³ ,
J. Cambon Bouzas⁴⁵ , P. Campana²⁶ , D.H. Campora Perez⁷⁶ ,
A.F. Campoverde Quezada⁷ , S. Capelli²⁹ , L. Capriotti²⁴ , R. Caravaca-Mora⁹ ,
A. Carbone^{23,j} , L. Carcedo Salgado⁴⁵ , R. Cardinale^{27,n} , A. Cardini³⁰ ,
P. Carniti^{29,p} , L. Carus²⁰ , A. Casais Vidal⁶³ , R. Caspary²⁰ , G. Casse⁵⁹ ,
J. Castro Godinez⁹ , M. Cattaneo⁴⁷ , G. Cavallero²⁴ , V. Cavallini^{24,l} , S. Celani²⁰ ,
J. Cerasoli¹² , D. Cervenkov⁶² , S. Cesare^{28,o} , A.J. Chadwick⁵⁹ , I. Chahrour⁸⁰ ,
M. Charles¹⁵ , Ph. Charpentier⁴⁷ , C.A. Chavez Barajas⁵⁹ , M. Chefdeville¹⁰ ,
C. Chen¹² , S. Chen⁵ , Z. Chen⁷ , A. Chernov³⁹ , S. Chernyshenko⁵¹ ,
V. Chobanova⁷⁸ , S. Cholak⁴⁸ , M. Chrzaszcz³⁹ , A. Chubykin⁴² , V. Chulikov⁴² ,
P. Ciambrone²⁶ , X. Cid Vidal⁴⁵ , G. Ciezarek⁴⁷ , P. Cifra⁴⁷ , P.E.L. Clarke⁵⁷ ,
M. Clemencic⁴⁷ , H.V. Cliff⁵⁴ , J. Closier⁴⁷ , C. Cocha Toapaxi²⁰ , V. Coco⁴⁷ ,
J. Cogan¹² , E. Cogneras¹¹ , L. Cojocariu⁴¹ , P. Collins⁴⁷ , T. Colombo⁴⁷ ,
A. Comerma-Montells⁴⁴ , L. Congedo²² , A. Contu³⁰ , N. Cooke⁵⁸ , I. Corredoira⁴⁵ ,
A. Correia¹⁵ , G. Corti⁴⁷ , J.J. Cottee Meldrum⁵³ , B. Couturier⁴⁷ , D.C. Craik⁴⁹ ,
M. Cruz Torres^{2,g} , E. Curras Rivera⁴⁸ , R. Currie⁵⁷ , C.L. Da Silva⁶⁶ , S. Dadabev⁴² ,
L. Dai⁶⁹ , X. Dai⁶ , E. Dall'Occo¹⁸ , J. Dalseno⁴⁵ , C. D'Ambrosio⁴⁷ , J. Daniel¹¹ ,
A. Danilina⁴² , P. d'Argent²² , A. Davidson⁵⁵ , J.E. Davies⁶¹ , A. Davis⁶¹ ,
O. De Aguiar Francisco⁶¹ , C. De Angelis^{30,k} , J. de Boer³⁶ , K. De Bruyn⁷⁵ ,
S. De Capua⁶¹ , M. De Cian^{20,47} , U. De Freitas Carneiro Da Graca^{2,b} , E. De Lucia²⁶ ,
J.M. De Miranda² , L. De Paula³ , M. De Serio^{22,h} , P. De Simone²⁶ , F. De Vellis¹⁸ ,

J.A. de Vries⁷⁶ , F. Debernardis²² , D. Decamp¹⁰ , V. Dedu¹² , L. Del Buono¹⁵ ,
 B. Delaney⁶³ , H.-P. Dembinski¹⁸ , J. Deng⁸ , V. Denysenko⁴⁹ , O. Deschamps¹¹ ,
 F. Dettori^{30,k} , B. Dey⁷⁴ , P. Di Nezza²⁶ , I. Diachkov⁴² , S. Didenko⁴² , S. Ding⁶⁷ ,
 L. Dittmann²⁰ , V. Dobishuk⁵¹ , A. D. Docheva⁵⁸ , A. Dolmatov⁴² , C. Dong⁴ ,
 A.M. Donohoe²¹ , F. Dordei³⁰ , A.C. dos Reis² , A. D. Dowling⁶⁷ , A.G. Downes¹⁰ ,
 W. Duan⁷⁰ , P. Duda⁷⁷ , M.W. Dudek³⁹ , L. Dufour⁴⁷ , V. Duk³² , P. Durante⁴⁷ , M.
 M. Duras⁷⁷ , J.M. Durham⁶⁶ , O. D. Durmus⁷⁴ , A. Dziurda³⁹ , A. Dzyuba⁴² ,
 S. Easo⁵⁶ , E. Eckstein¹⁷ , U. Egede¹ , A. Egorychev⁴² , V. Egorychev⁴² ,
 S. Eisenhardt⁵⁷ , E. Ejopu⁶¹ , S. Ek-In⁴⁸ , L. Eklund⁷⁹ , M. Elashri⁶⁴ ,
 J. Ellbracht¹⁸ , S. Ely⁶⁰ , A. Ene⁴¹ , E. Epple⁶⁴ , S. Escher¹⁶ , J. Eschle⁶⁷ ,
 S. Esen²⁰ , T. Evans⁶¹ , F. Fabiano^{30,k,47} , L.N. Falcao² , Y. Fan⁷ , B. Fang^{72,13} ,
 L. Fantini^{32,r} , M. Faria⁴⁸ , K. Farmer⁵⁷ , D. Fazzini^{29,p} , L. Felkowski⁷⁷ ,
 M. Feng^{5,7} , M. Feo^{18,47} , M. Fernandez Gomez⁴⁵ , A.D. Fernez⁶⁵ , F. Ferrari²³ ,
 F. Ferreira Rodrigues³ , S. Ferreres Sole³⁶ , M. Ferrillo⁴⁹ , M. Ferro-Luzzi⁴⁷ ,
 S. Filippov⁴² , R.A. Fini²² , M. Fiorini^{24,l} , K.M. Fischer⁶² , D.S. Fitzgerald⁸⁰ ,
 C. Fitzpatrick⁶¹ , F. Fleuret¹⁴ , M. Fontana²³ , L. F. Foreman⁶¹ , R. Forty⁴⁷ ,
 D. Foulds-Holt⁵⁴ , M. Franco Sevilla⁶⁵ , M. Frank⁴⁷ , E. Franzoso^{24,l} , G. Frau²⁰ ,
 C. Frei⁴⁷ , D.A. Friday⁶¹ , J. Fu⁷ , Q. Fuehring¹⁸ , Y. Fujii¹ , T. Fulghesu¹⁵ ,
 E. Gabriel³⁶ , G. Galati^{22,h} , M.D. Galati³⁶ , A. Gallas Torreira⁴⁵ , D. Galli^{23,j} ,
 S. Gambetta⁵⁷ , M. Gandelman³ , P. Gandini²⁸ , H. Gao⁷ , R. Gao⁶² , Y. Gao⁸ ,
 Y. Gao⁶ , M. Garau^{30,k} , L.M. Garcia Martin⁴⁸ , P. Garcia Moreno⁴⁴ ,
 J. García Pardiñas⁴⁷ , K. G. Garg⁸ , L. Garrido⁴⁴ , C. Gaspar⁴⁷ , R.E. Geertsema³⁶ ,
 L.L. Gerken¹⁸ , E. Gersabeck⁶¹ , M. Gersabeck⁶¹ , T. Gershon⁵⁵ ,
 Z. Ghorbanimoghaddam⁵³ , L. Giambastiani³¹ , F. I. Giasemis^{15,e} , V. Gibson⁵⁴ ,
 H.K. Giemza⁴⁰ , A.L. Gilman⁶² , M. Giovannetti²⁶ , A. Gioventù⁴⁴ ,
 P. Gironella Gironell⁴⁴ , C. Giugliano^{24,l} , M.A. Giza³⁹ , E.L. Gkoukousis⁶⁰ ,
 F.C. Glaser^{13,20} , V.V. Gligorov¹⁵ , C. Göbel⁶⁸ , E. Golobardes⁴³ , D. Golubkov⁴² ,
 A. Golutvin^{60,42,47} , A. Gomes^{2,a,†} , S. Gomez Fernandez⁴⁴ , F. Goncalves Abrantes⁶² ,
 M. Goncerz³⁹ , G. Gong⁴ , J. A. Gooding¹⁸ , I.V. Gorelov⁴² , C. Gotti²⁹ ,
 J.P. Grabowski¹⁷ , L.A. Granado Cardoso⁴⁷ , E. Graugés⁴⁴ , E. Graverini^{48,t} ,
 L. Gazette⁵⁵ , G. Graziani , A. T. Grecu⁴¹ , L.M. Greeven³⁶ , N.A. Grieser⁶⁴ ,
 L. Grillo⁵⁸ , S. Gromov⁴² , C. Gu¹⁴ , M. Guarise²⁴ , M. Guittiere¹³ , V. Guliaeva⁴² ,
 P. A. Günther²⁰ , A.-K. Guseinov⁴⁸ , E. Gushchin⁴² , Y. Guz^{6,42,47} , T. Gys⁴⁷ ,
 K. Habermann¹⁷ , T. Hadavizadeh¹ , C. Hadjivasiliou⁶⁵ , G. Haefeli⁴⁸ , C. Haen⁴⁷ ,
 J. Haimberger⁴⁷ , M. Hajheidari⁴⁷ , M.M. Halvorsen⁴⁷ , P.M. Hamilton⁶⁵ ,
 J. Hammerich⁵⁹ , Q. Han⁸ , X. Han²⁰ , S. Hansmann-Menzemer²⁰ , L. Hao⁷ ,
 N. Harnew⁶² , T. Harrison⁵⁹ , M. Hartmann¹³ , J. He^{7,c} , F. Hemmer⁴⁷ ,
 C. Henderson⁶⁴ , R.D.L. Henderson^{1,55} , A.M. Hennequin⁴⁷ , K. Hennessy⁵⁹ ,
 L. Henry⁴⁸ , J. Herd⁶⁰ , J. Herdieckerhoff¹⁸ , P. Herrero Gascon²⁰ , J. Heuel¹⁶ ,
 A. Hicheur³ , G. Hijano Mendizabal⁴⁹ , D. Hill⁴⁸ , S.E. Hollitt¹⁸ , J. Horswill⁶¹ ,
 R. Hou⁸ , Y. Hou¹¹ , N. Howarth⁵⁹ , J. Hu²⁰ , J. Hu⁷⁰ , W. Hu⁶ , X. Hu⁴ ,
 W. Huang⁷ , W. Hulsbergen³⁶ , R.J. Hunter⁵⁵ , M. Hushchyn⁴² , D. Hutchcroft⁵⁹ ,
 D. Ilin⁴² , P. Ilten⁶⁴ , A. Inglessi⁴² , A. Iniukhin⁴² , A. Ishteev⁴² , K. Ivshin⁴² ,
 R. Jacobsson⁴⁷ , H. Jage¹⁶ , S.J. Jaimes Elles^{46,73} , S. Jakobsen⁴⁷ , E. Jans³⁶ ,
 B.K. Jashal⁴⁶ , A. Jawahery^{65,47} , V. Jevtic¹⁸ , E. Jiang⁶⁵ , X. Jiang^{5,7} , Y. Jiang⁷ ,
 Y. J. Jiang⁶ , M. John⁶² , D. Johnson⁵² , C.R. Jones⁵⁴ , T.P. Jones⁵⁵ , S. Joshi⁴⁰ ,
 B. Jost⁴⁷ , N. Jurik⁴⁷ , I. Juszcak³⁹ , D. Kaminaris⁴⁸ , S. Kandybei⁵⁰ , Y. Kang⁴ ,
 M. Karacson⁴⁷ , D. Karpenkov⁴² , A. M. Kauniskangas⁴⁸ , J.W. Kautz⁶⁴ ,
 F. Keizer⁴⁷ , M. Kenzie⁵⁴ , T. Ketel³⁶ , B. Khanji⁶⁷ , A. Kharisova⁴² ,
 S. Kholodenko³³ , G. Khreich¹³ , T. Kirn¹⁶ , V.S. Kirsebom²⁹ , O. Kitouni⁶³ ,

S. Klaver³⁷ , N. Kleijne^{33,s} , K. Klimaszewski⁴⁰ , M.R. Kmiec⁴⁰ , S. Koliiev⁵¹ ,
 L. Kolk¹⁸ , A. Konoplyannikov⁴² , P. Kopciwicz^{38,47} , P. Koppenburg³⁶ ,
 M. Korolev⁴² , I. Kostiuk³⁶ , O. Kot⁵¹ , S. Kotriakhova , A. Kozachuk⁴² ,
 P. Kravchenko⁴² , L. Kravchuk⁴² , M. Kreps⁵⁵ , S. Kretzschmar¹⁶ , P. Krokovny⁴² ,
 W. Krupa⁶⁷ , W. Krzemien⁴⁰ , J. Kubat²⁰ , S. Kubis⁷⁷ , W. Kucewicz³⁹ ,
 M. Kucharczyk³⁹ , V. Kudryavtsev⁴² , E. Kulikova⁴² , A. Kupsc⁷⁹ , B. K. Kutsenko¹² ,
 D. Lacarrere⁴⁷ , A. Lai³⁰ , A. Lampis³⁰ , D. Lancierini⁵⁴ , C. Landesa Gomez⁴⁵ ,
 J.J. Lane¹ , R. Lane⁵³ , C. Langenbruch²⁰ , J. Langer¹⁸ , O. Lantwin⁴² ,
 T. Latham⁵⁵ , F. Lazzari^{33,t} , C. Lazzeroni⁵² , R. Le Gac¹² , R. Lefèvre¹¹ ,
 A. Leflat⁴² , S. Legotin⁴² , M. Lehuraux⁵⁵ , E. Lemos Cid⁴⁷ , O. Leroy¹² ,
 T. Lesiak³⁹ , B. Leverington²⁰ , A. Li⁴ , H. Li⁷⁰ , K. Li⁸ , L. Li⁶¹ , P. Li⁴⁷ ,
 P.-R. Li⁷¹ , S. Li⁸ , T. Li^{5,d} , T. Li⁷⁰ , Y. Li⁸ , Y. Li⁵ , Z. Li⁶⁷ , Z. Lian⁴ ,
 X. Liang⁶⁷ , S. Libralon⁴⁶ , C. Lin⁷ , T. Lin⁵⁶ , R. Lindner⁴⁷ , V. Lisovskyi⁴⁸ ,
 R. Litvinov³⁰ , F. L. Liu¹ , G. Liu⁷⁰ , K. Liu⁷¹ , Q. Liu⁷ , S. Liu^{5,7} , Y. Liu⁵⁷ ,
 Y. Liu⁷¹ , Y. L. Liu⁶⁰ , A. Lobo Salvia⁴⁴ , A. Loi³⁰ , J. Lomba Castro⁴⁵ , T. Long⁵⁴ ,
 J.H. Lopes³ , A. Lopez Huertas⁴⁴ , S. López Soliño⁴⁵ , C. Lucarelli^{25,m} ,
 D. Lucchesi^{31,q} , M. Lucio Martinez⁷⁶ , V. Lukashenko^{36,51} , Y. Luo⁶ , A. Lupato³¹ ,
 E. Luppi^{24,l} , K. Lynch²¹ , X.-R. Lyu⁷ , G. M. Ma⁴ , R. Ma⁷ , S. Maccolini¹⁸ ,
 F. Machefert¹³ , F. Maciuc⁴¹ , B. M. Mack⁶⁷ , I. Mackay⁶² , L. M. Mackey⁶⁷ ,
 L.R. Madhan Mohan⁵⁴ , M. M. Madurai⁵² , A. Maevskiy⁴² , D. Magdalinski³⁶ ,
 D. Maisuzenko⁴² , M.W. Majewski³⁸ , J.J. Malczewski³⁹ , S. Malde⁶² , B. Malecki³⁹ ,
 L. Malentacca⁴⁷ , A. Malinin⁴² , T. Maltsev⁴² , G. Manca^{30,k} , G. Mancinelli¹² ,
 C. Mancuso^{28,13,o} , R. Manera Escalero⁴⁴ , D. Manuzzi²³ , D. Marangotto^{28,o} ,
 J.F. Marchand¹⁰ , R. Marchevski⁴⁸ , U. Marconi²³ , S. Mariani⁴⁷ , C. Marin Benito⁴⁴ ,
 J. Marks²⁰ , A.M. Marshall⁵³ , P.J. Marshall⁵⁹ , G. Martelli^{32,r} , G. Martellotti³⁴ ,
 L. Martinazzoli⁴⁷ , M. Martinelli^{29,p} , D. Martinez Santos⁴⁵ , F. Martinez Vidal⁴⁶ ,
 A. Massafferri² , M. Materok¹⁶ , R. Matev⁴⁷ , A. Mathad⁴⁷ , V. Matiunin⁴² ,
 C. Matteuzzi⁶⁷ , K.R. Mattioli¹⁴ , A. Mauri⁶⁰ , E. Maurice¹⁴ , J. Mauricio⁴⁴ ,
 P. Mayencourt⁴⁸ , M. Mazurek⁴⁰ , M. McCann⁶⁰ , L. McConnell²¹ , T.H. McGrath⁶¹ ,
 N.T. McHugh⁵⁸ , A. McNab⁶¹ , R. McNulty²¹ , B. Meadows⁶⁴ , G. Meier¹⁸ ,
 D. Melnychuk⁴⁰ , M. Merk^{36,76} , A. Merli^{28,o} , L. Meyer Garcia³ , D. Miao^{5,7} ,
 H. Miao⁷ , M. Mikhasenko^{17,f} , D.A. Milanese⁷³ , A. Minotti^{29,p} , E. Minucci⁶⁷ ,
 T. Miralles¹¹ , B. Mitreska¹⁸ , D.S. Mitzel¹⁸ , A. Modak⁵⁶ , A. Mödden ¹⁸ ,
 R.A. Mohammed⁶² , R.D. Moise¹⁶ , S. Mokhnenko⁴² , T. Mombächer⁴⁷ , M. Monk^{55,1} ,
 S. Monteil¹¹ , A. Morcillo Gomez⁴⁵ , G. Morello²⁶ , M.J. Morello^{33,s} ,
 M.P. Morgenthaler²⁰ , A.B. Morris⁴⁷ , A.G. Morris¹² , R. Mountain⁶⁷ , H. Mu⁴ , Z. M.
 Mu⁶ , E. Muhammad⁵⁵ , F. Muheim⁵⁷ , M. Mulder⁷⁵ , K. Müller⁴⁹ ,
 F. Muñoz-Rojas⁹ , R. Murta⁶⁰ , P. Naik⁵⁹ , T. Nakada⁴⁸ , R. Nandakumar⁵⁶ ,
 T. Nanut⁴⁷ , I. Nasteva³ , M. Needham⁵⁷ , N. Neri^{28,o} , S. Neubert¹⁷ , N. Neufeld⁴⁷ ,
 P. Neustroev⁴² , J. Nicolini^{18,13} , D. Nicotra⁷⁶ , E.M. Niel⁴⁸ , N. Nikitin⁴² , P. Nogga¹⁷ ,
 N.S. Nolte⁶³ , C. Normand⁵³ , J. Novoa Fernandez⁴⁵ , G. Nowak⁶⁴ , C. Nunez⁸⁰ , H. N.
 Nur⁵⁸ , A. Oblakowska-Mucha³⁸ , V. Obraztsov⁴² , T. Oeser¹⁶ , S. Okamura^{24,l,47} ,
 A. Okhotnikov⁴² , R. Oldeman^{30,k} , F. Oliva⁵⁷ , M. Olocco¹⁸ , C.J.G. Onderwater⁷⁶ ,
 R.H. O'Neil⁵⁷ , J.M. Otalora Goicochea³ , P. Owen⁴⁹ , A. Oyanguren⁴⁶ , O. Ozcelik⁵⁷ ,
 K.O. Padeken¹⁷ , B. Pagare⁵⁵ , P.R. Pais²⁰ , T. Pajero⁶² , A. Palano²² ,
 M. Palutan²⁶ , G. Panshin⁴² , L. Paolucci⁵⁵ , A. Papanestis⁵⁶ , M. Pappagallo^{22,h} ,
 L.L. Pappalardo^{24,l} , C. Pappenheimer⁶⁴ , C. Parkes⁶¹ , B. Passalacqua²⁴ ,
 G. Passaleva²⁵ , D. Passaro^{33,s} , A. Pastore²² , M. Patel⁶⁰ , J. Patoc⁶² ,
 C. Patrignani^{23,j} , C.J. Pawley⁷⁶ , A. Pellegrino³⁶ , M. Pepe Altarelli²⁶ ,
 S. Perazzini²³ , D. Pereima⁴² , A. Pereiro Castro⁴⁵ , P. Perret¹¹ , A. Perro⁴⁷ ,

K. Petridis⁵³ , A. Petrolini^{27,n} , S. Petrucci⁵⁷ , J. P. Pfaller⁶⁴ , H. Pham⁶⁷ ,
 L. Pica^{33,s} , M. Piccini³² , B. Pietrzyk¹⁰ , G. Pietrzyk¹³ , D. Pinci³⁴ , F. Pisani⁴⁷ ,
 M. Pizzichemi^{29,p} , V. Placinta⁴¹ , M. Plo Casasus⁴⁵ , F. Polci^{15,47} , M. Poli Lener²⁶ ,
 A. Poluektov¹² , N. Polukhina⁴² , I. Polyakov⁴⁷ , E. Polycarpo³ , S. Ponce⁴⁷ ,
 D. Popov⁷ , S. Poslavskii⁴² , K. Prasanth³⁹ , C. Prouve⁴⁵ , V. Pugatch⁵¹ ,
 G. Punzi^{33,t} , W. Qian⁷ , N. Qin⁴ , S. Qu⁴ , R. Quagliani⁴⁸ , R.I. Rabadan Trejo⁵⁵ ,
 J.H. Rademacker⁵³ , M. Rama³³ , M. Ramírez García⁸⁰ , M. Ramos Pernas⁵⁵ ,
 M.S. Rangel³ , F. Ratnikov⁴² , G. Raven³⁷ , M. Rebollo De Miguel⁴⁶ , F. Redi^{28,i} ,
 J. Reich⁵³ , F. Reiss⁶¹ , Z. Ren⁷ , P.K. Resmi⁶² , R. Ribatti^{33,s} , G. R. Ricart^{14,81} ,
 D. Riccardi^{33,s} , S. Ricciardi⁵⁶ , K. Richardson⁶³ , M. Richardson-Slipper⁵⁷ ,
 K. Rinnert⁵⁹ , P. Robbe¹³ , G. Robertson⁵⁸ , E. Rodrigues⁵⁹ ,
 E. Rodriguez Fernandez⁴⁵ , J.A. Rodriguez Lopez⁷³ , E. Rodriguez Rodriguez⁴⁵ ,
 A. Rogovskiy⁵⁶ , D.L. Rolf⁴⁷ , P. Roloff⁴⁷ , V. Romanovskiy⁴² , M. Romero Lamas⁴⁵ ,
 A. Romero Vidal⁴⁵ , G. Romolini²⁴ , F. Ronchetti⁴⁸ , M. Rotondo²⁶ , S. R. Roy²⁰ ,
 M.S. Rudolph⁶⁷ , T. Ruf⁴⁷ , M. Ruiz Diaz²⁰ , R.A. Ruiz Fernandez⁴⁵ ,
 J. Ruiz Vidal^{79,z} , A. Ryzhikov⁴² , J. Ryzka³⁸ , J. J. Saavedra-Arias⁹ ,
 J.J. Saborido Silva⁴⁵ , R. Sadek¹⁴ , N. Sagidova⁴² , D. Sahoo⁷⁴ , N. Sahoo⁵² ,
 B. Saitta^{30,k} , M. Salnikova⁴² , M. Salomoni^{29,p} , C. Sanchez Gras³⁶ ,
 I. Sanderswood⁴⁶ , R. Santacesaria³⁴ , C. Santamarina Rios⁴⁵ , M. Santimaria²⁶ ,
 L. Santoro² , E. Santovetti³⁵ , A. Saputi^{24,47} , D. Saranin⁴² , G. Sarpis⁵⁷ ,
 M. Sarpis¹⁷ , A. Sarti³⁴ , C. Satriano^{34,u} , A. Satta³⁵ , M. Saur⁶ , D. Savrina⁴² ,
 H. Sazak¹⁶ , L.G. Scantlebury Smead⁶² , A. Scarabotto¹⁸ , S. Schael¹⁶ , S. Scherl⁵⁹ ,
 M. Schiller⁵⁸ , H. Schindler⁴⁷ , M. Schmelling¹⁹ , B. Schmidt⁴⁷ , S. Schmitt¹⁶ ,
 H. Schmitz¹⁷ , O. Schneider⁴⁸ , A. Schopper⁴⁷ , N. Schulte¹⁸ , S. Schulte⁴⁸ ,
 M.H. Schune¹³ , R. Schwemmer⁴⁷ , G. Schwering¹⁶ , B. Sciascia²⁶ , A. Sciuccati⁴⁷ ,
 S. Sellam⁴⁵ , A. Semennikov⁴² , T. Senger⁴⁹ , M. Senghi Soares³⁷ , A. Sergi²⁷ ,
 N. Serra⁴⁹ , L. Sestini³¹ , A. Seuthe¹⁸ , Y. Shang⁶ , D.M. Shangase⁸⁰ , M. Shapkin⁴² ,
 R. S. Sharma⁶⁷ , I. Shchemerov⁴² , L. Shchutka⁴⁸ , T. Shears⁵⁹ , L. Shekhtman⁴² ,
 Z. Shen⁶ , S. Sheng^{5,7} , V. Shevchenko⁴² , B. Shi⁷ , Q. Shi⁷ , E.B. Shields^{29,p} ,
 Y. Shimizu¹³ , E. Shmanin⁴² , R. Shorkin⁴² , J.D. Shupperd⁶⁷ , R. Silva Coutinho⁶⁷ ,
 G. Simi³¹ , S. Simone^{22,h} , N. Skidmore⁵⁵ , T. Skwarnicki⁶⁷ , M.W. Slater⁵² ,
 J.C. Smallwood⁶² , E. Smith⁶³ , K. Smith⁶⁶ , M. Smith⁶⁰ , A. Snoch³⁶ ,
 L. Soares Lavra⁵⁷ , M.D. Sokoloff⁶⁴ , F.J.P. Soler⁵⁸ , A. Solomin^{42,53} , A. Solovov⁴² ,
 I. Solovyev⁴² , R. Song¹ , Y. Song⁴⁸ , Y. Song⁴ , Y. S. Song⁶ ,
 F.L. Souza De Almeida⁶⁷ , B. Souza De Paula³ , E. Spadaro Norella^{28,o} ,
 E. Spedicato²³ , J.G. Speer¹⁸ , E. Spiridenkov⁴² , P. Spradlin⁵⁸ , V. Sriskaran⁴⁷ ,
 F. Stagni⁴⁷ , M. Stahl⁴⁷ , S. Stahl⁴⁷ , S. Stanislaus⁶² , E.N. Stein⁴⁷ , O. Steinkamp⁴⁹ ,
 O. Stenyakin⁴² , H. Stevens¹⁸ , D. Strelakina⁴² , Y. Su⁷ , F. Suljik⁶² , J. Sun³⁰ ,
 L. Sun⁷² , Y. Sun⁶⁵ , W. Sutcliffe⁴⁹ , P.N. Swallow⁵² , F. Swystun⁵⁴ , A. Szabelski⁴⁰ ,
 T. Szumlak³⁸ , Y. Tan⁴ , M.D. Tat⁶² , A. Terentev⁴⁹ , F. Terzuoli^{33,w} , F. Teubert⁴⁷ ,
 E. Thomas⁴⁷ , D.J.D. Thompson⁵² , H. Tilquin⁶⁰ , V. Tisserand¹¹ , S. T'Jampens¹⁰ ,
 M. Tobin⁵ , L. Tomassetti^{24,l} , G. Tonani^{28,o,47} , X. Tong⁶ , D. Torres Machado² ,
 L. Toscano¹⁸ , D.Y. Tou⁴ , C. Trippi⁴³ , G. Tuci²⁰ , N. Tuning³⁶ , L.H. Uecker²⁰ ,
 A. Ukleja³⁸ , D.J. Unverzagt²⁰ , E. Ursov⁴² , A. Usachov³⁷ , A. Ustyuzhanin⁴² ,
 U. Uwer²⁰ , V. Vagnoni²³ , A. Valassi⁴⁷ , G. Valenti²³ , N. Valls Canudas⁴⁷ ,
 H. Van Hecke⁶⁶ , E. van Herwijnen⁶⁰ , C.B. Van Hulse^{45,y} , R. Van Laak⁴⁸ ,
 M. van Veghel³⁶ , G. Vasquez⁴⁹ , R. Vazquez Gomez⁴⁴ , P. Vazquez Regueiro⁴⁵ ,
 C. Vázquez Sierra⁴⁵ , S. Vecchi²⁴ , J.J. Velthuis⁵³ , M. Veltri^{25,x} , A. Venkateswaran⁴⁸ ,
 M. Vesterinen⁵⁵ , M. Vieites Diaz⁴⁷ , X. Vilasis-Cardona⁴³ , E. Vilella Figueras⁵⁹ ,
 A. Villa²³ , P. Vincent¹⁵ , F.C. Volle⁵² , D. vom Bruch¹² , V. Vorobyev⁴²

N. Voropaev⁴² , K. Vos⁷⁶ , G. Vouters¹⁰, C. Vrahas⁵⁷ , J. Walsh³³ , E.J. Walton¹ ,
G. Wan⁶ , C. Wang²⁰ , G. Wang⁸ , J. Wang⁶ , J. Wang⁵ , J. Wang⁴ , J. Wang⁷² ,
M. Wang²⁸ , N. W. Wang⁷ , R. Wang⁵³ , X. Wang⁷⁰ , X. W. Wang⁶⁰ , Y. Wang⁸ ,
Z. Wang¹³ , Z. Wang⁴ , Z. Wang²⁸ , J.A. Ward^{55,1} , M. Waterlaet⁴⁷, N.K. Watson⁵² ,
D. Websdale⁶⁰ , Y. Wei⁶ , B.D.C. Westhenry⁵³ , D.J. White⁶¹ , M. Whitehead⁵⁸ ,
A.R. Wiederhold⁵⁵ , D. Wiedner¹⁸ , G. Wilkinson⁶² , M.K. Wilkinson⁶⁴ ,
M. Williams⁶³ , M.R.J. Williams⁵⁷ , R. Williams⁵⁴ , F.F. Wilson⁵⁶ , W. Wislicki⁴⁰ ,
M. Witek³⁹ , L. Witola²⁰ , C.P. Wong⁶⁶ , G. Wormser¹³ , S.A. Wotton⁵⁴ , H. Wu⁶⁷ ,
J. Wu⁸ , Y. Wu⁶ , K. Wyllie⁴⁷ , S. Xian⁷⁰, Z. Xiang⁵ , Y. Xie⁸ , A. Xu³³ , J. Xu⁷ ,
L. Xu⁴ , L. Xu⁴ , M. Xu⁵⁵ , Z. Xu¹¹ , Z. Xu⁷ , Z. Xu⁵ , D. Yang⁴ , S. Yang⁷ ,
X. Yang⁶ , Y. Yang^{27,n} , Z. Yang⁶ , Z. Yang⁶⁵ , V. Yeroshenko¹³ , H. Yeung⁶¹ ,
H. Yin⁸ , C. Y. Yu⁶ , J. Yu⁶⁹ , X. Yuan⁵ , E. Zaffaroni⁴⁸ , M. Zavertyaev¹⁹ ,
M. Zdybal³⁹ , M. Zeng⁴ , C. Zhang⁶ , D. Zhang⁸ , J. Zhang⁷ , L. Zhang⁴ ,
S. Zhang⁶⁹ , S. Zhang⁶ , Y. Zhang⁶ , Y. Z. Zhang⁴ , Y. Zhao²⁰ , A. Zharkova⁴² ,
A. Zhelezov²⁰ , X. Z. Zheng⁴ , Y. Zheng⁷ , T. Zhou⁶ , X. Zhou⁸ , Y. Zhou⁷ ,
V. Zhovkovska⁵⁵ , L. Z. Zhu⁷ , X. Zhu⁴ , X. Zhu⁸ , V. Zhukov¹⁶ , J. Zhuo⁴⁶ ,
Q. Zou^{5,7} , D. Zuliani³¹ , G. Zunica⁴⁸ .

¹*School of Physics and Astronomy, Monash University, Melbourne, Australia*

²*Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil*

³*Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil*

⁴*Center for High Energy Physics, Tsinghua University, Beijing, China*

⁵*Institute Of High Energy Physics (IHEP), Beijing, China*

⁶*School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China*

⁷*University of Chinese Academy of Sciences, Beijing, China*

⁸*Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China*

⁹*Consejo Nacional de Rectores (CONARE), San Jose, Costa Rica*

¹⁰*Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France*

¹¹*Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*

¹²*Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France*

¹³*Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France*

¹⁴*Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France*

¹⁵*LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France*

¹⁶*I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*

¹⁷*Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany*

¹⁸*Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*

¹⁹*Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*

²⁰*Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*

²¹*School of Physics, University College Dublin, Dublin, Ireland*

²²*INFN Sezione di Bari, Bari, Italy*

²³*INFN Sezione di Bologna, Bologna, Italy*

²⁴*INFN Sezione di Ferrara, Ferrara, Italy*

²⁵*INFN Sezione di Firenze, Firenze, Italy*

²⁶*INFN Laboratori Nazionali di Frascati, Frascati, Italy*

²⁷*INFN Sezione di Genova, Genova, Italy*

²⁸*INFN Sezione di Milano, Milano, Italy*

²⁹*INFN Sezione di Milano-Bicocca, Milano, Italy*

³⁰*INFN Sezione di Cagliari, Monserrato, Italy*

³¹*Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy*

³²*INFN Sezione di Perugia, Perugia, Italy*

³³*INFN Sezione di Pisa, Pisa, Italy*

³⁴*INFN Sezione di Roma La Sapienza, Roma, Italy*

³⁵*INFN Sezione di Roma Tor Vergata, Roma, Italy*

- ³⁶ *Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- ³⁷ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- ³⁸ *AGH - University of Krakow, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ³⁹ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ⁴⁰ *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ⁴¹ *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ⁴² *Affiliated with an institute covered by a cooperation agreement with CERN*
- ⁴³ *DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁴⁴ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁵ *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁶ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain*
- ⁴⁷ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴⁸ *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴⁹ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁵⁰ *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁵¹ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁵² *University of Birmingham, Birmingham, United Kingdom*
- ⁵³ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁵⁴ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁵ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵⁶ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵⁷ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵⁸ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁹ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁶⁰ *Imperial College London, London, United Kingdom*
- ⁶¹ *Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁶² *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁶³ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁶⁴ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶⁵ *University of Maryland, College Park, MD, United States*
- ⁶⁶ *Los Alamos National Laboratory (LANL), Los Alamos, NM, United States*
- ⁶⁷ *Syracuse University, Syracuse, NY, United States*
- ⁶⁸ *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ³*
- ⁶⁹ *School of Physics and Electronics, Hunan University, Changsha City, China, associated to ⁸*
- ⁷⁰ *Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to ⁴*
- ⁷¹ *Lanzhou University, Lanzhou, China, associated to ⁵*
- ⁷² *School of Physics and Technology, Wuhan University, Wuhan, China, associated to ⁴*
- ⁷³ *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to ¹⁵*
- ⁷⁴ *Eotvos Lorand University, Budapest, Hungary, associated to ⁴⁷*
- ⁷⁵ *Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to ³⁶*
- ⁷⁶ *Universiteit Maastricht, Maastricht, Netherlands, associated to ³⁶*
- ⁷⁷ *Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland, associated to ³⁹*
- ⁷⁸ *Universidade da Coruña, A Coruña, Spain, associated to ⁴³*
- ⁷⁹ *Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden, associated to ⁵⁸*
- ⁸⁰ *University of Michigan, Ann Arbor, MI, United States, associated to ⁶⁷*
- ⁸¹ *Departement de Physique Nucleaire (SPhN), Gif-Sur-Yvette, France*

^a *Universidade de Brasília, Brasília, Brazil*

^b *Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Rio De Janeiro, Brazil*

^c *Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China*

^d *School of Physics and Electronics, Henan University, Kaifeng, China*

^e *LIP6, Sorbonne Universite, Paris, France*

^f *Excellence Cluster ORIGINS, Munich, Germany*

^g *Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras*

^h *Università di Bari, Bari, Italy*

ⁱ *Università degli studi di Bergamo, Bergamo, Italy*

^j *Università di Bologna, Bologna, Italy*

^k *Università di Cagliari, Cagliari, Italy*

^l *Università di Ferrara, Ferrara, Italy*

^m *Università di Firenze, Firenze, Italy*

ⁿ *Università di Genova, Genova, Italy*

^o *Università degli Studi di Milano, Milano, Italy*

^p *Università di Milano Bicocca, Milano, Italy*

^q *Università di Padova, Padova, Italy*

^r *Università di Perugia, Perugia, Italy*

^s *Scuola Normale Superiore, Pisa, Italy*

^t *Università di Pisa, Pisa, Italy*

^u *Università della Basilicata, Potenza, Italy*

^v *Università di Roma Tor Vergata, Roma, Italy*

^w *Università di Siena, Siena, Italy*

^x *Università di Urbino, Urbino, Italy*

^y *Universidad de Alcalá, Alcalá de Henares, Spain*

^z *Department of Physics/Division of Particle Physics, Lund, Sweden*

[†] *Deceased*