



Search for decays of the Higgs boson into a pair of pseudoscalar particles decaying into $b\bar{b}\tau^+\tau^-$ using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector

The ATLAS Collaboration

This paper presents a search for exotic decays of the Higgs boson into a pair of new pseudoscalar particles, $H \rightarrow aa$, where one pseudoscalar decays into a b -quark pair and the other decays into a τ -lepton pair, in the mass range $12 \leq m_a \leq 60$ GeV. The analysis uses pp collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS detector at the LHC, corresponding to an integrated luminosity of 140 fb^{-1} . No significant excess above the Standard Model (SM) prediction is observed. Assuming the SM Higgs boson production cross-section, the search sets upper limits at 95% confidence level on the branching ratio of Higgs bosons decaying into $b\bar{b}\tau^+\tau^-$, $\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-)$, between 2.2% and 3.9% depending on the pseudoscalar mass.

1 Introduction

Following the observation of the Higgs boson H with mass m_H near 125 GeV by the ATLAS and CMS collaborations [1, 2], studies of its properties have been important programs of research. Global analyses of measurements of Higgs boson properties constrain the branching ratio of the Higgs boson into undetected Beyond the Standard Model (BSM) particles to approximately $\mathcal{B}_u \lesssim 12\%$ [3, 4]. Higgs boson decays are particularly sensitive to new physics due to the small total width ($\Gamma_H \approx 4$ MeV). Even very small couplings to new particles can give sizable branching ratios and can be compatible with available measurements [5].

Extensions of the Standard Model (SM) that include new light pseudoscalars, called a -bosons, can give rise to exotic Higgs boson decays $H \rightarrow aa$. Such new light particles appear in theories with an extended Higgs sector [6–10], dark matter models [11–15], models with a first-order electroweak phase transition [16, 17], and theories of neutral naturalness [18, 19]. Signatures of $H \rightarrow aa$ can also arise in models with hidden-sector particles that are singlets under the SM gauge transformations [5, 20–23]. In scenarios where the a -boson mixes with the SM Higgs boson and inherits its Yukawa couplings to fermions, decays of the a -boson into heavy fermions such as b -quarks and τ -leptons are favored, and the process $H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-$, shown in Figure 1, is expected to have a sizeable branching ratio in the mass range $2m_b < m_a < m_H/2$ [5, 24].

This paper presents a search for the exotic Higgs boson decay $H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-$ and uses the full Run-2 dataset of pp collisions at $\sqrt{s} = 13$ TeV recorded with the ATLAS detector corresponding to an integrated luminosity 140 fb^{-1} . The analysis is performed over the mass range $12 < m_a < 60$ GeV and targets the following production modes of the Higgs boson: gluon-gluon fusion (ggF), vector boson fusion (VBF), and associated production with a vector boson (VH). The associated production of a top-antitop-quark pair ($t\bar{t}$) with a Higgs boson provides a negligible signal contribution due to (i) the low cross-section and (ii) the additional b -quarks from the top-quark decays resulting in a reduced signal acceptance. Therefore, this production mode is not considered. Different analysis categories are defined depending on the τ -lepton decay modes, which can be into electrons (e), muons (μ), or hadrons (τ_{had} , also “hadronic taus”). Due to the relatively low mass of the Higgs boson and the four-body final state, the τ -leptons and b -quarks tend to have low transverse momentum (p_T), typically below $p_T \lesssim 50$ GeV. Therefore, the analysis relies on electron or muon triggers, which have lower p_T thresholds compared to triggers using hadronic activity, to select the events of interest and requires an electron or muon in the final state. The analysis defines different event categories depending on τ -lepton decay modes: $e\mu$, $e\tau_{\text{had}}$, and $\mu\tau_{\text{had}}$. The major backgrounds to the signal process are Drell–Yan production of τ -leptons produced in association with heavy-flavor jets, $t\bar{t}$, and non-prompt leptons plus τ_{had} .




Figure 1: Feynman diagram for the leading contribution to the $pp \rightarrow H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-$ process.

τ -lepton decays	$e\mu$	$(e\mu,1B)$	$(e\mu,1b)$	$(e\mu,2b)$
	$\mu\tau_{\text{had}}$	$(\mu\tau_{\text{had}},1B)$	$(\mu\tau_{\text{had}},1b)$	$(\mu\tau_{\text{had}},2b)$
	$e\tau_{\text{had}}$	$(e\tau_{\text{had}},1B)$	$(e\tau_{\text{had}},1b)$	$(e\tau_{\text{had}},2b)$
		1B,0b	0B,1b	0B,2b
		Heavy-flavor jets		

Figure 2: The analysis categories used in the search corresponding to different decay modes of the τ -leptons and strategies for identifying heavy-flavor jets, including merged b -jet pairs (B) and single b -jets (b). Note that the categories are exclusive.

For low m_a , the a -boson has a large Lorentz boost and its decay products can be collimated. Consequently, the $a \rightarrow b\bar{b}$ final state is reconstructed as a single jet that contains the hadronization products of the two b -quarks. While several techniques exist to resolve merged jets [25], most of them are only efficient in the case of high-mass and high- p_T jets. Due to the relatively low mass of the Higgs boson, the merged $a \rightarrow b\bar{b}$ jet will have low p_T . This analysis uses a novel, dedicated algorithm to identify low-mass, merged, “double b -quark” jets (B -jets) [26]. For high m_a , the b -quarks decays tend to be well separated and the reconstructed jets capture the hadronization of a single b -quark (b -jet). The analysis considers events with one B -jet, or one or two b -jets, resulting in nine analysis categories, as shown in Figure 2.

Similar searches in the $b\bar{b}\tau^+\tau^-$ decay channel were performed by the CMS Collaboration [27, 28]. The latest search has placed 95% CL upper limits on $\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-)$ in the range 1.7 – 7.7% for $12 \leq m_a \leq 60$ GeV using 138 fb^{-1} of Run 2 data at $\sqrt{s} = 13$ TeV. This analysis improves the sensitivity of previous results [28] in the low mass regime ($m_a < 20$ GeV) by targeting more final states and using a neural network discriminant to increase the separation of signal from background. This search is also complementary to other searches for $H \rightarrow aa$ decays performed by the ATLAS and CMS collaborations using both $\sqrt{s} = 8$ TeV and $\sqrt{s} = 13$ TeV data in several final states including $\mu^+\mu^-\mu^+\mu^-$ [29–31], $\mu^+\mu^-\tau^+\tau^-$ [32–35], $\tau^+\tau^-\tau^+\tau^-$ [34, 36], $b\bar{b}\mu^+\mu^-$ [37–40], $b\bar{b}b\bar{b}$ [41–43], $\gamma\gamma\gamma\gamma$ [44–46], and $\gamma\gamma g g$ [47].

2 ATLAS detector

The ATLAS detector [48] at the LHC covers nearly the entire solid angle around the collision point.¹ It consists of an inner tracking detector surrounded by a thin superconducting solenoid, electromagnetic

¹ ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the centre of the LHC ring, and the y -axis points upwards. Polar coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$ and is equal to the rapidity $y = \frac{1}{2} \ln \left(\frac{E+p_z c}{E-p_z c} \right)$ in the relativistic limit.

Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta y)^2 + (\Delta \phi)^2}$.

and hadronic calorimeters, and a muon spectrometer incorporating three large superconducting air-core toroidal magnets.

The inner-detector system (ID) is immersed in a 2 T axial magnetic field and provides charged-particle tracking in the range $|\eta| < 2.5$. The high-granularity silicon pixel detector covers the vertex region and typically provides four measurements per track, the first hit generally being in the insertable B-layer (IBL) installed before Run 2 [49, 50]. It is followed by the SemiConductor Tracker (SCT), which usually provides eight measurements per track. These silicon detectors are complemented by the transition radiation tracker (TRT), which enables radially extended track reconstruction up to $|\eta| = 2.0$. The TRT also provides electron identification information based on the fraction of hits (typically 30 in total) above a higher energy-deposit threshold corresponding to transition radiation.

The calorimeter system covers the pseudorapidity range $|\eta| < 4.9$. Within the region $|\eta| < 3.2$, electromagnetic calorimetry is provided by barrel and endcap high-granularity lead/liquid-argon (LAr) calorimeters, with an additional thin LAr presampler covering $|\eta| < 1.8$ to correct for energy loss in material upstream of the calorimeters. Hadronic calorimetry is provided by the steel/scintillator-tile calorimeter, segmented into three barrel structures within $|\eta| < 1.7$, and two copper/LAr hadronic endcap calorimeters. The solid angle coverage is completed with forward copper/LAr and tungsten/LAr calorimeter modules optimised for electromagnetic and hadronic energy measurements respectively.

The muon spectrometer (MS) comprises separate trigger and high-precision tracking chambers measuring the deflection of muons in a magnetic field generated by the superconducting air-core toroidal magnets. The field integral of the toroids ranges between 2.0 and 6.0 T m across most of the detector. Three layers of precision chambers, each consisting of layers of monitored drift tubes, cover the region $|\eta| < 2.7$, complemented by cathode-strip chambers in the forward region, where the background is highest. The muon trigger system covers the range $|\eta| < 2.4$ with resistive-plate chambers in the barrel, and thin-gap chambers in the endcap regions.

The luminosity is measured mainly by the LUCID-2 [51] detector that records Cherenkov light produced in the quartz windows of photomultipliers located close to the beam pipe. Events are selected by the first-level trigger system implemented in custom hardware, followed by selections made by algorithms implemented in software in the high-level trigger [52]. The first-level trigger accepts events from the 40 MHz bunch crossings at a rate below 100 kHz, which the high-level trigger further reduces in order to record complete events to disk at about 1 kHz. A software suite [53] is used in data simulation, in the reconstruction and analysis of real and simulated data, in detector operations, and in the trigger and data acquisition systems of the experiment.

3 Data and simulated event samples

This search uses pp collision data at $\sqrt{s} = 13$ TeV recorded by the ATLAS experiment from 2015 to 2018. Only events that satisfy data quality requirements that ensure the stable operation of the ATLAS detector [54] are considered. The resulting dataset corresponds to an integrated luminosity of $140.1 \pm 1.2 \text{ fb}^{-1}$ [55]. Data are selected using a combination of single-electron, and single-muon, and opposite-flavor dilepton ($e\mu$) triggers [56, 57].

Several Monte Carlo (MC) event generators are used to simulate the signal and background processes. The MC samples are used to optimize the event selection, evaluate efficiencies and acceptances, and to estimate yields. The main SM backgrounds are Drell–Yan production with decays into τ -leptons $Z/\gamma^* \rightarrow \tau^+ \tau^-$

produced in association with jets (Z +jets), top-quark production ($t\bar{t}$ or single top quarks) where at least one of the W bosons decay leptonically, and backgrounds where jets are misidentified as leptons. Other backgrounds from Higgs bosons decaying into τ -leptons $H \rightarrow \tau^+\tau^-$, $Z/\gamma^* \rightarrow \ell^+\ell^-$ ($\ell = e, \mu$), diboson production, and vector bosons produced in association with $t\bar{t}$ are also included.

Samples of Z +jets and W +jets events were produced using the SHERPA 2.2.11 [58] generator with next-to-leading-order (NLO) precision matrix elements for up to two partons, and leading-order (LO) precision for up to five partons in the five-flavor scheme calculated with the COMIX [59] and OPENLOOPS [60–62] libraries. The events were matched with the SHERPA parton shower [63] using the MEPS@NLO prescription [64–67] with a dedicated set of tuned parameters. The NNPDF3.0NNLO set of parton distribution functions (PDF) [68] was used in the sample generation and the samples were normalized to a next-to-next-to-leading-order (NNLO) prediction [69].

Samples of diboson (VV) events were produced with the SHERPA 2.2.1 generator for semileptonic final states or SHERPA 2.2.2 [58] generator for fully leptonic final states. Fully leptonic final states and semileptonic final states, where one boson decays leptonically and the other hadronically, were generated using matrix elements at NLO accuracy in QCD for up to one additional parton and at LO accuracy for up to three additional parton emissions. Samples for the loop-induced processes $gg \rightarrow VV$ were generated using LO-accurate matrix elements for up to one additional parton emission for both the cases of fully leptonic and semileptonic final states. The matrix element calculations were matched and merged with the SHERPA parton shower based on Catani–Seymour dipole factorization [59, 63] using the MEPS@NLO prescription. The virtual QCD corrections were provided by the OPENLOOPS library [60–62]. The NNPDF3.0NNLO PDF set was used [68], along with a dedicated set of tuned parton-shower parameters.

The production of $t\bar{t}$ events was modeled using the POWHEG BOX v2 [70–73] generator at NLO precision in QCD with the NNPDF3.0NLO PDF set and the h_{damp} ² parameter was set to $1.5 m_t$ [74], with the top-quark mass m_t set to 172.5 GeV. The parton shower, hadronization, and underlying event were modeled using PYTHIA 8.230 [75] with the A14 set of tuned parameters [76] and using the NNPDF2.3LO PDF set [77]. The decays of bottom and charm hadrons were performed by EVTGEN 1.6.0 [78].

Single-top-quark production events were modeled with dedicated samples covering s -channel, t -channel, or W -associated (tW) production. All three production modes were modeled using POWHEG BOX v2 [71–73, 79–81] at NLO in QCD with the NNPDF3.0NLO PDF set. For the s -channel and tW production, the calculation is performed in the five-flavor scheme, while the t -channel production uses the four-flavor scheme. Parton shower and hadronization of these events were modeled with PYTHIA 8.230 using the A14 tune and the NNPDF2.3LO PDF set.

The signal event samples include Higgs boson production via ggF, VBF, and VH . For the SM production of the $H \rightarrow \tau^+\tau^-$ background process, only the ggF and VBF production modes are considered. Top-quark-associated production modes of the SM $H \rightarrow \tau^+\tau^-$ background process are negligible.

The sample of Higgs boson production via ggF is generated at NLO accuracy in QCD using POWHEG BOX v2 [71, 72, 79, 82, 83]. The simulation achieves NNLO accuracy for $gg \rightarrow h$ observables by reweighting the Higgs boson rapidity spectrum in HJ-MiNLO [84–86] to that of HNNLO [87]. The Higgs boson production via VBF and VH were simulated with NLO precision using POWHEG BOX v2 [71, 72, 79, 88]. The PDF4LHC15NNLO PDF set [89] was used for all three production modes, while the parton shower and hadronization were performed using PYTHIA 8.244 with the A14 set of tuned parameters.

² The h_{damp} parameter regulates singularities in the emission of hard radiation in POWHEG. The value is chosen to provide good description of the $t\bar{t}$ system p_T .

Each of the simulated Higgs production samples are normalized using dedicated higher-order cross-section calculations. The ggF samples are normalized to the next-to-next-to-next-to-leading-order (N³LO) cross-section in QCD plus electroweak corrections at next-to-leading logarithm (NLL) [90–100]. The VBF samples are normalized to an approximate-NNLO QCD cross-section with NLO electroweak corrections [101–103]. Similarly, the VH samples are normalized to cross-sections calculated at NNLO in QCD with NLO electroweak corrections. The cross section estimate includes the $gg \rightarrow ZH$ contribution, even though it is not simulated explicitly. For the signal samples, the decay $H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-$ is performed using PYTHIA 8.244 and eight distinct mass points were simulated for the ggF, VBF, and the VH production modes in the range $m_a = 12 - 60$ GeV (12, 16, 20, 25, 30, 40, 50 and 60 GeV).

All simulated event samples were processed through a detailed simulation of the ATLAS detector based on GEANT4 [104] or a faster simulation where the full GEANT4 simulation of the calorimeter response is replaced by a detailed parameterization of the shower shapes [105]. The effects of multiple pp interactions in the same and neighboring bunch crossings (pileup) were modeled by overlaying the simulated hard-scattering event with inelastic pp (minimum-bias) events to reproduce the pileup distributions seen in the data. These inelastic events were generated with the soft QCD processes of PYTHIA 8.186 [106] using the NNPDF2.3LO PDF set and the A3 set of tuned parameters [107].

4 Object and event reconstruction

Events are required to contain at least one reconstructed pp collision vertex candidate with at least two associated ID tracks with $p_T > 0.5$ GeV [108]. The primary vertex (PV) is selected as the vertex with the highest sum of the squared transverse momentum of the associated tracks.

Electrons are reconstructed from a seed cluster in the EM calorimeter matched to a track in the ID [109] and are required to have $p_T > 7$ GeV and $|\eta| < 2.47$. Electrons in the calorimeter barrel–endcap transition region ($1.37 < |\eta| < 1.52$) are excluded from the analysis. Electrons from τ -lepton decays can be displaced from the PV due to the long τ -lepton lifetime, so no requirement on the transverse impact parameter (d_0) is applied. A requirement on the longitudinal impact parameter $|z_0 \sin \theta| < 0.5$ mm is applied to match the electron track to the PV of the event. Electrons must satisfy the *Medium* working point of the likelihood identification criteria [109]. Electrons satisfying these requirements are referred to as baseline electrons. Signal electrons are defined by additionally applying a tight isolation requirement using both tracking and calorimetry information with p_T -dependent ΔR cone radius [109]. This isolation variable rejects electrons that likely originated from light- or heavy-flavor hadrons. The isolation criterion is corrected for the presence of a nearby muon by subtracting the p_T of the muon track within the isolation cone of the electron from the isolation sum. This is particularly important for low mass signal samples ($m_a \lesssim 20$ GeV), where the electrons and muons from two τ -lepton decays in $a \rightarrow \tau^+\tau^-$ are often found to be within the isolation cone of each other.

Several methods are used to reconstruct muons, depending on the availability of tracks in the ID and the MS [110]. In order to benefit from the extended MS coverage up to $|\eta| = 2.7$, an ID track is not required in the region $2.5 < |\eta| < 2.7$. Muons are selected with $p_T > 7$ GeV and $|\eta| < 2.7$. As in the case of electrons, the only impact parameter requirement applied is $|z_0 \sin \theta| < 0.5$ mm. Muons must satisfy the *Medium* identification criterion [110], comprising the baseline muon selection. Signal muons are defined with an additional loose requirement on the isolation in both the inner tracking detector and calorimeters also with a p_T -dependent cone radius [110]. As above, an electron within the isolation cone of a muon is excluded from the calculation.

Jets are reconstructed using the anti- k_t algorithm [111] implemented in the FASTJET package [112, 113] with a radius parameter $R = 0.4$. A particle-flow approach [114] is used for the jet reconstruction. The jet energy scale is calibrated to the particle level using simulation and further corrected with in-situ methods [115]. The jet selection requires $p_T > 15$ GeV and $|\eta| < 2.5$. A multivariate jet vertex tagger (JVT) based on tracking information is used to identify a jet as originating from the PV and suppress jets from pileup interactions for jets with $p_T < 60$ GeV and $|\eta| < 2.4$ [116].

The strategy to identify jets containing b -hadrons depends on the kinematics of the signal. For low signal masses ($m_a \lesssim 20$ GeV), pairs of b -hadrons from $a \rightarrow b\bar{b}$ decays tend to be merged in the detector and are identified as a single reconstructed jet. DEXTER [26] is an end-to-end algorithm that classifies jets into three categories: merged b -jets, single b -jets, and other jet flavors. Track-jets associated to jets are reconstructed by reclustering jets with $p_T > 20$ GeV and $|\eta| < 2.0$, together with all ID tracks using an anti- k_t algorithm with radius parameter $R = 0.8$. When the track-jet contains a single $R = 0.4$ jet, the jet is called isolated. An exclusive- k_t clustering of the track-jet tracks into precisely two sub-jets (ex- $k_t^{(2)}$) [117] is used to reconstruct the flight axes of the two b -jets within a track-jet. Both ex- $k_t^{(2)}$ track-subjets are required to have $p_T > 5$ GeV, where the transverse momentum of the subjet is estimated by summing the four-momentum of the associated tracks. The DEXTER algorithm uses the tracks from the $R = 0.8$ track-jets to reconstruct secondary vertices, which provide a distinctive signature of merged b -jets when more than one is reconstructed inside the same jet or when they merge in a single secondary vertex with very high mass. Displaced tracks, secondary vertices, and the properties of the two ex- $k_t^{(2)}$ jets are used in a deep set neural network (NN) to classify the flavor of the jet. The NN exploits the presence of highly displaced tracks along the two flight axes as well as the reconstruction of multiple secondary vertices with large mass to resolve the two b -jets.

In the case of jets which are not merged, the jet flavor is identified with the DL1r algorithm [118] combining track impact parameter values with information from secondary vertices reconstructed within the jet. A working point corresponding to 60% efficiency for identifying B -jets is used for DEXTER, and a working point corresponding to 85% efficiency for identifying b -jets is used for DL1r.

The b -jet identification efficiency of the DL1r algorithm is measured in collider data by using $t\bar{t}$, Z +jets, W +jets, and multijet events [119–121]. A dedicated measurement of the DL1r algorithm identification efficiency for jets with $15 < p_T < 20$ GeV was performed for this search. The B -jet identification efficiency of the DEXTER algorithm is also measured in collider data by using $t\bar{t}$ and $Z+g(\rightarrow b\bar{b})$ events [26]. These measurements are used to correct the identification efficiency of heavy-flavored jets in simulation. The calibration procedure corrects the identification efficiency in simulation as a function of the jet p_T and η to match the one observed in data. Discrepancies between the identification efficiency in simulation and in data stem from both mismodeling of the detector response and of the underlying physics processes.

The reconstruction of τ_{had} candidates is seeded using jets reconstructed using the anti- k_t algorithm with distance parameter $R = 0.4$ whose inputs are topoclusters, three-dimensional clusters of calorimeter cells [122]. Reconstructed τ_{had} candidates have $p_T > 20$ GeV and $|\eta| < 2.5$, excluding the region $1.37 < |\eta| < 1.52$. The identification of τ_{had} uses a recurrent neural network (RNN) algorithm, which uses as inputs tracks and calorimeter clusters associated to τ_{had} candidates, as well as high-level discriminating variables [123]. Baseline τ_{had} candidates are required to pass the *Very Loose* working point, while signal τ_{had} candidates are further required to satisfy the *Medium* working point [123]. A dedicated multivariate electron veto is applied to τ_{had} candidates to reject electrons misidentified as τ_{had} . The *Medium* working point is used for the boosted-decision-tree-based electron veto [124].

A dedicated τ -jet-vertex algorithm (TJVA) is used to associate τ_{had} candidates to a PV [125]. The algorithm does not apply impact parameter requirements and finds the vertex with the largest fraction of the p_{T} from the tracks associated with the τ_{had} within a distance of $R = 0.2$ around the candidate. While TJVA does not apply impact parameter requirements, tracks must be sufficiently close to the τ_{had} vertex, so two selections are applied on the impact parameters of tracks: $|d_0^{\text{TJVA}}| < 1.0$ mm and $|z_0^{\text{TJVA}} \sin \theta| < 1.5$ mm [126].

An overlap removal procedure is applied to prevent double counting of objects. This procedure is applied to the objects described previously that satisfy the baseline criteria, with the exception of muons where candidates satisfying the *VeryLoose* identification criterion are used for the overlap removal and the *Medium* identification criteria is used after overlap removal for the baseline selection criteria. This reduces backgrounds from $Z/\gamma^* \rightarrow \mu^+ \mu^-$ events where a *VeryLoose* muon can be misidentified as a τ_{had} candidate.

The overlap removal procedure is executed as follows. The closest jet within a radius of $\Delta R = 0.2$ of a selected electron is removed. If the nearest jet is within $\Delta R = 0.4$ of an electron, the electron is excluded from the analysis. Muons are excluded from the analysis if they are separated from the nearest jet by $\Delta R < 0.4$, since this reduces the background from heavy-flavor decays inside jets. However, if the jet has fewer than three associated tracks, the muon is selected and the jet is excluded from the analysis instead. This avoids an inefficiency for high-energy muons undergoing significant energy loss in the calorimeter. Electrons are excluded from the analysis if they share their track with a muon.

From the remaining jets, isolated jets that satisfy the 60% DEXTER working point are classified as B -jets. Isolated jets that do not satisfy the 60% DEXTER working point and satisfy the 85% DL1r working point are classified as selected b -jets. Isolated jets that do not satisfy the 60% DEXTER working point and do not satisfy the 85% DL1r working point are not used in the analysis. Non-isolated jets that satisfy the 85% DL1r working point are also classified as b -jets, while those that fail are excluded from the analysis. Muons in the annular region $0.4 < \Delta R < 0.8$ region around a B -jet which have a relative calorimeter-based isolation larger than 0.1 are excluded. Electrons satisfying the same criteria are also removed from the analysis. Finally, any B -jet with either an electron or a muon in the annular $0.4 < \Delta R < 0.8$ region and relative calorimeter-based isolation less than 0.1 are excluded from the analysis.

Muons that are excluded in the overlap removal procedure, but that are within $\Delta R < 0.3$ of a b -jet or within $\Delta R < 0.3$ of either of the $ex\text{-}k_t^{(2)}$ track-subjets of a B -jet are classified as soft muons. The leading soft muon associated to a b -jet or to a $ex\text{-}k_t^{(2)}$ track-subjet is added to the jet four momentum to account for semileptonic decays of b -hadrons in the jet energy.

The missing transverse momentum $\vec{p}_{\text{T}}^{\text{miss}}$, with magnitude $E_{\text{T}}^{\text{miss}}$, is calculated using the magnitude of the vector sum of the p_{T} of all reconstructed objects and any additional tracks not associated to any reconstructed objects in the event [127]. The missing transverse energy ($E_{\text{T}}^{\text{miss}}$) measures the imbalance of the transverse momentum in the detector. In this search $E_{\text{T}}^{\text{miss}}$ is expected to arise from the neutrinos from τ -lepton decays.

5 Event selection

Events are placed in exclusive categories defined by the final state objects from the $\tau^+ \tau^-$ and $b\bar{b}$ systems, as shown in Figure 2. The trigger strategy used to select events depends on the leptons in the final state. For the $e\mu$ categories, three orthogonal trigger regions are defined. If the event has at least one electron with

$p_T > 27$ GeV, the single electron trigger is used and is required to match the selected electron. If the event has no such electron, but has a muon with $p_T > 27$ GeV, the single muon trigger is equivalently used and is required to match the selected muon. If there are no electrons nor muons satisfying the previous criteria, then opposite-flavor $e\mu$ triggers are used, also requiring trigger matching for both leptons. For the $e\tau_{\text{had}}$ and $\mu\tau_{\text{had}}$ categories, the single electron or muon trigger of the corresponding flavor with $p_T > 27$ GeV is used and matched to the electron or muon respectively.

Events in the $e\mu$ categories are required to have exactly one electron and one muon with opposite-sign (OS) charge that satisfy the signal selection criteria. In addition, events should have at most one baseline and no signal τ_{had} candidate. In the $e\tau_{\text{had}}$ and $\mu\tau_{\text{had}}$ categories, events are required to have exactly one electron and one muon, respectively, and exactly one signal τ_{had} with OS charge. For the $e\mu$ categories, the electron and muon are required to be separated by $\Delta R(e, \mu) > 0.1$ and for the $e\tau_{\text{had}}$ and $\mu\tau_{\text{had}}$ categories, this requirement is $\Delta R(e/\mu, \tau_{\text{had}}) > 0.2$, to reduce backgrounds from low-mass hadronic decays. The analysis categories are also defined by requiring one B -jet, or one or two b -jets.

Two additional criteria are applied in each category to define regions enhanced in backgrounds. These criteria are based on the kinematics of the visible τ -lepton decay products, i.e., excluding the neutrinos. The visible mass of the τ -lepton decay products $m^{\text{vis}}(\tau\tau)$ is lower for signal than for the $Z \rightarrow \tau^+\tau^-$ background, due to the lower mass of the a -boson. A requirement is thus applied on the visible mass for all categories: $m^{\text{vis}}(\tau\tau) = m^{\text{vis}}(e\mu) < 45$ GeV for the $e\mu$ categories and $m^{\text{vis}}(\tau\tau) = m^{\text{vis}}(e\tau_{\text{had}})$ or $m^{\text{vis}}(\mu\tau_{\text{had}}) < 60$ GeV for the $e\tau_{\text{had}}$ and $\mu\tau_{\text{had}}$ categories. The difference in the requirements is due to the smaller number of neutrinos in the categories with a hadronic τ -lepton compared to the $e\mu$ case. A low mass requirement on the visible mass of $m^{\text{vis}}(\tau\tau) > 4$ GeV is also applied to reduce backgrounds from hadronic decays. The sample obtained by inverting the higher requirement on the visible mass is used to correct the modeling of the Z +jets background (see Section 6), and is called the “ Z region.” Figure 3(a) demonstrates the ability of the background prediction to model the data in the most sensitive $\mu\tau_{\text{had}}$ category. The background model performs similarly well in the $e\tau_{\text{had}}$ and $e\mu$ categories.

When compared to the background, leptons from signal processes also have lower transverse mass calculated with the τ -lepton visible transverse momentum $p_T^{\text{vis}}(\tau)$ and E_T^{miss} . In the case of leptonic decays of the τ -lepton, the visible p_T is defined as the transverse momentum of the electron or muon. The transverse mass is defined as: $m_T(\tau) = \sqrt{2p_T^{\text{vis}}(\tau)E_T^{\text{miss}}(1 - \cos(\Delta\phi))}$, where $\Delta\phi$ is the difference in the azimuthal angle between $p_T^{\text{vis}}(\tau)$ and E_T^{miss} . A requirement is applied on the sum of the transverse mass calculated for the two τ -leptons $\Sigma m_T < 120$ GeV, where $\Sigma m_T = m_T(\tau^{\text{lead}}) + m_T(\tau^{\text{sublead}})$. Figure 3(b) shows the distribution of Σm_T in the $\mu\tau_{\text{had}}$ categories. The sample obtained by inverting the requirement on Σm_T and is used to correct the modeling of the $t\bar{t}$ background (see Section 6), and is called the “ $t\bar{t}$ region.” There is small signal contamination into the Z and $t\bar{t}$ regions, only ≈ 1 signal event is expected assuming $\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-) = 10\%$.

Finally a same-sign (SS) region, enhanced in backgrounds containing jets that are misidentified as electrons, muons or hadronic taus, is defined by applying the same selection criteria as the signal region but requiring the e , μ , and τ_{had} to have same-sign charge. This region is used to estimate backgrounds with non-prompt leptons (see Section 6). Table 1 summarizes the event selection used in the analysis.




Figure 3: (a) Visible mass $m^{\text{vis}}(\mu\tau_{\text{had}})$ and (b) sum of the transverse mass Σm_T distributions for signal and the expected background. Events with high $m^{\text{vis}}(\mu\tau_{\text{had}})$ and high Σm_T are included in the $t\bar{t}$ region. In order to compare the shapes, the expected signal distribution is shown assuming ten times the production cross section of the Higgs boson and a 100% branching ratio to $b\bar{b}\tau^+\tau^-$. The yields for the backgrounds correspond to the values obtained after applying the corrections to the background modeling described in Section 6. The cuts separating the signal region (SR) from the Z and $t\bar{t}$ regions are indicated by the vertical dashed line. The hashed area represents the total uncertainty of the background. Overflow events are included in the last bins.

Table 1: Event selection for the analysis categories and the background regions. The background regions only list the requirements that are different from the signal region. The definition of signal leptons can be found in Section 4.

Region	$e\mu$	$e\tau_{\text{had}}$ or $\mu\tau_{\text{had}}$
	1 OS signal $e\mu$ pair	1 OS signal $e\tau_{\text{had}}$ or $\mu\tau_{\text{had}}$ pair
	0 signal τ_{had}	1 signal τ_{had}
	$\Delta R(e, \mu) > 0.1$	$\Delta R(\ell, \tau) > 0.2$
Signal region	$4 < m^{\text{vis}}(\tau\tau) < 45 \text{ GeV}$	$4 < m^{\text{vis}}(\tau\tau) < 60 \text{ GeV}$
		$\Sigma m_T < 120 \text{ GeV}$
		1 B -jet or 1 or 2 b -jets
Z region	$m^{\text{vis}}(\tau\tau) > 45 \text{ GeV}$	$m^{\text{vis}}(\tau\tau) > 60 \text{ GeV}$
$t\bar{t}$ region	$\Sigma m_T > 120 \text{ GeV}$, no $m^{\text{vis}}(\tau\tau)$ requirement	
SS region	1 SS signal $e\mu$ pair	1 SS signal $e\tau_{\text{had}}$ or $\mu\tau_{\text{had}}$ pair

6 Background estimation

Several SM processes can produce final states that satisfy the object and event selections described previously. The contributions from these processes are estimated with simulation or data-driven methods. Background processes with all leptons originating from the decay of vector bosons are defined as prompt and described by simulation. The dominant sources of prompt backgrounds are top-quark production, from $t\bar{t}$ and single-top events, and Z +jets where the Z boson decays into pairs of τ -leptons. Other backgrounds estimated from simulation include $h \rightarrow \tau^+\tau^-$, diboson production, $t\bar{t}$ in association with a vector boson, and $Z/\gamma^* \rightarrow \ell^+\ell^-$ where one lepton is misidentified as a τ_{had} in the $e\tau_{\text{had}}$ and $\mu\tau_{\text{had}}$ categories.

The modeling of two of the main prompt backgrounds, Z +jets and $t\bar{t}$, in simulation is corrected to match the data in regions enhanced in these backgrounds (see Section 5). For the $t\bar{t}$ background, the simulation is reweighted to match the number of b -jets and the number of jets in the $t\bar{t}$ region. The $t\bar{t}$ region is devoid of Z +jets events and performing the $t\bar{t}$ correction first minimizes the need for further iterations. In the $e\mu$ categories, an additional correction is applied as a function of the variable H_T , defined as the scalar sum of the p_T of all jets and leptons in the event. The reweighting procedure improves the description of the p_T of the $t\bar{t}$ system and the description of the number of additional jets produced in association with the $t\bar{t}$ system. The same strategy was applied in other ATLAS searches with large $t\bar{t}$ background [128, 129]. After the $t\bar{t}$ reweighting, the Z +jets background is corrected with two normalization factors. The $Z \rightarrow \tau^+\tau^-$ and $Z \rightarrow e^+e^-$ backgrounds are scaled so that the predicted yields from simulation in two $m_{\ell\ell}$ bins match the observed number of events in the Z region.

Background processes with at least one lepton originating from the non-prompt decay of a hadron, from photon conversions, or from the misidentification of other particles are defined as non-prompt backgrounds. A tight-to-loose data-driven method [130] is used to estimate this background contribution. As described in Section 4, each lepton has two criteria baseline and signal. Leptons that satisfy the baseline selection but not the signal one are called loose (L), while those satisfying the signal selection are called tight (T). The estimates of the non-prompt backgrounds are obtained by assigning a weight $w_L = f\varepsilon/(\varepsilon - f)$ to each loose lepton and a weight $w_T = \varepsilon(1 - f)/(\varepsilon - f)$ to each tight lepton [130]. Here, f and ε are, respectively, the rate with which non-prompt and prompt leptons that satisfy the baseline selection also satisfy the signal criteria. The sign of the overall event weight is adjusted depending on the number of loose leptons to avoid double-counting of background events.

The non-prompt rates for electrons and muons are measured with events in the SS region that satisfy $\Delta R(e, \mu) > 1.4$. Events with one baseline muon (electron) and one signal electron (muon) are used to estimate the muon (electron) non-prompt rate f_μ (f_e). The electron and muon non-prompt rates are parameterized as a function of the lepton p_T , η , whether or not the lepton is matched to a trigger object, and the number of b -jets in the event. The prompt rates for electrons (muons) ϵ_e (ϵ_μ) are estimated as a function of the same variables from simulations of $Z \rightarrow \tau^+\tau^-$ events.

The condition $\Delta R(e, \mu) > 1.4$ ensures that the leptons do not interfere in their identification efficiency. When the weights are applied in the $e\mu$ signal region, a geometrical correction is necessary to account for the overlap of isolation cones with $\Delta R(e, \mu) < 0.6$. The correction is applied to events with two loose leptons and is parameterized as:

$$(-w_L^e w_L^\mu)^{\text{corr}} = (1 - f(\Delta R)) (-w_L^e w_L^\mu) + f(\Delta R) \times \frac{1}{2} (w_L^e w_T^\mu + w_T^e w_L^\mu),$$

$$f(\Delta R) = c \times \left(\frac{2}{\pi} \arccos\left(\frac{\Delta R}{2r}\right) - \frac{\Delta R}{\pi r^2} \sqrt{r^2 - \frac{(\Delta R)^2}{4}} \right),$$

where $f(\Delta R)$ is the fraction of the isolation cone that overlaps, c is an arbitrary constant fit from the data in the SS region, and $r = 0.3$ is the radius of the isolation cone.

For hadronic taus, the prompt rate is taken as $\varepsilon_\tau = 1$ in the weights w_L and w_T , and MC simulation is used to remove the contribution from prompt loose taus. The non-prompt rate is measured separately with three different processes: $Z(\rightarrow \mu^+ \mu^-) + \tau_{\text{had}}$, jet + τ_{had} where the τ_{had} candidate has high JVT score, and jet + τ_{had} with low JVT score. In each region, the non-prompt rate is parameterized as a function of the τ_{had} p_T , η , decay mode, and the number of b -jets in the event. The τ_{had} non-prompt rate used to estimate the non-prompt background is written as a linear combination of the three estimates:

$$f_\tau = [f_{Z \rightarrow \mu^+ \mu^-} (1 - r_{\text{QCD}}) + f_{\text{high JVT}} r_{\text{QCD}}] (1 - r_{\text{LJVT}}) + f_{\text{low JVT}} r_{\text{LJVT}},$$

where the coefficient r_{QCD} is the fraction of non-prompt background events with two non-prompt leptons in each $e\tau_{\text{had}}$ and $\mu\tau_{\text{had}}$ categories and the coefficient r_{LJVT} is determined by a maximum-likelihood fit to the baseline τ_{had} seed jet width distribution. The coefficients r_{QCD} and r_{LJVT} are determined separately for each category and region of the analysis, as well as for each τ -lepton decay mode.

7 Analysis strategy

In each analysis category, a NN is trained using the kinematic variables of the reconstructed heavy-flavor jets and τ -leptons. The NN is parameterized as a function of the a -boson mass in order to obtain an optimal discriminant for each different simulated mass [131]. During training, the background is assigned a random value as the value of m_a , while for signal the simulated mass is used. The signal events have different correlations between physical observables and the generated mass than the background, which the network exploits. Once the network is trained, the NN output score is calculated with the true mass replaced by the mass hypothesis under consideration for all events, be they signal, background, or data.

Due to the neutrinos in the decays of the τ -leptons, the four-momentum of the two τ -leptons cannot be directly reconstructed. An algorithm called missing-mass calculator (MMC) [132] uses a Markov-chain MC to perform a maximum likelihood estimate of the neutrinos four-momenta, which is then used to reconstruct the input variables of the parameterized NN (pNN).

Beyond the true a -boson mass $m^{\text{true}}(\tau\tau)$, the pNNs for all categories use the visible mass $m^{\text{vis}}(\tau\tau)$, the MMC-based mass of the neutrino pair $m^{\text{MMC}}(\nu\nu)$ in each $\tau \rightarrow e\nu_\tau \bar{\nu}_e$ or $\tau \rightarrow \mu\nu_\tau \bar{\nu}_\mu$ decay, the missing transverse energy E_T^{miss} , the two transverse masses calculated with the visible p_T of the final-state τ -leptons $m_T(\tau)$, the leading heavy-flavor jet $p_T(b^{\text{lead}})$, and the visible transverse momentum $p_T^{\text{vis}}(\tau\tau b^{\text{lead}})$. The variable D_ζ [133], defined as:

$$D_\zeta = [\vec{p}_T^{\text{miss}} - 0.85\vec{p}_T^{\text{vis}}(\tau\tau)] \cdot \vec{\zeta},$$

Table 2: Neural-network input variables with a summary of the final-state property it describes.


Feature	Description
$m^{\text{true}}(\tau\tau)$	During training: generated a -boson mass for signal MC. Background events are assigned a random value of the eight signal masses. During testing: the mass hypothesis under consideration.
$m^{\text{vis}}(\tau\tau)$	Visible mass of the $\tau\tau$ system.
$p_{\text{T}}(\tau\tau)$	p_{T} of the $\tau\tau$ system.
$m^{\text{MMC}}(\nu\nu)$	MMC-based mass of the two neutrinos in $\tau \rightarrow e\nu_{\tau}\bar{\nu}_e$ or $\tau \rightarrow e\nu_{\tau}\bar{\nu}_{\mu}$ decays.
$E_{\text{T}}^{\text{miss}}$	Missing transverse energy.
$m_{\text{T}}(\tau)$	Transverse mass calculated with the visible p_{T} of the final-state τ -leptons.
$p_{\text{T}}(b^{\text{lead}})$	Transverse momentum of the leading final-state b -jet.
$p_{\text{T}}^{\text{vis}}(\tau\tau b^{\text{lead}})$	Visible p_{T} of the $\tau\tau b^{\text{lead}}$ system.
D_{ζ}	Misalignment between the $\vec{E}_{\text{T}}^{\text{miss}}$ vector and the $\tau\tau$ system.
Categories with a B -jet or 2 b -jets	
$p_{\text{T}}(b^{\text{sublead}})$	Transverse momentum of the subleading final-state b -jet.
$p_{\text{T}}(bb)$	Transverse momentum of the bb system.
$m(bb)$	Mass of the bb system.
$m^{\text{vis}}(bb\tau\tau)$	Visible mass of the Higgs boson system.
$m^{\text{MMC}}(bb\tau\tau)$	MMC-based mass of the Higgs boson system.

where $\vec{\zeta}$ is a unit vector in the direction of the bisector between the two τ -leptons, is also used as input to the network in all categories. D_{ζ} provides an estimate of the misalignment between the missing transverse momentum and the visible $\tau\tau$ subsystem.


In categories with two b -jets or one B -jet, additional input variables are used: the subleading heavy-flavor jet $p_{\text{T}}(b^{\text{sublead}})$, the transverse momentum $p_{\text{T}}(bb)$ and mass $m(bb)$ of the bb subsystem, as well as the visible $m^{\text{vis}}(bb\tau\tau)$ and MMC-based $m^{\text{MMC}}(bb\tau\tau)$ mass estimates of the Higgs boson. In categories with a B -jet, the four-momentum of the two ex - $k_t^{(2)}$ track-subjets are used as proxies for individual heavy-flavor jets. Table 2 summarizes the variables used as input for the NN. Distributions of the pNN input variables for the $(\mu\tau_{\text{had}}, 2b)$ category are shown in Figure 4. Distributions of the pNN(m_a) output variable for a value of $m_a = 30$ GeV in the $(\mu\tau_{\text{had}}, 2b)$ category and for $m_a = 12$ GeV in the $(e\mu, 1B)$ category are shown in Figure 5.

Each pNN is a fully connected network with three hidden layers with 15 neurons each. The activation function is a leaky ReLU activation function with slope of 0.01 when the input is negative. The Objax [134] framework is used with the Adam optimization algorithm [135] and a binary cross entropy loss function. The signal sample used in the NN training contains all $H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-$ simulated samples, which are normalized so that each of them has the expected number of observed events. The background sample used in the NN training is composed of composed of top-quark, $Z(\rightarrow \tau\tau)$ + jets and non-prompt events.


For each a -boson hypothesis m_a , a statistical analysis is performed simultaneously in all nine categories of the analysis. To reduce the correlation among different m_a hypotheses, only events which satisfy $0.95m_a - 6 \text{ GeV} \leq m^{\text{MMC}}(\tau\tau) \leq 1.15m_a + 1 \text{ GeV}$ in the $e\tau_{\text{had}}$ and $\mu\tau_{\text{had}}$ categories, or $0.75m_a - 6 \text{ GeV} \leq m^{\text{MMC}}(\tau\tau) \leq 1.25m_a + 1 \text{ GeV}$ in the $e\mu$ categories are used. The pNN score calculated with this subset of events and $m^{\text{true}}(\tau\tau) = m_a$ is split into three bins with different signal-to-background ratios (S/B). These




(a)



(b)



(c)



(d)

Figure 4: The pNN input variables (a) visible mass $m^{\text{vis}}(\mu\tau_{\text{had}})$, (b) visible transverse momentum $p_{\text{T}}^{\text{vis}}(\mu\tau_{\text{had}})$, (c) transverse mass $m_{\text{T}}(\mu, E_{\text{T}}^{\text{miss}})$, and (d) MMC mass $m^{\text{MMC}}(bb\tau\tau)$ are shown in the SR with no cut on the pNN discriminant. The signal shape is normalized to the same integral as the total background prediction. The hashed area represents the total uncertainty of the background. Overflow events are included in the last bins.




Figure 5: Distribution of $\ln(\text{pNN})$ output score in the (a) $(\mu\tau_{\text{had}}, 2b)$ category with $m_a = 30$ GeV and (b) $(e\mu, 1b)$ category with $m_a = 12$ GeV. The signal shape is normalized to the integral of the total background model. The hashed area represents the total uncertainty of the background. Underflow and overflow events are included in the first and last bins, respectively.

bins are called CR, SR1, and SR2, in order of increasing S/B. Figure 6 shows the ranges for each mass hypothesis tested in this paper while Figure 7 shows the three regions used to search for a signal with $m_a = 30$ GeV.

A test statistic is built from a profile likelihood ratio calculated using the `pyhf` software [136]. The likelihood is given by the product of Poisson probability distributions for the three bins in each of the nine categories and the Gaussian distributions, which implement constraint terms for each source of systematic uncertainty as a function of nuisance parameters (NP).

$$L(\mu, \vec{\alpha}) = \prod_c^{N_{\text{cat}}} \prod_{j=1}^3 \frac{(\mu s_{c,j}(\vec{\alpha}) + b_{c,j}(\vec{\alpha}))^{n_{c,j}}}{n_{c,j}!} e^{-(\mu s_{c,j}(\vec{\alpha}) + b_{c,j}(\vec{\alpha}))} \prod_{k=1}^{N_{\text{syst}}} \frac{1}{\sqrt{2\pi}} e^{-\frac{\alpha_k^2}{2}},$$

where μ is the signal strength, $\vec{\alpha}$ is the vector of nuisance parameters, $s_{c,j}$ and $b_{c,j}$ are the expected number of signal and background events in the j^{th} bin of the category c , and $n_{c,j}$ is the observed number of events. Since the signal templates are normalized to the SM Higgs inclusive cross section, the signal strength is equal to $(\sigma(H)/\sigma_{\text{SM}}(H))\mathcal{B}(H \rightarrow aa \rightarrow bb\tau^+\tau^-)$. Note that the values $s_{c,j}$ and $b_{c,j}$ are themselves functions of the set of NPs. Each systematic uncertainty is fully correlated across all bins and categories.

Categories with very low signal acceptance are suppressed from the likelihood product. For the $m_a = 12$ GeV hypothesis, only the $e\mu$ categories are used. At this mass hypothesis, the cut $\Delta R > 0.2$ between the τ -leptons and lepton removes nearly all signal acceptance. For $m_a > 30$ GeV, the categories with B -jets are similarly not used. Removing these categories does not impact the observed limit due to the lack of signal acceptance. The one-sided alternative likelihood ratio test statistic for upper limits \tilde{q}_μ [137] was used.




Figure 6: Mass distribution of the $\tau\tau$ system for each of the generated MC signal mass points for the (a) $\mu\tau_{\text{had}}$ and (b) $e\mu$ categories. The rectangle above each histogram shows the domain used to test each mass hypothesis. Note that nearby mass hypotheses have overlapping regions. All distributions are normalized assuming $\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-) = 10\%$.




Figure 7: As an illustration, the binning used in the two-dimensional plane of pNN score and mass for the $\mu\tau_{\text{had}}$ categories and the $m_a = 30$ GeV mass hypothesis is shown.

8 Systematic uncertainties

Systematic uncertainties include experimental uncertainties in the reconstructed objects, uncertainties in the data-driven model for the non-prompt-lepton background, and modeling uncertainties in the simulated background and signal samples.

Experimental uncertainties include momentum scale and resolution uncertainties for all reconstructed objects: muons, electrons, hadronic τ -leptons, jets, and missing transverse energy. These uncertainties are estimated by using calibrations performed for each individual object and by comparing them among different simulation models. These uncertainties are considered in the E_T^{miss} reconstruction, and additional uncertainties for soft tracks are estimated from data [127].

Uncertainties in the reconstruction, identification and trigger efficiency of electrons [56, 109] and muons [57, 110] are determined from tag-and-probe efficiency measurements using $Z \rightarrow \ell^+ \ell^-$ and $J/\psi \rightarrow \ell^+ \ell^-$ events. Similarly, uncertainties in the reconstruction and identification efficiency of τ_{had} are determined from tag-and-probe efficiency measurements using $Z \rightarrow \tau_\mu \tau_{\text{had}}$ events (where τ_μ denotes a $\tau \rightarrow \mu \nu_\mu \nu_\tau$ decay) and by varying the physical models used for the simulation of hadronic interactions in the detector [123, 126]. An additional uncertainty is estimated for the electron-veto efficiency based on a tag-and-probe measurement using $Z \rightarrow ee$ events [124].

Uncertainties in the jet vertex tagger efficiency are estimated from efficiency measurements using $Z(\rightarrow \mu^+ \mu^-)$ events with a single jet recoiling against the di-muon pair [116]. A dedicated efficiency measurement for jets with $15 \leq p_T \leq 20$ GeV was performed for this search.

Uncertainties in the DL1r identification efficiency are estimated from a combination of several measurements sensitive to the true jet flavor. The efficiency is measured in several bins of jet p_T , η , and separately for true b -, c -, and light-jets [119–121]. A dedicated efficiency measurement for jets with $15 \leq p_T \leq 20$ GeV was performed for this search. The measurement uses $t\bar{t}$ and Z + jets events to measure the efficiency of the DL1r algorithm in different η bins for b - and non- b -jets (ie., c - and light-jets, which are considered together in this dedicated measurement). The uncertainties in the flavor-tagging efficiency of at low jet p_T are considered uncorrelated with the ones for jet $p_T > 20$ GeV, but the correlations between different jet η bins and between the different true jet flavors are accounted for.

Uncertainties in the DEXTER identification efficiency are estimated from the efficiency measurement using $t\bar{t}$ and $Z(\rightarrow \ell\ell) + g(\rightarrow bb)$ events [26]. The efficiency and associated uncertainties are determined in bins of jet p_T , η , and separately for b - and B -jets. The number of light-jets satisfying the DEXTER identification selection is negligible.

Modeling uncertainties for $t\bar{t}$ and Z +jets background processes and for the Higgs boson signal process are estimated by varying the hard-process renormalization and factorization scales, and by using the NNPDF replicas to estimate PDF uncertainties. For the $t\bar{t}$ background process, additional modeling uncertainties are estimated by comparing the prediction from POWHEG and MADGRAPH5_AMC@NLO [138], by varying the renormalization scale used to simulate the initial- and final-state radiation, and by comparing the prediction using PYTHIA and HERWIG [139] as parton showers. For the Z +jets background process, additional modeling uncertainties in the fraction of heavy-flavored associated jets are estimated by varying the resummation (QSF) and merging (CKKM) scales in SHERPA [58, 63]. For simplicity, the variations are estimated without detector simulation, but in a fiducial region identical to the one used in the search. In addition to generator-based estimates, uncertainties associated with the reweighting procedure in the $t\bar{t}$ and Z regions are propagated to the final result.

Several uncorrelated sources of systematic uncertainties are considered for background events with non-prompt leptons. Statistical uncertainties coming from the limited number of events in the regions used to measure the prompt and non-prompt lepton efficiencies are generally negligible. Statistical uncertainties coming from the limited number of data events with at least one loose lepton are propagated to the final result. In addition to uncertainties of a statistical source, the following uncertainties are considered for background events with non-prompt leptons. Each uncertainty is estimated separately for electrons, muons, and hadronic taus.

- Prompt-lepton efficiency: the uncertainty is estimated by comparing the efficiency determined from simulated Z +jets events with the one estimated from simulated $t\bar{t}$ events.
- Non-prompt-lepton composition: for electrons and muons, this uncertainty is estimated by comparing the non-prompt-lepton efficiency measured in the different analysis categories with the efficiency measured in a region with zero b - and B -jets. For hadronic taus, this uncertainty is obtained by using the same region without heavy-flavored jets to estimate variations on r_{QCD} and r_{LJVT} .
- Prompt-lepton subtraction: the uncertainty is estimated by varying the cross-section of processes with prompt leptons by 30% in the regions used to estimate non-prompt efficiencies.

The uncertainty in the combined 2015-2018 integrated luminosity is 0.83% [55], obtained using the LUCID-2 detector for the main luminosity measurement, and complemented by measurements using the ID and the calorimeters. The uncertainty in the pileup model is determined by varying the estimate of the inelastic pp cross-section between the value estimated by using only the LUCID-2 detector and the value estimated using only the ID.

Table 3: Impact of different group of systematic uncertainties on the upper limit on $(\sigma(H)/\sigma_{\text{SM}}(H))\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-)$. The different systematic uncertainties group are considered individually in each line of the table. The larger the difference relative to the expected limit without systematic uncertainties (stat-only), the more important the uncertainty group is for the final result. The entries with “–” change by less than 1% relative to the stat-only limit.

Uncertainty source	Expected limit on $(\sigma(H)/\sigma_{\text{SM}}(H))\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-)$		
	$m_a = 12 \text{ GeV}$	$m_a = 25 \text{ GeV}$	$m_a = 60 \text{ GeV}$
Stat-only limit	1.34	1.79	3.00
Observed limit	2.89	2.02	3.37
MC statistics	1.42	1.81	3.04
Experimental	2.72	1.94	3.21
Detector response	2.43	1.84	3.03
Luminosity and pileup	1.37	–	–
b -tagging	–	1.81	–
B -tagging	2.35	–	–
Jet and $E_{\text{T}}^{\text{miss}}$	–	1.83	–
Electrons	1.36	–	–
Muons	1.35	–	–
Taus	–	–	–
Data-driven normalization	1.58	1.94	3.19
Non-prompt leptons	1.58	1.85	3.16
Non-prompt taus	–	1.86	3.10
MC reweighting	–	–	–
Theoretical modeling	1.38	1.89	3.04
Signal	–	–	–
Background	1.37	1.87	3.03


9 Results

Different a -boson mass hypotheses are tested $m_a = (12, 16, 20, 25, 30, 40, 50, 60) \text{ GeV}$. Figures 8 and 9 show the pNN distribution for each hypothesis in the three pNN bins used this search for the $(e\mu, 1b)$ and $(e\tau_{\text{had}})$ categories, respectively. While the $m^{\text{MMC}}(\tau\tau)$ window reduces the correlation between different hypotheses, the events in the histogram for different masses are not completely independent. No significant excess over the background expectation is observed for any m_a hypothesis. The largest deviation is a local p -value of 0.055 at $m_a=20 \text{ GeV}$.


Upper limits on $(\sigma(H)/\sigma_{\text{SM}}(H))\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-)$ are determined at 95% confidence level (CL) using the CL_s technique [137] and the \tilde{q}_μ test statistic described in Section 7. The SM Higgs boson production cross-section used is $\sigma_{\text{SM}}(H) = 55.7 \text{ pb}$ [140]. Table 3 shows the impact of dominant sources of systematic uncertainties in the expected upper limit for different m_a hypotheses. The impact is estimated by the difference between the expected upper limit when only a group of uncertainties are considered and the case where no systematic uncertainty is considered. B -jet identification efficiency uncertainty is an important source of systematic uncertainty at low value of m_a due to the difficulty in measuring the identification efficiency of merged B -jets with low mass and low p_{T} . Figure 10(a) compares the

observed upper limits with the expected limits assuming the median value of \tilde{q}_μ under the background-only hypothesis. Figure 10(b) compares the expected limits for categories with heavy-flavor objects (B - or b -jets). The category using a B -jet is especially sensitive in the low-mass regime, where the $b\bar{b}$ system is collimated into a single object. Regarding the τ -lepton-decay mode, the $\mu\tau_{\text{had}}$ channel dominates the sensitivity, except for the low-mass regime where $e\mu$ dominates due to the isolation criterion, which is corrected for the presence of nearby leptons.

This is the first search in ATLAS for exotic decays of the Higgs boson in the final state $H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-$. The observed limit is up to a factor of two better at low masses when compared to previous results [28] due to the use of a novel identification algorithm for low-mass merged B -jets. In several BSM models, final state with b -quarks have large branching ratio and the result provided here can place strict limits on the existence of exotic decays of the Higgs boson.




(a)




(b)

Figure 8: The pNN spectrum in the three bins are shown separately for each mass hypothesis in the (a) $(e\mu, 1b)$ and (b) $(\mu\tau_{\text{had}}, 2b)$ category. The signal shape (normalized to $\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-) = 1$) for each corresponding mass hypothesis is overlaid on top of the SM prediction. The bins divided by vertical dashed lines are used for testing different m_a hypotheses indicated as a text in the figure. Bins used for different m_a hypotheses are not statistically independent. Note that the 12 GeV mass point is not shown for the $(\mu\tau_{\text{had}}, 2b)$ category because the signal acceptance is very low. The hashed area represents the total uncertainty of the background.



(a)



(b)

Figure 9: The pNN spectrum in the three bins are shown separately for each mass hypothesis in the (a) ($e\mu$, $1B$) and (b) ($e\tau_{\text{had}}$, $2b$) categories. The signal shape (normalized to $\mathcal{B}(H \rightarrow aa \rightarrow bb\tau\tau) = 1$) for each corresponding mass hypothesis is overlaid on top of the SM prediction. The bins divided by vertical dashed lines are used for testing different m_a hypotheses indicated as a text in the figure. Bins used for different m_a hypotheses are not statistically independent. The hashed area represents the total uncertainty of the background.




Figure 10: The observed (solid) 95% C.L. upper limits on $(\sigma(H)/\sigma_{\text{SM}}(H))\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-)$ as a function of m_a and the expected (dashed) limits under the background-only hypothesis when (a) combining all categories and (b) considering different categories based on the heavy-flavor objects separately. In the combined plot (a) the inner green and outer yellow shaded bands show the $\pm 1\sigma$ and $\pm 2\sigma$ uncertainties of the expected limits. The mass hypothesis m_a is probed between 12 and 60 GeV for the values shown with markers. A linear interpolation validated with MC simulation between adjacent mass points is used.

10 Conclusion

This paper presents the first search by the ATLAS experiment for exotic decays of the Higgs boson into a pair of pseudoscalar particles, $H \rightarrow aa$, where the a -boson decays into two b -quarks and two τ -leptons. The analysis is performed using the 140 fb^{-1} of proton–proton collision data at a center-of-mass energy of 13 TeV recorded by the ATLAS detector at the LHC between 2015 and 2018. The search makes use of heavy-flavor-tagging techniques to target collimated $a \rightarrow b\bar{b}$ decays and a mass-parameterized neural network to enhance the sensitivity. The analysis categorizes events depending on the τ -lepton decay modes: $e\mu$, $e\tau_{\text{had}}$, and $\mu\tau_{\text{had}}$, as well as the strategy for identifying heavy-flavor jets: a merged b -jet pair, B -jet, and one or two single b -jets. No significant excess above the SM background expectation is observed, and upper limits at 95% confidence level are set on $\mathcal{B}(H \rightarrow aa \rightarrow b\bar{b}\tau^+\tau^-)$ of 2.2% – 3.9% for pseudoscalar mass values in the range $12 \text{ GeV} \leq m_a \leq 60 \text{ GeV}$, assuming the SM Higgs boson production cross-section. These results contribute to the broad program of searches for $H \rightarrow aa$ decays in ATLAS and can be used to set constraints on a variety of BSM scenarios featuring such exotic Higgs decays.

Acknowledgements

We thank CERN for the very successful operation of the LHC and its injectors, as well as the support staff at CERN and at our institutions worldwide without whom ATLAS could not be operated efficiently.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF/SFU (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain),

RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [141].

We gratefully acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMFWF and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; D NRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ISF and Benozziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MNiSW, Poland; FCT, Portugal; MNE/IFA, Romania; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DSI/NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taipei; TENMAK, Türkiye; STFC, United Kingdom; DOE and NSF, United States of America.

Individual groups and members have received support from BCKDF, CANARIE, CRC and DRAC, Canada; CERN-CZ, FORTE and PRIMUS, Czech Republic; COST, ERC, ERDF, Horizon 2020, ICSC-NextGenerationEU and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex, Investissements d’Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

In addition, individual members wish to acknowledge support from Armenia: Yerevan Physics Institute (FAPERJ); CERN: European Organization for Nuclear Research (CERN PJAS); Chile: Agencia Nacional de Investigación y Desarrollo (FONDECYT 1230812, FONDECYT 1230987, FONDECYT 1240864); China: Chinese Ministry of Science and Technology (MOST-2023YFA1605700), National Natural Science Foundation of China (NSFC - 12175119, NSFC 12275265, NSFC-12075060); Czech Republic: Czech Science Foundation (GACR - 24-11373S), Ministry of Education Youth and Sports (FORTE CZ.02.01.01/00/22_008/0004632), PRIMUS Research Programme (PRIMUS/21/SCI/017); EU: H2020 European Research Council (ERC - 101002463); European Union: European Research Council (ERC - 948254, ERC 101089007), Horizon 2020 Framework Programme (MUCCA - CHIST-ERA-19-XAI-00), European Union, Future Artificial Intelligence Research (FAIR-NextGenerationEU PE00000013), Italian Center for High Performance Computing, Big Data and Quantum Computing (ICSC, NextGenerationEU); France: Agence Nationale de la Recherche (ANR-20-CE31-0013, ANR-21-CE31-0013, ANR-21-CE31-0022), Investissements d’Avenir Labex (ANR-11-LABX-0012); Germany: Baden-Württemberg Stiftung (BW Stiftung-Postdoc Eliteprogramme), Deutsche Forschungsgemeinschaft (DFG - 469666862, DFG - CR 312/5-2); Italy: Istituto Nazionale di Fisica Nucleare (ICSC, NextGenerationEU); Japan: Japan Society for the Promotion of Science (JSPS KAKENHI JP22H01227, JSPS KAKENHI JP22H04944, JSPS KAKENHI JP22KK0227, JSPS KAKENHI JP23KK0245); Netherlands: Netherlands Organisation for Scientific Research (NWO Veni 2020 - VI.Veni.202.179); Norway: Research Council of Norway (RCN-314472); Poland: Polish National Agency for Academic Exchange (PPN/PPO/2020/1/00002/U/00001), Polish National Science Centre (NCN 2021/42/E/ST2/00350, NCN OPUS nr 2022/47/B/ST2/03059, NCN UMO-2019/34/E/ST2/00393, UMO-2020/37/B/ST2/01043, UMO-2021/40/C/ST2/00187, UMO-2022/47/O/ST2/00148, UMO-2023/49/B/ST2/04085); Slovenia: Slovenian Research Agency (ARIS grant J1-3010); Spain: Generalitat Valenciana (Artemisa, FEDER, IDIFEDER/2018/048), Ministry of Science and Innovation (MCIN & NextGenEU PCI2022-135018-2, MICIN & FEDER PID2021-125273NB, RYC2019-028510-I, RYC2020-030254-I, RYC2021-031273-I, RYC2022-038164-I), PROMETEO and

GenT Programmes Generalitat Valenciana (CIDEAGENT/2019/027); Sweden: Swedish Research Council (Swedish Research Council 2023-04654, VR 2018-00482, VR 2022-03845, VR 2022-04683, VR 2023-03403, VR grant 2021-03651), Knut and Alice Wallenberg Foundation (KAW 2018.0157, KAW 2018.0458, KAW 2019.0447, KAW 2022.0358); Switzerland: Swiss National Science Foundation (SNSF - PCEFP2_194658); United Kingdom: Leverhulme Trust (Leverhulme Trust RPG-2020-004), Royal Society (NIF-R1-231091); United States of America: U.S. Department of Energy (ECA DE-AC02-76SF00515), Neubauer Family Foundation.

References

- [1] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1, arXiv: [1207.7214 \[hep-ex\]](#).
- [2] CMS Collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30, arXiv: [1207.7235 \[hep-ex\]](#).
- [3] ATLAS Collaboration, *A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery*, *Nature* **607** (2022) 52, arXiv: [2207.00092 \[hep-ex\]](#), Erratum: *Nature* **612** (2022) E24.
- [4] CMS Collaboration, *A portrait of the Higgs boson by the CMS experiment ten years after the discovery*, *Nature* **607** (2022) 60, arXiv: [2207.00043 \[hep-ex\]](#), Erratum: *Nature* **623** (2023) E4.
- [5] D. Curtin et al., *Exotic decays of the 125 GeV Higgs boson*, *Phys. Rev. D* **90** (2014) 075004, arXiv: [1312.4992 \[hep-ph\]](#).
- [6] B. A. Dobrescu and K. T. Matchev, *Light axion within the next-to-minimal supersymmetric standard model*, *JHEP* **09** (2000) 031, arXiv: [hep-ph/0008192 \[hep-ph\]](#).
- [7] U. Ellwanger, J. F. Gunion, C. Hugonie, and S. Moretti, *Towards a no lose theorem for NMSSM Higgs discovery at the LHC*, (2003), arXiv: [hep-ph/0305109](#).
- [8] R. Dermisek and J. F. Gunion, *Escaping the large fine tuning and little hierarchy problems in the next to minimal supersymmetric model and $h \rightarrow aa$ decays*, *Phys. Rev. Lett.* **95** (2005) 041801, arXiv: [hep-ph/0502105](#).
- [9] S. Chang, R. Dermisek, J. F. Gunion, and N. Weiner, *Nonstandard Higgs Boson Decays*, *Ann. Rev. Nucl. Part. Sci.* **58** (2008) 75, arXiv: [0801.4554 \[hep-ph\]](#).
- [10] D. E. Morrissey and A. Pierce, *Modified Higgs Boson Phenomenology from Gauge or Gaugino Mediation in the NMSSM*, *Phys. Rev. D* **78** (2008) 075029, arXiv: [0807.2259 \[hep-ph\]](#).
- [11] V. Silveira and A. Zee, *Scalar Phantoms*, *Phys. Lett. B* **161** (1985) 136.
- [12] M. Pospelov, A. Ritz, and M. B. Voloshin, *Secluded WIMP Dark Matter*, *Phys. Lett. B* **662** (2008) 53, arXiv: [0711.4866 \[hep-ph\]](#).
- [13] P. Draper, T. Liu, C. E. M. Wagner, L.-T. Wang, and H. Zhang, *Dark Light Higgs*, *Phys. Rev. Lett.* **106** (2011) 121805, arXiv: [1009.3963 \[hep-ph\]](#).
- [14] S. Ipek, D. McKeen, and A. E. Nelson, *A Renormalizable Model for the Galactic Center Gamma Ray Excess from Dark Matter Annihilation*, *Phys. Rev. D* **90** (2014) 055021, arXiv: [1404.3716 \[hep-ph\]](#).
- [15] A. Martin, J. Shelton, and J. Unwin, *Fitting the Galactic Center Gamma-Ray Excess with Cascade Annihilations*, *Phys. Rev. D* **90** (2014) 103513, arXiv: [1405.0272 \[hep-ph\]](#).

- [16] J. Kozaczuk, M. J. Ramsey-Musolf, and J. Shelton, *Exotic Higgs boson decays and the electroweak phase transition*, *Phys. Rev. D* **101** (2020) 115035, arXiv: [1911.10210 \[hep-ph\]](#).
- [17] M. Carena, Z. Liu, and Y. Wang, *Electroweak phase transition with spontaneous Z_2 -breaking*, *JHEP* **08** (2020) 107, arXiv: [1911.10206 \[hep-ph\]](#).
- [18] N. Craig, A. Katz, M. Strassler, and R. Sundrum, *Naturalness in the Dark at the LHC*, *JHEP* **07** (2015) 105, arXiv: [1501.05310 \[hep-ph\]](#).
- [19] D. Curtin and C. B. Verhaaren, *Discovering Uncolored Naturalness in Exotic Higgs Decays*, *JHEP* **12** (2015) 072, arXiv: [1506.06141 \[hep-ph\]](#).
- [20] T. Robens and T. Stefaniak, *Status of the Higgs Singlet Extension of the Standard Model after LHC Run 1*, *Eur. Phys. J. C* **75** (2015) 104, arXiv: [1501.02234 \[hep-ph\]](#).
- [21] T. Robens and T. Stefaniak, *LHC benchmark scenarios for the real Higgs singlet extension of the standard model*, *Eur. Phys. J. C* **76** (2016) 268, arXiv: [1601.07880 \[hep-ph\]](#).
- [22] T. Robens, T. Stefaniak, and J. Wittbrodt, *Two-real-scalar-singlet extension of the SM: LHC phenomenology and benchmark scenarios*, *Eur. Phys. J. C* **80** (2020) 151, arXiv: [1908.08554 \[hep-ph\]](#).
- [23] M. Bauer, M. Neubert, and A. Thamm, *Collider Probes of Axion-Like Particles*, *JHEP* **12** (2017) 044, arXiv: [1708.00443 \[hep-ph\]](#).
- [24] U. Haisch, J. F. Kamenik, A. Malinauskas, and M. Spira, *Collider constraints on light pseudoscalars*, *JHEP* **03** (2018) 178, arXiv: [1802.02156 \[hep-ph\]](#).
- [25] A. J. Larkoski, I. Moulton, and B. Nachman, *Jet Substructure at the Large Hadron Collider: A Review of Recent Advances in Theory and Machine Learning*, *Phys. Rept.* **841** (2020) 1, arXiv: [1709.04464 \[hep-ph\]](#).
- [26] ATLAS Collaboration, *DeXTer: Deep Sets based Neural Networks for Low- p_T $X \rightarrow b\bar{b}$ Identification in ATLAS*, ATL-PHYS-PUB-2022-042, 2022, URL: <https://cds.cern.ch/record/2825434>.
- [27] CMS Collaboration, *Search for an exotic decay of the Higgs boson to a pair of light pseudoscalars in the final state with two b quarks and two τ leptons in proton–proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **785** (2018) 462, arXiv: [1805.10191 \[hep-ex\]](#).
- [28] CMS Collaboration, *Search for exotic decays of the Higgs boson to a pair of pseudoscalars in the $\mu\mu b\bar{b}$ and $\tau\tau b\bar{b}$ final states*, *Eur. Phys. J. C* **84** (2024) 493, arXiv: [2402.13358 \[hep-ex\]](#).
- [29] ATLAS Collaboration, *Search for new light gauge bosons in Higgs boson decays to four-lepton final states in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector at the LHC*, *Phys. Rev. D* **92** (2015) 092001, arXiv: [1505.07645 \[hep-ex\]](#).
- [30] ATLAS Collaboration, *Search for Higgs boson decays to beyond-the-Standard-Model light bosons in four-lepton events with the ATLAS detector at $\sqrt{s} = 13$ TeV*, *JHEP* **06** (2018) 166, arXiv: [1802.03388 \[hep-ex\]](#).
- [31] ATLAS Collaboration, *Search for Higgs bosons decaying into new spin-0 or spin-1 particles in four-lepton final states with the ATLAS detector with 139fb^{-1} of pp collision data at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2022) 041, arXiv: [2110.13673 \[hep-ex\]](#).

- [32] ATLAS Collaboration, *Search for Higgs bosons decaying to aa in the $\mu\mu\tau\tau$ final state in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS experiment*, *Phys. Rev. D* **92** (2015) 052002, arXiv: 1505.01609 [hep-ex].
- [33] CMS Collaboration, *Search for an exotic decay of the Higgs boson to a pair of light pseudoscalars in the final state of two muons and two τ leptons in proton–proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **11** (2018) 018, arXiv: 1805.04865 [hep-ex].
- [34] CMS Collaboration, *Search for light pseudoscalar boson pairs produced from decays of the 125 GeV Higgs boson in final states with two muons and two nearby tracks in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **800** (2020) 135087, arXiv: 1907.07235 [hep-ex].
- [35] CMS Collaboration, *Search for a light pseudoscalar Higgs boson in the boosted $\mu\mu\tau\tau$ final state in proton–proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **08** (2020) 139, arXiv: 2005.08694 [hep-ex].
- [36] CMS Collaboration, *Search for a very light NMSSM Higgs boson produced in decays of the 125 GeV scalar boson and decaying into τ leptons in pp collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **01** (2016) 079, arXiv: 1510.06534 [hep-ex].
- [37] ATLAS Collaboration, *Search for Higgs boson decays into a pair of light bosons in the $bb\mu\mu$ final state in pp collision at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **790** (2019) 1, arXiv: 1807.00539 [hep-ex].
- [38] ATLAS Collaboration, *Search for Higgs boson decays into a pair of pseudoscalar particles in the $bb\mu\mu$ final state with the ATLAS detector in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **105** (2022) 012006, arXiv: 2110.00313 [hep-ex].
- [39] CMS Collaboration, *Search for light bosons in decays of the 125 GeV Higgs boson in proton–proton collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **10** (2017) 076, arXiv: 1701.02032 [hep-ex].
- [40] CMS Collaboration, *Search for an exotic decay of the Higgs boson to a pair of light pseudoscalars in the final state with two muons and two b quarks in pp collisions at 13 TeV*, *Phys. Lett. B* **795** (2019) 398, arXiv: 1812.06359 [hep-ex].
- [41] ATLAS Collaboration, *Search for the Higgs boson produced in association with a W boson and decaying to four b -quarks via two spin-zero particles in pp collisions at 13 TeV with the ATLAS detector*, *Eur. Phys. J. C* **76** (2016) 605, arXiv: 1606.08391 [hep-ex].
- [42] ATLAS Collaboration, *Search for the Higgs boson produced in association with a vector boson and decaying into two spin-zero particles in the $H \rightarrow aa \rightarrow 4b$ channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **10** (2018) 031, arXiv: 1806.07355 [hep-ex].
- [43] ATLAS Collaboration, *Search for Higgs boson decays into two new low-mass spin-0 particles in the $4b$ channel with the ATLAS detector using pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **102** (2020) 112006, arXiv: 2005.12236 [hep-ex].
- [44] ATLAS Collaboration, *Search for new phenomena in events with at least three photons collected in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **76** (2016) 210, arXiv: 1509.05051 [hep-ex].
- [45] CMS Collaboration, *Search for the exotic decay of the Higgs boson into two light pseudoscalars with four photons in the final state in proton–proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2023) 148, arXiv: 2208.01469 [hep-ex].

- [46] CMS Collaboration, *Search for exotic Higgs boson decays $H \rightarrow \mathcal{A}\mathcal{A} \rightarrow 4\gamma$ with events containing two merged diphotons in proton–proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **131** (2023) 101801, arXiv: 2209.06197 [hep-ex].
- [47] ATLAS Collaboration, *Search for Higgs boson decays into pairs of light (pseudo)scalar particles in the $\gamma\gamma jj$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **782** (2018) 750, arXiv: 1803.11145 [hep-ex].
- [48] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, *JINST* **3** (2008) S08003.
- [49] ATLAS Collaboration, *ATLAS Insertable B-Layer: Technical Design Report*, ATLAS-TDR-19; CERN-LHCC-2010-013, 2010, URL: <https://cds.cern.ch/record/1291633>, Addendum: ATLAS-TDR-19-ADD-1; CERN-LHCC-2012-009, 2012, URL: <https://cds.cern.ch/record/1451888>.
- [50] B. Abbott et al., *Production and integration of the ATLAS Insertable B-Layer*, *JINST* **13** (2018) T05008, arXiv: 1803.00844 [physics.ins-det].
- [51] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*, *JINST* **13** (2018) P07017.
- [52] ATLAS Collaboration, *Performance of the ATLAS trigger system in 2015*, *Eur. Phys. J. C* **77** (2017) 317, arXiv: 1611.09661 [hep-ex].
- [53] ATLAS Collaboration, *Software and computing for Run 3 of the ATLAS experiment at the LHC*, (2024), arXiv: 2404.06335 [hep-ex].
- [54] ATLAS Collaboration, *ATLAS data quality operations and performance for 2015–2018 data-taking*, *JINST* **15** (2020) P04003, arXiv: 1911.04632 [physics.ins-det].
- [55] ATLAS Collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC*, *Eur. Phys. J. C* **83** (2023) 982, arXiv: 2212.09379 [hep-ex].
- [56] ATLAS Collaboration, *Performance of electron and photon triggers in ATLAS during LHC Run 2*, *Eur. Phys. J. C* **80** (2020) 47, arXiv: 1909.00761 [hep-ex].
- [57] ATLAS Collaboration, *Performance of the ATLAS muon triggers in Run 2*, *JINST* **15** (2020) P09015, arXiv: 2004.13447 [physics.ins-det].
- [58] E. Bothmann et al., *Event generation with Sherpa 2.2*, *SciPost Phys.* **7** (2019) 034, arXiv: 1905.09127 [hep-ph].
- [59] T. Gleisberg and S. Höche, *Comix, a new matrix element generator*, *JHEP* **12** (2008) 039, arXiv: 0808.3674 [hep-ph].
- [60] F. Buccioni et al., *OpenLoops 2*, *Eur. Phys. J. C* **79** (2019) 866, arXiv: 1907.13071 [hep-ph].
- [61] F. Cascioli, P. Maierhöfer, and S. Pozzorini, *Scattering Amplitudes with Open Loops*, *Phys. Rev. Lett.* **108** (2012) 111601, arXiv: 1111.5206 [hep-ph].
- [62] A. Denner, S. Dittmaier, and L. Hofer, *COLLIER: A fortran-based complex one-loop library in extended regularizations*, *Comput. Phys. Commun.* **212** (2017) 220, arXiv: 1604.06792 [hep-ph].

- [63] S. Schumann and F. Krauss,
A parton shower algorithm based on Catani–Seymour dipole factorisation, **JHEP** **03** (2008) 038,
arXiv: [0709.1027 \[hep-ph\]](#).
- [64] S. Höche, F. Krauss, M. Schönherr, and F. Siegert,
A critical appraisal of NLO+PS matching methods, **JHEP** **09** (2012) 049,
arXiv: [1111.1220 \[hep-ph\]](#).
- [65] S. Höche, F. Krauss, M. Schönherr, and F. Siegert,
QCD matrix elements + parton showers. The NLO case, **JHEP** **04** (2013) 027,
arXiv: [1207.5030 \[hep-ph\]](#).
- [66] S. Catani, F. Krauss, B. R. Webber, and R. Kuhn, *QCD Matrix Elements + Parton Showers*,
JHEP **11** (2001) 063, arXiv: [hep-ph/0109231](#).
- [67] S. Höche, F. Krauss, S. Schumann, and F. Siegert, *QCD matrix elements and truncated showers*,
JHEP **05** (2009) 053, arXiv: [0903.1219 \[hep-ph\]](#).
- [68] NNPDF Collaboration, R. D. Ball, et al., *Parton distributions for the LHC run II*,
JHEP **04** (2015) 040, arXiv: [1410.8849 \[hep-ph\]](#).
- [69] C. Anastasiou, L. Dixon, K. Melnikov, and F. Petriello, *High-precision QCD at hadron colliders: Electroweak gauge boson rapidity distributions at next-to-next-to leading order*,
Phys. Rev. D **69** (2004) 094008, arXiv: [hep-ph/0312266](#).
- [70] S. Frixione, G. Ridolfi, and P. Nason,
A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction,
JHEP **09** (2007) 126, arXiv: [0707.3088 \[hep-ph\]](#).
- [71] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*,
JHEP **11** (2004) 040, arXiv: [hep-ph/0409146](#).
- [72] S. Frixione, P. Nason, and C. Oleari,
Matching NLO QCD computations with parton shower simulations: the POWHEG method,
JHEP **11** (2007) 070, arXiv: [0709.2092 \[hep-ph\]](#).
- [73] S. Alioli, P. Nason, C. Oleari, and E. Re,
NLO single-top production matched with shower in POWHEG: s- and t-channel contributions,
JHEP **09** (2009) 111, arXiv: [0907.4076 \[hep-ph\]](#), Erratum: **JHEP** **02** (2010) 011.
- [74] ATLAS Collaboration, *Studies on top-quark Monte Carlo modelling for Top2016*,
ATL-PHYS-PUB-2016-020, 2016, URL: <https://cds.cern.ch/record/2216168>.
- [75] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, **Comput. Phys. Commun.** **191** (2015) 159,
arXiv: [1410.3012 \[hep-ph\]](#).
- [76] ATLAS Collaboration, *ATLAS Pythia 8 tunes to 7 TeV data*, ATL-PHYS-PUB-2014-021, 2014,
URL: <https://cds.cern.ch/record/1966419>.
- [77] NNPDF Collaboration, R. D. Ball, et al., *Parton distributions with LHC data*,
Nucl. Phys. B **867** (2013) 244, arXiv: [1207.1303 \[hep-ph\]](#).
- [78] D. J. Lange, *The EvtGen particle decay simulation package*,
Nucl. Instrum. Meth. A **462** (2001) 152.
- [79] S. Alioli, P. Nason, C. Oleari, and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, **JHEP** **06** (2010) 043,
arXiv: [1002.2581 \[hep-ph\]](#).

- [80] R. Frederix, E. Re, and P. Torrielli, *Single-top t -channel hadroproduction in the four-flavour scheme with POWHEG and aMC@NLO*, [JHEP **09** \(2012\) 130](#), arXiv: [1207.5391 \[hep-ph\]](#).
- [81] E. Re, *Single-top Wt -channel production matched with parton showers using the POWHEG method*, [Eur. Phys. J. C **71** \(2011\) 1547](#), arXiv: [1009.2450 \[hep-ph\]](#).
- [82] K. Hamilton, P. Nason, E. Re, and G. Zanderighi, *NNLOPS simulation of Higgs boson production*, [JHEP **10** \(2013\) 222](#), arXiv: [1309.0017 \[hep-ph\]](#).
- [83] K. Hamilton, P. Nason, and G. Zanderighi, *Finite quark-mass effects in the NNLOPS POWHEG+MiNLO Higgs generator*, [JHEP **05** \(2015\) 140](#), arXiv: [1501.04637 \[hep-ph\]](#).
- [84] K. Hamilton, P. Nason, and G. Zanderighi, *MINLO: multi-scale improved NLO*, [JHEP **10** \(2012\) 155](#), arXiv: [1206.3572 \[hep-ph\]](#).
- [85] J. M. Campbell et al., *NLO Higgs boson production plus one and two jets using the POWHEG BOX, MadGraph4 and MCFM*, [JHEP **07** \(2012\) 092](#), arXiv: [1202.5475 \[hep-ph\]](#).
- [86] K. Hamilton, P. Nason, C. Oleari, and G. Zanderighi, *Merging $H/W/Z + 0$ and 1 jet at NLO with no merging scale: a path to parton shower + NNLO matching*, [JHEP **05** \(2013\) 082](#), arXiv: [1212.4504 \[hep-ph\]](#).
- [87] S. Catani and M. Grazzini, *Next-to-Next-to-Leading-Order Subtraction Formalism in Hadron Collisions and its Application to Higgs-Boson Production at the Large Hadron Collider*, [Phys. Rev. Lett. **98** \(2007\) 222002](#), arXiv: [hep-ph/0703012 \[hep-ph\]](#).
- [88] P. Nason and C. Oleari, *NLO Higgs boson production via vector-boson fusion matched with shower in POWHEG*, [JHEP **02** \(2010\) 037](#), arXiv: [0911.5299 \[hep-ph\]](#).
- [89] J. Butterworth et al., *PDF4LHC recommendations for LHC Run II*, [J. Phys. G **43** \(2016\) 023001](#), arXiv: [1510.03865 \[hep-ph\]](#).
- [90] D. de Florian et al., *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, (2017), arXiv: [1610.07922 \[hep-ph\]](#).
- [91] C. Anastasiou et al., *High precision determination of the gluon fusion Higgs boson cross-section at the LHC*, [JHEP **05** \(2016\) 058](#), arXiv: [1602.00695 \[hep-ph\]](#).
- [92] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog, and B. Mistlberger, *Higgs Boson Gluon-Fusion Production in QCD at Three Loops*, [Phys. Rev. Lett. **114** \(2015\) 212001](#), arXiv: [1503.06056 \[hep-ph\]](#).
- [93] F. Dulat, A. Lazopoulos, and B. Mistlberger, *iHixs 2 – Inclusive Higgs cross sections*, [Comput. Phys. Commun. **233** \(2018\) 243](#), arXiv: [1802.00827 \[hep-ph\]](#).
- [94] R. V. Harlander and K. J. Ozeren, *Finite top mass effects for hadronic Higgs production at next-to-next-to-leading order*, [JHEP **11** \(2009\) 088](#), arXiv: [0909.3420 \[hep-ph\]](#).
- [95] R. V. Harlander and K. J. Ozeren, *Top mass effects in Higgs production at next-to-next-to-leading order QCD: Virtual corrections*, [Phys. Lett. B **679** \(2009\) 467](#), arXiv: [0907.2997 \[hep-ph\]](#).

- [96] R. V. Harlander, H. Mantler, S. Marzani, and K. J. Ozeren, *Higgs production in gluon fusion at next-to-next-to-leading order QCD for finite top mass*, [Eur. Phys. J. C **66** \(2010\) 359](#), arXiv: [0912.2104 \[hep-ph\]](#).
- [97] A. Pak, M. Rogal, and M. Steinhauser, *Finite top quark mass effects in NNLO Higgs boson production at LHC*, [JHEP **02** \(2010\) 025](#), arXiv: [0911.4662 \[hep-ph\]](#).
- [98] S. Actis, G. Passarino, C. Sturm, and S. Uccirati, *NLO electroweak corrections to Higgs boson production at hadron colliders*, [Phys. Lett. B **670** \(2008\) 12](#), arXiv: [0809.1301 \[hep-ph\]](#).
- [99] S. Actis, G. Passarino, C. Sturm, and S. Uccirati, *NNLO computational techniques: The cases $H \rightarrow \gamma\gamma$ and $H \rightarrow gg$* , [Nucl. Phys. B **811** \(2009\) 182](#), arXiv: [0809.3667 \[hep-ph\]](#).
- [100] M. Bonetti, K. Melnikov, and L. Tancredi, *Higher order corrections to mixed QCD-EW contributions to Higgs boson production in gluon fusion*, [Phys. Rev. D **97** \(2018\) 056017](#), arXiv: [1801.10403 \[hep-ph\]](#), Erratum: [Phys. Rev. D **97** \(2018\) 099906\(E\)](#).
- [101] M. Ciccolini, A. Denner, and S. Dittmaier, *Strong and Electroweak Corrections to the Production of a Higgs Boson + 2 Jets via Weak Interactions at the Large Hadron Collider*, [Phys. Rev. Lett. **99** \(2007\) 161803](#), arXiv: [0707.0381 \[hep-ph\]](#).
- [102] M. Ciccolini, A. Denner, and S. Dittmaier, *Electroweak and QCD corrections to Higgs production via vector-boson fusion at the CERN LHC*, [Phys. Rev. D **77** \(2008\) 013002](#), arXiv: [0710.4749 \[hep-ph\]](#).
- [103] P. Bolzoni, F. Maltoni, S.-O. Moch, and M. Zaro, *Higgs Boson Production via Vector-Boson Fusion at Next-to-Next-to-Leading Order in QCD*, [Phys. Rev. Lett. **105** \(2010\) 011801](#), arXiv: [1003.4451 \[hep-ph\]](#).
- [104] GEANT4 Collaboration, *GEANT4—a simulation toolkit*, [Nucl. Instrum. Meth. A **506** \(2003\) 250](#).
- [105] ATLAS Collaboration, *The ATLAS Simulation Infrastructure*, [Eur. Phys. J. C **70** \(2010\) 823](#), arXiv: [1005.4568 \[physics.ins-det\]](#).
- [106] T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, [Comput. Phys. Commun. **178** \(2008\) 852](#), arXiv: [0710.3820 \[hep-ph\]](#).
- [107] ATLAS Collaboration, *The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie–Landshoff diffractive model*, ATL-PHYS-PUB-2016-017, 2016, URL: <https://cds.cern.ch/record/2206965>.
- [108] ATLAS Collaboration, *Vertex Reconstruction Performance of the ATLAS Detector at $\sqrt{s} = 13$ TeV*, ATL-PHYS-PUB-2015-026, 2015, URL: <https://cds.cern.ch/record/2037717>.
- [109] ATLAS Collaboration, *Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton–proton collision data*, [JINST **14** \(2019\) P12006](#), arXiv: [1908.00005 \[hep-ex\]](#).
- [110] ATLAS Collaboration, *Muon reconstruction and identification efficiency in ATLAS using the full Run 2 pp collision data set at $\sqrt{s} = 13$ TeV*, [Eur. Phys. J. C **81** \(2021\) 578](#), arXiv: [2012.00578 \[hep-ex\]](#).
- [111] M. Cacciari, G. P. Salam, and G. Soyez, *The anti- k_t jet clustering algorithm*, [JHEP **04** \(2008\) 063](#), arXiv: [0802.1189 \[hep-ph\]](#).

- [112] M. Cacciari, G. P. Salam, and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896, arXiv: [1111.6097 \[hep-ph\]](https://arxiv.org/abs/1111.6097).
- [113] M. Cacciari and G. P. Salam, *Dispelling the N^3 myth for the k_t jet-finder*, *Phys. Lett. B* **641** (2006) 57, arXiv: [hep-ph/0512210](https://arxiv.org/abs/hep-ph/0512210).
- [114] ATLAS Collaboration, *Jet reconstruction and performance using particle flow with the ATLAS Detector*, *Eur. Phys. J. C* **77** (2017) 466, arXiv: [1703.10485 \[hep-ex\]](https://arxiv.org/abs/1703.10485).
- [115] ATLAS Collaboration, *Jet energy scale and resolution measured in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2021) 689, arXiv: [2007.02645 \[hep-ex\]](https://arxiv.org/abs/2007.02645).
- [116] ATLAS Collaboration, *Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector*, *Eur. Phys. J. C* **76** (2016) 581, arXiv: [1510.03823 \[hep-ex\]](https://arxiv.org/abs/1510.03823).
- [117] S. D. Ellis and D. E. Soper, *Successive combination jet algorithm for hadron collisions*, *Phys. Rev. D* **48** (1993) 3160, arXiv: [hep-ph/9305266](https://arxiv.org/abs/hep-ph/9305266).
- [118] ATLAS Collaboration, *ATLAS flavour-tagging algorithms for the LHC Run 2 pp collision dataset*, *Eur. Phys. J. C* **83** (2023) 681, arXiv: [2211.16345 \[physics.data-an\]](https://arxiv.org/abs/2211.16345).
- [119] ATLAS Collaboration, *ATLAS b-jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 970, arXiv: [1907.05120 \[hep-ex\]](https://arxiv.org/abs/1907.05120).
- [120] ATLAS Collaboration, *Measurement of the c-jet mistagging efficiency in $t\bar{t}$ events using pp collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS detector*, *Eur. Phys. J. C* **82** (2022) 95, arXiv: [2109.10627 \[hep-ex\]](https://arxiv.org/abs/2109.10627).
- [121] ATLAS Collaboration, *Calibration of the light-flavour jet mistagging efficiency of the b-tagging algorithms with Z+jets events using 139 fb^{-1} of ATLAS proton–proton collision data at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **83** (2023) 728, arXiv: [2301.06319 \[hep-ex\]](https://arxiv.org/abs/2301.06319).
- [122] ATLAS Collaboration, *Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1*, *Eur. Phys. J. C* **77** (2017) 490, arXiv: [1603.02934 \[hep-ex\]](https://arxiv.org/abs/1603.02934).
- [123] ATLAS Collaboration, *Identification of hadronic tau lepton decays using neural networks in the ATLAS experiment*, ATL-PHYS-PUB-2019-033, 2019, URL: <https://cds.cern.ch/record/2688062>.
- [124] ATLAS Collaboration, *Reconstruction, Energy Calibration, and Identification of Hadronically Decaying Tau Leptons in the ATLAS Experiment for Run-2 of the LHC*, ATL-PHYS-PUB-2015-045, 2015, URL: <https://cds.cern.ch/record/2064383>.
- [125] ATLAS Collaboration, *Identification and energy calibration of hadronically decaying tau leptons with the ATLAS experiment in pp collisions at $\sqrt{s} = 8$ TeV*, *Eur. Phys. J. C* **75** (2015) 303, arXiv: [1412.7086 \[hep-ex\]](https://arxiv.org/abs/1412.7086).
- [126] ATLAS Collaboration, *Measurement of the tau lepton reconstruction and identification performance in the ATLAS experiment using pp collisions at $\sqrt{s} = 13$ TeV*, ATLAS-CONF-2017-029, 2017, URL: <https://cds.cern.ch/record/2261772>.

- [127] ATLAS Collaboration, *The performance of missing transverse momentum reconstruction and its significance with the ATLAS detector using 140 fb^{-1} of $\sqrt{s} = 13 \text{ TeV}$ pp collisions*, (2024), arXiv: [2402.05858 \[hep-ex\]](#).
- [128] ATLAS Collaboration, *Search for charged Higgs bosons decaying into a top quark and a bottom quark at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector*, *JHEP* **06** (2021) 145, arXiv: [2102.10076 \[hep-ex\]](#).
- [129] ATLAS Collaboration, *Search for a new scalar resonance in flavour-changing neutral-current top-quark decays $t \rightarrow qX$ ($q = u, c$), with $X \rightarrow b\bar{b}$, in proton–proton collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector*, *JHEP* **07** (2023) 199, arXiv: [2301.03902 \[hep-ex\]](#).
- [130] ATLAS Collaboration, *Tools for estimating fake/non-prompt lepton backgrounds with the ATLAS detector at the LHC*, *JINST* **18** (2023) T11004, arXiv: [2211.16178 \[hep-ex\]](#).
- [131] P. Baldi, K. Cranmer, T. Faucett, P. Sadowski, and D. Whiteson, *Parameterized neural networks for high-energy physics*, *Eur. Phys. J. C* **76** (2016) 235, arXiv: [1601.07913 \[hep-ex\]](#).
- [132] A. Elagin, P. Murat, A. Pranko, and A. Safonov, *A New Mass Reconstruction Technique for Resonances Decaying to di-tau*, *Nucl. Instrum. Meth. A* **654** (2011) 481, arXiv: [1012.4686 \[hep-ex\]](#).
- [133] CMS Collaboration, *Search for neutral MSSM Higgs bosons decaying to a pair of tau leptons in pp collisions*, *JHEP* **10** (2014) 160, arXiv: [1408.3316 \[hep-ex\]](#).
- [134] Objax Developers, *Objax*, version 1.2.0, 2020, URL: <https://github.com/google/objax>.
- [135] D. P. Kingma and J. Ba, *Adam: A Method for Stochastic Optimization*, (2014), arXiv: [1412.6980 \[cs.LG\]](#).
- [136] L. Heinrich, M. Feickert, G. Stark, and K. Cranmer, *pyhf: pure-Python implementation of HistFactory statistical models*, *J. Open Source Softw.* **6** (2021) 2823.
- [137] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554, arXiv: [1007.1727 \[physics.data-an\]](#), Erratum: *Eur. Phys. J. C* **73** (2013) 2501.
- [138] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079, arXiv: [1405.0301 \[hep-ph\]](#).
- [139] J. Bellm et al., *Herwig 7.0/Herwig++ 3.0 release note*, *Eur. Phys. J. C* **76** (2016) 196, arXiv: [1512.01178 \[hep-ph\]](#).
- [140] J. R. Andersen et al., *Handbook of LHC Higgs Cross Sections: 3. Higgs Properties*, (2013), arXiv: [1307.1347 \[hep-ph\]](#).
- [141] ATLAS Collaboration, *ATLAS Computing Acknowledgements*, ATL-SOFT-PUB-2023-001, 2023, URL: <https://cds.cern.ch/record/2869272>.

The ATLAS Collaboration

G. Aad ¹⁰⁴, E. Aakvaag ¹⁷, B. Abbott ¹²³, S. Abdelhameed ^{119a}, K. Abeling ⁵⁶, N.J. Abicht ⁵⁰, S.H. Abidi ³⁰, M. Aboeela ⁴⁵, A. Aboulhorma ^{36e}, H. Abramowicz ¹⁵⁴, H. Abreu ¹⁵³, Y. Abulaiti ¹²⁰, B.S. Acharya ^{70a,70b,1}, A. Ackermann ^{64a}, C. Adam Bourdarios ⁴, L. Adamczyk ^{87a}, S.V. Addepalli ²⁷, M.J. Addison ¹⁰³, J. Adelman ¹¹⁸, A. Adiguzel ^{22c}, T. Adye ¹³⁷, A.A. Affolder ¹³⁹, Y. Afik ⁴⁰, M.N. Agaras ¹³, J. Agarwala ^{74a,74b}, A. Aggarwal ¹⁰², C. Agheorghiesei ^{28c}, F. Ahmadov ^{39,x}, W.S. Ahmed ¹⁰⁶, S. Ahuja ⁹⁷, X. Ai ^{63e}, G. Aielli ^{77a,77b}, A. Aikot ¹⁶⁶, M. Ait Tamliah ^{36e}, B. Aitbenkikh ^{36a}, M. Akbiyik ¹⁰², T.P.A. Åkesson ¹⁰⁰, A.V. Akimov ³⁸, D. Akiyama ¹⁷¹, N.N. Akolkar ²⁵, S. Aktas ^{22a}, K. Al Houry ⁴², G.L. Alberghi ^{24b}, J. Albert ¹⁶⁸, P. Albicocco ⁵⁴, G.L. Albouy ⁶¹, S. Alderweireldt ⁵³, Z.L. Alegria ¹²⁴, M. Aleksa ³⁷, I.N. Aleksandrov ³⁹, C. Alexa ^{28b}, T. Alexopoulos ¹⁰, F. Alfonsi ^{24b}, M. Algren ⁵⁷, M. Alhroob ¹⁷⁰, B. Ali ¹³⁵, H.M.J. Ali ⁹³, S. Ali ³², S.W. Alibocus ⁹⁴, M. Aliev ^{34c}, G. Alimonti ^{72a}, W. Alkahi ⁵⁶, C. Allaire ⁶⁷, B.M.M. Allbrooke ¹⁴⁹, J.F. Allen ⁵³, C.A. Allendes Flores ^{140f}, P.P. Allport ²¹, A. Aloisio ^{73a,73b}, F. Alonso ⁹², C. Alpigiani ¹⁴¹, Z.M.K. Alsolami ⁹³, M. Alvarez Estevez ¹⁰¹, A. Alvarez Fernandez ¹⁰², M. Alves Cardoso ⁵⁷, M.G. Alvigi ^{73a,73b}, M. Aly ¹⁰³, Y. Amaral Coutinho ^{84b}, A. Ambler ¹⁰⁶, C. Amelung ³⁷, M. Amerl ¹⁰³, C.G. Ames ¹¹¹, D. Amidei ¹⁰⁸, B. Amini ⁵⁵, K.J. Amirie ¹⁵⁸, S.P. Amor Dos Santos ^{133a}, K.R. Amos ¹⁶⁶, D. Amperiadou ¹⁵⁵, S. An ⁸⁵, V. Ananiev ¹²⁸, C. Anastopoulos ¹⁴², T. Andeen ¹¹, J.K. Anders ³⁷, A.C. Anderson ⁶⁰, S.Y. Andrean ^{48a,48b}, A. Andreazza ^{72a,72b}, S. Angelidakis ⁹, A. Angerami ⁴², A.V. Anisenkov ³⁸, A. Annovi ^{75a}, C. Antel ⁵⁷, E. Antipov ¹⁴⁸, M. Antonelli ⁵⁴, F. Anulli ^{76a}, M. Aoki ⁸⁵, T. Aoki ¹⁵⁶, M.A. Aparo ¹⁴⁹, L. Aperio Bella ⁴⁹, C. Appelt ¹⁹, A. Apyan ²⁷, S.J. Arbiol Val ⁸⁸, C. Arcangeletti ⁵⁴, A.T.H. Arce ⁵², J-F. Arguin ¹¹⁰, S. Argyropoulos ⁵⁵, J.-H. Arling ⁴⁹, O. Arnaez ⁴, H. Arnold ¹⁴⁸, G. Artoni ^{76a,76b}, H. Asada ¹¹³, K. Asai ¹²¹, S. Asai ¹⁵⁶, N.A. Asbah ³⁷, R.A. Ashby Pickering ¹⁷⁰, K. Assamagan ³⁰, R. Astalos ^{29a}, K.S.V. Astrand ¹⁰⁰, S. Atashi ¹⁶², R.J. Atkin ^{34a}, M. Atkinson ¹⁶⁵, H. Atmani ^{36f}, P.A. Atmasiddha ¹³¹, K. Augsten ¹³⁵, S. Auricchio ^{73a,73b}, A.D. Auriol ²¹, V.A. Austrup ¹⁰³, G. Avolio ³⁷, K. Axiotis ⁵⁷, G. Azuelos ^{110,ac}, D. Babal ^{29b}, H. Bachacou ¹³⁸, K. Bachas ^{155,p}, A. Bachi ³⁵, F. Backman ^{48a,48b}, A. Badea ⁴⁰, T.M. Baer ¹⁰⁸, P. Bagnaia ^{76a,76b}, M. Bahmani ¹⁹, D. Bahner ⁵⁵, K. Bai ¹²⁶, J.T. Baines ¹³⁷, L. Baines ⁹⁶, O.K. Baker ¹⁷⁵, E. Bakos ¹⁶, D. Bakshi Gupta ⁸, L.E. Balabram Filho ^{84b}, V. Balakrishnan ¹²³, R. Balasubramanian ¹¹⁷, E.M. Baldin ³⁸, P. Balek ^{87a}, E. Ballabene ^{24b,24a}, F. Balli ¹³⁸, L.M. Baltes ^{64a}, W.K. Balunas ³³, J. Balz ¹⁰², I. Bamwidhi ^{119b}, E. Banas ⁸⁸, M. Bandieramonte ¹³², A. Bandyopadhyay ²⁵, S. Bansal ²⁵, L. Barak ¹⁵⁴, M. Barakat ⁴⁹, E.L. Barberio ¹⁰⁷, D. Barberis ^{58b,58a}, M. Barbero ¹⁰⁴, M.Z. Barel ¹¹⁷, T. Barillari ¹¹², M-S. Barisits ³⁷, T. Barklow ¹⁴⁶, P. Baron ¹²⁵, D.A. Baron Moreno ¹⁰³, A. Baroncelli ^{63a}, A.J. Barr ¹²⁹, J.D. Barr ⁹⁸, F. Barreiro ¹⁰¹, J. Barreiro Guimarães da Costa ¹⁴, U. Barron ¹⁵⁴, M.G. Barros Teixeira ^{133a}, S. Barsov ³⁸, F. Bartels ^{64a}, R. Bartoldus ¹⁴⁶, A.E. Barton ⁹³, P. Bartos ^{29a}, A. Basan ¹⁰², M. Baselga ⁵⁰, A. Bassalat ^{67,b}, M.J. Basso ^{159a}, S. Bataju ⁴⁵, R. Bate ¹⁶⁷, R.L. Bates ⁶⁰, S. Batlamous ¹⁰¹, B. Batool ¹⁴⁴, M. Battaglia ¹³⁹, D. Battulga ¹⁹, M. Baucé ^{76a,76b}, M. Bauer ⁸⁰, P. Bauer ²⁵, L.T. Bazzano Hurrell ³¹, J.B. Beacham ⁵², T. Beau ¹³⁰, J.Y. Beaucamp ⁹², P.H. Beauchemin ¹⁶¹, P. Bechtel ²⁵, H.P. Beck ^{20,o}, K. Becker ¹⁷⁰, A.J. Beddall ⁸³, V.A. Bednyakov ³⁹, C.P. Bee ¹⁴⁸, L.J. Beemster ¹⁶, T.A. Beermann ³⁷, M. Begalli ^{84d}, M. Begel ³⁰, A. Behera ¹⁴⁸, J.K. Behr ⁴⁹, J.F. Beirer ³⁷, F. Beisiegel ²⁵, M. Belfkir ^{119b}, G. Bella ¹⁵⁴, L. Bellagamba ^{24b}, A. Bellerive ³⁵, P. Bellos ²¹, K. Beloborodov ³⁸, D. Benchechroun ^{36a}, F. Bendebba ^{36a}, Y. Benhammou ¹⁵⁴,

K.C. Benkendorfer ⁶², L. Beresford ⁴⁹, M. Beretta ⁵⁴, E. Bergeaas Kuutmann ¹⁶⁴, N. Berger ⁴,
 B. Bergmann ¹³⁵, J. Beringer ^{18a}, G. Bernardi ⁵, C. Bernius ¹⁴⁶, F.U. Bernlochner ²⁵,
 F. Bernon ^{37,104}, A. Berrocal Guardia ¹³, T. Berry ⁹⁷, P. Berta ¹³⁶, A. Berthold ⁵¹, S. Bethke ¹¹²,
 A. Betti ^{76a,76b}, A.J. Bevan ⁹⁶, N.K. Bhalla ⁵⁵, S. Bhatta ¹⁴⁸, D.S. Bhattacharya ¹⁶⁹,
 P. Bhattarai ¹⁴⁶, K.D. Bhide ⁵⁵, V.S. Bhopatkar ¹²⁴, R.M. Bianchi ¹³², G. Bianco ^{24b,24a},
 O. Biebel ¹¹¹, R. Bielski ¹²⁶, M. Biglietti ^{78a}, C.S. Billingsley ⁴⁵, Y. Bimgdi ^{36f}, M. Bindi ⁵⁶,
 A. Bingul ^{22b}, C. Bini ^{76a,76b}, G.A. Bird ³³, M. Birman ¹⁷², M. Biros ¹³⁶, S. Biryukov ¹⁴⁹,
 T. Bisanz ⁵⁰, E. Bisceglie ^{44b,44a}, J.P. Biswal ¹³⁷, D. Biswas ¹⁴⁴, I. Bloch ⁴⁹, A. Blue ⁶⁰,
 U. Blumenschein ⁹⁶, J. Blumenthal ¹⁰², V.S. Bobrovnikov ³⁸, M. Boehler ⁵⁵, B. Boehm ¹⁶⁹,
 D. Bogavac ³⁷, A.G. Bogdanchikov ³⁸, L.S. Boggia ¹³⁰, C. Bohm ^{48a}, V. Boisvert ⁹⁷,
 P. Bokan ³⁷, T. Bold ^{87a}, M. Bomben ⁵, M. Bona ⁹⁶, M. Boonekamp ¹³⁸, C.D. Booth ⁹⁷,
 A.G. Borbély ⁶⁰, I.S. Bordulev ³⁸, G. Borissov ⁹³, D. Bortoletto ¹²⁹, D. Boscherini ^{24b},
 M. Bosman ¹³, J.D. Bossio Sola ³⁷, K. Bouaouda ^{36a}, N. Bouchhar ¹⁶⁶, L. Boudet ⁴,
 J. Boudreau ¹³², E.V. Bouhova-Thacker ⁹³, D. Boumediene ⁴¹, R. Bouquet ^{58b,58a}, A. Boveia ¹²²,
 J. Boyd ³⁷, D. Boye ³⁰, I.R. Boyko ³⁹, L. Bozianu ⁵⁷, J. Bracinek ²¹, N. Brahimi ⁴,
 G. Brandt ¹⁷⁴, O. Brandt ³³, F. Braren ⁴⁹, B. Brau ¹⁰⁵, J.E. Brau ¹²⁶, R. Brenner ¹⁷²,
 L. Brenner ¹¹⁷, R. Brenner ¹⁶⁴, S. Bressler ¹⁷², G. Brianti ^{79a,79b}, D. Britton ⁶⁰, D. Britzger ¹¹²,
 I. Brock ²⁵, G. Brooijmans ⁴², E.M. Brooks ^{159b}, E. Brost ³⁰, L.M. Brown ¹⁶⁸, L.E. Bruce ⁶²,
 T.L. Bruckler ¹²⁹, P.A. Bruckman de Renstrom ⁸⁸, B. Brüers ⁴⁹, A. Bruni ^{24b}, G. Bruni ^{24b},
 M. Bruschi ^{24b}, N. Bruscinò ^{76a,76b}, T. Buanes ¹⁷, Q. Buat ¹⁴¹, D. Buchin ¹¹², A.G. Buckley ⁶⁰,
 O. Bulekov ³⁸, B.A. Bullard ¹⁴⁶, S. Burdin ⁹⁴, C.D. Burgard ⁵⁰, A.M. Burger ³⁷,
 B. Burghgrave ⁸, O. Burlayenko ⁵⁵, J. Burleson ¹⁶⁵, J.T.P. Burr ³³, J.C. Burzynski ¹⁴⁵,
 E.L. Busch ⁴², V. Büscher ¹⁰², P.J. Bussey ⁶⁰, J.M. Butler ²⁶, C.M. Buttar ⁶⁰,
 J.M. Butterworth ⁹⁸, W. Buttinger ¹³⁷, C.J. Buxo Vazquez ¹⁰⁹, A.R. Buzykaev ³⁸,
 S. Cabrera Urbán ¹⁶⁶, L. Cadamuro ⁶⁷, D. Caforio ⁵⁹, H. Cai ¹³², Y. Cai ^{14,114c}, Y. Cai ^{114a},
 V.M.M. Cairo ³⁷, O. Cakir ^{3a}, N. Calace ³⁷, P. Calafiura ^{18a}, G. Calderini ¹³⁰, P. Calfayan ⁶⁹,
 G. Callea ⁶⁰, L.P. Caloba ^{84b}, D. Calvet ⁴¹, S. Calvet ⁴¹, M. Calvetti ^{75a,75b}, R. Camacho Toro ¹³⁰,
 S. Camarda ³⁷, D. Camarero Munoz ²⁷, P. Camarri ^{77a,77b}, M.T. Camerlingo ^{73a,73b},
 D. Cameron ³⁷, C. Camincher ¹⁶⁸, M. Campanelli ⁹⁸, A. Camplani ⁴³, V. Canale ^{73a,73b},
 A.C. Canbay ^{3a}, E. Canonero ⁹⁷, J. Cantero ¹⁶⁶, Y. Cao ¹⁶⁵, F. Capocasa ²⁷, M. Capua ^{44b,44a},
 A. Carbone ^{72a,72b}, R. Cardarelli ^{77a}, J.C.J. Cardenas ⁸, G. Carducci ^{44b,44a}, T. Carli ³⁷,
 G. Carlino ^{73a}, J.I. Carlotto ¹³, B.T. Carlson ^{132,q}, E.M. Carlson ^{168,159a}, J. Carmignani ⁹⁴,
 L. Carminati ^{72a,72b}, A. Carnelli ¹³⁸, M. Carnesale ^{76a,76b}, S. Caron ¹¹⁶, E. Carquin ^{140f},
 S. Carrá ^{72a}, G. Carratta ^{24b,24a}, A.M. Carroll ¹²⁶, T.M. Carter ⁵³, M.P. Casado ^{13,i},
 M. Caspar ⁴⁹, F.L. Castillo ⁴, L. Castillo Garcia ¹³, V. Castillo Gimenez ¹⁶⁶, N.F. Castro ^{133a,133e},
 A. Catinaccio ³⁷, J.R. Catmore ¹²⁸, T. Cavaliere ⁴, V. Cavaliere ³⁰, N. Cavalli ^{24b,24a},
 L.J. Caviedes Betancourt ^{23b}, Y.C. Cekmecelioglu ⁴⁹, E. Celebi ⁸³, S. Cella ³⁷, F. Celli ¹²⁹,
 M.S. Centonze ^{71a,71b}, V. Cepaitis ⁵⁷, K. Cerny ¹²⁵, A.S. Cerqueira ^{84a}, A. Cerri ¹⁴⁹,
 L. Cerrito ^{77a,77b}, F. Cerutti ^{18a}, B. Cervato ¹⁴⁴, A. Cervelli ^{24b}, G. Cesarini ⁵⁴, S.A. Cetin ⁸³,
 D. Chakraborty ¹¹⁸, J. Chan ^{18a}, W.Y. Chan ¹⁵⁶, J.D. Chapman ³³, E. Chapon ¹³⁸,
 B. Chargeishvili ^{152b}, D.G. Charlton ²¹, M. Chatterjee ²⁰, C. Chauhan ¹³⁶, Y. Che ^{114a},
 S. Chekanov ⁶, S.V. Chekulaev ^{159a}, G.A. Chelkov ^{39,a}, A. Chen ¹⁰⁸, B. Chen ¹⁵⁴, B. Chen ¹⁶⁸,
 H. Chen ^{114a}, H. Chen ³⁰, J. Chen ^{63c}, J. Chen ¹⁴⁵, M. Chen ¹²⁹, S. Chen ¹⁵⁶, S.J. Chen ^{114a},
 X. Chen ^{63c}, X. Chen ^{15,ab}, Y. Chen ^{63a}, C.L. Cheng ¹⁷³, H.C. Cheng ^{65a}, S. Cheong ¹⁴⁶,
 A. Cheplakov ³⁹, E. Cheremushkina ⁴⁹, E. Cherepanova ¹¹⁷, R. Cherkaoui El Moursli ^{36e},
 E. Cheu ⁷, K. Cheung ⁶⁶, L. Chevalier ¹³⁸, V. Chiarella ⁵⁴, G. Chiarelli ^{75a}, N. Chiedde ¹⁰⁴,
 G. Chiodini ^{71a}, A.S. Chisholm ²¹, A. Chitan ^{28b}, M. Chitishvili ¹⁶⁶, M.V. Chizhov ³⁹,

K. Choi ¹¹, Y. Chou ¹⁴¹, E.Y.S. Chow ¹¹⁶, K.L. Chu ¹⁷², M.C. Chu ^{65a}, X. Chu ^{14,114c},
 Z. Chubinidze ⁵⁴, J. Chudoba ¹³⁴, J.J. Chwastowski ⁸⁸, D. Cieri ¹¹², K.M. Ciesla ^{87a},
 V. Cindro ⁹⁵, A. Ciocio ^{18a}, F. Cirotto ^{73a,73b}, Z.H. Citron ¹⁷², M. Citterio ^{72a}, D.A. Ciubotaru ^{28b},
 A. Clark ⁵⁷, P.J. Clark ⁵³, N. Clarke Hall ⁹⁸, C. Clarry ¹⁵⁸, J.M. Clavijo Columbie ⁴⁹,
 S.E. Clawson ⁴⁹, C. Clement ^{48a,48b}, Y. Coadou ¹⁰⁴, M. Cobal ^{70a,70c}, A. Coccaro ^{58b},
 R.F. Coelho Barrue ^{133a}, R. Coelho Lopes De Sa ¹⁰⁵, S. Coelli ^{72a}, B. Cole ⁴², J. Collot ⁶¹,
 P. Conde Muiño ^{133a,133g}, M.P. Connell ^{34c}, S.H. Connell ^{34c}, E.I. Conroy ¹²⁹, F. Conventi ^{73a,ad},
 H.G. Cooke ²¹, A.M. Cooper-Sarkar ¹²⁹, F.A. Corchia ^{24b,24a}, A. Cordeiro Oudot Choi ¹³⁰,
 L.D. Corpe ⁴¹, M. Corradi ^{76a,76b}, F. Corriveau ^{106,w}, A. Cortes-Gonzalez ¹⁹, M.J. Costa ¹⁶⁶,
 F. Costanza ⁴, D. Costanzo ¹⁴², B.M. Cote ¹²², J. Couthures ⁴, G. Cowan ⁹⁷, K. Cranmer ¹⁷³,
 D. Cremonini ^{24b,24a}, S. Crépe-Renaudin ⁶¹, F. Crescioli ¹³⁰, M. Cristinziani ¹⁴⁴,
 M. Cristoforetti ^{79a,79b}, V. Croft ¹¹⁷, J.E. Crosby ¹²⁴, G. Crosetti ^{44b,44a}, A. Cueto ¹⁰¹, H. Cui ⁹⁸,
 Z. Cui ⁷, W.R. Cunningham ⁶⁰, F. Curcio ¹⁶⁶, J.R. Curran ⁵³, P. Czodrowski ³⁷,
 M.J. Da Cunha Sargedas De Sousa ^{58b,58a}, J.V. Da Fonseca Pinto ^{84b}, C. Da Via ¹⁰³,
 W. Dabrowski ^{87a}, T. Dado ³⁷, S. Dahbi ¹⁵¹, T. Dai ¹⁰⁸, D. Dal Santo ²⁰, C. Dallapiccola ¹⁰⁵,
 M. Dam ⁴³, G. D'amen ³⁰, V. D'Amico ¹¹¹, J. Damp ¹⁰², J.R. Dandoy ³⁵, D. Dannheim ³⁷,
 M. Danninger ¹⁴⁵, V. Dao ¹⁴⁸, G. Darbo ^{58b}, S.J. Das ^{30,ae}, F. Dattola ⁴⁹, S. D'Auria ^{72a,72b},
 A. D'Avanzo ^{73a,73b}, C. David ^{34a}, T. Davidek ¹³⁶, I. Dawson ⁹⁶, H.A. Day-hall ¹³⁵, K. De ⁸,
 R. De Asmundis ^{73a}, N. De Biase ⁴⁹, S. De Castro ^{24b,24a}, N. De Groot ¹¹⁶, P. de Jong ¹¹⁷,
 H. De la Torre ¹¹⁸, A. De Maria ^{114a}, A. De Salvo ^{76a}, U. De Sanctis ^{77a,77b}, F. De Santis ^{71a,71b},
 A. De Santo ¹⁴⁹, J.B. De Vivie De Regie ⁶¹, J. Debevc ⁹⁵, D.V. Dedovich ³⁹, J. Degens ⁹⁴,
 A.M. Deiana ⁴⁵, F. Del Corso ^{24b,24a}, J. Del Peso ¹⁰¹, L. Delagrangé ¹³⁰, F. Deliot ¹³⁸,
 C.M. Delitzsch ⁵⁰, M. Della Pietra ^{73a,73b}, D. Della Volpe ⁵⁷, A. Dell'Acqua ³⁷,
 L. Dell'Asta ^{72a,72b}, M. Delmastro ⁴, P.A. Delsart ⁶¹, S. Demers ¹⁷⁵, M. Demichev ³⁹,
 S.P. Denisov ³⁸, L. D'Eramo ⁴¹, D. Derendarz ⁸⁸, F. Derue ¹³⁰, P. Dervan ⁹⁴, K. Desch ²⁵,
 C. Deutsch ²⁵, F.A. Di Bello ^{58b,58a}, A. Di Ciaccio ^{77a,77b}, L. Di Ciaccio ⁴,
 A. Di Domenico ^{76a,76b}, C. Di Donato ^{73a,73b}, A. Di Girolamo ³⁷, G. Di Gregorio ³⁷,
 A. Di Luca ^{79a,79b}, B. Di Micco ^{78a,78b}, R. Di Nardo ^{78a,78b}, K.F. Di Petrillo ⁴⁰,
 M. Diamantopoulou ³⁵, F.A. Dias ¹¹⁷, T. Dias Do Vale ¹⁴⁵, M.A. Diaz ^{140a,140b},
 F.G. Diaz Capriles ²⁵, A.R. Didenko ³⁹, M. Didenko ¹⁶⁶, E.B. Diehl ¹⁰⁸, S. Díez Cornell ⁴⁹,
 C. Díez Pardos ¹⁴⁴, C. Dimitriadi ¹⁶⁴, A. Dimitrievska ²¹, J. Dingfelder ²⁵, T. Dingley ¹²⁹,
 I-M. Dinu ^{28b}, S.J. Dittmeier ^{64b}, F. Dittus ³⁷, M. Divisek ¹³⁶, B. Dixit ⁹⁴, F. Djama ¹⁰⁴,
 T. Djobava ^{152b}, C. Doglioni ^{103,100}, A. Dohalova ^{29a}, J. Dolejsi ¹³⁶, Z. Dolezal ¹³⁶,
 K. Domijan ^{87a}, K.M. Dona ⁴⁰, M. Donadelli ^{84d}, B. Dong ¹⁰⁹, J. Donini ⁴¹,
 A. D'Onofrio ^{73a,73b}, M. D'Onofrio ⁹⁴, J. Dopke ¹³⁷, A. Doria ^{73a}, N. Dos Santos Fernandes ^{133a},
 P. Dougan ¹⁰³, M.T. Dova ⁹², A.T. Doyle ⁶⁰, M.A. Draguet ¹²⁹, E. Dreyer ¹⁷²,
 I. Drivas-koulouris ¹⁰, M. Drnevich ¹²⁰, M. Drozdova ⁵⁷, D. Du ^{63a}, T.A. du Pree ¹¹⁷,
 F. Dubinin ³⁸, M. Dubovsky ^{29a}, E. Duchovni ¹⁷², G. Duckeck ¹¹¹, O.A. Ducu ^{28b}, D. Duda ⁵³,
 A. Dudarev ³⁷, E.R. Duden ²⁷, M. D'uffizi ¹⁰³, L. Duflot ⁶⁷, M. Dührssen ³⁷, I. Duminica ^{28g},
 A.E. Dumitriu ^{28b}, M. Dunford ^{64a}, S. Dungs ⁵⁰, K. Dunne ^{48a,48b}, A. Duperrin ¹⁰⁴,
 H. Duran Yildiz ^{3a}, M. Düren ⁵⁹, A. Durglishvili ^{152b}, B.L. Dwyer ¹¹⁸, G.I. Dyckes ^{18a},
 M. Dyndal ^{87a}, B.S. Dziedzic ³⁷, Z.O. Earnshaw ¹⁴⁹, G.H. Eberwein ¹²⁹, B. Eckerova ^{29a},
 S. Eggebrecht ⁵⁶, E. Egidio Purcino De Souza ^{84c}, L.F. Ehrke ⁵⁷, G. Eigen ¹⁷, K. Einsweiler ^{18a},
 T. Ekelof ¹⁶⁴, P.A. Ekman ¹⁰⁰, S. El Farkh ^{36b}, Y. El Ghazali ^{63a}, H. El Jarrari ³⁷,
 A. El Moussaouy ^{36a}, V. Ellajosyula ¹⁶⁴, M. Ellert ¹⁶⁴, F. Ellinghaus ¹⁷⁴, N. Ellis ³⁷,
 J. Elmsheuser ³⁰, M. Elsayy ^{119a}, M. Elsing ³⁷, D. Emelianov ¹³⁷, Y. Enari ⁸⁵, I. Ene ^{18a},
 S. Epari ¹³, P.A. Erland ⁸⁸, D. Ernani Martins Neto ⁸⁸, M. Errenst ¹⁷⁴, M. Escalier ⁶⁷,

C. Escobar ¹⁶⁶, E. Etzion ¹⁵⁴, G. Evans ^{133a}, H. Evans ⁶⁹, L.S. Evans ⁹⁷, A. Ezhilov ³⁸,
 S. Ezzarqtouni ^{36a}, F. Fabbri ^{24b,24a}, L. Fabbri ^{24b,24a}, G. Facini ⁹⁸, V. Fadeyev ¹³⁹,
 R.M. Fakhrutdinov ³⁸, D. Fakoudis ¹⁰², S. Falciano ^{76a}, L.F. Falda Ulhoa Coelho ³⁷,
 F. Fallavollita ¹¹², G. Falsetti ^{44b,44a}, J. Faltova ¹³⁶, C. Fan ¹⁶⁵, K.Y. Fan ^{65b}, Y. Fan ¹⁴,
 Y. Fang ^{14,114c}, M. Fanti ^{72a,72b}, M. Faraj ^{70a,70b}, Z. Farazpay ⁹⁹, A. Farbin ⁸, A. Farilla ^{78a},
 T. Farooque ¹⁰⁹, S.M. Farrington ⁵³, F. Fassi ^{36e}, D. Fassouliotis ⁹, M. Faucci Giannelli ^{77a,77b},
 W.J. Fawcett ³³, L. Fayard ⁶⁷, P. Federic ¹³⁶, P. Federicova ¹³⁴, O.L. Fedin ^{38,a}, M. Feickert ¹⁷³,
 L. Feligioni ¹⁰⁴, D.E. Fellers ¹²⁶, C. Feng ^{63b}, Z. Feng ¹¹⁷, M.J. Fenton ¹⁶², L. Ferencz ⁴⁹,
 R.A.M. Ferguson ⁹³, S.I. Fernandez Luengo ^{140f}, P. Fernandez Martinez ¹³, M.J.V. Fernoux ¹⁰⁴,
 J. Ferrando ⁹³, A. Ferrari ¹⁶⁴, P. Ferrari ^{117,116}, R. Ferrari ^{74a}, D. Ferrere ⁵⁷, C. Ferretti ¹⁰⁸,
 D. Fiacco ^{76a,76b}, F. Fiedler ¹⁰², P. Fiedler ¹³⁵, A. Filipčić ⁹⁵, E.K. Filmer ¹, F. Filthaut ¹¹⁶,
 M.C.N. Fiolhais ^{133a,133c,c}, L. Fiorini ¹⁶⁶, W.C. Fisher ¹⁰⁹, T. Fitschen ¹⁰³, P.M. Fitzhugh ¹³⁸,
 I. Fleck ¹⁴⁴, P. Fleischmann ¹⁰⁸, T. Flick ¹⁷⁴, M. Flores ^{34d,z}, L.R. Flores Castillo ^{65a},
 L. Flores Sanz De Acedo ³⁷, F.M. Follega ^{79a,79b}, N. Fomin ³³, J.H. Foo ¹⁵⁸, A. Formica ¹³⁸,
 A.C. Forti ¹⁰³, E. Fortin ³⁷, A.W. Fortman ^{18a}, M.G. Foti ^{18a}, L. Fountas ^{9j}, D. Fournier ⁶⁷,
 H. Fox ⁹³, P. Francavilla ^{75a,75b}, S. Francescato ⁶², S. Franchellucci ⁵⁷, M. Franchini ^{24b,24a},
 S. Franchino ^{64a}, D. Francis ³⁷, L. Franco ¹¹⁶, V. Franco Lima ³⁷, L. Franconi ⁴⁹, M. Franklin ⁶²,
 G. Frattari ²⁷, Y.Y. Frid ¹⁵⁴, J. Friend ⁶⁰, N. Fritzsche ³⁷, A. Froch ⁵⁵, D. Froidevaux ³⁷,
 J.A. Frost ¹²⁹, Y. Fu ^{63a}, S. Fuenzalida Garrido ^{140f}, M. Fujimoto ¹⁰⁴, K.Y. Fung ^{65a},
 E. Furtado De Simas Filho ^{84e}, M. Furukawa ¹⁵⁶, J. Fuster ¹⁶⁶, A. Gaa ⁵⁶, A. Gabrielli ^{24b,24a},
 A. Gabrielli ¹⁵⁸, P. Gadow ³⁷, G. Gagliardi ^{58b,58a}, L.G. Gagnon ^{18a}, S. Gaid ¹⁶³,
 S. Galantzan ¹⁵⁴, J. Gallagher ¹, E.J. Gallas ¹²⁹, B.J. Gallop ¹³⁷, K.K. Gan ¹²², S. Ganguly ¹⁵⁶,
 Y. Gao ⁵³, F.M. Garay Walls ^{140a,140b}, B. Garcia ³⁰, C. García ¹⁶⁶, A. Garcia Alonso ¹¹⁷,
 A.G. Garcia Caffaro ¹⁷⁵, J.E. García Navarro ¹⁶⁶, M. Garcia-Sciveres ^{18a}, G.L. Gardner ¹³¹,
 R.W. Gardner ⁴⁰, N. Garelli ¹⁶¹, D. Garg ⁸¹, R.B. Garg ¹⁴⁶, J.M. Gargan ⁵³, C.A. Garner ¹⁵⁸,
 C.M. Garvey ^{34a}, V.K. Gassmann ¹⁶¹, G. Gaudio ^{74a}, V. Gautam ¹³, P. Gauzzi ^{76a,76b},
 J. Gavranovic ⁹⁵, I.L. Gavrilenko ³⁸, A. Gavriluk ³⁸, C. Gay ¹⁶⁷, G. Gaycken ¹²⁶,
 E.N. Gazis ¹⁰, A.A. Geanta ^{28b}, C.M. Gee ¹³⁹, A. Gekow ¹²², C. Gemme ^{58b}, M.H. Genest ⁶¹,
 A.D. Gentry ¹¹⁵, S. George ⁹⁷, W.F. George ²¹, T. Geralis ⁴⁷, P. Gessinger-Befurt ³⁷,
 M.E. Geyik ¹⁷⁴, M. Ghani ¹⁷⁰, K. Ghorbanian ⁹⁶, A. Ghosal ¹⁴⁴, A. Ghosh ¹⁶², A. Ghosh ⁷,
 B. Giacobbe ^{24b}, S. Giagu ^{76a,76b}, T. Giani ¹¹⁷, A. Giannini ^{63a}, S.M. Gibson ⁹⁷, M. Gignac ¹³⁹,
 D.T. Gil ^{87b}, A.K. Gilbert ^{87a}, B.J. Gilbert ⁴², D. Gillberg ³⁵, G. Gilles ¹¹⁷, L. Ginabat ¹³⁰,
 D.M. Gingrich ^{2,ac}, M.P. Giordani ^{70a,70c}, P.F. Giraud ¹³⁸, G. Giugliarelli ^{70a,70c}, D. Giugni ^{72a},
 F. Giuli ³⁷, I. Gkialas ^{9j}, L.K. Gladilin ³⁸, C. Glasman ¹⁰¹, G.R. Gledhill ¹²⁶, G. Glemža ⁴⁹,
 M. Glisic ¹²⁶, I. Gnesi ^{44b,e}, Y. Go ³⁰, M. Goblirsch-Kolb ³⁷, B. Gocke ⁵⁰, D. Godin ¹¹⁰,
 B. Gokturk ^{22a}, S. Goldfarb ¹⁰⁷, T. Golling ⁵⁷, M.G.D. Gololo ^{34g}, D. Golubkov ³⁸,
 J.P. Gombas ¹⁰⁹, A. Gomes ^{133a,133b}, G. Gomes Da Silva ¹⁴⁴, A.J. Gomez Delegido ¹⁶⁶,
 R. Gonçalves ^{133a}, L. Gonella ²¹, A. Gongadze ^{152c}, F. Gonnella ²¹, J.L. Gonski ¹⁴⁶,
 R.Y. González Andana ⁵³, S. González de la Hoz ¹⁶⁶, R. Gonzalez Lopez ⁹⁴,
 C. Gonzalez Renteria ^{18a}, M.V. Gonzalez Rodrigues ⁴⁹, R. Gonzalez Suarez ¹⁶⁴,
 S. Gonzalez-Sevilla ⁵⁷, L. Goossens ³⁷, B. Gorini ³⁷, E. Gorini ^{71a,71b}, A. Gorišek ⁹⁵,
 T.C. Gosart ¹³¹, A.T. Goshaw ⁵², M.I. Gostkin ³⁹, S. Goswami ¹²⁴, C.A. Gottardo ³⁷,
 S.A. Gotz ¹¹¹, M. Goughri ^{36b}, V. Goumarre ⁴⁹, A.G. Goussiou ¹⁴¹, N. Govender ^{34c},
 R.P. Grabarczyk ¹²⁹, I. Grabowska-Bold ^{87a}, K. Graham ³⁵, E. Gramstad ¹²⁸,
 S. Grancagnolo ^{71a,71b}, C.M. Grant ^{1,138}, P.M. Gravila ^{28f}, F.G. Gravili ^{71a,71b}, H.M. Gray ^{18a},
 M. Greco ^{71a,71b}, M.J. Green ¹, C. Grefe ²⁵, A.S. Grefsrud ¹⁷, I.M. Gregor ⁴⁹, K.T. Greif ¹⁶²,
 P. Grenier ¹⁴⁶, S.G. Grewe ¹¹², A.A. Grillo ¹³⁹, K. Grimm ³², S. Grinstein ^{13,s}, J.-F. Grivaz ⁶⁷,

E. Gross ¹⁷², J. Grosse-Knetter ⁵⁶, L. Guan ¹⁰⁸, J.G.R. Guerrero Rojas ¹⁶⁶, G. Guerrieri ³⁷,
 R. Gugel ¹⁰², J.A.M. Guhit ¹⁰⁸, A. Guida ¹⁹, E. Guilloton ¹⁷⁰, S. Guindon ³⁷, F. Guo ^{14,114c},
 J. Guo ^{63c}, L. Guo ⁴⁹, Y. Guo ¹⁰⁸, R. Gupta ¹³², S. Gurbuz ²⁵, S.S. Gurdasani ⁵⁵,
 G. Gustavino ^{76a,76b}, P. Gutierrez ¹²³, L.F. Gutierrez Zagazeta ¹³¹, M. Gutsche ⁵¹,
 C. Gutschow ⁹⁸, C. Gwenlan ¹²⁹, C.B. Gwilliam ⁹⁴, E.S. Haaland ¹²⁸, A. Haas ¹²⁰,
 M. Habedank ⁴⁹, C. Haber ^{18a}, H.K. Hadavand ⁸, A. Hadeif ⁵¹, S. Hadzic ¹¹², A.I. Hagan ⁹³,
 J.J. Hahn ¹⁴⁴, E.H. Haines ⁹⁸, M. Haleem ¹⁶⁹, J. Haley ¹²⁴, J.J. Hall ¹⁴², G.D. Hallewell ¹⁰⁴,
 L. Halser ²⁰, K. Hamano ¹⁶⁸, M. Hamer ²⁵, G.N. Hamity ⁵³, E.J. Hampshire ⁹⁷, J. Han ^{63b},
 K. Han ^{63a}, L. Han ^{114a}, L. Han ^{63a}, S. Han ^{18a}, Y.F. Han ¹⁵⁸, K. Hanagaki ⁸⁵, M. Hance ¹³⁹,
 D.A. Hangal ⁴², H. Hanif ¹⁴⁵, M.D. Hank ¹³¹, J.B. Hansen ⁴³, P.H. Hansen ⁴³, D. Harada ⁵⁷,
 T. Harenberg ¹⁷⁴, S. Harkusha ³⁸, M.L. Harris ¹⁰⁵, Y.T. Harris ²⁵, J. Harrison ¹³,
 N.M. Harrison ¹²², P.F. Harrison ¹⁷⁰, N.M. Hartman ¹¹², N.M. Hartmann ¹¹¹, R.Z. Hasan ^{97,137},
 Y. Hasegawa ¹⁴³, F. Haslbeck ¹²⁹, S. Hassan ¹⁷, R. Hauser ¹⁰⁹, C.M. Hawkes ²¹,
 R.J. Hawkings ³⁷, Y. Hayashi ¹⁵⁶, D. Hayden ¹⁰⁹, C. Hayes ¹⁰⁸, R.L. Hayes ¹¹⁷, C.P. Hays ¹²⁹,
 J.M. Hays ⁹⁶, H.S. Hayward ⁹⁴, F. He ^{63a}, M. He ^{14,114c}, Y. He ⁴⁹, Y. He ⁹⁸, N.B. Heatley ⁹⁶,
 V. Hedberg ¹⁰⁰, A.L. Heggelund ¹²⁸, N.D. Hehir ^{96,*}, C. Heidegger ⁵⁵, K.K. Heidegger ⁵⁵,
 J. Heilman ³⁵, S. Heim ⁴⁹, T. Heim ^{18a}, J.G. Heinlein ¹³¹, J.J. Heinrich ¹²⁶, L. Heinrich ^{112,aa},
 J. Hejbal ¹³⁴, A. Held ¹⁷³, S. Hellesund ¹⁷, C.M. Helling ¹⁶⁷, S. Hellman ^{48a,48b},
 R.C.W. Henderson ⁹³, L. Henkelmann ³³, A.M. Henriques Correia ³⁷, H. Herde ¹⁰⁰,
 Y. Hernández Jiménez ¹⁴⁸, L.M. Herrmann ²⁵, T. Herrmann ⁵¹, G. Herten ⁵⁵, R. Hertenberger ¹¹¹,
 L. Hervas ³⁷, M.E. Hesping ¹⁰², N.P. Hessey ^{159a}, M. Hidaoui ^{36b}, N. Hidic ¹³⁶, E. Hill ¹⁵⁸,
 S.J. Hillier ²¹, J.R. Hinds ¹⁰⁹, F. Hinterkeuser ²⁵, M. Hirose ¹²⁷, S. Hirose ¹⁶⁰,
 D. Hirschbuehl ¹⁷⁴, T.G. Hitchings ¹⁰³, B. Hiti ⁹⁵, J. Hobbs ¹⁴⁸, R. Hobincu ^{28c}, N. Hod ¹⁷²,
 M.C. Hodgkinson ¹⁴², B.H. Hodgkinson ¹²⁹, A. Hoecker ³⁷, D.D. Hofer ¹⁰⁸, J. Hofer ⁴⁹,
 T. Holm ²⁵, M. Holzbock ³⁷, L.B.A.H. Hommels ³³, B.P. Honan ¹⁰³, J.J. Hong ⁶⁹, J. Hong ^{63c},
 T.M. Hong ¹³², B.H. Hooberman ¹⁶⁵, W.H. Hopkins ⁶, M.C. Hoppesch ¹⁶⁵, Y. Horii ¹¹³,
 S. Hou ¹⁵¹, A.S. Howard ⁹⁵, J. Howarth ⁶⁰, J. Hoya ⁶, M. Hrabovsky ¹²⁵, A. Hrynevich ⁴⁹,
 T. Hryn'ova ⁴, P.J. Hsu ⁶⁶, S.-C. Hsu ¹⁴¹, T. Hsu ⁶⁷, M. Hu ^{18a}, Q. Hu ^{63a}, S. Huang ^{65b},
 X. Huang ^{14,114c}, Y. Huang ¹⁴², Y. Huang ¹⁰², Y. Huang ¹⁴, Z. Huang ¹⁰³, Z. Hubacek ¹³⁵,
 M. Huebner ²⁵, F. Huegging ²⁵, T.B. Huffman ¹²⁹, C.A. Hugli ⁴⁹, M. Huhtinen ³⁷,
 S.K. Huiberts ¹⁷, R. Hulsken ¹⁰⁶, N. Huseynov ^{12,g}, J. Huston ¹⁰⁹, J. Huth ⁶², R. Hyneman ¹⁴⁶,
 G. Iacobucci ⁵⁷, G. Iakovidis ³⁰, L. Iconomidou-Fayard ⁶⁷, J.P. Iddon ³⁷, P. Iengo ^{73a,73b},
 R. Iguchi ¹⁵⁶, Y. Iiyama ¹⁵⁶, T. Iizawa ¹²⁹, Y. Ikegami ⁸⁵, N. Ilic ¹⁵⁸, H. Imam ^{84c},
 G. Inacio Goncalves ^{84d}, M. Ince Lezki ⁵⁷, T. Ingebretsen Carlson ^{48a,48b}, J.M. Inglis ⁹⁶,
 G. Introzzi ^{74a,74b}, M. Iodice ^{78a}, V. Ippolito ^{76a,76b}, R.K. Irwin ⁹⁴, M. Ishino ¹⁵⁶, W. Islam ¹⁷³,
 C. Issever ^{19,49}, S. Istin ^{22a,ag}, H. Ito ¹⁷¹, R. Iuppa ^{79a,79b}, A. Ivina ¹⁷², J.M. Izen ⁴⁶,
 V. Izzo ^{73a}, P. Jacka ¹³⁴, P. Jackson ¹, C.S. Jagfeld ¹¹¹, G. Jain ^{159a}, P. Jain ⁴⁹, K. Jakobs ⁵⁵,
 T. Jakoubek ¹⁷², J. Jamieson ⁶⁰, W. Jang ¹⁵⁶, M. Javurkova ¹⁰⁵, P. Jawahar ¹⁰³, L. Jeanty ¹²⁶,
 J. Jejelava ^{152a,y}, P. Jenni ^{55,f}, C.E. Jessiman ³⁵, C. Jia ^{63b}, J. Jia ¹⁴⁸, X. Jia ^{14,114c}, Z. Jia ^{114a},
 C. Jiang ⁵³, S. Jiggins ⁴⁹, J. Jimenez Pena ¹³, S. Jin ^{114a}, A. Jinaru ^{28b}, O. Jinnouchi ¹⁵⁷,
 P. Johansson ¹⁴², K.A. Johns ⁷, J.W. Johnson ¹³⁹, F.A. Jolly ⁴⁹, D.M. Jones ¹⁴⁹, E. Jones ⁴⁹,
 K.S. Jones ⁸, P. Jones ³³, R.W.L. Jones ⁹³, T.J. Jones ⁹⁴, H.L. Joos ^{56,37}, R. Joshi ¹²²,
 J. Jovicevic ¹⁶, X. Ju ^{18a}, J.J. Junggeburth ¹⁰⁵, T. Junkermann ^{64a}, A. Juste Rozas ^{13,s},
 M.K. Juzek ⁸⁸, S. Kabana ^{140e}, A. Kaczmarska ⁸⁸, M. Kado ¹¹², H. Kagan ¹²², M. Kagan ¹⁴⁶,
 A. Kahn ¹³¹, C. Kahra ¹⁰², T. Kaji ¹⁵⁶, E. Kajomovitz ¹⁵³, N. Kakati ¹⁷², I. Kalaitzidou ⁵⁵,
 C.W. Kalderon ³⁰, N.J. Kang ¹³⁹, D. Kar ^{34g}, K. Karava ¹²⁹, M.J. Kareem ^{159b}, E. Karentzos ⁵⁵,
 O. Karkout ¹¹⁷, S.N. Karpov ³⁹, Z.M. Karpova ³⁹, V. Kartvelishvili ⁹³, A.N. Karyukhin ³⁸,

E. Kasimi ¹⁵⁵, J. Katzy ⁴⁹, S. Kaur ³⁵, K. Kawade ¹⁴³, M.P. Kawale ¹²³, C. Kawamoto ⁸⁹,
 T. Kawamoto ^{63a}, E.F. Kay ³⁷, F.I. Kaya ¹⁶¹, S. Kazakos ¹⁰⁹, V.F. Kazanin ³⁸, Y. Ke ¹⁴⁸,
 J.M. Keaveney ^{34a}, R. Keeler ¹⁶⁸, G.V. Kehris ⁶², J.S. Keller ³⁵, A.S. Kelly ⁹⁸, J.J. Kempster ¹⁴⁹,
 P.D. Kennedy ¹⁰², O. Kepka ¹³⁴, B.P. Kerridge ¹³⁷, S. Kersten ¹⁷⁴, B.P. Kerševan ⁹⁵,
 L. Keszeghova ^{29a}, S. Ketabchi Haghghat ¹⁵⁸, R.A. Khan ¹³², A. Khanov ¹²⁴, A.G. Kharlamov ³⁸,
 T. Kharlamova ³⁸, E.E. Khoda ¹⁴¹, M. Kholodenko ^{133a}, T.J. Khoo ¹⁹, G. Khorauli ¹⁶⁹,
 J. Khubua ^{152b,*}, Y.A.R. Khwaira ¹³⁰, B. Kibirige ^{34g}, D. Kim ⁶, D.W. Kim ^{48a,48b}, Y.K. Kim ⁴⁰,
 N. Kimura ⁹⁸, M.K. Kingston ⁵⁶, A. Kirchhoff ⁵⁶, C. Kirfel ²⁵, F. Kirfel ²⁵, J. Kirk ¹³⁷,
 A.E. Kiryunin ¹¹², C. Kitsaki ¹⁰, O. Kivernyk ²⁵, M. Klassen ¹⁶¹, C. Klein ³⁵, L. Klein ¹⁶⁹,
 M.H. Klein ⁴⁵, S.B. Klein ⁵⁷, U. Klein ⁹⁴, P. Klimek ³⁷, A. Klimentov ³⁰, T. Klioutchnikova ³⁷,
 P. Kluit ¹¹⁷, S. Kluth ¹¹², E. Kneringer ⁸⁰, T.M. Knight ¹⁵⁸, A. Knue ⁵⁰, D. Kobylanskii ¹⁷²,
 S.F. Koch ¹²⁹, M. Kocian ¹⁴⁶, P. Kodyš ¹³⁶, D.M. Koeck ¹²⁶, P.T. Koenig ²⁵, T. Koffas ³⁵,
 O. Kolay ⁵¹, I. Koletsou ⁴, T. Komarek ⁸⁸, K. Köneke ⁵⁵, A.X.Y. Kong ¹, T. Kono ¹²¹,
 N. Konstantinidis ⁹⁸, P. Kontaxakis ⁵⁷, B. Konya ¹⁰⁰, R. Kopeliansky ⁴², S. Koperny ^{87a},
 K. Korcyl ⁸⁸, K. Kordas ^{155,d}, A. Korn ⁹⁸, S. Korn ⁵⁶, I. Korolkov ¹³, N. Korotkova ³⁸,
 B. Kortman ¹¹⁷, O. Kortner ¹¹², S. Kortner ¹¹², W.H. Kostecka ¹¹⁸, V.V. Kostyukhin ¹⁴⁴,
 A. Kotsokechagia ³⁷, A. Kotwal ⁵², A. Koulouris ³⁷, A. Kourkoumeli-Charalampidi ^{74a,74b},
 C. Kourkoumelis ⁹, E. Kourlitis ^{112,aa}, O. Kovanda ¹²⁶, R. Kowalewski ¹⁶⁸, W. Kozanecki ¹²⁶,
 A.S. Kozhin ³⁸, V.A. Kramarenko ³⁸, G. Kramberger ⁹⁵, P. Kramer ¹⁰², M.W. Krasny ¹³⁰,
 A. Krasznahorkay ³⁷, A.C. Kraus ¹¹⁸, J.W. Kraus ¹⁷⁴, J.A. Kremer ⁴⁹, T. Kresse ⁵¹,
 L. Kretschmann ¹⁷⁴, J. Kretschmar ⁹⁴, K. Kreul ¹⁹, P. Krieger ¹⁵⁸, M. Krivos ¹³⁶, K. Krizka ²¹,
 K. Kroeninger ⁵⁰, H. Kroha ¹¹², J. Kroll ¹³⁴, J. Kroll ¹³¹, K.S. Krowpman ¹⁰⁹, U. Kruchonak ³⁹,
 H. Krüger ²⁵, N. Krumnack ⁸², M.C. Kruse ⁵², O. Kuchinskaia ³⁸, S. Kuday ^{3a}, S. Kuehn ³⁷,
 R. Kuesters ⁵⁵, T. Kuhl ⁴⁹, V. Kukhtin ³⁹, Y. Kulchitsky ^{38,a}, S. Kuleshov ^{140d,140b},
 M. Kumar ^{34g}, N. Kumari ⁴⁹, P. Kumari ^{159b}, A. Kupco ¹³⁴, T. Kupfer ⁵⁰, A. Kupich ³⁸,
 O. Kuprash ⁵⁵, H. Kurashige ⁸⁶, L.L. Kurchaninov ^{159a}, O. Kurdysh ⁶⁷, Y.A. Kurochkin ³⁸,
 A. Kurova ³⁸, M. Kuze ¹⁵⁷, A.K. Kvam ¹⁰⁵, J. Kvitá ¹²⁵, T. Kwan ¹⁰⁶, N.G. Kyriacou ¹⁰⁸,
 L.A.O. Laatu ¹⁰⁴, C. Lacasta ¹⁶⁶, F. Lacava ^{76a,76b}, H. Lacker ¹⁹, D. Lacour ¹³⁰, N.N. Lad ⁹⁸,
 E. Ladygin ³⁹, A. Lafarge ⁴¹, B. Laforge ¹³⁰, T. Lagouri ¹⁷⁵, F.Z. Lahbabi ^{36a}, S. Lai ⁵⁶,
 J.E. Lambert ¹⁶⁸, S. Lammers ⁶⁹, W. Lampl ⁷, C. Lampoudis ^{155,d}, G. Lamprinoudis ¹⁰²,
 A.N. Lancaster ¹¹⁸, E. Lançon ³⁰, U. Landgraf ⁵⁵, M.P.J. Landon ⁹⁶, V.S. Lang ⁵⁵,
 O.K.B. Langrekken ¹²⁸, A.J. Lankford ¹⁶², F. Lanni ³⁷, K. Lantzsch ²⁵, A. Lanza ^{74a},
 M. Lanzac Berrocal ¹⁶⁶, J.F. Laporte ¹³⁸, T. Lari ^{72a}, F. Lasagni Manghi ^{24b}, M. Lassnig ³⁷,
 V. Latonova ¹³⁴, A. Laurier ¹⁵³, S.D. Lawlor ¹⁴², Z. Lawrence ¹⁰³, R. Lazaridou ¹⁷⁰,
 M. Lazzaroni ^{72a,72b}, B. Le ¹⁰³, H.D.M. Le ¹⁰⁹, E.M. Le Boulicaut ⁵², L.T. Le Pottier ^{18a},
 B. Leban ^{24b,24a}, A. Lebedev ⁸², M. LeBlanc ¹⁰³, F. Ledroit-Guillon ⁶¹, S.C. Lee ¹⁵¹,
 S. Lee ^{48a,48b}, T.F. Lee ⁹⁴, L.L. Leeuw ^{34c}, H.P. Lefebvre ⁹⁷, M. Lefebvre ¹⁶⁸, C. Leggett ^{18a},
 G. Lehmann Miotto ³⁷, M. Leigh ⁵⁷, W.A. Leight ¹⁰⁵, W. Leinonen ¹¹⁶, A. Leisos ^{155,r},
 M.A.L. Leite ^{84c}, C.E. Leitgeb ¹⁹, R. Leitner ¹³⁶, K.J.C. Leney ⁴⁵, T. Lenz ²⁵, S. Leone ^{75a},
 C. Leonidopoulos ⁵³, A. Leopold ¹⁴⁷, R. Les ¹⁰⁹, C.G. Lester ³³, M. Levchenko ³⁸,
 J. Levêque ⁴, L.J. Levinson ¹⁷², G. Levrini ^{24b,24a}, M.P. Lewicki ⁸⁸, C. Lewis ¹⁴¹, D.J. Lewis ⁴,
 L. Lewitt ¹⁴², A. Li ⁵, B. Li ^{63b}, C. Li ^{63a}, C-Q. Li ¹¹², H. Li ^{63a}, H. Li ^{63b}, H. Li ^{114a}, H. Li ¹⁵,
 H. Li ^{63b}, J. Li ^{63c}, K. Li ¹⁴¹, L. Li ^{63c}, M. Li ^{14,114c}, S. Li ^{14,114c}, S. Li ^{63d,63c}, T. Li ⁵,
 X. Li ¹⁰⁶, Z. Li ¹²⁹, Z. Li ¹⁵⁶, Z. Li ^{14,114c}, Z. Li ^{63a}, S. Liang ^{14,114c}, Z. Liang ¹⁴,
 M. Liberatore ¹³⁸, B. Liberti ^{77a}, K. Lie ^{65c}, J. Lieber Marin ^{84e}, H. Lien ⁶⁹, H. Lin ¹⁰⁸,
 K. Lin ¹⁰⁹, R.E. Lindley ⁷, J.H. Lindon ², J. Ling ⁶², E. Lipeles ¹³¹, A. Lipniacka ¹⁷,
 A. Lister ¹⁶⁷, J.D. Little ⁶⁹, B. Liu ¹⁴, B.X. Liu ^{114b}, D. Liu ^{63d,63c}, E.H.L. Liu ²¹, J.B. Liu ^{63a},

J.K.K. Liu ³³, K. Liu ^{63d}, K. Liu ^{63d,63c}, M. Liu ^{63a}, M.Y. Liu ^{63a}, P. Liu ¹⁴, Q. Liu ^{63d,141,63c}, X. Liu ^{63a}, X. Liu ^{63b}, Y. Liu ^{114b,114c}, Y.L. Liu ^{63b}, Y.W. Liu ^{63a}, S.L. Lloyd ⁹⁶, E.M. Lobodzinska ⁴⁹, P. Loch ⁷, T. Lohse ¹⁹, K. Lohwasser ¹⁴², E. Loiacono ⁴⁹, M. Lokajicek ^{134,*}, J.D. Lomas ²¹, J.D. Long ¹⁶⁵, I. Longarini ¹⁶², R. Longo ¹⁶⁵, I. Lopez Paz ⁶⁸, A. Lopez Solis ⁴⁹, N.A. Lopez-canelas ⁷, N. Lorenzo Martinez ⁴, A.M. Lory ¹¹¹, M. Losada ^{119a}, G. Löschcke Centeno ¹⁴⁹, O. Loseva ³⁸, X. Lou ^{48a,48b}, X. Lou ^{14,114c}, A. Lounis ⁶⁷, P.A. Love ⁹³, G. Lu ^{14,114c}, M. Lu ⁶⁷, S. Lu ¹³¹, Y.J. Lu ⁶⁶, H.J. Lubatti ¹⁴¹, C. Luci ^{76a,76b}, F.L. Lucio Alves ^{114a}, F. Luehring ⁶⁹, I. Luise ¹⁴⁸, O. Lukianchuk ⁶⁷, O. Lundberg ¹⁴⁷, B. Lund-Jensen ^{147,*}, N.A. Luongo ⁶, M.S. Lutz ³⁷, A.B. Lux ²⁶, D. Lynn ³⁰, R. Lysak ¹³⁴, E. Lytken ¹⁰⁰, V. Lyubushkin ³⁹, T. Lyubushkina ³⁹, M.M. Lyukova ¹⁴⁸, M.Firdaus M. Soberi ⁵³, H. Ma ³⁰, K. Ma ^{63a}, L.L. Ma ^{63b}, W. Ma ^{63a}, Y. Ma ¹²⁴, J.C. MacDonald ¹⁰², P.C. Machado De Abreu Farias ^{84e}, R. Madar ⁴¹, T. Madula ⁹⁸, J. Maeda ⁸⁶, T. Maeno ³⁰, H. Maguire ¹⁴², V. Maiboroda ¹³⁸, A. Maio ^{133a,133b,133d}, K. Maj ^{87a}, O. Majersky ⁴⁹, S. Majewski ¹²⁶, N. Makovec ⁶⁷, V. Maksimovic ¹⁶, B. Malaescu ¹³⁰, Pa. Malecki ⁸⁸, V.P. Maleev ³⁸, F. Malek ^{61,n}, M. Mali ⁹⁵, D. Malito ⁹⁷, U. Mallik ⁸¹, S. Maltezos ¹⁰, S. Malyukov ³⁹, J. Mamuzic ¹³, G. Mancini ⁵⁴, M.N. Mancini ²⁷, G. Manco ^{74a,74b}, J.P. Mandalia ⁹⁶, S.S. Mandarray ¹⁴⁹, I. Mandić ⁹⁵, L. Manhaes de Andrade Filho ^{84a}, I.M. Maniatis ¹⁷², J. Manjarres Ramos ⁹¹, D.C. Mankad ¹⁷², A. Mann ¹¹¹, S. Manzoni ³⁷, L. Mao ^{63c}, X. Mapekula ^{34c}, A. Marantis ^{155,r}, G. Marchiori ⁵, M. Marcisovsky ¹³⁴, C. Marcon ^{72a}, M. Marinescu ²¹, S. Marium ⁴⁹, M. Marjanovic ¹²³, A. Markhoos ⁵⁵, M. Markovitch ⁶⁷, E.J. Marshall ⁹³, Z. Marshall ^{18a}, S. Marti-Garcia ¹⁶⁶, J. Martin ⁹⁸, T.A. Martin ¹³⁷, V.J. Martin ⁵³, B. Martin dit Latour ¹⁷, L. Martinelli ^{76a,76b}, M. Martinez ^{13,s}, P. Martinez Agullo ¹⁶⁶, V.I. Martinez Outschoorn ¹⁰⁵, P. Martinez Suarez ¹³, S. Martin-Haugh ¹³⁷, G. Martinovicova ¹³⁶, V.S. Martoiu ^{28b}, A.C. Martyniuk ⁹⁸, A. Marzin ³⁷, D. Mascione ^{79a,79b}, L. Masetti ¹⁰², J. Masik ¹⁰³, A.L. Maslennikov ³⁸, P. Massarotti ^{73a,73b}, P. Mastrandrea ^{75a,75b}, A. Mastroberardino ^{44b,44a}, T. Masubuchi ¹²⁷, T. Mathisen ¹⁶⁴, J. Matousek ¹³⁶, J. Maurer ^{28b}, T. Maurin ⁶⁰, A.J. Maury ⁶⁷, B. Maček ⁹⁵, D.A. Maximov ³⁸, A.E. May ¹⁰³, R. Mazini ¹⁵¹, I. Maznas ¹¹⁸, M. Mazza ¹⁰⁹, S.M. Mazza ¹³⁹, E. Mazzeo ^{72a,72b}, C. Mc Ginn ³⁰, J.P. Mc Gowan ¹⁶⁸, S.P. Mc Kee ¹⁰⁸, C.C. McCracken ¹⁶⁷, E.F. McDonald ¹⁰⁷, A.E. McDougall ¹¹⁷, J.A. Mcfayden ¹⁴⁹, R.P. McGovern ¹³¹, R.P. Mckenzie ^{34g}, T.C. McLachlan ⁴⁹, D.J. McLaughlin ⁹⁸, S.J. McMahon ¹³⁷, C.M. Mcpartland ⁹⁴, R.A. McPherson ^{168,w}, S. Mehlhase ¹¹¹, A. Mehta ⁹⁴, D. Melini ¹⁶⁶, B.R. Mellado Garcia ^{34g}, A.H. Melo ⁵⁶, F. Meloni ⁴⁹, A.M. Mendes Jacques Da Costa ¹⁰³, H.Y. Meng ¹⁵⁸, L. Meng ⁹³, S. Menke ¹¹², M. Mentink ³⁷, E. Meoni ^{44b,44a}, G. Mercado ¹¹⁸, S. Merianos ¹⁵⁵, C. Merlassino ^{70a,70c}, L. Merola ^{73a,73b}, C. Meroni ^{72a,72b}, J. Metcalfe ⁶, A.S. Mete ⁶, E. Meuser ¹⁰², C. Meyer ⁶⁹, J-P. Meyer ¹³⁸, R.P. Middleton ¹³⁷, L. Mijović ⁵³, G. Mikenberg ¹⁷², M. Mikestikova ¹³⁴, M. Mikuž ⁹⁵, H. Mildner ¹⁰², A. Milic ³⁷, D.W. Miller ⁴⁰, E.H. Miller ¹⁴⁶, L.S. Miller ³⁵, A. Milov ¹⁷², D.A. Milstead ^{48a,48b}, T. Min ^{114a}, A.A. Minaenko ³⁸, I.A. Minashvili ^{152b}, L. Mince ⁶⁰, A.I. Mincer ¹²⁰, B. Mindur ^{87a}, M. Mineev ³⁹, Y. Mino ⁸⁹, L.M. Mir ¹³, M. Miralles Lopez ⁶⁰, M. Mironova ^{18a}, M.C. Missio ¹¹⁶, A. Mitra ¹⁷⁰, V.A. Mitsou ¹⁶⁶, Y. Mitsumori ¹¹³, O. Miu ¹⁵⁸, P.S. Miyagawa ⁹⁶, T. Mkrtchyan ^{64a}, M. Mlinarevic ⁹⁸, T. Mlinarevic ⁹⁸, M. Mlynarikova ³⁷, S. Mobius ²⁰, P. Mogg ¹¹¹, M.H. Mohamed Farook ¹¹⁵, A.F. Mohammed ^{14,114c}, S. Mohapatra ⁴², G. Mokgatitwane ^{34g}, L. Moleri ¹⁷², B. Mondal ¹⁴⁴, S. Mondal ¹³⁵, K. Mönig ⁴⁹, E. Monnier ¹⁰⁴, L. Monsonis Romero ¹⁶⁶, J. Montejo Berlingen ¹³, A. Montella ^{48a,48b}, M. Montella ¹²², F. Montekali ^{78a,78b}, F. Monticelli ⁹², S. Monzani ^{70a,70c}, A. Morancho Tarda ⁴³, N. Morange ⁶⁷, A.L. Moreira De Carvalho ⁴⁹, M. Moreno Llácer ¹⁶⁶, C. Moreno Martinez ⁵⁷,

J.M. Moreno Perez^{23b}, P. Morettini^{58b}, S. Morgenstern³⁷, M. Morii⁶², M. Morinaga¹⁵⁶,
 F. Morodei^{76a,76b}, L. Morvaj³⁷, P. Moschovakos³⁷, B. Moser¹²⁹, M. Mosidze^{152b},
 T. Moskalets⁴⁵, P. Moskvitina¹¹⁶, J. Moss^{32,k}, P. Moszkowicz^{87a}, A. Moussa^{36d},
 E.J.W. Moyse¹⁰⁵, O. Mtintsilana^{34g}, S. Muanza¹⁰⁴, J. Mueller¹³², D. Muenstermann⁹³,
 R. Müller³⁷, G.A. Mullier¹⁶⁴, A.J. Mullin³³, J.J. Mullin¹³¹, D.P. Mungo¹⁵⁸, D. Munoz Perez¹⁶⁶,
 F.J. Munoz Sanchez¹⁰³, M. Murin¹⁰³, W.J. Murray^{170,137}, M. Muškinja⁹⁵, C. Mwewa³⁰,
 A.G. Myagkov^{38,a}, A.J. Myers⁸, G. Myers¹⁰⁸, M. Myska¹³⁵, B.P. Nachman^{18a},
 O. Nackenhorst⁵⁰, K. Nagai¹²⁹, K. Nagano⁸⁵, J.L. Nagle^{30,ae}, E. Nagy¹⁰⁴, A.M. Nairz³⁷,
 Y. Nakahama⁸⁵, K. Nakamura⁸⁵, K. Nakkalil⁵, H. Nanjo¹²⁷, E.A. Narayanan¹¹⁵,
 I. Naryshkin³⁸, L. Nasella^{72a,72b}, M. Naseri³⁵, S. Nasri^{119b}, C. Nass²⁵, G. Navarro^{23a},
 J. Navarro-Gonzalez¹⁶⁶, R. Nayak¹⁵⁴, A. Nayaz¹⁹, P.Y. Nechaeva³⁸, S. Nechaeva^{24b,24a},
 F. Nechansky⁴⁹, L. Nedic¹²⁹, T.J. Neep²¹, A. Negri^{74a,74b}, M. Negrini^{24b}, C. Nellist¹¹⁷,
 C. Nelson¹⁰⁶, K. Nelson¹⁰⁸, S. Nemecek¹³⁴, M. Nessi^{37,h}, M.S. Neubauer¹⁶⁵, F. Neuhaus¹⁰²,
 J. Neundorf⁴⁹, P.R. Newman²¹, C.W. Ng¹³², Y.W.Y. Ng⁴⁹, B. Ngair^{119a}, H.D.N. Nguyen¹¹⁰,
 R.B. Nickerson¹²⁹, R. Nicolaidou¹³⁸, J. Nielsen¹³⁹, M. Niemeyer⁵⁶, J. Niermann⁵⁶,
 N. Nikiforou³⁷, V. Nikolaenko^{38,a}, I. Nikolic-Audit¹³⁰, K. Nikolopoulos²¹, P. Nilsson³⁰,
 I. Ninca⁴⁹, G. Ninio¹⁵⁴, A. Nisati^{76a}, N. Nishu², R. Nisius¹¹², J-E. Nitschke⁵¹,
 E.K. Nkadimeng^{34g}, T. Nobe¹⁵⁶, T. Nommensen¹⁵⁰, M.B. Norfolk¹⁴², B.J. Norman³⁵,
 M. Noury^{36a}, J. Novak⁹⁵, T. Novak⁹⁵, L. Novotny¹³⁵, R. Novotny¹¹⁵, L. Nozka¹²⁵,
 K. Ntekas¹⁶², N.M.J. Nunes De Moura Junior^{84b}, J. Ocariz¹³⁰, A. Ochi⁸⁶, I. Ochoa^{133a},
 S. Oerdek^{49,t}, J.T. Offermann⁴⁰, A. Ogrodnik¹³⁶, A. Oh¹⁰³, C.C. Ohm¹⁴⁷, H. Oide⁸⁵,
 R. Oishi¹⁵⁶, M.L. Ojeda⁴⁹, Y. Okumura¹⁵⁶, L.F. Oleiro Seabra^{133a}, I. Oleksiyuk⁵⁷,
 S.A. Olivares Pino^{140d}, G. Oliveira Correa¹³, D. Oliveira Damazio³⁰, J.L. Oliver¹⁶²,
 Ö.O. Öncel⁵⁵, A.P. O'Neill²⁰, A. Onofre^{133a,133e}, P.U.E. Onyisi¹¹, M.J. Oreglia⁴⁰,
 G.E. Orellana⁹², D. Orestano^{78a,78b}, N. Orlando¹³, R.S. Orr¹⁵⁸, L.M. Osojnak¹³¹,
 R. Ospanov^{63a}, G. Otero y Garzon³¹, H. Otono⁹⁰, P.S. Ott^{64a}, G.J. Ottino^{18a}, M. Ouchrif^{36d},
 F. Ould-Saada¹²⁸, T. Ovsianikova¹⁴¹, M. Owen⁶⁰, R.E. Owen¹³⁷, V.E. Ozcan^{22a},
 F. Ozturk⁸⁸, N. Ozturk⁸, S. Ozturk⁸³, H.A. Pacey¹²⁹, A. Pacheco Pages¹³,
 C. Padilla Aranda¹³, G. Padovano^{76a,76b}, S. Pagan Griso^{18a}, G. Palacino⁶⁹, A. Palazzo^{71a,71b},
 J. Pampel²⁵, J. Pan¹⁷⁵, T. Pan^{65a}, D.K. Panchal¹¹, C.E. Pandini¹¹⁷, J.G. Panduro Vazquez¹³⁷,
 H.D. Pandya¹, H. Pang¹⁵, P. Pani⁴⁹, G. Panizzo^{70a,70c}, L. Panwar¹³⁰, L. Paolozzi⁵⁷,
 S. Parajuli¹⁶⁵, A. Paramonov⁶, C. Paraskevopoulos⁵⁴, D. Paredes Hernandez^{65b},
 A. Pareti^{74a,74b}, K.R. Park⁴², T.H. Park¹⁵⁸, M.A. Parker³³, F. Parodi^{58b,58a}, E.W. Parrish¹¹⁸,
 V.A. Parrish⁵³, J.A. Parsons⁴², U. Parzefall⁵⁵, B. Pascual Dias¹¹⁰, L. Pascual Dominguez¹⁰¹,
 E. Pasqualucci^{76a}, S. Passaggio^{58b}, F. Pastore⁹⁷, P. Patel⁸⁸, U.M. Patel⁵², J.R. Pater¹⁰³,
 T. Pauly³⁷, C.I. Pazos¹⁶¹, J. Pearkes¹⁴⁶, M. Pedersen¹²⁸, R. Pedro^{133a}, S.V. Peleganchuk³⁸,
 O. Penc³⁷, E.A. Pender⁵³, G.D. Penn¹⁷⁵, K.E. Pensi¹¹¹, M. Penzin³⁸, B.S. Peralva^{84d},
 A.P. Pereira Peixoto¹⁴¹, L. Pereira Sanchez¹⁴⁶, D.V. Perepelitsa^{30,ae}, G. Perera¹⁰⁵,
 E. Perez Codina^{159a}, M. Perganti¹⁰, H. Pernegger³⁷, S. Perrella^{76a,76b}, O. Perrin⁴¹,
 K. Peters⁴⁹, R.F.Y. Peters¹⁰³, B.A. Petersen³⁷, T.C. Petersen⁴³, E. Petit¹⁰⁴, V. Petousis¹³⁵,
 C. Petridou^{155,d}, T. Petru¹³⁶, A. Petrukhin¹⁴⁴, M. Pettee^{18a}, A. Petukhov³⁸, K. Petukhova³⁷,
 R. Pezoa^{140f}, L. Pezzotti³⁷, G. Pezzullo¹⁷⁵, T.M. Pham¹⁷³, T. Pham¹⁰⁷, P.W. Phillips¹³⁷,
 G. Piacquadio¹⁴⁸, E. Pianori^{18a}, F. Piazza¹²⁶, R. Piegai³¹, D. Pietreanu^{28b},
 A.D. Pilkington¹⁰³, M. Pinamonti^{70a,70c}, J.L. Pinfold², B.C. Pinheiro Pereira^{133a},
 J. Pinol Bel¹³, A.E. Pinto Pinoargote^{138,138}, L. Pintucci^{70a,70c}, K.M. Piper¹⁴⁹, A. Pirttikoski⁵⁷,
 D.A. Pizzi³⁵, L. Pizzimento^{65b}, A. Pizzini¹¹⁷, M.-A. Pleier³⁰, V. Pleskot¹³⁶, E. Plotnikova³⁹,
 G. Poddar⁹⁶, R. Poettgen¹⁰⁰, L. Poggioli¹³⁰, I. Pokharel⁵⁶, S. Polacek¹³⁶, G. Polesello^{74a},

A. Poley ^{145,159a}, A. Polini ^{24b}, C.S. Pollard ¹⁷⁰, Z.B. Pollock ¹²², E. Pompa Pacchi ^{76a,76b},
 N.I. Pond ⁹⁸, D. Ponomarenko ⁶⁹, L. Pontecorvo ³⁷, S. Popa ^{28a}, G.A. Popeneciu ^{28d},
 A. Poreba ³⁷, D.M. Portillo Quintero ^{159a}, S. Pospisil ¹³⁵, M.A. Postill ¹⁴², P. Postolache ^{28c},
 K. Potamianos ¹⁷⁰, P.A. Potepa ^{87a}, I.N. Potrap ³⁹, C.J. Potter ³³, H. Potti ¹⁵⁰, J. Poveda ¹⁶⁶,
 M.E. Pozo Astigarraga ³⁷, A. Prades Ibanez ^{77a,77b}, J. Pretel ¹⁶⁸, D. Price ¹⁰³, M. Primavera ^{71a},
 L. Primomo ^{70a,70c}, M.A. Principe Martin ¹⁰¹, R. Privara ¹²⁵, T. Procter ⁶⁰, M.L. Proffitt ¹⁴¹,
 N. Proklova ¹³¹, K. Prokofiev ^{65c}, G. Proto ¹¹², J. Proudfoot ⁶, M. Przybycien ^{87a},
 W.W. Przygoda ^{87b}, A. Psallidas ⁴⁷, J.E. Puddefoot ¹⁴², D. Pudzha ⁵⁵, D. Pyatiizbyantseva ³⁸,
 J. Qian ¹⁰⁸, D. Qichen ¹⁰³, Y. Qin ¹³, T. Qiu ⁵³, A. Quadt ⁵⁶, M. Queitsch-Maitland ¹⁰³,
 G. Quetant ⁵⁷, R.P. Quinn ¹⁶⁷, G. Rabanal Bolanos ⁶², D. Rafanoharana ⁵⁵, F. Raffaelli ^{77a,77b},
 F. Ragusa ^{72a,72b}, J.L. Rainbolt ⁴⁰, J.A. Raine ⁵⁷, S. Rajagopalan ³⁰, E. Ramakoti ³⁸,
 L. Rambelli ^{58b,58a}, I.A. Ramirez-Berend ³⁵, K. Ran ^{49,114c}, D.S. Rankin ¹³¹, N.P. Rapheeha ^{34g},
 H. Rasheed ^{28b}, V. Raskina ¹³⁰, D.F. Rassloff ^{64a}, A. Rastogi ^{18a}, S. Rave ¹⁰², S. Ravera ^{58b,58a},
 B. Ravina ⁵⁶, I. Ravinovich ¹⁷², M. Raymond ³⁷, A.L. Read ¹²⁸, N.P. Readioff ¹⁴²,
 D.M. Rebuzzi ^{74a,74b}, G. Redlinger ³⁰, A.S. Reed ¹¹², K. Reeves ²⁷, J.A. Reidelsturz ¹⁷⁴,
 D. Reikher ¹²⁶, A. Rej ⁵⁰, C. Rembser ³⁷, M. Renda ^{28b}, F. Renner ⁴⁹, A.G. Rennie ¹⁶²,
 A.L. Rescia ⁴⁹, S. Resconi ^{72a}, M. Ressegotti ^{58b,58a}, S. Rettie ³⁷, J.G. Reyes Rivera ¹⁰⁹,
 E. Reynolds ^{18a}, O.L. Rezanova ³⁸, P. Reznicek ¹³⁶, H. Riani ^{36d}, N. Ribaric ⁹³, E. Ricci ^{79a,79b},
 R. Richter ¹¹², S. Richter ^{48a,48b}, E. Richter-Was ^{87b}, M. Ridel ¹³⁰, S. Ridouani ^{36d}, P. Rieck ¹²⁰,
 P. Riedler ³⁷, E.M. Riefel ^{48a,48b}, J.O. Rieger ¹¹⁷, M. Rijssenbeek ¹⁴⁸, M. Rimoldi ³⁷,
 L. Rinaldi ^{24b,24a}, P. Rincke ^{56,164}, T.T. Rinn ³⁰, M.P. Rinnagel ¹¹¹, G. Ripellino ¹⁶⁴, I. Riu ¹³,
 J.C. Rivera Vergara ¹⁶⁸, F. Rizatdinova ¹²⁴, E. Rizvi ⁹⁶, B.R. Roberts ^{18a}, S.S. Roberts ¹³⁹,
 S.H. Robertson ^{106,w}, D. Robinson ³³, M. Robles Manzano ¹⁰², A. Robson ⁶⁰, A. Rocchi ^{77a,77b},
 C. Roda ^{75a,75b}, S. Rodriguez Bosca ³⁷, Y. Rodriguez Garcia ^{23a}, A. Rodriguez Rodriguez ⁵⁵,
 A.M. Rodríguez Vera ¹¹⁸, S. Roe ³⁷, J.T. Roemer ³⁷, A.R. Roepe-Gier ¹³⁹, O. Røhne ¹²⁸,
 R.A. Rojas ¹⁰⁵, C.P.A. Roland ¹³⁰, J. Roloff ³⁰, A. Romaniouk ³⁸, E. Romano ^{74a,74b},
 M. Romano ^{24b}, A.C. Romero Hernandez ¹⁶⁵, N. Rompotis ⁹⁴, L. Roos ¹³⁰, S. Rosati ^{76a},
 B.J. Rosser ⁴⁰, E. Rossi ¹²⁹, E. Rossi ^{73a,73b}, L.P. Rossi ⁶², L. Rossini ⁵⁵, R. Rosten ¹²²,
 M. Rotaru ^{28b}, B. Rottler ⁵⁵, C. Rougier ⁹¹, D. Rousseau ⁶⁷, D. Rousso ⁴⁹, A. Roy ¹⁶⁵,
 S. Roy-Garand ¹⁵⁸, A. Rozanov ¹⁰⁴, Z.M.A. Rozario ⁶⁰, Y. Rozen ¹⁵³, A. Rubio Jimenez ¹⁶⁶,
 A.J. Ruby ⁹⁴, V.H. Ruelas Rivera ¹⁹, T.A. Ruggeri ¹, A. Ruggiero ¹²⁹, A. Ruiz-Martinez ¹⁶⁶,
 A. Rummler ³⁷, Z. Rurikova ⁵⁵, N.A. Rusakovich ³⁹, H.L. Russell ¹⁶⁸, G. Russo ^{76a,76b},
 J.P. Rutherford ⁷, S. Rutherford Colmenares ³³, M. Rybar ¹³⁶, E.B. Rye ¹²⁸, A. Ryzhov ⁴⁵,
 J.A. Sabater Iglesias ⁵⁷, H.F.W. Sadrozinski ¹³⁹, F. Safai Tehrani ^{76a}, B. Safarzadeh Samani ¹³⁷,
 S. Saha ¹, M. Sahinsoy ⁸³, A. Saibel ¹⁶⁶, M. Saimpert ¹³⁸, M. Saito ¹⁵⁶, T. Saito ¹⁵⁶,
 A. Sala ^{72a,72b}, D. Salamani ³⁷, A. Salnikov ¹⁴⁶, J. Salt ¹⁶⁶, A. Salvador Salas ¹⁵⁴,
 D. Salvatore ^{44b,44a}, F. Salvatore ¹⁴⁹, A. Salzburger ³⁷, D. Sammel ⁵⁵, E. Sampson ⁹³,
 D. Sampsonidis ^{155,d}, D. Sampsonidou ¹²⁶, J. Sánchez ¹⁶⁶, V. Sanchez Sebastian ¹⁶⁶,
 H. Sandaker ¹²⁸, C.O. Sander ⁴⁹, J.A. Sandesara ¹⁰⁵, M. Sandhoff ¹⁷⁴, C. Sandoval ^{23b},
 L. Sanfilippo ^{64a}, D.P.C. Sankey ¹³⁷, T. Sano ⁸⁹, A. Sansoni ⁵⁴, L. Santi ^{37,76b}, C. Santoni ⁴¹,
 H. Santos ^{133a,133b}, A. Santra ¹⁷², E. Sanzani ^{24b,24a}, K.A. Saoucha ¹⁶³, J.G. Saraiva ^{133a,133d},
 J. Sardain ⁷, O. Sasaki ⁸⁵, K. Sato ¹⁶⁰, C. Sauer ^{64b}, E. Sauvan ⁴, P. Savard ^{158,ac}, R. Sawada ¹⁵⁶,
 C. Sawyer ¹³⁷, L. Sawyer ⁹⁹, C. Sbarra ^{24b}, A. Sbrizzi ^{24b,24a}, T. Scanlon ⁹⁸,
 J. Schaarschmidt ¹⁴¹, U. Schäfer ¹⁰², A.C. Schaffer ^{67,45}, D. Schaile ¹¹¹, R.D. Schamberger ¹⁴⁸,
 C. Scharf ¹⁹, M.M. Schefer ²⁰, V.A. Schegelsky ³⁸, D. Scheirich ¹³⁶, M. Schernau ¹⁶²,
 C. Scheulen ⁵⁶, C. Schiavi ^{58b,58a}, M. Schioppa ^{44b,44a}, B. Schlag ^{146,m}, K.E. Schleicher ⁵⁵,
 S. Schlenker ³⁷, J. Schmeing ¹⁷⁴, M.A. Schmidt ¹⁷⁴, K. Schmieden ¹⁰², C. Schmitt ¹⁰²,

N. Schmitt ¹⁰², S. Schmitt ⁴⁹, L. Schoeffel ¹³⁸, A. Schoening ^{64b}, P.G. Scholer ³⁵, E. Schopf ¹²⁹,
 M. Schott ²⁵, J. Schovancova ³⁷, S. Schramm ⁵⁷, T. Schroer ⁵⁷, H-C. Schultz-Coulon ^{64a},
 M. Schumacher ⁵⁵, B.A. Schumm ¹³⁹, Ph. Schune ¹³⁸, A.J. Schuy ¹⁴¹, H.R. Schwartz ¹³⁹,
 A. Schwartzman ¹⁴⁶, T.A. Schwarz ¹⁰⁸, Ph. Schwemling ¹³⁸, R. Schwienhorst ¹⁰⁹,
 F.G. Sciacca ²⁰, A. Sciandra ³⁰, G. Sciolla ²⁷, F. Scuri ^{75a}, C.D. Sebastiani ⁹⁴, K. Sedlaczek ¹¹⁸,
 S.C. Seidel ¹¹⁵, A. Seiden ¹³⁹, B.D. Seidlitz ⁴², C. Seitz ⁴⁹, J.M. Seixas ^{84b}, G. Sekhniaidze ^{73a},
 L. Selem ⁶¹, N. Semprini-Cesari ^{24b,24a}, D. Sengupta ⁵⁷, V. Senthilkumar ¹⁶⁶, L. Serin ⁶⁷,
 M. Sessa ^{77a,77b}, H. Severini ¹²³, F. Sforza ^{58b,58a}, A. Sfyrla ⁵⁷, Q. Sha ¹⁴, E. Shabalina ⁵⁶,
 A.H. Shah ³³, R. Shaheen ¹⁴⁷, J.D. Shahinian ¹³¹, D. Shaked Renous ¹⁷², L.Y. Shan ¹⁴,
 M. Shapiro ^{18a}, A. Sharma ³⁷, A.S. Sharma ¹⁶⁷, P. Sharma ⁸¹, P.B. Shatalov ³⁸, K. Shaw ¹⁴⁹,
 S.M. Shaw ¹⁰³, Q. Shen ^{63c}, D.J. Sheppard ¹⁴⁵, P. Sherwood ⁹⁸, L. Shi ⁹⁸, X. Shi ¹⁴,
 S. Shimizu ⁸⁵, C.O. Shimmin ¹⁷⁵, J.D. Shinner ⁹⁷, I.P.J. Shipsey ¹²⁹, S. Shirabe ⁹⁰,
 M. Shiyakova ^{39,u}, M.J. Shochet ⁴⁰, D.R. Shope ¹²⁸, B. Shrestha ¹²³, S. Shrestha ^{122,af},
 M.J. Shroff ¹⁶⁸, P. Sicho ¹³⁴, A.M. Sickles ¹⁶⁵, E. Sideras Haddad ^{34g}, A.C. Sidley ¹¹⁷,
 A. Sidoti ^{24b}, F. Siegert ⁵¹, Dj. Sijacki ¹⁶, F. Sili ⁹², J.M. Silva ⁵³, I. Silva Ferreira ^{84b},
 M.V. Silva Oliveira ³⁰, S.B. Silverstein ^{48a}, S. Simion ⁶⁷, R. Simoniello ³⁷, E.L. Simpson ¹⁰³,
 H. Simpson ¹⁴⁹, L.R. Simpson ¹⁰⁸, N.D. Simpson ¹⁰⁰, S. Simsek ⁸³, S. Sindhu ⁵⁶, P. Sinervo ¹⁵⁸,
 S. Singh ¹⁵⁸, S. Sinha ⁴⁹, S. Sinha ¹⁰³, M. Sioli ^{24b,24a}, I. Siral ³⁷, E. Sitnikova ⁴⁹,
 J. Sjölin ^{48a,48b}, A. Skaf ⁵⁶, E. Skorda ²¹, P. Skubic ¹²³, M. Slawinska ⁸⁸, V. Smakhtin ¹⁷²,
 B.H. Smart ¹³⁷, S. Yu. Smirnov ³⁸, Y. Smirnov ³⁸, L.N. Smirnova ^{38,a}, O. Smirnova ¹⁰⁰,
 A.C. Smith ⁴², D.R. Smith ¹⁶², E.A. Smith ⁴⁰, J.L. Smith ¹⁰³, R. Smith ¹⁴⁶, M. Smizanska ⁹³,
 K. Smolek ¹³⁵, A.A. Snesarev ³⁸, S.R. Snider ¹⁵⁸, H.L. Snoek ¹¹⁷, S. Snyder ³⁰, R. Sobie ^{168,w},
 A. Soffer ¹⁵⁴, C.A. Solans Sanchez ³⁷, E. Yu. Soldatov ³⁸, U. Soldevila ¹⁶⁶, A.A. Solodkov ³⁸,
 S. Solomon ²⁷, A. Soloshenko ³⁹, K. Solovieva ⁵⁵, O.V. Solovyanov ⁴¹, P. Sommer ⁵¹,
 A. Sonay ¹³, W.Y. Song ^{159b}, A. Sopczak ¹³⁵, A.L. Sopio ⁹⁸, F. Sopkova ^{29b}, J.D. Sorenson ¹¹⁵,
 I.R. Sotarriva Alvarez ¹⁵⁷, V. Sothilingam ^{64a}, O.J. Soto Sandoval ^{140c,140b}, S. Sottocornola ⁶⁹,
 R. Soualah ¹⁶³, Z. Soumami ^{36e}, D. South ⁴⁹, N. Soybelman ¹⁷², S. Spagnolo ^{71a,71b},
 M. Spalla ¹¹², D. Sperlich ⁵⁵, G. Spigo ³⁷, B. Spisso ^{73a,73b}, D.P. Spiteri ⁶⁰, M. Spousta ¹³⁶,
 E.J. Staats ³⁵, R. Stamen ^{64a}, A. Stampekis ²¹, M. Standke ²⁵, E. Stanecka ⁸⁸,
 W. Stanek-Maslouska ⁴⁹, M.V. Stange ⁵¹, B. Stanislaus ^{18a}, M.M. Stanitzki ⁴⁹, B. Stapf ⁴⁹,
 E.A. Starchenko ³⁸, G.H. Stark ¹³⁹, J. Stark ⁹¹, P. Staroba ¹³⁴, P. Starovoitov ^{64a}, S. Stärz ¹⁰⁶,
 R. Staszewski ⁸⁸, G. Stavropoulos ⁴⁷, P. Steinberg ³⁰, B. Stelzer ^{145,159a}, H.J. Stelzer ¹³²,
 O. Stelzer-Chilton ^{159a}, H. Stenzel ⁵⁹, T.J. Stevenson ¹⁴⁹, G.A. Stewart ³⁷, J.R. Stewart ¹²⁴,
 M.C. Stockton ³⁷, G. Stoicea ^{28b}, M. Stolarski ^{133a}, S. Stonjek ¹¹², A. Straessner ⁵¹,
 J. Strandberg ¹⁴⁷, S. Strandberg ^{48a,48b}, M. Stratmann ¹⁷⁴, M. Strauss ¹²³, T. Streblor ¹⁰⁴,
 P. Strizenc ^{29b}, R. Ströhmer ¹⁶⁹, D.M. Strom ¹²⁶, R. Stroynowski ⁴⁵, A. Strubig ^{48a,48b},
 S.A. Stucci ³⁰, B. Stugu ¹⁷, J. Stupak ¹²³, N.A. Styles ⁴⁹, D. Su ¹⁴⁶, S. Su ^{63a}, W. Su ^{63d},
 X. Su ^{63a}, D. Suchy ^{29a}, K. Sugizaki ¹⁵⁶, V.V. Sulin ³⁸, M.J. Sullivan ⁹⁴, D.M.S. Sultan ¹²⁹,
 L. Sultanaliyeva ³⁸, S. Sultansoy ^{3b}, T. Sumida ⁸⁹, S. Sun ¹⁷³, O. Sunneborn Gudnadottir ¹⁶⁴,
 N. Sur ¹⁰⁴, M.R. Sutton ¹⁴⁹, H. Suzuki ¹⁶⁰, M. Svatos ¹³⁴, M. Swiatlowski ^{159a}, T. Swirski ¹⁶⁹,
 I. Sykora ^{29a}, M. Sykora ¹³⁶, T. Sykora ¹³⁶, D. Ta ¹⁰², K. Tackmann ^{49,t}, A. Taffard ¹⁶²,
 R. Tafirout ^{159a}, J.S. Tafoya Vargas ⁶⁷, Y. Takubo ⁸⁵, M. Talby ¹⁰⁴, A.A. Talyshev ³⁸,
 K.C. Tam ^{65b}, N.M. Tamir ¹⁵⁴, A. Tanaka ¹⁵⁶, J. Tanaka ¹⁵⁶, R. Tanaka ⁶⁷, M. Tanasini ¹⁴⁸,
 Z. Tao ¹⁶⁷, S. Tapia Araya ^{140f}, S. Tapprogge ¹⁰², A. Tarek Abouelfadl Mohamed ¹⁰⁹,
 S. Tarem ¹⁵³, K. Tariq ¹⁴, G. Tarna ^{28b}, G.F. Tartarelli ^{72a}, M.J. Tartarin ⁹¹, P. Tas ¹³⁶,
 M. Tasevsky ¹³⁴, E. Tassi ^{44b,44a}, A.C. Tate ¹⁶⁵, G. Tateno ¹⁵⁶, Y. Tayalati ^{36e,v}, G.N. Taylor ¹⁰⁷,
 W. Taylor ^{159b}, R. Teixeira De Lima ¹⁴⁶, P. Teixeira-Dias ⁹⁷, J.J. Teoh ¹⁵⁸, K. Terashi ¹⁵⁶,

J. Terron ¹⁰¹, S. Terzo ¹³, M. Testa ⁵⁴, R.J. Teuscher ^{158,w}, A. Thaler ⁸⁰, O. Theiner ⁵⁷,
 N. Themistokleous ⁵³, T. Thevenaux-Pelzer ¹⁰⁴, O. Thielmann ¹⁷⁴, D.W. Thomas ⁹⁷,
 J.P. Thomas ²¹, E.A. Thompson ^{18a}, P.D. Thompson ²¹, E. Thomson ¹³¹, R.E. Thornberry ⁴⁵,
 C. Tian ^{63a}, Y. Tian ⁵⁶, V. Tikhomirov ^{38,a}, Yu.A. Tikhonov ³⁸, S. Timoshenko ³⁸,
 D. Timoshyn ¹³⁶, E.X.L. Ting ¹, P. Tipton ¹⁷⁵, A. Tishelman-Charny ³⁰, S.H. Tlou ^{34g},
 K. Todome ¹⁵⁷, S. Todorova-Nova ¹³⁶, S. Todt ⁵¹, L. Toffolin ^{70a,70c}, M. Togawa ⁸⁵, J. Tojo ⁹⁰,
 S. Tokár ^{29a}, K. Tokushuku ⁸⁵, O. Toldaiev ⁶⁹, M. Tomoto ^{85,113}, L. Tompkins ^{146,m},
 K.W. Topolnicki ^{87b}, E. Torrence ¹²⁶, H. Torres ⁹¹, E. Torró Pastor ¹⁶⁶, M. Toscani ³¹,
 C. Toscirri ⁴⁰, M. Tost ¹¹, D.R. Tovey ¹⁴², I.S. Trandaafir ^{28b}, T. Trefzger ¹⁶⁹, A. Tricoli ³⁰,
 I.M. Trigger ^{159a}, S. Trincaz-Duvoid ¹³⁰, D.A. Trischuk ²⁷, B. Trocmé ⁶¹, A. Tropina ³⁹,
 L. Truong ^{34c}, M. Trzebinski ⁸⁸, A. Trzupiek ⁸⁸, F. Tsai ¹⁴⁸, M. Tsai ¹⁰⁸, A. Tsiamis ¹⁵⁵,
 P.V. Tsiareshka ³⁸, S. Tsigaridas ^{159a}, A. Tsirigotis ^{155,r}, V. Tsiskaridze ¹⁵⁸, E.G. Tskhadadze ^{152a},
 M. Tsopoulou ¹⁵⁵, Y. Tsujikawa ⁸⁹, I.I. Tsukerman ³⁸, V. Tsulaia ^{18a}, S. Tsuno ⁸⁵, K. Tsuru ¹²¹,
 D. Tsybychev ¹⁴⁸, Y. Tu ^{65b}, A. Tudorache ^{28b}, V. Tudorache ^{28b}, A.N. Tuna ⁶²,
 S. Turchikhin ^{58b,58a}, I. Turk Cakir ^{3a}, R. Turra ^{72a}, T. Turtuvshin ³⁹, P.M. Tuts ⁴²,
 S. Tzamarias ^{155,d}, E. Tzovara ¹⁰², F. Ukegawa ¹⁶⁰, P.A. Ulloa Poblete ^{140c,140b}, E.N. Umaka ³⁰,
 G. Unal ³⁷, A. Undrus ³⁰, G. Unel ¹⁶², J. Urban ^{29b}, P. Urrejola ^{140a}, G. Usai ⁸, R. Ushioda ¹⁵⁷,
 M. Usman ¹¹⁰, F. Ustuner ⁵³, Z. Uysal ⁸³, V. Vacek ¹³⁵, B. Vachon ¹⁰⁶, T. Vafeiadis ³⁷,
 A. Vaitkus ⁹⁸, C. Valderanis ¹¹¹, E. Valdes Santurio ^{48a,48b}, M. Valente ^{159a}, S. Valentinetti ^{24b,24a},
 A. Valero ¹⁶⁶, E. Valiente Moreno ¹⁶⁶, A. Vallier ⁹¹, J.A. Valls Ferrer ¹⁶⁶, D.R. Van Arneeman ¹¹⁷,
 T.R. Van Daalen ¹⁴¹, A. Van Der Graaf ⁵⁰, P. Van Gemmeren ⁶, M. Van Rijnbach ³⁷,
 S. Van Stroud ⁹⁸, I. Van Vulpen ¹¹⁷, P. Vana ¹³⁶, M. Vanadia ^{77a,77b}, W. Vandelli ³⁷,
 E.R. Vandewall ¹²⁴, D. Vannicola ¹⁵⁴, L. Vannoli ⁵⁴, R. Vari ^{76a}, E.W. Varnes ⁷, C. Varni ^{18b},
 T. Varol ¹⁵¹, D. Varouchas ⁶⁷, L. Varriale ¹⁶⁶, K.E. Varvell ¹⁵⁰, M.E. Vasile ^{28b}, L. Vaslin ⁸⁵,
 G.A. Vasquez ¹⁶⁸, A. Vasyukov ³⁹, L.M. Vaughan ¹²⁴, R. Vavricka ¹⁰², T. Vazquez Schroeder ³⁷,
 J. Veatch ³², V. Vecchio ¹⁰³, M.J. Veen ¹⁰⁵, I. Veliscek ³⁰, L.M. Veloce ¹⁵⁸, F. Veloso ^{133a,133c},
 S. Veneziano ^{76a}, A. Ventura ^{71a,71b}, S. Ventura Gonzalez ¹³⁸, A. Verbytskyi ¹¹²,
 M. Verducci ^{75a,75b}, C. Vergis ⁹⁶, M. Verissimo De Araujo ^{84b}, W. Verkerke ¹¹⁷,
 J.C. Vermeulen ¹¹⁷, C. Vernieri ¹⁴⁶, M. Vessella ¹⁰⁵, M.C. Vetterli ^{145,ac}, A. Vgenopoulos ¹⁰²,
 N. Viaux Maira ^{140f}, T. Vickey ¹⁴², O.E. Vickey Boeriu ¹⁴², G.H.A. Viehhauser ¹²⁹, L. Vignani ^{64b},
 M. Vigl ¹¹², M. Villa ^{24b,24a}, M. Villaplana Perez ¹⁶⁶, E.M. Villhauer ⁵³, E. Vilucchi ⁵⁴,
 M.G. Vincter ³⁵, A. Visibile ¹¹⁷, C. Vittori ³⁷, I. Vivarelli ^{24b,24a}, E. Voevodina ¹¹², F. Vogel ¹¹¹,
 J.C. Voigt ⁵¹, P. Vokac ¹³⁵, Yu. Volkotrub ^{87b}, J. Von Ahnen ⁴⁹, E. Von Toerne ²⁵,
 B. Vormwald ³⁷, V. Vorobel ¹³⁶, K. Vorobev ³⁸, M. Vos ¹⁶⁶, K. Voss ¹⁴⁴, M. Vozak ¹¹⁷,
 L. Vozdecky ¹²³, N. Vranjes ¹⁶, M. Vranjes Milosavljevic ¹⁶, M. Vreeswijk ¹¹⁷, N.K. Vu ^{63d,63c},
 R. Vuillermet ³⁷, O. Vujinovic ¹⁰², I. Vukotic ⁴⁰, S. Wada ¹⁶⁰, C. Wagner ¹⁰⁵, J.M. Wagner ^{18a},
 W. Wagner ¹⁷⁴, S. Wahdan ¹⁷⁴, H. Wahlberg ⁹², J. Walder ¹³⁷, R. Walker ¹¹¹, W. Walkowiak ¹⁴⁴,
 A. Wall ¹³¹, E.J. Wallin ¹⁰⁰, T. Wamorkar ⁶, A.Z. Wang ¹³⁹, C. Wang ¹⁰², C. Wang ¹¹,
 H. Wang ^{18a}, J. Wang ^{65c}, P. Wang ⁹⁸, R. Wang ⁶², R. Wang ⁶, S.M. Wang ¹⁵¹, S. Wang ^{63b},
 S. Wang ¹⁴, T. Wang ^{63a}, W.T. Wang ⁸¹, W. Wang ¹⁴, X. Wang ^{114a}, X. Wang ¹⁶⁵,
 X. Wang ^{63c}, Y. Wang ^{63d}, Y. Wang ^{114a}, Y. Wang ^{63a}, Z. Wang ¹⁰⁸, Z. Wang ^{63d,52,63c},
 Z. Wang ¹⁰⁸, A. Warburton ¹⁰⁶, R.J. Ward ²¹, N. Warrack ⁶⁰, S. Waterhouse ⁹⁷, A.T. Watson ²¹,
 H. Watson ⁶⁰, M.F. Watson ²¹, E. Watton ^{60,137}, G. Watts ¹⁴¹, B.M. Waugh ⁹⁸, J.M. Webb ⁵⁵,
 C. Weber ³⁰, H.A. Weber ¹⁹, M.S. Weber ²⁰, S.M. Weber ^{64a}, C. Wei ^{63a}, Y. Wei ⁵⁵,
 A.R. Weidberg ¹²⁹, E.J. Weik ¹²⁰, J. Weingarten ⁵⁰, C. Weiser ⁵⁵, C.J. Wells ⁴⁹, T. Wenaus ³⁰,
 B. Wendland ⁵⁰, T. Wengler ³⁷, N.S. Wenke ¹¹², N. Wermes ²⁵, M. Wessels ^{64a}, A.M. Wharton ⁹³,
 A.S. White ⁶², A. White ⁸, M.J. White ¹, D. Whiteson ¹⁶², L. Wickremasinghe ¹²⁷,

W. Wiedenmann ¹⁷³, M. Wielers ¹³⁷, C. Wigglesworth ⁴³, D.J. Wilbern ¹²³, H.G. Wilkens ³⁷, J.J.H. Wilkinson ³³, D.M. Williams ⁴², H.H. Williams ¹³¹, S. Williams ³³, S. Willocq ¹⁰⁵, B.J. Wilson ¹⁰³, P.J. Windischhofer ⁴⁰, F.I. Winkel ³¹, F. Winklmeier ¹²⁶, B.T. Winter ⁵⁵, J.K. Winter ¹⁰³, M. Wittgen ¹⁴⁶, M. Wobisch ⁹⁹, T. Wojtkowski ⁶¹, Z. Wolffs ¹¹⁷, J. Wollrath ¹⁶², M.W. Wolter ⁸⁸, H. Wolters ^{133a,133c}, M.C. Wong ¹³⁹, E.L. Woodward ⁴², S.D. Worm ⁴⁹, B.K. Wosiek ⁸⁸, K.W. Woźniak ⁸⁸, S. Wozniowski ⁵⁶, K. Wraight ⁶⁰, C. Wu ²¹, M. Wu ^{114b}, M. Wu ¹¹⁶, S.L. Wu ¹⁷³, X. Wu ⁵⁷, Y. Wu ^{63a}, Z. Wu ⁴, J. Wuerzinger ^{112,aa}, T.R. Wyatt ¹⁰³, B.M. Wynne ⁵³, S. Xella ⁴³, L. Xia ^{114a}, M. Xia ¹⁵, M. Xie ^{63a}, S. Xin ^{14,114c}, A. Xiong ¹²⁶, J. Xiong ^{18a}, D. Xu ¹⁴, H. Xu ^{63a}, L. Xu ^{63a}, R. Xu ¹³¹, T. Xu ¹⁰⁸, Y. Xu ¹⁵, Z. Xu ⁵³, Z. Xu ^{114a}, B. Yabsley ¹⁵⁰, S. Yacoob ^{34a}, Y. Yamaguchi ⁸⁵, E. Yamashita ¹⁵⁶, H. Yamauchi ¹⁶⁰, T. Yamazaki ^{18a}, Y. Yamazaki ⁸⁶, J. Yan ^{63c}, S. Yan ⁶⁰, Z. Yan ¹⁰⁵, H.J. Yang ^{63c,63d}, H.T. Yang ^{63a}, S. Yang ^{63a}, T. Yang ^{65c}, X. Yang ³⁷, X. Yang ¹⁴, Y. Yang ⁴⁵, Y. Yang ^{63a}, Z. Yang ^{63a}, W-M. Yao ^{18a}, H. Ye ^{114a}, H. Ye ⁵⁶, J. Ye ¹⁴, S. Ye ³⁰, X. Ye ^{63a}, Y. Yeh ⁹⁸, I. Yeletsikh ³⁹, B.K. Yeo ^{18b}, M.R. Yexley ⁹⁸, T.P. Yildirim ¹²⁹, P. Yin ⁴², K. Yorita ¹⁷¹, S. Younas ^{28b}, C.J.S. Young ³⁷, C. Young ¹⁴⁶, C. Yu ^{14,114c}, Y. Yu ^{63a}, J. Yuan ^{14,114c}, M. Yuan ¹⁰⁸, R. Yuan ^{63d,63c}, L. Yue ⁹⁸, M. Zaazoua ^{63a}, B. Zabinski ⁸⁸, E. Zaid ⁵³, Z.K. Zak ⁸⁸, T. Zakareishvili ¹⁶⁶, S. Zambito ⁵⁷, J.A. Zamora Saa ^{140d,140b}, J. Zang ¹⁵⁶, D. Zanzi ⁵⁵, O. Zaplatilek ¹³⁵, C. Zeitnitz ¹⁷⁴, H. Zeng ¹⁴, J.C. Zeng ¹⁶⁵, D.T. Zenger Jr ²⁷, O. Zenin ³⁸, T. Ženiš ^{29a}, S. Zenz ⁹⁶, S. Zerradi ^{36a}, D. Zerwas ⁶⁷, M. Zhai ^{14,114c}, D.F. Zhang ¹⁴², J. Zhang ^{63b}, J. Zhang ⁶, K. Zhang ^{14,114c}, L. Zhang ^{63a}, L. Zhang ^{114a}, P. Zhang ^{14,114c}, R. Zhang ¹⁷³, S. Zhang ¹⁰⁸, S. Zhang ⁹¹, T. Zhang ¹⁵⁶, X. Zhang ^{63c}, X. Zhang ^{63b}, Y. Zhang ^{63c}, Y. Zhang ⁹⁸, Y. Zhang ^{114a}, Z. Zhang ^{18a}, Z. Zhang ^{63b}, Z. Zhang ⁶⁷, H. Zhao ¹⁴¹, T. Zhao ^{63b}, Y. Zhao ¹³⁹, Z. Zhao ^{63a}, Z. Zhao ^{63a}, A. Zhemchugov ³⁹, J. Zheng ^{114a}, K. Zheng ¹⁶⁵, X. Zheng ^{63a}, Z. Zheng ¹⁴⁶, D. Zhong ¹⁶⁵, B. Zhou ¹⁰⁸, H. Zhou ⁷, N. Zhou ^{63c}, Y. Zhou ¹⁵, Y. Zhou ^{114a}, Y. Zhou ⁷, C.G. Zhu ^{63b}, J. Zhu ¹⁰⁸, X. Zhu ^{63d}, Y. Zhu ^{63c}, Y. Zhu ^{63a}, X. Zhuang ¹⁴, K. Zhukov ⁶⁹, N.I. Zimine ³⁹, J. Zinsser ^{64b}, M. Ziolkowski ¹⁴⁴, L. Živković ¹⁶, A. Zoccoli ^{24b,24a}, K. Zoch ⁶², T.G. Zorbas ¹⁴², O. Zormpa ⁴⁷, W. Zou ⁴², L. Zwalinski ³⁷.

¹Department of Physics, University of Adelaide, Adelaide; Australia.

²Department of Physics, University of Alberta, Edmonton AB; Canada.

³(^a)Department of Physics, Ankara University, Ankara; (^b)Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye.

⁴LAPP, Université Savoie Mont Blanc, CNRS/IN2P3, Annecy; France.

⁵APC, Université Paris Cité, CNRS/IN2P3, Paris; France.

⁶High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America.

⁷Department of Physics, University of Arizona, Tucson AZ; United States of America.

⁸Department of Physics, University of Texas at Arlington, Arlington TX; United States of America.

⁹Physics Department, National and Kapodistrian University of Athens, Athens; Greece.

¹⁰Physics Department, National Technical University of Athens, Zografou; Greece.

¹¹Department of Physics, University of Texas at Austin, Austin TX; United States of America.

¹²Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.

¹³Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain.

¹⁴Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; China.

¹⁵Physics Department, Tsinghua University, Beijing; China.

¹⁶Institute of Physics, University of Belgrade, Belgrade; Serbia.

- ¹⁷Department for Physics and Technology, University of Bergen, Bergen; Norway.
- ¹⁸(^a)Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA; (^b)University of California, Berkeley CA; United States of America.
- ¹⁹Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany.
- ²⁰Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland.
- ²¹School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom.
- ²²(^a)Department of Physics, Bogazici University, Istanbul; (^b)Department of Physics Engineering, Gaziantep University, Gaziantep; (^c)Department of Physics, Istanbul University, Istanbul; Türkiye.
- ²³(^a)Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá; (^b)Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia.
- ²⁴(^a)Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna; (^b)INFN Sezione di Bologna; Italy.
- ²⁵Physikalisches Institut, Universität Bonn, Bonn; Germany.
- ²⁶Department of Physics, Boston University, Boston MA; United States of America.
- ²⁷Department of Physics, Brandeis University, Waltham MA; United States of America.
- ²⁸(^a)Transilvania University of Brasov, Brasov; (^b)Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; (^c)Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi; (^d)National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca; (^e)National University of Science and Technology Politehnica, Bucharest; (^f)West University in Timisoara, Timisoara; (^g)Faculty of Physics, University of Bucharest, Bucharest; Romania.
- ²⁹(^a)Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava; (^b)Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic.
- ³⁰Physics Department, Brookhaven National Laboratory, Upton NY; United States of America.
- ³¹Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina.
- ³²California State University, CA; United States of America.
- ³³Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom.
- ³⁴(^a)Department of Physics, University of Cape Town, Cape Town; (^b)iThemba Labs, Western Cape; (^c)Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg; (^d)National Institute of Physics, University of the Philippines Diliman (Philippines); (^e)University of South Africa, Department of Physics, Pretoria; (^f)University of Zululand, KwaDlangezwa; (^g)School of Physics, University of the Witwatersrand, Johannesburg; South Africa.
- ³⁵Department of Physics, Carleton University, Ottawa ON; Canada.
- ³⁶(^a)Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca; (^b)Faculté des Sciences, Université Ibn-Tofail, Kénitra; (^c)Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech; (^d)LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda; (^e)Faculté des sciences, Université Mohammed V, Rabat; (^f)Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco.
- ³⁷CERN, Geneva; Switzerland.
- ³⁸Affiliated with an institute covered by a cooperation agreement with CERN.
- ³⁹Affiliated with an international laboratory covered by a cooperation agreement with CERN.
- ⁴⁰Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America.
- ⁴¹LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France.
- ⁴²Nevis Laboratory, Columbia University, Irvington NY; United States of America.
- ⁴³Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark.

- ^{44(a)}Dipartimento di Fisica, Università della Calabria, Rende; ^(b)INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy.
- ⁴⁵Physics Department, Southern Methodist University, Dallas TX; United States of America.
- ⁴⁶Physics Department, University of Texas at Dallas, Richardson TX; United States of America.
- ⁴⁷National Centre for Scientific Research "Demokritos", Agia Paraskevi; Greece.
- ^{48(a)}Department of Physics, Stockholm University; ^(b)Oskar Klein Centre, Stockholm; Sweden.
- ⁴⁹Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany.
- ⁵⁰Fakultät Physik, Technische Universität Dortmund, Dortmund; Germany.
- ⁵¹Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany.
- ⁵²Department of Physics, Duke University, Durham NC; United States of America.
- ⁵³SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom.
- ⁵⁴INFN e Laboratori Nazionali di Frascati, Frascati; Italy.
- ⁵⁵Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany.
- ⁵⁶II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany.
- ⁵⁷Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.
- ^{58(a)}Dipartimento di Fisica, Università di Genova, Genova; ^(b)INFN Sezione di Genova; Italy.
- ⁵⁹II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany.
- ⁶⁰SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom.
- ⁶¹LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France.
- ⁶²Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America.
- ^{63(a)}Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; ^(b)Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao; ^(c)School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai; ^(d)Tsung-Dao Lee Institute, Shanghai; ^(e)School of Physics and Microelectronics, Zhengzhou University; China.
- ^{64(a)}Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; ^(b)Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany.
- ^{65(a)}Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong; ^(b)Department of Physics, University of Hong Kong, Hong Kong; ^(c)Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China.
- ⁶⁶Department of Physics, National Tsing Hua University, Hsinchu; Taiwan.
- ⁶⁷IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France.
- ⁶⁸Centro Nacional de Microelectrónica (IMB-CNM-CSIC), Barcelona; Spain.
- ⁶⁹Department of Physics, Indiana University, Bloomington IN; United States of America.
- ^{70(a)}INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; ^(b)ICTP, Trieste; ^(c)Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy.
- ^{71(a)}INFN Sezione di Lecce; ^(b)Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy.
- ^{72(a)}INFN Sezione di Milano; ^(b)Dipartimento di Fisica, Università di Milano, Milano; Italy.
- ^{73(a)}INFN Sezione di Napoli; ^(b)Dipartimento di Fisica, Università di Napoli, Napoli; Italy.
- ^{74(a)}INFN Sezione di Pavia; ^(b)Dipartimento di Fisica, Università di Pavia, Pavia; Italy.
- ^{75(a)}INFN Sezione di Pisa; ^(b)Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy.
- ^{76(a)}INFN Sezione di Roma; ^(b)Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy.
- ^{77(a)}INFN Sezione di Roma Tor Vergata; ^(b)Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy.
- ^{78(a)}INFN Sezione di Roma Tre; ^(b)Dipartimento di Matematica e Fisica, Università Roma Tre, Roma;

Italy.

^{79(a)} INFN-TIFPA; ^(b) Università degli Studi di Trento, Trento; Italy.

⁸⁰ Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck; Austria.

⁸¹ University of Iowa, Iowa City IA; United States of America.

⁸² Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America.

⁸³ İstinye University, Sariyer, Istanbul; Türkiye.

^{84(a)} Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; ^(b) Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; ^(c) Instituto de Física, Universidade de São Paulo, São Paulo; ^(d) Rio de Janeiro State University, Rio de Janeiro; ^(e) Federal University of Bahia, Bahia; Brazil.

⁸⁵ KEK, High Energy Accelerator Research Organization, Tsukuba; Japan.

⁸⁶ Graduate School of Science, Kobe University, Kobe; Japan.

^{87(a)} AGH University of Krakow, Faculty of Physics and Applied Computer Science, Krakow; ^(b) Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland.

⁸⁸ Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland.

⁸⁹ Faculty of Science, Kyoto University, Kyoto; Japan.

⁹⁰ Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka ; Japan.

⁹¹ L2IT, Université de Toulouse, CNRS/IN2P3, UPS, Toulouse; France.

⁹² Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina.

⁹³ Physics Department, Lancaster University, Lancaster; United Kingdom.

⁹⁴ Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom.

⁹⁵ Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia.

⁹⁶ School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom.

⁹⁷ Department of Physics, Royal Holloway University of London, Egham; United Kingdom.

⁹⁸ Department of Physics and Astronomy, University College London, London; United Kingdom.

⁹⁹ Louisiana Tech University, Ruston LA; United States of America.

¹⁰⁰ Fysiska institutionen, Lunds universitet, Lund; Sweden.

¹⁰¹ Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain.

¹⁰² Institut für Physik, Universität Mainz, Mainz; Germany.

¹⁰³ School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom.

¹⁰⁴ CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France.

¹⁰⁵ Department of Physics, University of Massachusetts, Amherst MA; United States of America.

¹⁰⁶ Department of Physics, McGill University, Montreal QC; Canada.

¹⁰⁷ School of Physics, University of Melbourne, Victoria; Australia.

¹⁰⁸ Department of Physics, University of Michigan, Ann Arbor MI; United States of America.

¹⁰⁹ Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.

¹¹⁰ Group of Particle Physics, University of Montreal, Montreal QC; Canada.

¹¹¹ Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany.

¹¹² Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany.

¹¹³ Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan.

^{114(a)} Department of Physics, Nanjing University, Nanjing; ^(b) School of Science, Shenzhen Campus of Sun Yat-sen University; ^(c) University of Chinese Academy of Science (UCAS), Beijing; China.

¹¹⁵ Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America.

- ¹¹⁶Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen; Netherlands.
- ¹¹⁷Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands.
- ¹¹⁸Department of Physics, Northern Illinois University, DeKalb IL; United States of America.
- ¹¹⁹^(a)New York University Abu Dhabi, Abu Dhabi; ^(b)United Arab Emirates University, Al Ain; United Arab Emirates.
- ¹²⁰Department of Physics, New York University, New York NY; United States of America.
- ¹²¹Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan.
- ¹²²Ohio State University, Columbus OH; United States of America.
- ¹²³Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America.
- ¹²⁴Department of Physics, Oklahoma State University, Stillwater OK; United States of America.
- ¹²⁵Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic.
- ¹²⁶Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America.
- ¹²⁷Graduate School of Science, Osaka University, Osaka; Japan.
- ¹²⁸Department of Physics, University of Oslo, Oslo; Norway.
- ¹²⁹Department of Physics, Oxford University, Oxford; United Kingdom.
- ¹³⁰LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France.
- ¹³¹Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America.
- ¹³²Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America.
- ¹³³^(a)Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisboa; ^(b)Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa; ^(c)Departamento de Física, Universidade de Coimbra, Coimbra; ^(d)Centro de Física Nuclear da Universidade de Lisboa, Lisboa; ^(e)Departamento de Física, Universidade do Minho, Braga; ^(f)Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain); ^(g)Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal.
- ¹³⁴Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic.
- ¹³⁵Czech Technical University in Prague, Prague; Czech Republic.
- ¹³⁶Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic.
- ¹³⁷Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom.
- ¹³⁸IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France.
- ¹³⁹Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America.
- ¹⁴⁰^(a)Departamento de Física, Pontificia Universidad Católica de Chile, Santiago; ^(b)Millennium Institute for Subatomic physics at high energy frontier (SAPHIR), Santiago; ^(c)Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena; ^(d)Universidad Andres Bello, Department of Physics, Santiago; ^(e)Instituto de Alta Investigación, Universidad de Tarapacá, Arica; ^(f)Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile.
- ¹⁴¹Department of Physics, University of Washington, Seattle WA; United States of America.
- ¹⁴²Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom.
- ¹⁴³Department of Physics, Shinshu University, Nagano; Japan.
- ¹⁴⁴Department Physik, Universität Siegen, Siegen; Germany.
- ¹⁴⁵Department of Physics, Simon Fraser University, Burnaby BC; Canada.
- ¹⁴⁶SLAC National Accelerator Laboratory, Stanford CA; United States of America.

- ¹⁴⁷Department of Physics, Royal Institute of Technology, Stockholm; Sweden.
- ¹⁴⁸Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America.
- ¹⁴⁹Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom.
- ¹⁵⁰School of Physics, University of Sydney, Sydney; Australia.
- ¹⁵¹Institute of Physics, Academia Sinica, Taipei; Taiwan.
- ¹⁵²^(a) E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi; ^(b) High Energy Physics Institute, Tbilisi State University, Tbilisi; ^(c) University of Georgia, Tbilisi; Georgia.
- ¹⁵³Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel.
- ¹⁵⁴Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel.
- ¹⁵⁵Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece.
- ¹⁵⁶International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan.
- ¹⁵⁷Department of Physics, Tokyo Institute of Technology, Tokyo; Japan.
- ¹⁵⁸Department of Physics, University of Toronto, Toronto ON; Canada.
- ¹⁵⁹^(a) TRIUMF, Vancouver BC; ^(b) Department of Physics and Astronomy, York University, Toronto ON; Canada.
- ¹⁶⁰Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan.
- ¹⁶¹Department of Physics and Astronomy, Tufts University, Medford MA; United States of America.
- ¹⁶²Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America.
- ¹⁶³University of Sharjah, Sharjah; United Arab Emirates.
- ¹⁶⁴Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden.
- ¹⁶⁵Department of Physics, University of Illinois, Urbana IL; United States of America.
- ¹⁶⁶Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia; Spain.
- ¹⁶⁷Department of Physics, University of British Columbia, Vancouver BC; Canada.
- ¹⁶⁸Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada.
- ¹⁶⁹Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany.
- ¹⁷⁰Department of Physics, University of Warwick, Coventry; United Kingdom.
- ¹⁷¹Waseda University, Tokyo; Japan.
- ¹⁷²Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel.
- ¹⁷³Department of Physics, University of Wisconsin, Madison WI; United States of America.
- ¹⁷⁴Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany.
- ¹⁷⁵Department of Physics, Yale University, New Haven CT; United States of America.
- ^a Also Affiliated with an institute covered by a cooperation agreement with CERN.
- ^b Also at An-Najah National University, Nablus; Palestine.
- ^c Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America.
- ^d Also at Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Thessaloniki; Greece.
- ^e Also at Centro Studi e Ricerche Enrico Fermi; Italy.
- ^f Also at CERN, Geneva; Switzerland.
- ^g Also at CMD-AC UNEC Research Center, Azerbaijan State University of Economics (UNEC); Azerbaijan.
- ^h Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.

- ⁱ Also at Departament de Fisica de la Universitat Autònoma de Barcelona, Barcelona; Spain.
- ^j Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece.
- ^k Also at Department of Physics, California State University, Sacramento; United States of America.
- ^l Also at Department of Physics, King's College London, London; United Kingdom.
- ^m Also at Department of Physics, Stanford University, Stanford CA; United States of America.
- ⁿ Also at Department of Physics, Stellenbosch University; South Africa.
- ^o Also at Department of Physics, University of Fribourg, Fribourg; Switzerland.
- ^p Also at Department of Physics, University of Thessaly; Greece.
- ^q Also at Department of Physics, Westmont College, Santa Barbara; United States of America.
- ^r Also at Hellenic Open University, Patras; Greece.
- ^s Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain.
- ^t Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany.
- ^u Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia; Bulgaria.
- ^v Also at Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco.
- ^w Also at Institute of Particle Physics (IPP); Canada.
- ^x Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.
- ^y Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia.
- ^z Also at National Institute of Physics, University of the Philippines Diliman (Philippines); Philippines.
- ^{aa} Also at Technical University of Munich, Munich; Germany.
- ^{ab} Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China.
- ^{ac} Also at TRIUMF, Vancouver BC; Canada.
- ^{ad} Also at Università di Napoli Parthenope, Napoli; Italy.
- ^{ae} Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America.
- ^{af} Also at Washington College, Chestertown, MD; United States of America.
- ^{ag} Also at Yeditepe University, Physics Department, Istanbul; Türkiye.
- * Deceased