


The Compact Muon Solenoid Experiment
Conference Report

Mailing address: CMS CERN, CH-1211 GENEVA 23, Switzerland



24 October 2024 (v3, 29 October 2024)

Searches for Vector Like Leptons in CMS

Charis Kleio Koraka for the CMS Collaboration

Abstract

We present results from two distinct searches for Vector-Like Leptons (VLLs) conducted by the CMS experiment using data collected during LHC Run 2. The first analysis uses a model-independent approach probing multilepton final states, while the second one focuses on hadronic final states with multiple tau leptons, leveraging model-dependent techniques. Both searches target SU(2) doublet VLL scenarios with couplings to third-generation Standard Model (SM) leptons while the multilepton analysis also targets singlet VLL scenarios. The first analysis yields results that exclude VLL masses up to 1040 GeV in the doublet scenario and between 125-170 GeV in the singlet scenario. In the second analysis, the maximum likelihood fit prefers the presence of signal at the level of 2.8 standard deviations, for a representative VLL mass point of 600 GeV. Finally, projections for the High-Luminosity LHC (HL-LHC) suggest sensitivity improvements, extending the excluded mass range further into the multi-TeV region.

Presented at *LHCP2024 12th Large Hadron Collider Physics Conference*

Searches for Vector Like Leptons in CMS

Charis Kleio Koraka¹

¹*On behalf of the CMS collaboration*

E-mail: charis.kleio.koraka@cern.ch

We present results from two distinct searches for Vector-Like Leptons (VLLs) conducted by the CMS experiment using data collected during LHC Run 2. The first analysis uses a model-independent approach probing multilepton final states, while the second one focuses on hadronic final states with multiple tau leptons, leveraging model-dependent techniques. Both searches target SU(2) doublet VLL scenarios with couplings to third-generation Standard Model (SM) leptons while the multilepton analysis also targets singlet VLL scenarios. The first analysis yields results that exclude VLL masses up to 1040 GeV in the doublet scenario and between 125-170 GeV in the singlet scenario. In the second analysis, the maximum likelihood fit prefers the presence of signal at the level of 2.8 standard deviations, for a representative VLL mass point of 600 GeV. Finally, projections for the High-Luminosity LHC (HL-LHC) suggest sensitivity improvements, extending the excluded mass range further into the multi-TeV region.

*12th Large Hadron Collider Physics Conference (LHCP2024)
3-7 June 2024
Boston, USA*

1. Introduction

Vector-Like Leptons (VLLs) are predicted in various Beyond Standard Model (BSM) theories as solutions to unresolved issues in the Standard Model (SM), such as the hierarchy problem, the existence of dark matter and tensions in muon $g-2$ measurements [3]. These non-chiral particles can be SU(2) doublets or singlets and arise in models such as supersymmetry [12–14], grand unified theories (GUTs) [15, 16], and extra-dimension models [17, 18]. VLLs are further motivated by observed anomalies in B-physics, particularly in lepton flavor universality tests, such as $R(D^*)$ and $R(D)$ measurements, which deviate from SM predictions by approximately 3σ [4]. The CMS experiment [1, 2] has conducted two key searches for VLLs using data from LHC Run 2. One search [7] adopts a model-independent strategy considering a minimal SM extension, while the other [5] focuses on hadronic signatures, primarily involving tau leptons, within the context of the 4321 model [8–10]. Finally, projections for the sensitivity of VLL searches in the High-Luminosity LHC (HL-LHC) [6] are presented.




Figure 1: Left: Values of a_μ from different experiments compared to the SM expectation [3]. Right: Measurements of $R(D)$ versus $R(D^*)$ from various experiments compared against the SM expectation [4].

2. Model-Independent Multilepton Analysis

This model-independent search [7] is performed considering a minimal extension of the SM. VLLs can decay into SM boson-lepton pairs as seen in Figure 2, with their decay rates governed by their mass. This analysis investigates multilepton final states, focusing on events involving light leptons and hadronically decaying tau leptons (τ_h). The events are categorized into seven orthogonal channels, as shown in Figure 3.

2.1 Background Estimation

The dominant backgrounds in this search arise from SM processes that generate three or more leptons, such as WZ , ZZ , $t\bar{t}Z$, and $t\bar{t}W$. These irreducible backgrounds are estimated using simulations and validated in dedicated control regions (CRs). Contributions from misidentified leptons, including photon conversions, are estimated using data-driven methods.




Figure 2: Example processes illustrating production and decay of doublet (left) and singlet (right) VLL pairs at the LHC in the minima SM extension scenarios [7].

Trilepton channels	Quadlepton channels*
<ul style="list-style-type: none"> • 3L: 3 light leptons, 0 had. taus • 2L1T: 2 light leptons, 1 had. tau • 1L2T: 1 light lepton, 2 had. taus 	<ul style="list-style-type: none"> • 4L: 4 light leptons, 0 had. tau • 3L1T: 3 light leptons, 1 had. tau • 2L2T: 2 light lepton, 2 had. taus • 1L3T: 2 light lepton, 2 had. taus <p>* No veto on additional leptons → only lead. four objects are considered</p>

Figure 3: Event categorization into orthogonal channels based on the number of light leptons and τ_h leptons.

2.2 Signal Regions

The signal regions (SRs) are defined in two ways: a model-independent scheme and a BDT-based model-dependent method. The model-independent SRs rely on kinematic properties of the final states (e.g. L_T , p_T^{miss}) to cover a broad phase space for VLL signals, an example of which is shown in Figure 4. In contrast, the BDTs are trained on object and event level variables to enhance sensitivity for specific VLL mass ranges as shown in Figure 5.

2.3 Results

The search excludes VLL masses up to 1040 GeV in the doublet scenario, while in the singlet scenario, masses between 125-170 GeV are excluded. Results are shown in Figure 6. No significant deviations from the SM background were observed.

3. Hadronic Signature Analysis in the 4321 Model

The second analysis searches for VLLs in the context of the 4321 model [5], where VLLs couple primarily to third-generation fermions through interactions with leptoquarks. This is shown in the diagram of Figure 7. The analysis targets hadronic final states with at least three b-tagged jets and multiple tau leptons. Events are categorized based on the multiplicity of τ_h leptons (0, 1, or 2) as shown in Figure 8 and uses a combination of graph neural networks (GNNs) and data-driven methods to distinguish signal from background.




Figure 4: The L_T distribution is shown for events in the trilepton channels. The lower panel shows the ratio of observed events to the total expected background prediction. The gray band on the ratio represents the quadratic sum of statistical and systematic uncertainties in the SM background prediction shown after fitting the data under the background-only hypothesis. An example signal hypothesis for the production of the vector-like tau lepton in the doublet scenario with a mass of 1 TeV, before the fit, is overlaid. Taken from [6].




Figure 5: BDT region defined for the four-lepton channels for the full Run-2 data set. The lower panel shows the ratio of observed events to the total expected background prediction. The gray band on the ratio represents the quadratic sum of statistical and systematic uncertainties in the SM background prediction after fitting the data under the background-only hypothesis. An example signal hypothesis for the production of the vector-like tau lepton in the doublet scenario for a VLL mass of 900 GeV, before the fit, is overlaid. Taken from [6].

3.1 Machine learning classifiers

A machine learning based strategy is employed to construct the observables for signal and background discrimination. A graph neural network approach is selected, in particular the use of ABCNet [11]. Up to 10 objects per event are utilized including jets and hadronic τ leptons to construct the graph. Two classifiers are trained targeting the QCD and $t\bar{t}$ backgrounds considering several kinematical features for the training shown in Figure 9.




Figure 6: Observed and expected upper limits at 95% CL on the production cross section for the vector-like tau leptons in the doublet model (left) and singlet model (right). For the doublet vector-like lepton model, to the left of the vertical dashed gray line, the limits are shown from the model-independent scheme, while to the right the limits are shown from the model dependent BDT regions. For the singlet vector-like lepton model, the limit is shown from the model-independent scheme for all masses [6].




Figure 7: Feynman diagrams showing (left) electroweak production of VLL pairs through s-channel bosons where L represents the VLL, either neutral (N) or charged (E) and (right) vector-like lepton decays mediated by a vector leptoquark, U, decaying primarily to third-generation leptons and quarks. Taken from [5].

3.2 Background Estimation

In the $0\text{-}\tau_h$ category, the main background arises from QCD multijet events, and is estimated using an ABCD method based on the missing transverse energy and the output of the relevant GNN. The $1\text{-}\tau_h$ and $2\text{-}\tau_h$ regions are dominated by $t\bar{t}$ events which are modeled using simulation. Finally, misidentified τ_h leptons contribute significantly to the background in this region and are estimated using a data-driven method. Distributions of the post-fit observables in the different signal extraction regions are shown in Figure 10.

3.3 Results

This search observes a broad mild excess in both the $1\tau_h$ and $2\tau_h$ regions with a local significance of 2.8σ at a representative VLL mass point of 600 GeV. The exclusion limits for the VLL pair production cross section, shown in Figure 11, are set between 10-30 fb, depending on the assumed mass hypothesis.




Figure 8: Diagram of the event categorization and the various signal and control regions used in the analysis. The regions within the solid box are used for the signal extraction while the regions in the dashed box are used for the data driven background determination. Taken from [5].

Variable	Description
η	Pseudorapidity
ϕ	Azimuthal angle
$\log\left(\frac{p_T}{\text{GeV}}\right)$	Logarithm of the p_T
$\log\left(\frac{m}{\text{GeV}}\right)$	Logarithm of the mass
Q	Charge
DEEPIJET score	b tagging discriminant

Figure 9: List of input features used during training of the ABCNet model. Taken from [5].

4. Projections for the HL-LHC

Looking ahead, searches at the High-Luminosity LHC (HL-LHC) will provide greater sensitivity to VLL production. Projections indicate that the excluded mass range for VLLs will extend beyond 1 TeV, reaching into the multi-TeV region. The increased luminosity will allow for further exploration of VLL scenarios, including those involving couplings to first- and second-generation SM leptons. Results of the projection studies are shown in Figure 12.

5. Conclusion

The CMS experiment has conducted two complementary searches for Vector-Like Leptons using LHC Run 2 data. In the first, a minimal SM extension scenario is considered, resulting to VLL masses up to 1040 GeV being excluded in the doublet and between 125-170 GeV in the singlet scenario. The second search is performed in the context of the 4321 model, and hadronic final states are explored. Results show a broad mild excess, with a maximum local significance of 2.8 sigma at a representative VLL mass of 600 GeV. Ongoing and future analyses at the HL-LHC will significantly improve sensitivity, providing critical insights into these BSM candidates.

References

- [1] CMS Collaboration, *The CMS experiment at the CERN LHC*, JINST **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.




Figure 10: Post-fit plots for 2017 (top row) and 2018 (bottom row) showing the fitted distributions in the signal region for the different τ_h multiplicity channels (from left to right: $0\text{-}\tau_h$, $1\text{-}\tau_h$, $2\text{-}\tau_h$). Taken from [5].

- [2] CMS Collaboration, *Development of the CMS detector for the CERN LHC Run 3*, JINST **19** (2024) P05064, doi:10.1088/1748-0221/19/05/P05064.
- [3] Muon g-2 Collaboration, *Measurement of the anomalous magnetic moment of the muon*, Fermilab, 2023. Available at: <https://muon-g-2.fnal.gov/result2023.pdf>
- [4] HFLAV Collaboration, “Combined results for $R(D)$ and $R(D^*)$ ”, available at: <https://hflav-eos.web.cern.ch/hflav-eos/semi/moriond24/html/RDsDstar/RDRDs.html>
- [5] CMS Collaboration, “Search for pair-produced vector-like leptons in final states with third-generation leptons and at least three b quark jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Lett. B* **846** (2023) 137713, doi:10.1016/j.physletb.2023.137713, arXiv:2208.09700.
- [6] CMS Collaboration, *Review of searches for vector-like quarks, vector-like leptons, and heavy neutral leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV at the CMS experiment*, arXiv preprint arXiv:2405.17605 (2024). Available at: <https://arxiv.org/abs/2405.17605>
- [7] CMS Collaboration, *Inclusive nonresonant multilepton probes of new phenomena $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **105** (2022) 112007, doi:10.1103/PhysRevD.105.112007.




Figure 11: Expected and observed 95%CL upper limits on the product of the VLL pair production cross section and the branching fraction to third-generation quarks and leptons, combining the 2017 and 2018 data and all τ_h multiplicity channels. The theoretical prediction in the 4321 model for EW production of VLLs is also shown [5].

- [8] L. Di Luzio, A. Greljo, and M. Nardecchia, “Gauge leptoquark as the origin of B-physics anomalies”, *Phys. Rev. D* **96** (2017) 115011, doi:10.1103/physrevd.96.115011, arXiv:1708.08450.
- [9] L. Di Luzio et al., “Maximal flavour violation: a Cabibbo mechanism for leptoquarks”, *JHEP* **11** (2018) 081, doi:10.1007/jhep11(2018)081, arXiv:1808.00942.
- [10] A. Greljo and B. A. Stefanek, “Third family quark–lepton unification at the TeV scale”, *Phys. Lett. B* **782** (2018) 131, doi:10.1016/j.physletb.2018.05.033, arXiv:1802.04274.
- [11] Zhang, H., *Generalization Bounds for Machine Learning*, arXiv preprint arXiv:2001.05311 (2020). Available at: <https://arxiv.org/abs/2001.05311>
- [12] K. Agashe, R. Contino, and A. Pomarol, “The minimal composite Higgs model”, *Nucl. Phys. B* **719** (2005) 165, doi:10.1016/j.nuclphysb.2005.04.035, arXiv:hep-ph/0412089.
- [13] M. Perelstein, “Little Higgs models and their phenomenology”, *Prog. Part. Nucl. Phys.* **58** (2007) 247, doi:10.1016/j.pnpnp.2006.04.001, arXiv:hep-ph/0512128.
- [14] S. P. Martin, “Extra vectorlike matter and the lightest Higgs scalar boson mass in low-energy supersymmetry”, *Phys. Rev. D* **81** (2010) 035004, doi:10.1103/PhysRevD.81.035004, arXiv:0910.2732.
- [15] J. A. Aguilar-Saavedra, D. E. López-Fogliani, and C. Muñoz, “Novel signatures for vector-like quarks”, *JHEP* **06** (2017) 095, doi:10.1007/JHEP06(2017)095, arXiv:1705.02526.
- [16] G.-Y. Huang, K. Kong, and S. C. Park, “Bounds on the fermion-bulk masses in models with universal extra dimensions”, *JHEP* **06** (2012) 099, doi:10.1007/JHEP06(2012)099, arXiv:1204.05.




Figure 12: Expected HL-LHC exclusion limits for vector-like electrons (upper row), muons (middle row), and tau leptons (lower row) in the doublet model (left) and the singlet model (right). In both cases, limits are calculated using $L_T + p_T^{miss}$ from the model independent SRs described in Sec. 2.2 for all masses [6].

[17] R. Nevzorov, “E6 inspired supersymmetric models with exact custodial symmetry”, *Phys. Rev. D* **87** (2013) 015029, doi:10.1103/PhysRevD.87.015029, arXiv:1205.5967.

[18] I. Dorsner, S. Fajfer, and I. Mustać, “Light vector-like fermions in a minimal SU(5) setup”, *Phys. Rev. D* **89** (2014) 115004, doi:10.1103/PhysRevD.89.115004, arXiv:1401.6870.