



Recent Electroweak Measurements with ATLAS

Brigitte Vachon
McGill University

On behalf of the ATLAS Collaboration

XXXI Cracow Epiphany Conference on the recent LHC Results
13-17 January 2025

The Electroweak Theory

- Rich phenomenology arises from the non-Abelian gauge group structure and spontaneous EW symmetry breaking in the Standard Model

$$SU(2)_L \times U(1)_Y \longrightarrow W^+, W^-, Z^0, \gamma$$


- Mass of electroweak gauge bosons and interactions strength predicted precisely from: g, g', v, λ

$$\rho = \frac{m_W^2}{m_Z^2 \cos^2 \theta_W}$$

- Four input parameters: e.g. $\alpha_{\text{QED}}, G_F, m_Z, m_H$
- LHC offers unique environment to test EW theory!

Electroweak Tests at LHC

Precision Frontier




Radiation corrections modify propagators and decay vertices

$$m_W^2 \left(1 - \frac{m_W^2}{m_Z^2} \right) = \frac{\pi\alpha}{\sqrt{2}G_F} (1 + \Delta r)$$

- Sensitivity to a wide range of physics through quantum loops

Measurements of SM parameters

Energy Frontier



from Nucl. Phys. B525 (1998) 27-50


- Delicate gauge cancellations at high energy
- Enhanced sensitivity to non-SM contributions

Investigation of EW gauge structure

W Boson Mass and Width

Eur. Phys. J. C 84 (2024) 1309, arXiv:2403.15085

- Re-analysis of 2011 data
 - Favourable experimental environment for m_W measurement
 - Consolidate earlier ATLAS results, in the perspective of the latest measurement by CDF.
- Fit p_T^ℓ and m_T^W distributions in $W \rightarrow \ell\nu$ decays
 - Improved statistic based on the profile likelihood
 - Updated to more modern PDF sets
- Measurement requires exquisite precision in lepton energy/momentum calibration ($O(10^{-4})$) and recoil response (\sim few %).



W Boson Mass and Width

Eur. Phys. J. C 84 (2024) 1309, arXiv:2403.15085




W boson mass

$$m_W = 80,366.5 \pm 15.9 \text{ MeV}$$

W boson width

$$\Gamma_W = 2202 \pm 47 \text{ MeV}$$

Simultaneous extraction of m_W and Γ_W




- Total uncertainty reduced by $\sim 15\%$

- **First measurement at LHC**
- **Most precise single-experiment measurements.**

Exclusive W Boson hadronic decays


Phys. Rev. Lett. 133 (2024) 161804, arXiv:2309.15887

- Test bench for the QCD factorization framework
- Could enable W boson mass measurement with a fully reconstructed at future colliders




95% CL upper limits

Branching fraction	Expected $\times 10^{-6}$	Observed $\times 10^{-6}$	
$B(W^\pm \rightarrow \pi^\pm \gamma)$	$1.2^{+0.5}_{-0.3}$	1.9	x4 improvement
$B(W^\pm \rightarrow K^\pm \gamma)$	$1.1^{+0.4}_{-0.3}$	1.7	
$B(W^\pm \rightarrow \rho^\pm \gamma)$	$6.0^{+2.3}_{-1.7}$	5.2	First upper limits



W and Z Boson Production



- First step in the study of gauge bosons at 13.6TeV
 - Total/fiducial/differential cross-section measurements

$pp \rightarrow W$ and $pp \rightarrow Z/\gamma^*$

PLB 854 (2024) 138725, arXiv:2403.12902

- Uncertainty \sim 2-3% dominated by luminosity uncertainty
- Ratio measurements for increased sensitivity to PDF
- Good agreement with NNLO+NNLL QCD and NLO EW predictions

$pp \rightarrow ZZ$

PLB 855 (2024) 138764, arXiv:2311.09715

- Uncertainty \sim 6% (with comparable stat./syst. uncertainties)
- Good agreement with NNLO QCD + NLO EW predictions


ATL-PHYS-PUB-2024-011

Z Boson Invisible Width

Phys. Lett. B 854 (2024) 138705, arXiv:2312.02789


- Sensitive to number of light neutrinos coupling to Z boson and potential non-SM contributions
- Measurements via different final states and analysis strategies test the consistency of SM

$$\Gamma(Z \rightarrow \text{inv}) = R^{\text{miss}} \cdot \Gamma(Z \rightarrow \ell\ell)$$




=

Z(→ inv) + jets



Z(→ ℓℓ) + jets




Z Boson Invisible Width

Phys. Lett. B 854 (2024) 138705, arXiv:2312.02789


$$\Gamma(Z \rightarrow \text{inv}) = 506 \pm 2(\text{stat}) \pm 12(\text{stat})$$

- Single most precise recoil-based measurement ($\sim 2.5\%$)
- Dominated by systematic uncertainties (leptons)
- Result is in agreement with LEP combination and SM predictions



WZ Polarization and RAZ

Phys. Rev. Lett. 133 (2024) 101802, arXiv:2402.16365




- Study of diboson polarization probes **gauge symmetry structure** and **electroweak symmetry breaking mechanism**.
- Radiation-amplitude-zero (RAZ): At LO, $W_T Z_T$ amplitudes predicted to be exactly zero for specific W scattering angle in the WZ rest frame
- Experiments gaining sensitivity to $V_L V_L$ production and starting to study energy dependence of cross-section.

WZ Polarization and RAZ

Phys. Rev. Lett. 133 (2024) 101802, arXiv:2402.16365

- Use leptonic decays: $WZ \rightarrow \ell\nu\ell'\ell'$ ($\ell, \ell' = e, \mu$)
- Longitudinally polarized final state $W_L Z_L$ only make up $\sim 6\%$ of total WZ cross-section.
- Measure polarization fractions in longitudinally-enriched regions.
 - $p_T^{WZ} < 70$ GeV : Reduce jet activity
 - High p_T^Z : Enhance s-channel contributions




- Measurements compatible with SM predictions.

	Measurement		Prediction	
	$100 < p_T^Z \leq 200$ GeV	$p_T^Z > 200$ GeV	$100 < p_T^Z \leq 200$ GeV	$p_T^Z > 200$ GeV
f_{00}	$0.17 \pm_{0.02}^{0.02} \text{ (stat)} \pm_{0.02}^{0.01} \text{ (syst)}$	$0.16 \pm_{0.05}^{0.05} \text{ (stat)} \pm_{0.03}^{0.02} \text{ (syst)}$	0.152 ± 0.006	0.234 ± 0.007
f_{XX}	$0.83 \pm_{0.02}^{0.02} \text{ (stat)} \pm_{0.01}^{0.02} \text{ (syst)}$	$0.84 \pm_{0.05}^{0.05} \text{ (stat)} \pm_{0.02}^{0.03} \text{ (syst)}$	0.120 ± 0.002	0.062 ± 0.002
f_{00} obs (exp) sig.	7.7 (6.9) σ	3.2 (4.2) σ	0.109 ± 0.001	0.058 ± 0.001
			0.619 ± 0.007	0.646 ± 0.008


WZ Polarization and RAZ

Phys. Rev. Lett. 133 (2024) 101802, arXiv:2402.16365

- Variables sensitive to RAZ effect: $\Delta Y(WZ), \Delta Y(\ell_W Z)$
- Extra parton emissions dilute RAZ effect
 - Define 3 regions with reduced jet activity: $p_T^{WZ} < 20, 40, 70$ GeV
- RAZ arises primarily from $W_T Z_T$ amplitude
 - Subtract measured $W_L Z_L$ fractions and background
- Unfolded distributions compatible with SM prediction




Transversely-polarized $W_T Z_T$




Quartic Electroweak Couplings

- Some of the rarest processes experimentally accessible at LHC
 - Vector-boson scattering observed in most channels
 - Increasing sensitivity to triboson production



Diboson
 $W\gamma jj, WZjj$








Triboson
 $W\gamma\gamma, VVZ$ **NEW**

Diboson: $W\gamma jj$

Eur. Phys. J. C 84 (2024) 1064, arXiv:2403.02809




- Observation of EW $W\gamma jj$ with $> 6\sigma$ established using ML techniques
- Study modelling of several key kinematic observables.



Diboson: $WZjj$

JHEP 06 (2024) 192, arXiv:2403.15296

- Multivariate discriminant used to separate EW and strong production modes
- First study of modelling of several key kinematic observables



Triboson



VVZ

arXiv:2412.15123

- BDT discriminants trained in each event category to enhance the separation between signal and background

Process	Signal strength	Cross section (fb)	Observed (expected) sensitivity
VVZ	$1.43 \pm 0.20(\text{stat.})^{+0.21}_{-0.19}(\text{syst.})$	$660^{+93}_{-90}(\text{stat.})^{+88}_{-81}(\text{syst.})$	6.4 (4.7) σ
WWZ	$1.33 \pm 0.28(\text{stat.})^{+0.21}_{-0.17}(\text{syst.})$	$442 \pm 94(\text{stat.})^{+60}_{-52}(\text{syst.})$	4.4 (3.6) σ
WZZ	$2.13^{+1.18}_{-0.96}(\text{stat.})^{+0.76}_{-0.41}(\text{syst.})$	$200^{+111}_{-91}(\text{stat.})^{+65}_{-37}(\text{syst.})$	2.8 (1.6) σ


Observation

Evidence

$W\gamma\gamma$

Phys. Lett. B 848 (2024) 138400

Observation with 5.6σ significance



Anomalous Quartic Gauge Couplings

- Deviations from the SM are quantified in an Effective Field Theory approach

$$\mathcal{L}_{\text{SMEFT}} = \mathcal{L}_{\text{SM}} + \sum_i \frac{f_i^{(6)}}{\Lambda^2} O_i^{(6)} + \sum_i \frac{f_i^{(8)}}{\Lambda^4} O_i^{(8)} + \dots$$

- Considering only operators affecting QGC at dimension-8
- Limits on Wilson coefficients reported without and with unitarity preservation using clipping technique.

W γ jj

Eur. Phys. J. C 84 (2024) 1064, arXiv:2403.02809


Coefficients [TeV ⁻⁴]	Observable	Expected [TeV ⁻⁴]	Observed [TeV ⁻⁴]
f_{T0}/Λ^4	p_T^{jj}	[-2.4, 2.4]	[-1.8, 1.8]
f_{T1}/Λ^4	p_T^{jj}	[-1.5, 1.6]	[-1.1, 1.2]
f_{T2}/Λ^4	p_T^{jj}	[-4.4, 4.7]	[-3.1, 3.5]
f_{T3}/Λ^4	p_T^{jj}	[-3.3, 3.5]	[-2.4, 2.6]
f_{T4}/Λ^4	p_T^{jj}	[-3.0, 3.0]	[-2.2, 2.2]
f_{T5}/Λ^4	p_T^{jj}	[-1.7, 1.7]	[-1.2, 1.3]
f_{T6}/Λ^4	p_T^{jj}	[-1.5, 1.5]	[-1.0, 1.1]
f_{T7}/Λ^4	p_T^{jj}	[-3.8, 3.9]	[-2.7, 2.8]
f_{M0}/Λ^4	p_T^l	[-28, 28]	[-24, 24]
f_{M1}/Λ^4	p_T^l	[-43, 44]	[-37, 38]
f_{M2}/Λ^4	p_T^l	[-10, 10]	[-8.6, 8.5]
f_{M3}/Λ^4	p_T^l	[-16, 16]	[-13, 14]
f_{M4}/Λ^4	p_T^l	[-18, 18]	[-15, 15]
f_{M5}/Λ^4	p_T^l	[-17, 14]	[-14, 12]
f_{M7}/Λ^4	p_T^l	[-78, 77]	[-66, 65]

First LHC constraints

VVZ

arXiv:2412.15123

Most sensitive to $f_{M2}, f_{M3}, f_{M4}, f_{M5}$



Summary

- LHC offers unique environment to test EW theory
- Wealth of electroweak measurements made by ATLAS in recent months.
 - Precision measurements of key Standard Model parameters
 - Investigation of electroweak gauge structure
- Measurements are in good agreement with SM predictions.
- As datasets grow, new opportunities arise to explore increasingly rare physics processes.
- New physics could be just around the corner – the search continues.
- All ATLAS results available at <https://twiki.cern.ch/twiki/bin/view/AtlasPublic>