



CERN-EP-2025-033
24 February 2025

Measurement of ω meson production in pp and p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV

ALICE Collaboration*

Abstract

We present the measurement of the p_{T} -differential production cross section of ω mesons in pp and p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV at midrapidity by ALICE. In addition, the first measurement of the nuclear modification factor R_{pPb} for ω mesons at LHC energies is presented, complementing the existing measurements of lighter neutral mesons such as the π^0 and η . Within the measured p_{T} -range, the R_{pPb} of ω mesons shows no cold nuclear matter effects within the uncertainties, consistent with previous measurements at lower energies. The ω/π^0 ratio is presented for both collision systems, showing no collision system dependence within the uncertainties. The comparison to previously published ω/π^0 ratios at lower and higher collision energies in pp collisions suggests a decreasing trend of the ratio above $p_{\text{T}} = 4$ GeV/ c with increasing collision energy. The data in both collision systems are compared to predictions from PYTHIA 8, EPOS LHC and DPMJET event generators, revealing significant shortcomings in these models' ability to describe the production of ω mesons.

arXiv:2502.19956v1 [nucl-ex] 27 Feb 2025

© 2025 CERN for the benefit of the ALICE Collaboration.

Reproduction of this article or parts of it is allowed as specified in the CC-BY-4.0 license.

*See Appendix A for the list of collaboration members

1 Introduction

Measurements of hadron production in ultra-relativistic nuclear collisions have substantially improved the understanding of Quantum Chromodynamics (QCD), the theory of the strong interaction [1, 2]. While pp collisions serve as a vacuum baseline, analyses of p–A and A–A collisions reveal emergent QCD phenomena, such as collective flow and energy loss in cold and hot nuclear matter [2]. Theoretically, particle production in such collisions can be divided into the hard and soft regimes, denoting scattering processes with large and small momentum transfer Q^2 , respectively. Hard processes, if not modified by energy loss in the QCD medium, can be calculated using perturbative QCD (pQCD), however, this relies on experimental input for the Parton Distribution Function (PDF) and Fragmentation Function (FF) [3–5]. On the other hand, soft processes are not calculable perturbatively, and phenomenological models and Monte-Carlo (MC) event generators are used to describe them.

Studying particle production in collisions of protons and heavy nuclei, such as p–Pb collisions, gives insights into the influence of Cold Nuclear Matter (CNM). Modifications of particle production with respect to the vacuum baseline, namely pp collisions, can be estimated using the nuclear modification factor, which, in the case of p–Pb collisions, is defined as

$$R_{\text{pPb}} = \frac{1}{A_{\text{Pb}}} \left(E \frac{d^3\sigma_{\text{pPb}}}{dp^3} \right) / \left(E \frac{d^3\sigma_{\text{pp}}}{dp^3} \right), \quad (1)$$

where $E \frac{d^3\sigma_{\text{pPb}}}{dp^3}$ denotes the measured production cross section of a certain particle species in p–Pb (pp) collisions, and $A_{\text{Pb}} = 208$ represents the number of nucleons in a lead nucleus. It was found in measurements at the Relativistic Heavy Ion Collider (RHIC) and the Large Hadron Collider (LHC), that the nuclear modification factor is in agreement with unity above $p_{\text{T}} \approx 2$ GeV/c for various mesons and baryons [6–14]. At lower transverse momenta ($p_{\text{T}} < 2$ GeV/c), a significant depletion is observed, showcasing the presence of CNM effects [6–8].

There are different approaches to explain the modification theoretically: nuclear PDFs (nPDFs) are used to model the PDF for single nucleons in a nucleus. Since the first observations of modification of the PDF in such environments [15, 16], the understanding of those nPDFs has been refined, leading to very precise knowledge of their evolution with Bjorken x and momentum transfer Q [17]. Nuclear shadowing, a depletion of the PDF when measured for a nucleon inside a nucleus for $x \lesssim 0.01$, is the cause for a reduction of the observed production cross section of particles below $p_{\text{T}} \approx 3$ GeV/c. Furthermore, a slight enhancement of the PDF is predicted at $x \approx 0.1$, known as anti-shadowing.

In the Color Glass Condensate (CGC) model, suppression arises from high parton densities, resulting in gluon saturation in the low- p_{T} regime [18]. Calculations using the CGC successfully describe the suppression of particle production at low transverse momenta. High- p_{T} measurements at midrapidity, however, probe a region far from the saturation scale, for which those calculations are not applicable.

An alternative to modifications of the initial state is to explain the depletion of particle production at low p_{T} in p–Pb collisions via fully coherent energy loss (FCEL) induced by gluon radiation in the CNM [19].

While measurements of light mesons and baryons are abundant, measurements of heavier particles, like the ω meson, are scarce, mostly due to more complicated decay chains compared to lighter particles, like the neutral pion. However, measurements of these heavier particles give insights into the mass dependence of particle production and the nuclear modification factor. In the case of the ω meson, the comparison to the neutral pion also reveals a possible spin dependence (spin 1 and spin 0, respectively), as the quark content of the π^0 and ω is almost the same [20].

This paper is structured as follows: first, a brief overview of the ALICE experiment and the detector systems involved in the analysis will be presented in Sec. 2, followed by an overview of the datasets in Sec. 3. The reconstruction of charged pions, photons and neutral pions is given in Sec. 4. The ω meson

reconstruction is described in section Sec. 5, followed by a discussion of the systematic uncertainties in Sec. 6. Finally, the results are being discussed in Sec. 7, and the conclusion is presented in Sec. 8.

2 ALICE detector

The ω mesons are reconstructed via their decay into three pions and the subsequent decay of the neutral pion into a pair of photons: $\omega \rightarrow \pi^+\pi^-\pi^0 \rightarrow \pi^+\pi^-\gamma\gamma$. The charged pions are identified using the Time Projection Chamber (TPC) and Time Of Flight (TOF) detector information of the associated tracks reconstructed in ALICE’s central tracking detectors, the Inner Tracking System (ITS) and the TPC. The photons are measured in the Electromagnetic Calorimeter (EMCal), or, in the case the photon converts into a dielectron pair in the detector material, they are reconstructed from their e^+e^- tracks in the tracking detectors. The following paragraphs briefly describe the five key detectors utilized in the analysis, in their configuration during LHC Run 2 (2015–2018). Comprehensive details on the ALICE detectors and their performance are available in Refs. [21] and [22].

The ITS [23, 24] consists of six cylindrical layers of silicon detectors surrounding the beam pipe, going from the innermost to the outermost layer: two layers of Silicon Pixel Detectors (SPD), two layers of Silicon Drift Detectors (SDD), and two layers of Silicon Strip Detectors (SSD). This analysis utilizes the SPD and SSD, covering the full azimuthal angle and a pseudorapidity range of $|\eta| < 0.9$, to locate the collision vertex and to provide the innermost clusters for tracking charged particles.

The TPC [25] is a cylindrical drift detector surrounding the ITS. Charged particles ionize the gas atoms along their trajectory within a pseudorapidity coverage of $|\eta| < 0.9$ and full azimuthal coverage. Due to an applied electric field, the freed electrons and ions then drift to the electrodes on either end of the TPC, multiplied using Multi Wire Proportional Chambers, and finally the signal is read out using cathode pads [26].

The V0 detector [27] is made up of two scintillation counters V0A and V0C, placed on either side of the interaction point along the beam axis, covering the full azimuthal angle for $-3.7 < \eta < -1.7$ as well as $2.8 < \eta < 5.1$, respectively. The V0 detector measures the event multiplicity in the forward region and also serve as ALICE’s Minimum Bias (MB) trigger detector. The adjacent T0 detectors T0A and T0C [28] consist of twelve Cherenkov detectors, covering the full azimuthal angle for $-3.3 < \eta < -2.9$ and $4.5 < \eta < 5.0$. With a time resolution of 50 ps, the T0 detectors are used to determine the collision time as a reference for other detectors, such as the EMCal and TOF.

The Time Of Flight (TOF) detector [29] surrounds the beam pipe with an array of Multigap Resistive Plate Chamber strips at a distance of about 3.7 m, thereby covering the full azimuthal angle within $|\eta| < 0.9$. In combination with the collision time signal provided by the T0 detectors, the TOF measures the flight time of charged particles.

The EMCal [30] is a sampling calorimeter with alternating layers of lead and scintillation material. Within $|\eta| < 0.7$, it covers the azimuthal angles between $80 < \varphi (^{\circ}) < 187$ and $261 < \varphi (^{\circ}) < 319$. Photons can be reconstructed using the EMCal from their electromagnetic showers, starting with a minimum energy of $E_{\text{min}} = 0.7$ GeV [31].

3 Datasets and event selection

The analysis utilizes all datasets of pp and p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV recorded by ALICE in the second run of the LHC. The collisions used for this analysis fulfill the minimum bias (MB) trigger condition of a coincidental signal in both arms of the V0 detector. This includes approximately one billion pp collisions recorded in November 2015 and 2017 and around half as many p–Pb collisions collected in late November 2016. These datasets correspond to an inspected integrated luminosity in

pp (p–Pb) collisions of $\mathcal{L}_{\text{int}} = N_{\text{evt}}/\sigma_{\text{MB}} = 18 \text{ nb}^{-1}$ (0.27 nb^{-1}). The respective visible minimum bias cross sections σ_{MB} were determined via Van der Meer scans [32] for $\sqrt{s_{\text{NN}}} = 5.02$ TeV in pp [33] and p–Pb [34] collisions.

The primary vertex of each collision is reconstructed via tracks in the TPC and ITS, as described in Ref. [22]. The distance in the beam direction (z) between the vertex and the nominal interaction point is required to be within $|z_{\text{vtx}}| < 10$ cm, to ensure good vertex resolution and uniform detector acceptance. To account for triggered collisions in which no vertex is reconstructed, the luminosity used for the normalization was corrected upward by 1.4% (0.5%) in pp (p–Pb) collisions, assuming no ω mesons are produced at mid-rapidity in such events. Pileup collisions with more than one collision vertex reconstructed in the SPD in the same bunch crossing are rejected; this is the case for 0.2% (3.7%) of pp (p–Pb) collisions.

In p–Pb collisions, the nucleon-nucleon center-of-mass system is boosted with $\Delta y_{\text{NN}} = 0.465$ in the direction of the proton beam due to the 2-in-1 magnet design of the LHC [35], which uses the same magnetic field for both beams, accelerating the protons to higher velocities because of their larger charge-to-mass ratio.

4 Pion reconstruction

As already mentioned, in the presented measurement, ω mesons are reconstructed via their dominant decay into $\pi^+\pi^-\pi^0$ ($\mathcal{B} \approx 89.2\%$ [20]). This section discusses the reconstruction of the charged pions, followed by the reconstruction of the neutral pions. The selection criteria and reconstruction methods are the same for the measurement in pp and p–Pb collisions.

4.1 Charged pion selection

Charged particles can be reconstructed from clusters along their trajectory using a Kalman filter, with their momentum measured through their curvature in ALICE’s magnetic field; see Ref. [22] for details. With a lifetime of $\tau^{\pi^\pm} = 7.8 \text{ m}/c$ [20], the vast majority of charged pions traverse the tracking detectors, ITS and TPC, so they can be reconstructed from their tracks. To ensure good quality of the tracks, they are required to have a transverse momentum of $p_{\text{T}} > 100 \text{ MeV}/c$, contain at least one hit in the SPD, include at least 80 TPC clusters and $\chi^2/N_{\text{cluster}}^{\text{TPC}} < 4$.

The charged pions are identified based on each track’s specific energy loss dE/dx in the TPC, quantified via the deviation of the measured energy loss from the expected energy loss relative to the detector resolution. This relative deviation is required to be $|n\sigma_{\text{TPC}}^{\pi^\pm}| < 3$. For charged tracks that include a TOF cluster, the relative deviation from the expected flight time for pions relative to the detector resolution is examined to increase the purity of the pion selection further. Thus, charged particles that the TOF classified as unlikely to be a charged pion ($|n\sigma_{\text{TOF}}^{\pi^\pm}| > 5$) and likely to be either a proton or a kaon ($|n\sigma_{\text{TOF}}^{p/K}| < 3$) were removed from the sample of selected pion tracks.

With the selection criteria described above, the reconstruction efficiency of charged pions is around $\epsilon^{\pi^\pm} \approx 80\%$ for $p_{\text{T}} > 500 \text{ MeV}$, with a purity of $P^{\pi^\pm} = 99\%$ at $p_{\text{T}} \approx 500 \text{ MeV}/c$, decreasing to $P^{\pi^\pm} = 80\%$ at $p_{\text{T}} \approx 7 \text{ GeV}/c$.

4.2 Neutral pion reconstruction

Neutral pions decay at the primary vertex with a branching ratio of $\mathcal{B} \approx 99.8\%$ [20] into a photon pair. The decay photons are reconstructed using either the EMCAL or the Photon Conversion Method (PCM), as laid out below.

Photons reaching the EMCAL produce electromagnetic showers that typically deposit their energy in multiple nearby cells. Clusters are formed by grouping adjacent cells, starting with seed cells that have

an energy $E_{\text{seed}} > 500$ MeV and then aggregating adjacent cells with an energy $E_{\text{agg}} > 100$ MeV, as outlined in Ref. [31]. The cluster energy E_{cls} is obtained as the sum of the individual cell energies, with an additional correction to account for the non-linearity of the energy response [31]. Furthermore, the signal time of each cluster relative to the collision time has to fulfill $-20 \leq t_{\text{cls}} \text{ (ns)} < 25$ to reduce out-of-bunch pileup. Only clusters with an energy of $E_{\text{cls}} > 0.7$ GeV are considered in the analysis to reduce the number of clusters induced by minimum ionizing particles. The shape of clusters is characterized by σ_{long}^2 , quantifying the elongation of the underlying electromagnetic shower; for more details, see Ref. [31]. The elongation of clusters containing more than one cell must be within $0.1 < \sigma_{\text{long}}^2 < 0.7$ to reduce contributions from the merged clusters of π^0 decay photons and those from the low- p_{T} electrons and hadrons. In a final selection step, clusters are rejected if a charged particle track, reconstructed in TPC and ITS, is found within a p_{T} -dependent distance in η and φ from the cluster, as described in Ref. [36].

Approximately 8.5% [22] of photons passing through the ALICE inner detector material convert into an e^+e^- pair within a radial distance of 180 cm from the beam axis. The first step in reconstructing these photons is to select the electron and positron tracks measured in the central tracking detectors. Tracks are required to have a transverse momentum of $p_{\text{T}} > 40$ MeV/ c and to contain at least 60% of the TPC clusters that are expected from the track's trajectory. The track's measured energy loss dE/dx in the TPC is required to be close to that expected for electrons ($-3 < n\sigma_{\text{TPC}}^e < 4$). In order to reduce contamination from charged pions, the measured energy loss is furthermore required to be more than one sigma away from the expected energy loss of pions ($|n\sigma_{\text{TPC}}^{\pi^\pm}| > 1$) for transverse momenta above $p_{\text{T}} > 0.4$ GeV/ c . The distinct V-shaped topology of two tracks with opposite curvatures originating from a common secondary vertex is identified with a V^0 finding algorithm [22]. The distance of the conversion point from the beam axis is required to be $R > 5$ cm to reduce contamination from Dalitz decays but within $R < 180$ cm to ensure good track quality. In simulations, each V^0 candidate is weighted according to [37], to correct for a residual mismatch of the material budget in simulation compared to the data. Further selections that enhance the purity of conversion photons in this V^0 candidate sample, such as requiring the momentum vector of the V^0 to point to the primary vertex, have been applied as described in Ref. [38].

These two photon-reconstruction methods provide three methods of reconstructing neutral pions. These are called PCM or EMCal if both of the decay photons are reconstructed using the same method, or PCM-EMCal, if one photon is measured in the calorimeter while the other is reconstructed with PCM. Neutral pion candidates are accepted if the invariant mass of the photon pair is within $M_{\text{rec}}^{\pi^0}(p_{\text{T}}) \pm n\sigma_{\text{rec}}^{\pi^0}(p_{\text{T}})$, where $M_{\text{rec}}^{\pi^0}$ and $\sigma_{\text{rec}}^{\pi^0}$ are the p_{T} -dependent mass and width of the reconstructed neutral pion taken from Ref. [38]. The parameter n is chosen to be 3 for PCM, 2.5 for PCM-EMCal, and 2 for EMCal to increase the available statistics for the PCM analysis while prioritizing the enhanced purity for the EMCal method.

5 Reconstruction of ω mesons

The transverse momentum and invariant mass of ω meson candidates is calculated from the four-momenta of all possible $\pi^+\pi^-\pi^0$ combinations of selected pions in each event. As the ω meson decays via the strong interaction, with a lifetime of $\tau^\omega \approx 23.2$ fm/ c [20], all decay products are assumed to originate from the collision vertex.

The energy resolution of the involved detectors causes a smearing of the reconstructed mass of the ω meson candidates, thereby reducing the signal-to-background ratio in an invariant-mass-based signal extraction. The dominant fraction of the smearing comes from the reconstruction of the π^0 , whose mass is similarly affected by the detector resolution. As demonstrated in previous publications [39, 40], this effect can be mitigated by shifting the reconstructed invariant mass of an ω candidate by the difference of the reconstructed π^0 invariant mass to the literature value of $M_{\text{PDG}}^{\pi^0} = 134.98$ MeV/ c^2 [20]. In the

presented measurement, a novel approach is introduced to further refine this correction by determining the correlation between a given discrepancy of the reconstructed π^0 mass (M^{π^0}) and the corresponding discrepancy of the reconstructed ω mass (M^ω), determined from the average of this ratio using Monte-Carlo simulations: $\lambda = \langle (M_{\text{rec}}^\omega - M_{\text{gen}}^\omega) / (M_{\text{rec}}^{\pi^0} - M_{\text{PDG}}^{\pi^0}) \rangle$. Here, M_{gen}^ω corresponds to the reference mass that the ω meson was generated with and is extracted from the simulation. This correlation λ was found to depend on the opening angle between the decay photons of the π^0 due to the interplay of the photon energy resolution and the resulting position resolution of the neutral pion candidate. Depending on the reconstruction method and opening angle, its value ranges between $1.3 < \lambda < 2.3$. For further details and figures, see the supplemental note [41]. This correlation is then used to shift the reconstructed mass of each $\pi^+\pi^-\pi^0$ combination individually based on the deviation of the contained π^0 from the PDG mass: $M^{\pi^+\pi^-\pi^0} = M_{\text{rec}}^{\pi^+\pi^-\pi^0} - \lambda (M_{\text{rec}}^{\pi^0} - M_{\text{PDG}}^{\pi^0})$.

The obtained invariant-mass distributions for one p_T interval for each π^0 reconstruction method and collision system are shown in Fig. 1 as black markers. A signal peak around the nominal mass of the ω meson is visible on top of the combinatorial background. The background is described using a third-order polynomial parametrization, shown as a blue line in Fig. 1. It is obtained from a fit of the invariant mass distribution excluding a signal region of $M^\omega \pm 3\sigma^\omega$. This signal region is defined using a preliminary, p_T -independent mass and width, extracted via Gaussian parametrizations of the invariant mass distributions for each reconstruction method. The signal invariant-mass distribution, shown in red, is extracted by subtracting the parametrized combinatorial background from the invariant mass distribution. The signal-to-background ratio at low p_T is approximately three times higher in pp collisions than in p–Pb collisions, which coincides with an approximately three times larger event multiplicity in the latter [42]. Due to the better energy resolution of PCM compared to the EMCal, the width of the peaks, and thereby the signal-to-background ratio, also vary between the reconstruction methods. The raw ω yield for each p_T interval is then calculated as the sum of bin contents within the aforementioned signal region, denoted by black dashed lines in Fig. 1.

The reconstruction efficiency (ϵ_{rec}) and geometrical acceptance (A), needed to correct the raw yield, are determined by applying the reconstruction and selections to simulated events using PYTHIA 8.2 Monash 2013 [43] in pp and DPMJET [44] in p–Pb collisions. The propagation of particles through the detector is simulated using GEANT 3 [45], taking into account the detector conditions during the data taking. An additional correction is applied to the cluster energies in the simulation, referred to as MC cluster energy fine tuning, to accurately reproduce the cluster energies observed in data, as described in Ref. [31]. The resulting mass and width of the ω meson extracted in data and simulations are in agreement within the respective statistical uncertainties of each π^0 reconstruction method; see supplemental note [41].

The geometrical acceptance is calculated as the ratio of the number of accepted ω mesons (i.e., where all decay particles are within the geometrical limits of the respective detector) to all generated ω mesons within $|y| < 0.85$. The reconstruction efficiency is defined as the ratio of reconstructed to accepted ω mesons in the simulation: $\epsilon_{\text{rec}} = N_{\text{rec}}^\omega / N_{\text{acc}}^\omega$. Here, the number of reconstructed ω mesons can either be obtained from the background-subtracted signal or by counting $\pi^+\pi^-\pi^0$ triplets that are validated to come from the same ω meson, checked at generator level in the simulation, thereby avoiding the need for background subtraction. Since both efficiencies were found to agree within their uncertainties and the latter significantly decreases the statistical uncertainties, the *validated* efficiency is utilized in this analysis. Figure 2 shows the total correction factor applied to the raw spectra, consisting of a product of the geometrical acceptance, validated efficiency, as well as solid angle normalization, where $\Delta\phi = 2\pi$ and $\Delta y = 1.7$ represent the azimuthal and rapidity ranges in which ω mesons were produced in the simulations.

While the total correction factors do not differ between the two collision systems, the requirement of either one or both photons converting for the PCM-EMCal and PCM methods reduces their overall




Figure 1: Invariant mass distributions of $\pi^+\pi^-\pi^0$ triplets in the indicated p_T ranges for π^0 's reconstructed with PCM (top), PCM-EMCal (middle), and EMCal (bottom) in pp (left) and p-Pb (right) collisions. Subtraction of the third-order polynomial background description (blue) from the $\pi^+\pi^-\pi^0$ candidates (black) results in the signal of ω candidates (red). In p-Pb collisions, this signal is scaled by a factor of three for better visibility. The vertical dashed lines denote the invariant mass region where the raw yield was obtained through bin counting, as described in Sec. 5.




Figure 2: Correction factors ε applied to the raw ω yields for each π^0 reconstruction method in pp and p–Pb collisions. The correction factor comprises the geometrical acceptance (A), the reconstruction efficiency (ε_{rec}), and normalization to azimuthal and rapidity ranges.

reconstruction efficiency by a factor of $1/8.5\%$. Although this limits the p_T reach of PCM at high p_T , the good energy resolution of PCM photons enables the reconstruction of ω mesons down to lower p_T .

For the calculation of the nuclear modification factor R_{pPb} , a correction factor is applied to the p_T spectrum of the ω measured in pp collisions to account for the boost of the center-of-mass frame in p–Pb collisions. Using a PYTHIA 8 [43] simulation, the production is found to be 0.8% (1.2%) smaller in the shifted rapidity regime of $-1.315 < y < 0.385$ compared to the central rapidity range of $|y| < 0.85$ for low (high) p_T .

6 Systematic uncertainties

The systematic uncertainties of the ω meson cross sections and the nuclear modification factor were evaluated as a function of p_T by variations of the selection criteria. Table 1 displays all sources of systematic uncertainties and their magnitude for the three π^0 reconstruction methods.

For p_T -dependent sources, the relative uncertainty is given in a low p_T interval ($3.2 < p_T$ (GeV/ c) < 4.0) and an intermediate p_T interval ($8.0 < p_T$ (GeV/ c) < 10.0). The total systematic uncertainty for each reconstruction method is calculated as the quadratic sum, assuming no correlation between the different uncertainty sources. Due to the small signal-to-background ratio, the signal extraction represents the dominant contribution of systematic uncertainty for all measured spectra and reconstruction methods, as also found in previous measurements of the ω meson [39, 40]. Six parameters were identified as possible sources of this systematic uncertainty: the use of the validated efficiency, the integration and parametrization ranges, the description of the background and signal, and the choice of the λ parameter in the applied ω meson mass resolution correction, introduced in Sec. 5. The signal extraction is performed using all possible combinations of reasonable settings for these parameters to estimate this uncertainty accurately. The resulting 198 variations of the spectra were found to be normally distributed, allowing for the extraction of the 1σ uncertainty.

The material budget uncertainty quoted in Table 1 accounts for the accuracy of the detector materials description in the simulation. The material uncertainties per EMCal photon and conversion photon are 2.1% and 2.5% , respectively [36, 37].

Uncertainties stemming from the reconstruction of conversion and calorimeter photons (see conv. and calo. photons in Tab. 1), as well as from neutral and charged pions, were estimated as described in Refs. [38, 40] through variations of the selection criteria introduced in Sec. 4.

π^0 reconstruction	ω cross section in pp						ω cross section in p-Pb						ωR_{pPb}					
	PCM		PCM-EMCaI		EMCaI		PCM		PCM-EMCaI		EMCaI		PCM		PCM-EMCaI		EMCaI	
	3.6	9.0	3.6	9.0	3.6	9.0	3.6	9.0	3.6	9.0	3.6	9.0	3.6	9.0	3.6	9.0	3.6	9.0
Signal extraction	8.6	16.3	8.4	14.3	8.0	10.6	14.4	11.7	16.5	12.7	8.1	9.6	11.9	14.4	12.7	14.9	7.8	11.5
Material budget	5.0	5.0	4.6	4.6	4.2	4.2	5.0	5.0	4.6	4.6	4.2	4.2	-	-	-	-	-	-
Conv. photons	5.8	12.7	1.7	6.5	-	-	9.4	8.9	1.6	4.4	-	-	10.5	13.3	2.6	7.6	-	-
Calo. photons	-	-	3.1	4.3	3.6	5.6	-	-	3.7	4.3	6.5	3.4	-	-	4.4	4.7	5.4	7.5
Neutral pions	5.1	8.3	5.1	11.7	3.3	3.3	4.3	5.6	5.9	7.9	6.5	3.4	6.1	6.6	6.8	14.4	4.4	4.5
Charged pions	5.7	17.7	3.5	4.3	4.4	6.3	7.2	9.8	4.2	3.9	3.4	3.4	8.3	13.4	6.8	6.1	5.4	7.5
Cross section	1.8	1.8	1.8	1.8	1.8	1.8	3.5	3.5	3.5	3.5	3.5	3.5	3.9	3.9	3.9	3.9	3.9	3.9
Branching ratio	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	-	-	-	-	-	-
Rapidity shift	-	-	-	-	-	-	-	-	-	-	-	-	0.1	0.1	0.1	0.1	0.1	0.1
Total systematic	14.0	29.0	12.1	21.0	11.3	14.7	20.1	19.5	19.3	17.6	12.9	12.9	19.7	25.2	17.6	23.9	12.7	16.5
Statistical	11.8	32.2	7.6	16.5	6.1	9.6	11.8	32.2	8.5	9.9	7.1	6.7	19.5	32.2	11.5	19.5	9.5	11.9
Combined stat.	4.6	8.1	8.9	13.7	8.9	13.7	5.3	5.4	5.3	5.4	7.1	9.9	7.1	9.9	7.1	9.9	7.1	9.9
Combined sys.	8.9	13.7	8.9	13.7	8.9	13.7	11.3	11.4	11.3	11.4	11.3	11.4	11.2	15.3	11.2	15.3	11.2	15.3

Table 1: Compilation of the relative systematic uncertainties (%) from the various sources for the pp and p-Pb cross section and the R_{pPb} using the three π^0 reconstruction methods. For p_T -dependent uncertainties, the relative value is given for the two intervals $3.2 < p_T$ (GeV/c) < 4.0 and $8.0 < p_T$ (GeV/c) < 10.0 , allowing for a comparison between the reconstruction methods in the low- and mid- p_T regime. The systematic uncertainties from different sources are added in quadrature, yielding the total systematic uncertainty. Together with the statistical uncertainties, the cross sections and nuclear modification factors are combined using the BLUE method [46]. The two last rows compile the resulting combined statistical and systematic uncertainties. The two bottom rows show the statistical and systematic uncertainty of the cross sections and the R_{pPb} after combining the results from the three π^0 reconstruction methods for the low- and mid- p_T regime.




Figure 3: Ratio of the ω meson production cross sections extracted using the three different π^0 reconstruction methods to a combined Lévy-Tsallis parametrization. Both in pp (left) and in p–Pb (right), agreement between the reconstruction methods is observed within the statistical (bars) and systematic (boxes) uncertainties.

For R_{pPb} , correlated uncertainties between the two collision systems cancel out by simultaneously varying the selection criteria and signal extraction in the two contributing cross sections.

The uncertainties of the visible MB cross section in pp and p–Pb of 1.8% [33] and 3.5% [34], respectively, were taken from their corresponding publications. For the uncertainty in the branching ratio $\omega \rightarrow \pi^+\pi^-\gamma\gamma$, the uncertainties of the two consecutive decays of the ω and π^0 meson quoted by the Particle Data Group (PDG) [20] are added in quadrature. The rapidity shift applied to the pp reference measurement for the R_{pPb} is assumed to introduce a constant uncertainty of 0.1%, which corresponds to 10% of the shift, taking into consideration the statistical uncertainty and introduced MC dependence.

7 Results

7.1 Production cross section of ω mesons

The p_{T} differential Lorentz-invariant production cross section of ω mesons in pp and p–Pb collisions is calculated for all three π^0 reconstruction methods as

$$E \frac{d^3\sigma^\omega}{d^3p} = \frac{1}{\mathcal{L}_{\text{int}}} \frac{1}{2\pi\Delta y A \epsilon_{\text{rec}}} \frac{1}{\mathcal{B}} \frac{N^\omega}{p_{\text{T}} \Delta p_{\text{T}}}, \quad (2)$$

where \mathcal{L}_{int} is the integrated luminosity in the respective collision system introduced in Sec. 3. The number of reconstructed ω mesons N^ω is furthermore normalized by the total correction factor $\epsilon = 2\pi\Delta y A \epsilon_{\text{rec}}$ discussed in Sec. 5, its p_{T} value and the width of the p_{T} interval Δp_{T} , as well as the branching ratio $\mathcal{B}(\omega \rightarrow \pi^+\pi^-\gamma\gamma) = (88.15 \pm 0.70)\%$.

Figure 3 shows the ratio of the ω meson cross sections extracted using the three π^0 reconstruction methods to a combined Lévy-Tsallis parametrization, as defined in Ref. [40]. Agreement is observed between the extracted cross sections within their respective statistical and systematic uncertainties. While the overlap of different π^0 reconstruction methods reduces the statistical and systematic uncertainties, Fig. 3 highlights their complementarity, extending the p_{T} coverage of the measurement. The three production cross sections, and the respective R_{pPb} 's, for each collision system are combined using the best linear unbiased estimate (BLUE) method [46], taking into consideration the correlation between the uncertainties, as done in previous measurements [39, 40].

The combined p_{T} differential Lorentz-invariant production cross section of ω mesons is extracted in pp and p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV within the transverse momentum range of $1.8 \leq p_{\text{T}} \text{ (GeV/c)} < 20$




Figure 4: Lorentz-invariant ω meson production cross section in pp (open markers) and p–Pb (closed markers). Statistical uncertainties are represented by vertical error bars, while boxes show the systematic uncertainties. Furthermore shown are Lévy-Tsallis parametrizations of the two cross sections and two predictions from simulations of the ω meson production per collision system. The lower panels contain the ratios of the data and simulations to the parametrization of the data points in the respective collision system. A gray box in each ratio panel depicts the normalization uncertainty of the visible minimum bias cross section.

and $2.2 \leq p_T$ (GeV/ c) < 20 , respectively. The pp and p–Pb measurements are shown in Fig. 4, covering the rapidity interval of $|y| < 0.85$ and $-1.315 < y < 0.385$, respectively. To account for the finite width of the p_T -intervals, the horizontal positions of the data points shown in Fig. 4 are shifted towards lower p_T , according to the method outlined in Ref. [47]. The description of the data points using a Lévy-Tsallis parametrization, analogous to Ref. [40], yielded p_T -shifts on the order of 1% towards lower p_T . This bin-shift is performed in y -coordinates for the ω/π^0 ratios and R_{pPb} where, due to the similar shape of the spectra, the shift is below 1% for all p_T .

Figure 4 also contains four predictions from simulations using different MC generators and a Lévy-Tsallis parametrization of the data for each collision system. The χ^2/ndf of the parametrization is 0.49 (0.48) in pp (p–Pb), taking into account the total uncertainties. The lower panels depict the ratio of data and the predictions from the event generators to the Lévy-Tsallis parametrization. The DPMJET simulation [44], based on the Dual Parton Model (DPM), roughly describes the p_T dependence of the cross section in p–Pb, but it underestimates the production of ω mesons by approximately 30%. On the other hand, the production of ω mesons in p–Pb collisions is well described by EPOS LHC [48]. The EPOS model describes hadronic interactions following a quantum mechanical multiple-scattering approach based on partons and strings with built-in collective hadronization [49]. The EPOS LHC adaptation [48] includes modifications, aiming to describe the first LHC results. However, when applied to pp collisions, the EPOS LHC simulations do not describe the data, with discrepancies of up to 100%, as shown in the lower panel. While the other pp simulation, PYTHIA 8.2 with the Monash 2013 tune [43], describes the shape of the production cross section in pp collisions, it overestimates the production of ω




Figure 5: Production ratio ω/π^0 in pp and p–Pb collisions calculated using the ω meson production measurements and the corresponding charged [50] and neutral pion [6] references, respectively. The charged pion reference was scaled up by 3.3% (see text for details). Also shown are two predictions from simulations per collision system, with bands representing their statistical uncertainties, as well as a m_T scaling curve, which converges towards a high- p_T constant fitted to the ratio for $p_T > 4$ GeV/ c .

mesons by about 30%. A similar deviation between these PYTHIA 8 predictions and the respective measured cross section was observed in previous ω meson analyses at different collision energies [39, 40], hinting at a possible energy-independent overestimation of the ω meson production in the Monash 2013 tune of PYTHIA 8.

7.2 ω/π^0 ratios

In p–Pb collisions, the ratio of produced ω and π^0 mesons is calculated from the measured ω meson cross section and the corresponding π^0 spectrum at $\sqrt{s_{\text{NN}}} = 5.02$ TeV [6]. To obtain a π^0 reference for the pp measurement, the published spectrum of charged pions [50] is used as a proxy. To account for the slightly higher production yield of neutral pions mainly due to the isospin symmetry breaking $\eta \rightarrow \pi^0\pi^0\pi^0$ decay, the measured charged pion spectrum is scaled up by 3.3%. This surplus is determined from the $(\pi^+ + \pi^-)/(2\pi^0)$ ratio in pp collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV simulated using PYTHIA 8 [43] via a constant parametrization for $1.8 \leq p_T$ (GeV/ c) < 20 . An additional relative systematic uncertainty of 3.3% is added to the ω/π^0 ratio in pp collisions to account for the model dependence introduced. Correlated systematic uncertainties of the visible MB cross section, the $\pi^0 \rightarrow \gamma\gamma$ branching ratio, and in the case of pp the material budget, are hereby removed. All other systematic and statistical uncertainties are assumed to be uncorrelated. Figure 5 shows the measured ω/π^0 ratio in the two collision systems; neither of them display a significant p_T dependence.

Taking into consideration the bin-by-bin correlation between the systematic uncertainties, an average ω/π^0 constant for $p_T > 4$ GeV/ c is evaluated following the procedure described in Ref. [46], yielding $C_{\text{pp}}^{\omega/\pi^0} = 0.48 \pm 0.02$ (stat.) ± 0.06 (sys.) in pp collisions, and $C_{\text{pPb}}^{\omega/\pi^0} = 0.52 \pm 0.01$ (stat.) ± 0.04 (sys.) in p–Pb collisions. Within the given uncertainties and p_T reach, the ω/π^0 production ratios agree with one another, suggesting the production ratio to be independent of CNM effects within the given uncertainties and inspected p_T range.

The extracted high- p_T constants can furthermore be used as input for the transverse mass scaling prediction of the ω/π^0 ratio. This empirical scaling relation describes the production ratio by assuming that the production of all mesons follows the same fundamental underlying function of the particle’s transverse mass ($m_T = \sqrt{p_T^2 + m_{\text{inv}}^2}$), scaled by a constant parameter C [51]. From the extracted high- p_T constants, this assumption is used to derive the p_T dependence of the particle ratio, as shown as purple bands in




Figure 6: Compilation of measurements of the ω/π^0 ratio (left) in pp collisions at various center-of-mass energies covering $0.062 < \sqrt{s}$ (TeV) < 13 at the ISR [54], PHENIX [55] and ALICE [39, 40], including the pp measurement presented herein. Vertical error bars represent total uncertainties, high- p_T data points at $\sqrt{s} = 13$ TeV up to $p_T = 50$ GeV/c are omitted here for better visibility. The right-hand side displays values obtained from parametrizations of the ω/π^0 ratios above $p_T > 4$ GeV/c, as well as predictions of this high- p_T constant for different lower bounds using PYTHIA 8 [43].

Fig. 5. In both collision systems, a slight tension between the ω/π^0 ratio prediction assuming m_T scaling and the data is observed at low transverse momenta $p_T \approx 3$ GeV/c, where m_T scaling suggests a decrease of the ω/π^0 ratio, which is not observed in the data within uncertainties. A slight discrepancy is expected, as m_T scaling was found to be broken for LHC energies due to the feed-down of decays into the π^0 [51]. However, the increased number of neutral pions from the feed-down of higher mass particles would decrease the ω/π^0 ratio compared to the m_T scaling prediction. The excess of the ω/π^0 ratio at low p_T compared to the m_T scaling was also observed at $\sqrt{s} = 13$ TeV, but its underlying cause remains unexplained. This discrepancy shows the need for ω meson production measurements such as this one, as studies of direct photons or dileptons commonly resort back to this m_T scaling prediction to evaluate hadronic feed-down contributions due to a lack of experimental data [52, 53].

In addition to the measured production ratios, Fig. 5 also includes predictions of the ω/π^0 ratio by three different event generators. The predictions of the ω/π^0 ratio by EPOS LHC [48], shown in red, are very similar between pp and p–Pb, suggesting the system size does not have a strong impact on the relative hadronization fractions into ω mesons and neutral pions. However, the production ratio is overestimated by about 100% in both collision systems. So, while EPOS LHC describes the production of ω mesons in p–Pb and not in pp collisions, see Fig. 4, the π^0 production is described only in pp and not in p–Pb collisions. This inadequate collision system and particle species dependence of the EPOS LHC predictions possibly hints at a lack of experimental data for tuning the collective hadronization implemented in EPOS LHC [49]. The similar production ratio predictions by PYTHIA 8 [43] in pp and DPMJET [44] in p–Pb can be explained by the fact that the hadronization implemented in DPMJET is based on the Lund string model, also used by the PYTHIA 8 event generator. Both generators using the Lund string model overestimate the ω/π^0 ratio by about 10 – 20%. It is, however, not evident whether the different hadronization descriptions cause this more accurate description of the ω/π^0 ratio compared to EPOS LHC, or whether this difference should be attributed to the different tunes and experimental input used.

Figure 6 shows the measured ω/π^0 ratio in pp collisions, compared to previous measurements, covering almost three orders of magnitude in center-of-mass energies. For the data points of the ISR measurements, only a minimum p_{T} was defined in the measurement. The p_{T} positions of these data points in a given p_{T} interval were therefore set to the expectation value of the Lévy-Tsallis parametrization of the ω cross section in pp collisions at $\sqrt{s} = 62$ GeV, taken from PYTHIA 8 [43].

The ω/π^0 ratios measured at different energies are compatible within the given total uncertainties represented by vertical bars. A slight tension is visible when considering the entire p_{T} range of the measurement at $\sqrt{s_{\text{NN}}} = 200$ GeV, depicted with green markers, and at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, shown with red markers. To better visualize this possible difference of the ω/π^0 ratio for different center-of-mass energies, the ratios shown in Fig. 6 are parametrized with a constant for $p_{\text{T}} > 4$ GeV/c, with the resulting values of these high- p_{T} fits of the ω/π^0 ratio compiled in Fig. 6, right panel. As the bin-by-bin correlation of the systematic uncertainty for the cited measurements is unknown, their systematic uncertainties are assumed to be fully correlated between the bins for calculating the uncertainty of the high- p_{T} constant. This approach provides an upper-limit estimation of the total uncertainties shown in Fig. 6. To isolate possible effects of the center-of-mass energy, only ω/π^0 ratios measured in pp collisions are considered for this comparison. The resulting high- p_{T} ω/π^0 constant as a function of the center-of-mass energy presented in Fig. 6 displays a slight tendency towards a lower ω/π^0 ratio with rising center-of-mass energy. The significance of this effect is estimated by calculating an averaged ratio at low center-of-mass energy ($62 < \sqrt{s}$ (GeV) < 200) and one at high center-of-mass energy ($5.02 < \sqrt{s}$ (TeV) < 13). By assuming all uncertainties between the different analyses to be fully uncorrelated, the averaged ratios are $\langle \omega/\pi^0 \rangle_{\text{low } \sqrt{s}} = 0.78 \pm 0.09$ and $\langle \omega/\pi^0 \rangle_{\text{high } \sqrt{s}} = 0.56 \pm 0.04$, which are shown as colored bands in Fig. 6. This corresponds to a difference of the ratios at low and high center-of-mass energies with a significance of 2.2σ .

The trend of a decreasing ω/π^0 ratio with rising center-of-mass energy is also observed in PYTHIA 8 simulations [43], shown in blue, cyan, and purple lines for different p_{T} -thresholds in Fig. 6. In the simulations, the production of heavier particles at larger center-of-mass energies is found to result in more hadronic feed-down into primary neutral pions, thereby decreasing the ω/π^0 ratio. Further precision measurements at both low and high collision energies are needed to investigate this trend.

7.3 Nuclear modification factor R_{pPb}

The nuclear modification factor of the ω meson production at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, extracted in the transverse momentum range of $2.2 \leq p_{\text{T}}$ (GeV/c) < 20 and the rapidity interval of $-1.315 < y < 0.385$, is shown with red markers in Fig. 7. This represents the first measurement of the nuclear modification factor of the ω meson at LHC energies. Within the statistical and systematic uncertainties, represented by vertical bars and boxes, respectively, the nuclear modification factor is found to be compatible with unity over the measured p_{T} range. Consequently, no CNM effects on the ω meson production are observed within the given uncertainties.

Figure 7 also displays the nuclear modification factor for neutral pions, measured at the same center-of-mass energy of $\sqrt{s_{\text{NN}}} = 5.02$ TeV [6]. The visible agreement between the nuclear modification factors for the light π^0 and the six times heavier ω meson [20] implies that, within the p_{T} range and the given uncertainties of the two measurements, no mass dependence of the nuclear modification factor is observed.

Finally, Fig. 7 also includes the nuclear modification factors R_{dAu} for the production of ω mesons at $\sqrt{s_{\text{NN}}} = 200$ GeV [56] in d–Au collisions, measured in different decay channels. These nuclear modification factors R_{dAu} are compatible with the herein presented R_{pPb} within the respective statistical and systematic uncertainties.




Figure 7: Measured nuclear modification factor R_{pPb} of the ω meson at $\sqrt{s_{\text{NN}}} = 5.02$ TeV shown with red markers, as well as a nuclear modification factor of neutral pions at the same center-of-mass energy [6] and the measurements for ω mesons at $\sqrt{s_{\text{NN}}} = 200$ GeV [56]. Vertical error bars and boxes represent statistical and systematic uncertainties, while the two solid boxes show the normalization uncertainty of the ALICE measurements.

8 Conclusion

The p_{T} -differential invariant cross sections of ω mesons in pp and p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV have been presented. The measurements, performed using the $\omega \rightarrow \pi^+ \pi^- \pi^0$ decay channel, cover a transverse momentum range of $1.8 \leq p_{\text{T}} \text{ (GeV/c)} < 20$ and $2.2 \leq p_{\text{T}} \text{ (GeV/c)} < 20$, respectively. Three partially independent reconstruction methods were used for the reconstruction of the neutral pion, significantly reducing the systematic uncertainties of the combined ω cross section. Predictions from MC event generators in both pp and p–Pb collisions do not describe the ω production cross section within the given uncertainties. While EPOS LHC describes the spectrum in p–Pb reasonably well, it overestimates the data by up to 100% in pp collisions. PYTHIA 8 overestimates the spectrum in pp as previously seen in pp at $\sqrt{s} = 13$ TeV, and predictions from DPMJET in p–Pb collisions underestimate the spectrum. The ω/π^0 ratio agrees between pp and p–Pb collisions, indicating no system size dependence of the ratio, as also observed for the η/π^0 ratio [6]. Comparisons between measurements of the ω/π^0 in pp collisions for energies up to $\sqrt{s} = 200$ GeV and LHC energies suggest an energy dependence of the ratio for $p_{\text{T}} > 4$ GeV/c with a significance of 2.2σ . In PYTHIA 8, showing qualitatively the same dependence, this trend is attributed to a rising feed-down contribution into the π^0 spectrum with rising collision energy. Finally, the first R_{pPb} for ω mesons at LHC energies was presented for $2.2 \leq p_{\text{T}} \text{ (GeV/c)} < 20$. Agreeing with unity over the full p_{T} range, the data shows no modification due to cold nuclear matter. Furthermore, the data agrees with measurements of π^0 at the same collision energy, as well as with a measurement of ω mesons in d–Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics In-

stitute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N36] and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (Finep), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Universidade Federal do Rio Grande do Sul (UFRGS), Brazil; Bulgarian Ministry of Education and Science, within the National Roadmap for Research Infrastructures 2020-2027 (object CERN), Bulgaria; Ministry of Education of China (MOEC), Ministry of Science & Technology of China (MSTC) and National Natural Science Foundation of China (NSFC), China; Ministry of Science and Education and Croatian Science Foundation, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research | Natural Sciences, the VILLUM FONDEN and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l’Energie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS), France; Bundesministerium für Bildung und Forschung (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; General Secretariat for Research and Technology, Ministry of Education, Research and Religions, Greece; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; National Research and Innovation Agency - BRIN, Indonesia; Istituto Nazionale di Fisica Nucleare (INFN), Italy; Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Académico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Pontificia Universidad Católica del Perú, Peru; Ministry of Science and Higher Education, National Science Centre and WUT ID-UB, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics, Ministry of Research and Innovation and Institute of Atomic Physics and Universitatea Nationala de Stiinta si Tehnologie Politehnica Bucuresti, Romania; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; Suranaree University of Technology (SUT), National Science and Technology Development Agency (NSTDA) and National Science, Research and Innovation Fund (NSRF via PMU-B B05F650021), Thailand; Turkish Energy, Nuclear and Mineral Research Agency (TENMAK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), United States of America. In addition, individual groups or members have received support from: Czech Science Foundation (grant no. 23-07499S), Czech Republic; FORTE project, reg. no. CZ.02.01.01/00/22_008/0004632, Czech Republic, co-funded by the European Union, Czech Republic; European Research Council (grant no. 950692), European Union; Deutsche Forschungs Gemeinschaft (DFG, German Research Foundation) “Neutrinos and Dark Matter in Astro- and Particle Physics” (grant no. SFB 1258), Germany; ICSC - National Research Center for High Performance Computing, Big Data and Quantum Computing and FAIR - Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy).

References

- [1] M. R. M. Jacob and K. Johnsen, “A review of accelerator and particle physics at the CERN Intersecting Storage Rings”, *CERN Yellow Report: CERN-84-13* (1984) .
- [2] ALICE Collaboration, S. Acharya *et al.*, “The ALICE experiment: a journey through QCD”, *Eur. Phys. J. C* **84** (2024) 813.
- [3] nCTEQ Collaboration, P. Duwentäster, L. A. Husová, T. Ježo, M. Klasen, K. Kovařík, A. Kusina, K. F. Muzakka, F. I. Olness, I. Schienbein, and J. Y. Yu, “Impact of inclusive hadron production data on nuclear gluon pdfs”, *Phys. Rev. D* **104** (2021) 094005.
- [4] V. Bertone, S. Carrazza, N. P. Hartland, E. R. Nocera, and J. Rojo, “A determination of the fragmentation functions of pions, kaons, and protons with faithful uncertainties”, *Eur. Phys. J. C* **77** (2017) 516.
- [5] H. Saveetha and D. Indumathi, “Fragmentation of ω and ϕ mesons in e^+e^- and pp collisions at NLO”, *Int. J. Mod. Phys. A* **32** (2017) 1750199.
- [6] ALICE Collaboration, S. Acharya *et al.*, “Neutral pion and η meson production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Eur. Phys. J. C* **78** (2018) 624.
- [7] ATLAS Collaboration, G. Aad *et al.*, “Transverse momentum, rapidity, and centrality dependence of inclusive charged-particle production in $\sqrt{s_{NN}} = 5.02$ TeV p+Pb collisions measured by the ATLAS experiment”, *Phys. Lett. B* **763** (2016) 313–336.
- [8] ALICE Collaboration, B. Abelev *et al.*, “Transverse momentum distribution and nuclear modification factor of charged particles in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Phys. Rev. Lett.* **110** (2013) 082302.
- [9] ALICE Collaboration, S. Acharya *et al.*, “Nuclear modification factor of light neutral-meson spectra up to high transverse momentum in p–Pb collisions at $\sqrt{s_{NN}} = 8.16$ TeV”, *Phys. Lett. B* **827** (2022) 136943.
- [10] PHENIX Collaboration, S. S. Adler *et al.*, “Centrality dependence of π^0 and η production at large transverse momentum in $\sqrt{s_{NN}} = 200$ GeV d + Au collisions”, *Phys. Rev. Lett.* **98** (2007) 172302.
- [11] STAR Collaboration, J. Adams *et al.*, “Identified hadron spectra at large transverse momentum in p+p and d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV”, *Phys. Lett. B* **637** (2006) 161–169.
- [12] CMS Collaboration, S. Chatrchyan *et al.*, “Study of the production of charged pions, kaons, and protons in p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Eur. Phys. J. C* **74** (2014) .
- [13] ALICE Collaboration, B. B. Abelev *et al.*, “Multiplicity Dependence of Pion, Kaon, Proton and Lambda Production in p-Pb Collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Phys. Lett. B* **728** (2014) 25–38.
- [14] ALICE Collaboration, J. Adam *et al.*, “Multiplicity dependence of charged pion, kaon, and (anti)proton production at large transverse momentum in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Phys. Lett. B* **760** (2016) 720–735.
- [15] EMC Collaboration, J. Aubert *et al.*, “The ratio of the nucleon structure functions f_2^n for iron and deuterium”, *Phys. Lett. B* **123** (1983) 275–278.
- [16] EMC Collaboration, M. Arneodo *et al.*, “Shadowing in deep inelastic muon scattering from nuclear targets”, *Phys. Lett. B* **211** (1988) 493–499.

- [17] M. Klasen and H. Paukkunen, “Nuclear Parton Distribution Functions After the First Decade of LHC Data”, *Annu. Rev. Nucl. Part. Sci.* **74** (2024) 49–87.
- [18] F. Gelis, E. Iancu, J. Jalilian-Marian, and R. Venugopalan, “The Color Glass Condensate”, *Annu. Rev. Nucl. Part. Sci.* **60** (2010) 463–489.
- [19] F. Arleo, F. Cougoulic, and S. Peigné, “Fully coherent energy loss effects on light hadron production in pA collisions.”, *J. High Energ. Phys.* **2020** (2020) .
- [20] **Particle Data Group** Collaboration, S. Navas *et al.*, “The Review of Particle Physics”, *Phys. Rev. D* **110** (2024) 030001.
- [21] **ALICE** Collaboration, K. Aamodt *et al.*, “The ALICE experiment at the CERN LHC”, *JINST* **3** (2008) S08002.
- [22] **ALICE** Collaboration, B. B. Abelev *et al.*, “Performance of the ALICE Experiment at the CERN LHC”, *Int. J. Mod. Phys. A* **29** (2014) 1430044.
- [23] **ALICE** Collaboration, A. Andronic *et al.*, “ALICE Inner Tracking System (ITS): Technical Design Report”, *CERN-LHCC-99-012* ; *ALICE-TDR-4* (1999) .
- [24] **ALICE** Collaboration, K. Aamodt *et al.*, “Alignment of the ALICE Inner Tracking System with cosmic-ray tracks”, *JINST* **5** (2010) P03003.
- [25] **ALICE** Collaboration, G. Dellacasa *et al.*, “ALICE time projection chamber: Technical Design Report”, *CERN-LHCC-2000-001* ; *CERN-OPEN-2000-183* ; *ALICE-TDR-7* (2000) .
- [26] J. Alme *et al.*, “The ALICE TPC, a large 3-dimensional tracking device with fast readout for ultra-high multiplicity events”, *Nucl. Instrum. Methods Phys. Res. A* **622** (2010) 316–367.
- [27] **ALICE** Collaboration, E. Abbas *et al.*, “Performance of the ALICE VZERO system”, *JINST* **8** (2013) P10016.
- [28] M. Bondila *et al.*, “ALICE T0 detector”, *IEEE Trans. Nucl. Sci.* **52** (2005) 1705–1711.
- [29] **ALICE** Collaboration, P. Cortese *et al.*, “ALICE Time-Of-Flight system (TOF): Technical Design Report”, *CERN-LHCC-2000-012* ; *ALICE-TDR-8* (2000) .
- [30] **ALICE** Collaboration, P. Cortese, G. Dellacasa, *et al.*, “ALICE Electromagnetic Calorimeter Technical Design Report”, *CERN-LHCC-2008-014*, *ALICE-TDR-14* (2008) .
- [31] **ALICE** Collaboration, S. Acharya *et al.*, “Performance of the ALICE Electromagnetic Calorimeter”, *JINST* **18** (2023) P08007.
- [32] S. van der Meer, “Calibration of the effective beam height in the ISR”, *CERN-ISR-PO-68-31*, *ISR-PO-68-31* (1968) .
- [33] **ALICE** Collaboration, S. Acharya *et al.*, “ALICE 2017 luminosity determination for pp collisions at $\sqrt{s} = 5$ TeV”, *ALICE-PUBLIC-2018-014* (2018) .
- [34] **ALICE** Collaboration, B. B. Abelev *et al.*, “Measurement of visible cross sections in proton-lead collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV in van der Meer scans with the ALICE detector”, *JINST* **9** (2014) P11003.
- [35] L. Evans and P. Bryant, “LHC machine”, *JINST* **3** (2008) S08001.

- [36] ALICE Collaboration, S. Acharya *et al.*, “ π^0 and η meson production in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *Eur. Phys. J. C* **78** (2018) 263.
- [37] ALICE Collaboration, S. Acharya *et al.*, “Data-driven precision determination of the material budget in ALICE”, *JINST* **18** (2023) P11032.
- [38] ALICE Collaboration, S. Acharya *et al.*, “Light neutral-meson production in pp collisions at $\sqrt{s} = 13$ TeV”, arXiv:2411.09560 [hep-ex].
- [39] ALICE Collaboration, S. Acharya *et al.*, “Production of ω mesons in pp collisions at $\sqrt{s} = 7$ TeV”, *Eur. Phys. J. C* **80** (2020) 1130.
- [40] ALICE Collaboration, S. Acharya *et al.*, “Measurement of ω meson production in pp collisions at $\sqrt{s} = 13$ TeV”, arXiv:2411.09432 [hep-ex].
- [41] ALICE Collaboration, S. Acharya *et al.*, “Supplemental figures: “Measurement of ω meson production in pp and p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV””, ALICE-PUBLIC-2025-003.
- [42] ALICE Collaboration, S. Acharya *et al.*, “Multiplicity dependence of charged-particle production in pp, p–Pb, Xe–Xe and Pb–Pb collisions at the LHC”, *Phys. Lett. B* **845** (2023) 138110.
- [43] C. Bierlich *et al.*, “A comprehensive guide to the physics and usage of PYTHIA 8.3”, *LU-TP 22-16; MCNET-22-04* (2022), arXiv:2203.11601 [hep-ph].
- [44] S. Roesler, R. Engel, and J. Ranft, “The Monte Carlo Event Generator DPMJET-III”, in *Advanced Monte Carlo for Radiation Physics, Particle Transport Simulation and Applications*, pp. 1033–1038. Springer Berlin Heidelberg, 2001.
- [45] R. Brun, F. Bruyant, M. Maire, A. C. McPherson, and P. Zancarini, “GEANT 3: user’s guide Geant 3.10, Geant 3.11; rev. version”, *CERN-DD-EE-84-01* (1987).
- [46] L. Lista, “Combination of measurements and the BLUE method”, *EPJ Web Conf.* **137** (2017) 11006.
- [47] G. Lafferty and T. Wyatt, “Where to stick your data points: The treatment of measurements within wide bins”, *Nucl. Instrum. Methods Phys. Res. A* **355** (1995) 541–547.
- [48] T. Pierog, I. Karpenko, J. M. Katzy, E. Yatsenko, and K. Werner, “EPOS LHC: Test of collective hadronization with data measured at the CERN large hadron collider”, *Phys. Rev. C* **92** (2015) 034906.
- [49] S. Porteboeuf, T. Pierog, and K. Werner, “Producing Hard Processes Regarding the Complete Event: The EPOS Event Generator”, in *45th Rencontres de Moriond on QCD and High Energy Interactions*, pp. 135–140. Gioi Publishers, 2010. arXiv:1006.2967 [hep-ph].
- [50] ALICE Collaboration, S. Acharya *et al.*, “Production of charged pions, kaons, and (anti-)protons in Pb–Pb and inelastic pp collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Phys. Rev. C* **101** (2020) 044907.
- [51] L. Altenkämper, F. Bock, C. Loizides, and N. Schmidt, “Applicability of transverse mass scaling in hadronic collisions at energies available at the CERN Large Hadron Collider”, *Phys. Rev. C* **96** (2017).
- [52] ALICE Collaboration, J. Adam *et al.*, “Direct photon production in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, *Phys. Lett. B* **754** (2016) 235–248.
- [53] ALICE Collaboration, S. Acharya *et al.*, “Dielectron and heavy-quark production in inelastic and high-multiplicity proton–proton collisions at $\sqrt{s_{NN}} = 13$ TeV”, *Phys. Lett. B* **788** (2019) 505–518.

- [54] M. Diakonou, C. Kourkouvelis, *et al.*, “Inclusive high- p_T ω^0 and η' production at the ISR”, *Phys. Lett. B* **89** (1980) 432–436.
- [55] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of neutral mesons in $p + p$ collisions at $\sqrt{s} = 200$ GeV and scaling properties of hadron production”, *Phys. Rev. D* **83** (2011) 052004.
- [56] **PHENIX** Collaboration, A. Adare *et al.*, “Production of ω mesons in $p + p$, $d + Au$, $Cu + Cu$, and $Au + Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV”, *Phys. Rev. C* **84** (2011) 044902, [arXiv:1105.3467](https://arxiv.org/abs/1105.3467) [nucl-ex].

A The ALICE Collaboration

S. Acharya⁵⁰, A. Agarwal¹³³, G. Aglieri Rinella³², L. Aglietta²⁴, M. Agnello²⁹, N. Agrawal²⁵, Z. Ahammed¹³³, S. Ahmad¹⁵, S.U. Ahn⁷¹, I. Ahuja³⁶, A. Akindinov¹³⁹, V. Akishina³⁸, M. Al-Turany⁹⁶, D. Aleksandrov¹³⁹, B. Alessandro⁵⁶, H.M. Alfanda⁶, R. Alfaro Molina⁶⁷, B. Ali¹⁵, A. Alici²⁵, N. Alizadehvandchali¹¹⁴, A. Alkin¹⁰³, J. Alme²⁰, G. Alocco²⁴, T. Alt⁶⁴, A.R. Altamura⁵⁰, I. Altsybeev⁹⁴, J.R. Alvarado⁴⁴, M.N. Anaam⁶, C. Andrei⁴⁵, N. Andreou¹¹³, A. Andronic¹²⁴, E. Andronov¹³⁹, V. Anguelov⁹³, F. Antinori⁵⁴, P. Antonioli⁵¹, N. Apadula⁷³, H. Appelshäuser⁶⁴, C. Arata⁷², S. Arcelli²⁵, R. Arnaldi⁵⁶, J.G.M.C.A. Arneiro¹⁰⁹, I.C. Arsene¹⁹, M. Arslanok¹³⁶, A. Augustinus³², R. Averbeck⁹⁶, D. Averyanov¹³⁹, M.D. Azmi¹⁵, H. Baba¹²², A. Badalà⁵³, J. Bae¹⁰³, Y. Bae¹⁰³, Y.W. Baek⁴⁰, X. Bai¹¹⁸, R. Bailhache⁶⁴, Y. Bailung⁴⁸, R. Bala⁹⁰, A. Baldisseri¹²⁸, B. Balis², S. Bangalia¹¹⁶, Z. Banoo⁹⁰, V. Barbasova³⁶, F. Barile³¹, L. Barioglio⁵⁶, M. Barlou⁷⁷, B. Barman⁴¹, G.G. Barnaföldi⁴⁶, L.S. Barnby¹¹³, E. Barreau¹⁰², V. Barret¹²⁵, L. Barreto¹⁰⁹, K. Barth³², E. Bartsch⁶⁴, N. Bastid¹²⁵, S. Basu⁷⁴, G. Batigne¹⁰², D. Battistini⁹⁴, B. Batyunya¹⁴⁰, D. Bauri⁴⁷, J.L. Bazo Alba¹⁰⁰, I.G. Bearden⁸², P. Becht⁹⁶, D. Behera⁴⁸, I. Belikov¹²⁷, A.D.C. Bell Hechavarría¹²⁴, F. Bellini²⁵, R. Bellwied¹¹⁴, S. Belokurova¹³⁹, L.G.E. Beltran¹⁰⁸, Y.A.V. Beltran⁴⁴, G. Bencedi⁴⁶, A. Bensaoula¹¹⁴, S. Beole²⁴, Y. Berdnikov¹³⁹, A. Berdnikova⁹³, L. Bergmann⁹³, L. Bernardinis²³, L. Betev³², P.P. Bhaduri¹³³, A. Bhasin⁹⁰, B. Bhattacharjee⁴¹, S. Bhattarai¹¹⁶, L. Bianchi²⁴, J. Bielčik³⁴, J. Bielčíková⁸⁵, A.P. Bigot¹²⁷, A. Bilandzic⁹⁴, A. Binoy¹¹⁶, G. Biro⁴⁶, S. Biswas⁴, N. Bize¹⁰², J.T. Blair¹⁰⁷, D. Blau¹³⁹, M.B. Blidaru⁹⁶, N. Bluhme³⁸, C. Blume⁶⁴, F. Bock⁸⁶, T. Bodova²⁰, J. Bok¹⁶, L. Boldizsár⁴⁶, M. Bombara³⁶, P.M. Bond³², G. Bonomi^{132,55}, H. Borel¹²⁸, A. Borissov¹³⁹, A.G. Borquez Carcamo⁹³, E. Botta²⁴, Y.E.M. Bouziani⁶⁴, D.C. Brandibur⁶³, L. Bratrud⁶⁴, P. Braun-Munzinger⁹⁶, M. Bregant¹⁰⁹, M. Broz³⁴, G.E. Bruno^{95,31}, V.D. Buchakchiev³⁵, M.D. Buckland⁸⁴, D. Budnikov¹³⁹, H. Buesching⁶⁴, S. Bufalino²⁹, P. Buhler¹⁰¹, N. Burmasov¹³⁹, Z. Buthelezi^{68,121}, A. Bylinkin²⁰, S.A. Bysiak¹⁰⁶, J.C. Cabanillas Noris¹⁰⁸, M.F.T. Cabrera¹¹⁴, H. Caines¹³⁶, A. Caliva²⁸, E. Calvo Villar¹⁰⁰, J.M.M. Camacho¹⁰⁸, P. Camerini²³, M.T. Camerlingo⁵⁰, F.D.M. Canedo¹⁰⁹, S. Cannito²³, S.L. Cantway¹³⁶, M. Carabas¹¹², F. Carnesecchi³², L.A.D. Carvalho¹⁰⁹, J. Castillo Castellanos¹²⁸, M. Castoldi³², F. Catalano³², S. Cattaruzzi²³, R. Cerri²⁴, I. Chakaberia⁷³, P. Chakraborty¹³⁴, S. Chandra¹³³, S. Chapeland³², M. Chartier¹¹⁷, S. Chattopadhyay¹³³, M. Chen³⁹, T. Cheng⁶, C. Cheshkov¹²⁶, D. Chiappara²⁷, V. Chibante Barroso³², D.D. Chinellato¹⁰¹, F. Chinu²⁴, E.S. Chizzali^{11,94}, J. Cho⁵⁸, S. Cho⁵⁸, P. Chochula³², Z.A. Chochulska¹³⁴, D. Choudhury⁴¹, S. Choudhury⁹⁸, P. Christakoglou⁸³, C.H. Christensen⁸², P. Christiansen⁷⁴, T. Chujo¹²³, M. Ciacco²⁹, C. Cicalo⁵², G. Cimdador²⁴, F. Cindolo⁵¹, M.R. Ciupek⁹⁶, G. Clai^{III,51}, F. Colamaria⁵⁰, J.S. Colburn⁹⁹, D. Colella³¹, A. Colelli³¹, M. Colocci²⁵, M. Concas³², G. Conesa Balbastre⁷², Z. Conesa del Valle¹²⁹, G. Contin²³, J.G. Contreras³⁴, M.L. Coquet¹⁰², P. Cortese^{131,56}, M.R. Cosentino¹¹¹, F. Costa³², S. Costanza²¹, P. Crochet¹²⁵, M.M. Czarnynoga¹³⁴, A. Dainese⁵⁴, G. Dange³⁸, M.C. Danisch⁹³, A. Danu⁶³, P. Das^{32,79}, S. Das⁴, A.R. Dash¹²⁴, S. Dash⁴⁷, A. De Caro²⁸, G. de Cataldo⁵⁰, J. de Cuveland³⁸, A. De Falco²², D. De Gruttola²⁸, N. De Marco⁵⁶, C. De Martin²³, S. De Pasquale²⁸, R. Deb¹³², R. Del Grande⁹⁴, L. Dello Stritto³², K.C. Devereaux¹⁸, G.G.A. de Souza¹⁰⁹, P. Dhankher¹⁸, D. Di Bari³¹, M. Di Costanzo²⁹, A. Di Mauro³², B. Di Ruzza¹³⁰, B. Diab¹²⁸, R.A. Diaz^{140,7}, Y. Ding⁶, J. Ditzel⁶⁴, R. Divià³², Ø. Djuvsland²⁰, U. Dmitrieva¹³⁹, A. Dobrin⁶³, B. Dönigus⁶⁴, J.M. Dubinski¹³⁴, A. Dubla⁹⁶, P. Dupieux¹²⁵, N. Dzalaiova¹³, T.M. Eder¹²⁴, R.J. Ehlers⁷³, F. Eisenhut⁶⁴, R. Ejima⁹¹, D. Elia⁵⁰, B. Erasmus¹⁰², F. Ercolessi²⁵, B. Espagnon¹²⁹, G. Eulisse³², D. Evans⁹⁹, S. Evdokimov¹³⁹, L. Fabbietti⁹⁴, M. Faggin³², J. Faivre⁷², F. Fan⁶, W. Fan⁷³, A. Fantoni⁴⁹, M. Fasel⁸⁶, G. Feofilov¹³⁹, A. Fernández Téllez⁴⁴, L. Ferrandi¹⁰⁹, M.B. Ferrer³², A. Ferrero¹²⁸, C. Ferrero^{IV,56}, A. Ferretti²⁴, V.J.G. Feuillard⁹³, V. Filova³⁴, D. Finogeev¹³⁹, F.M. Fionda⁵², F. Flor¹³⁶, A.N. Flores¹⁰⁷, S. Foertsch⁶⁸, I. Fokin⁹³, S. Fokin¹³⁹, U. Follo^{IV,56}, E. Fragiaco⁵⁷, E. Frajna⁴⁶, H. Fribert⁹⁴, U. Fuchs³², N. Funicello²⁸, C. Furget⁷², A. Furs¹³⁹, T. Fusayasu⁹⁷, J.J. Gaardhøje⁸², M. Gagliardi²⁴, A.M. Gago¹⁰⁰, T. Gahlaut⁴⁷, C.D. Galvan¹⁰⁸, S. Gami⁷⁹, D.R. Gangadharan¹¹⁴, P. Ganoti⁷⁷, C. Garabatos⁹⁶, J.M. Garcia⁴⁴, T. García Chávez⁴⁴, E. García-Solis⁹, S. Garetti¹²⁹, C. Gargiulo³², P. Gasik⁹⁶, H.M. Gaur³⁸, A. Gautam¹¹⁶, M.B. Gay Ducati⁶⁶, M. Germain¹⁰², R.A. Gernhaeuser⁹⁴, C. Ghosh¹³³, M. Giacalone⁵¹, G. Gioachin²⁹, S.K. Giri¹³³, P. Giubellino^{96,56}, P. Giubilato²⁷, A.M.C. Glaenzer¹²⁸, P. Glässel⁹³, E. Glimos¹²⁰, D.J.Q. Goh⁷⁵, V. Gonzalez¹³⁵, P. Gordeev¹³⁹, M. Gorgon², K. Goswami⁴⁸, S. Gotovac³³, V. Grabski⁶⁷, L.K. Graczykowski¹³⁴, E. Grecka⁸⁵,

A. Grelli ⁵⁹, C. Grigoras ³², V. Grigoriev ¹³⁹, S. Grigoryan ^{140,1}, O.S. Groettvik ³², F. Grosa ³², J.F. Grosse-Oetringhaus ³², R. Grosso ⁹⁶, D. Grund ³⁴, N.A. Grunwald⁹³, R. Guernane ⁷², M. Guilbaud ¹⁰², K. Gulbrandsen ⁸², J.K. Gumprecht ¹⁰¹, T. Gündem ⁶⁴, T. Gunji ¹²², J. Guo¹⁰, W. Guo ⁶, A. Gupta ⁹⁰, R. Gupta ⁹⁰, R. Gupta ⁴⁸, K. Gwizdziel ¹³⁴, L. Gyulai ⁴⁶, C. Hadjidakis ¹²⁹, F.U. Haider ⁹⁰, S. Haidlova ³⁴, M. Haldar⁴, H. Hamagaki ⁷⁵, Y. Han ¹³⁸, B.G. Hanley ¹³⁵, R. Hannigan ¹⁰⁷, J. Hansen ⁷⁴, J.W. Harris ¹³⁶, A. Harton ⁹, M.V. Hartung ⁶⁴, H. Hassan ¹¹⁵, D. Hatzifotiadou ⁵¹, P. Hauer ⁴², L.B. Havener ¹³⁶, E. Hellbär ³², H. Helstrup ³⁷, M. Hemmer ⁶⁴, T. Herman ³⁴, S.G. Hernandez¹¹⁴, G. Herrera Corral ⁸, S. Herrmann ¹²⁶, K.F. Hetland ³⁷, B. Heybeck ⁶⁴, H. Hillemanns ³², B. Hippolyte ¹²⁷, I.P.M. Hobus ⁸³, F.W. Hoffmann ⁷⁰, B. Hofman ⁵⁹, M. Horst ⁹⁴, A. Horzyk ², Y. Hou ⁶, P. Hristov ³², P. Huhn⁶⁴, L.M. Huhta ¹¹⁵, T.J. Humanic ⁸⁷, A. Hutson ¹¹⁴, D. Hutter ³⁸, M.C. Hwang ¹⁸, R. Ilkaev¹³⁹, M. Inaba ¹²³, M. Ippolitov ¹³⁹, A. Isakov ⁸³, T. Isidori ¹¹⁶, M.S. Islam ^{47,98}, S. Iurchenko ¹³⁹, M. Ivanov ⁹⁶, M. Ivanov¹³, V. Ivanov ¹³⁹, K.E. Iversen ⁷⁴, M. Jablonski ², B. Jacak ^{18,73}, N. Jacazio ²⁵, P.M. Jacobs ⁷³, S. Jadlovská¹⁰⁵, J. Jadlovsky¹⁰⁵, S. Jaelani ⁸¹, C. Jahnke ¹¹⁰, M.J. Jakubowska ¹³⁴, M.A. Janik ¹³⁴, S. Ji ¹⁶, S. Jia ¹⁰, T. Jiang ¹⁰, A.A.P. Jimenez ⁶⁵, F. Jonas ⁷³, D.M. Jones ¹¹⁷, J.M. Jowett ^{32,96}, J. Jung ⁶⁴, M. Jung ⁶⁴, A. Junique ³², A. Jusko ⁹⁹, J. Kaewjai¹⁰⁴, P. Kalinak ⁶⁰, A. Kalweit ³², A. Karasu Uysal ¹³⁷, D. Karatovic ⁸⁸, N. Karatzenis⁹⁹, O. Karavichev ¹³⁹, T. Karavicheva ¹³⁹, E. Karpechev ¹³⁹, M.J. Karwowska ¹³⁴, U. Keschull ⁷⁰, M. Keil ³², B. Ketzer ⁴², J. Keul ⁶⁴, S.S. Khade ⁴⁸, A.M. Khan ¹¹⁸, S. Khan ¹⁵, A. Khanzadeev ¹³⁹, Y. Kharlov ¹³⁹, A. Khatun ¹¹⁶, A. Khuntia ³⁴, Z. Khuranova ⁶⁴, B. Kileng ³⁷, B. Kim ¹⁰³, C. Kim ¹⁶, D.J. Kim ¹¹⁵, D. Kim ¹⁰³, E.J. Kim ⁶⁹, J. Kim ¹³⁸, J. Kim ⁵⁸, J. Kim ^{32,69}, M. Kim ¹⁸, S. Kim ¹⁷, T. Kim ¹³⁸, K. Kimura ⁹¹, S. Kirsch ⁶⁴, I. Kisel ³⁸, S. Kiselev ¹³⁹, A. Kisiel ¹³⁴, J.L. Klay ⁵, J. Klein ³², S. Klein ⁷³, C. Klein-Bösing ¹²⁴, M. Kleiner ⁶⁴, T. Klemenz ⁹⁴, A. Kluge ³², C. Kobdaj ¹⁰⁴, R. Kohara ¹²², T. Kollegger⁹⁶, A. Kondratyev ¹⁴⁰, N. Kondratyeva ¹³⁹, J. König ⁶⁴, S.A. Königstorfer ⁹⁴, P.J. Konopka ³², G. Kornakov ¹³⁴, M. Korwieser ⁹⁴, S.D. Koryciak ², C. Koster ⁸³, A. Kotliarov ⁸⁵, N. Kovacic ⁸⁸, V. Kovalenko ¹³⁹, M. Kowalski ¹⁰⁶, V. Kozuharov ³⁵, G. Kozlov³⁸, I. Králik ⁶⁰, A. Kravčáková ³⁶, L. Krcal ³², M. Krivda ^{99,60}, F. Krizek ⁸⁵, K. Krizkova Gajdosova ³⁴, C. Krug ⁶⁶, M. Krüger ⁶⁴, D.M. Krupova ³⁴, E. Kryshen ¹³⁹, V. Kučera ⁵⁸, C. Kuhn ¹²⁷, P.G. Kuijer ⁸³, T. Kumaoka¹²³, D. Kumar¹³³, L. Kumar ⁸⁹, N. Kumar⁸⁹, S. Kumar ⁵⁰, S. Kundu ³², M. Kuo¹²³, P. Kurashvili ⁷⁸, A.B. Kurepin ¹³⁹, A. Kuryakin ¹³⁹, S. Kushpil ⁸⁵, V. Kuskov ¹³⁹, M. Kutyla¹³⁴, A. Kuznetsov ¹⁴⁰, M.J. Kweon ⁵⁸, Y. Kwon ¹³⁸, S.L. La Pointe ³⁸, P. La Rocca ²⁶, A. Lakrathok¹⁰⁴, M. Lamanna ³², S. Lambert¹⁰², A.R. Landou ⁷², R. Langoy ¹¹⁹, P. Larionov ³², E. Laudi ³², L. Lautner ⁹⁴, R.A.N. Laveaga¹⁰⁸, R. Lavicka ¹⁰¹, R. Lea ^{132,55}, H. Lee ¹⁰³, I. Legrand ⁴⁵, G. Legras ¹²⁴, A.M. Lejeune ³⁴, T.M. Lelek ², R.C. Lemmon ^{1,84}, I. León Monzón ¹⁰⁸, M.M. Lesch ⁹⁴, P. Lévai ⁴⁶, M. Li⁶, P. Li¹⁰, X. Li¹⁰, B.E. Liang-gilman ¹⁸, J. Lien ¹¹⁹, R. Lietava ⁹⁹, I. Likmeta ¹¹⁴, B. Lim ²⁴, H. Lim ¹⁶, S.H. Lim ¹⁶, S. Lin¹⁰, V. Lindenstruth ³⁸, C. Lippmann ⁹⁶, D. Liskova ¹⁰⁵, D.H. Liu ⁶, J. Liu ¹¹⁷, G.S.S. Liveraro ¹¹⁰, I.M. Lofnes ²⁰, C. Loizides ⁸⁶, S. Lokos ¹⁰⁶, J. Lömker ⁵⁹, X. Lopez ¹²⁵, E. López Torres ⁷, C. Lotteau¹²⁶, P. Lu ^{96,118}, W. Lu ⁶, Z. Lu ¹⁰, F.V. Lugo ⁶⁷, J. Luo³⁹, G. Luparello ⁵⁷, Y.G. Ma ³⁹, M. Mager ³², A. Maire ¹²⁷, E.M. Majerz ², M.V. Makariev ³⁵, M. Malaev ¹³⁹, G. Malfattore ^{51,25}, N.M. Malik ⁹⁰, N. Malik ¹⁵, S.K. Malik ⁹⁰, D. Mallick ¹²⁹, N. Mallick ^{115,48}, G. Mandaglio ^{30,53}, S.K. Mandal ⁷⁸, A. Manea ⁶³, V. Manko ¹³⁹, A.K. Manna⁴⁸, F. Manso ¹²⁵, G. Mantzaridis ⁹⁴, V. Manzari ⁵⁰, Y. Mao ⁶, R.W. Marcjan ², G.V. Margagliotti ²³, A. Margotti ⁵¹, A. Marín ⁹⁶, C. Markert ¹⁰⁷, P. Martinengo ³², M.I. Martínez ⁴⁴, G. Martínez García ¹⁰², M.P.P. Martins ^{32,109}, S. Masciocchi ⁹⁶, M. Maserà ²⁴, A. Masoni ⁵², L. Massacrier ¹²⁹, O. Massen ⁵⁹, A. Mastroserio ^{130,50}, L. Mattei ^{24,125}, S. Mattiazzo ²⁷, A. Matyja ¹⁰⁶, F. Mazzaschi ^{32,24}, M. Mazzilli ¹¹⁴, Y. Melikyan ⁴³, M. Melo ¹⁰⁹, A. Menchaca-Rocha ⁶⁷, J.E.M. Mendez ⁶⁵, E. Meninno ¹⁰¹, A.S. Menon ¹¹⁴, M.W. Menzel^{32,93}, M. Meres ¹³, L. Micheletti ³², D. Mihai¹¹², D.L. Mihaylov ⁹⁴, A.U. Mikalsen ²⁰, K. Mikhaylov ^{140,139}, N. Minafra ¹¹⁶, D. Miśkowiec ⁹⁶, A. Modak ^{57,132}, B. Mohanty ⁷⁹, M. Mohisin Khan ^{V,15}, M.A. Molander ⁴³, M.M. Mondal ⁷⁹, S. Monira ¹³⁴, C. Mordasini ¹¹⁵, D.A. Moreira De Godoy ¹²⁴, I. Morozov ¹³⁹, A. Morsch ³², T. Mrnjavac ³², V. Muccifora ⁴⁹, S. Muhuri ¹³³, A. Mulliri ²², M.G. Munhoz ¹⁰⁹, R.H. Munzer ⁶⁴, H. Murakami ¹²², L. Musa ³², J. Musinsky ⁶⁰, J.W. Myrcha ¹³⁴, N.B. Sundstrom⁵⁹, B. Naik ¹²¹, A.I. Nambrath ¹⁸, B.K. Nandi ⁴⁷, R. Nania ⁵¹, E. Nappi ⁵⁰, A.F. Nassirpour ¹⁷, V. Nastase¹¹², A. Nath ⁹³, N.F. Nathanson⁸², C. Nattrass ¹²⁰, K. Naumov¹⁸, M.N. Naydenov ³⁵, A. Neagu¹⁹, L. Nellen ⁶⁵, R. Nepeivoda ⁷⁴, S. Nese ¹⁹, N. Nicassio ³¹, B.S. Nielsen ⁸², E.G. Nielsen ⁸², S. Nikolaev ¹³⁹, V. Nikulin ¹³⁹, F. Noferini ⁵¹, S. Noh ¹², P. Nomokonov ¹⁴⁰, J. Norman ¹¹⁷,

N. Novitzky ⁸⁶, A. Nyanin ¹³⁹, J. Nystrand ²⁰, M.R. Ockleton ¹¹⁷, M. Ogino ⁷⁵, S. Oh ¹⁷, A. Ohlson ⁷⁴, V.A. Okorokov ¹³⁹, J. Oleniacz ¹³⁴, A. Onnerstad ¹¹⁵, C. Oppedisano ⁵⁶, A. Ortiz Velasquez ⁶⁵, J. Otwinowski ¹⁰⁶, M. Oya ⁹¹, K. Oyama ⁷⁵, S. Padhan ⁴⁷, D. Pagano ^{132,55}, G. Paic ⁶⁵, S. Paisano-Guzmán ⁴⁴, A. Palasciano ⁵⁰, I. Panasenko ⁷⁴, S. Panebianco ¹²⁸, P. Panigrahi ⁴⁷, C. Pantouvakis ²⁷, H. Park ¹²³, J. Park ¹²³, S. Park ¹⁰³, J.E. Parkkila ³², Y. Patley ⁴⁷, R.N. Patra ⁵⁰, P. Paudel ¹¹⁶, B. Paul ¹³³, H. Pei ⁶, T. Peitzmann ⁵⁹, X. Peng ¹¹, M. Pennisi ²⁴, S. Perciballi ²⁴, D. Peresunko ¹³⁹, G.M. Perez ⁷, Y. Pestov ¹³⁹, M.T. Petersen ⁸², V. Petrov ¹³⁹, M. Petrovici ⁴⁵, S. Piano ⁵⁷, M. Pikna ¹³, P. Pillot ¹⁰², O. Pinazza ^{51,32}, L. Pinsky ¹¹⁴, C. Pinto ³², S. Pisano ⁴⁹, M. Płoskoń ⁷³, M. Planinic ⁸⁸, D.K. Plociennik ², M.G. Poghosyan ⁸⁶, B. Polichtchouk ¹³⁹, S. Politano ^{32,24}, N. Poljak ⁸⁸, A. Pop ⁴⁵, S. Porteboeuf-Houssais ¹²⁵, V. Pozdniakov ^{1,140}, I.Y. Pozos ⁴⁴, K.K. Pradhan ⁴⁸, S.K. Prasad ⁴, S. Prasad ⁴⁸, R. Preghenella ⁵¹, F. Prino ⁵⁶, C.A. Pruneau ¹³⁵, I. Pshenichnov ¹³⁹, M. Puccio ³², S. Pucillo ²⁴, S. Qiu ⁸³, L. Quaglia ²⁴, A.M.K. Radhakrishnan ⁴⁸, S. Ragoni ¹⁴, A. Rai ¹³⁶, A. Rakotozafindrabe ¹²⁸, N. Ramasubramanian ¹²⁶, L. Ramello ^{131,56}, C.O. Ramirez-Alvarez ⁴⁴, M. Rasa ²⁶, S.S. Räsänen ⁴³, R. Rath ⁵¹, M.P. Rauch ²⁰, I. Ravasenga ³², K.F. Read ^{86,120}, C. Reckziegel ¹¹¹, A.R. Redelbach ³⁸, K. Redlich ^{VI,78}, C.A. Reetz ⁹⁶, H.D. Regules-Medel ⁴⁴, A. Rehman ²⁰, F. Reidt ³², H.A. Reme-Ness ³⁷, K. Reygers ⁹³, A. Riabov ¹³⁹, V. Riabov ¹³⁹, R. Ricci ²⁸, M. Richter ²⁰, A.A. Riedel ⁹⁴, W. Riegler ³², A.G. Riffero ²⁴, M. Rignanese ²⁷, C. Ripoli ²⁸, C. Ristea ⁶³, M.V. Rodriguez ³², M. Rodríguez Cahuantzi ⁴⁴, S.A. Rodríguez Ramírez ⁴⁴, K. Røed ¹⁹, R. Rogalev ¹³⁹, E. Rogochaya ¹⁴⁰, T.S. Rogoschinski ⁶⁴, D. Rohr ³², D. Röhrich ²⁰, S. Rojas Torres ³⁴, P.S. Rokita ¹³⁴, G. Romanenko ²⁵, F. Ronchetti ³², D. Rosales Herrera ⁴⁴, E.D. Rosas ⁶⁵, K. Roslon ¹³⁴, A. Rossi ⁵⁴, A. Roy ⁴⁸, S. Roy ⁴⁷, N. Rubini ⁵¹, J.A. Rudolph ⁸³, D. Ruggiano ¹³⁴, R. Rui ²³, P.G. Russek ², R. Russo ⁸³, A. Rustamov ⁸⁰, E. Ryabinkin ¹³⁹, Y. Ryabov ¹³⁹, A. Rybicki ¹⁰⁶, L.C.V. Ryder ¹¹⁶, J. Ryu ¹⁶, W. Rzesza ¹³⁴, B. Sabiu ⁵¹, S. Sadhu ⁴², S. Sadovsky ¹³⁹, J. Saetre ²⁰, S. Saha ⁷⁹, B. Sahoo ⁴⁸, R. Sahoo ⁴⁸, D. Sahu ⁴⁸, P.K. Sahu ⁶¹, J. Saini ¹³³, K. Sajdakova ³⁶, S. Sakai ¹²³, S. Sambyal ⁹⁰, D. Samitz ¹⁰¹, I. Sanna ^{32,94}, T.B. Saramela ¹⁰⁹, D. Sarkar ⁸², P. Sarma ⁴¹, V. Sarritzu ²², V.M. Sarti ⁹⁴, M.H.P. Sas ³², S. Sawan ⁷⁹, E. Scapparone ⁵¹, J. Schambach ⁸⁶, H.S. Scheid ^{32,64}, C. Schiaua ⁴⁵, R. Schicker ⁹³, F. Schlepfer ^{32,93}, A. Schmah ⁹⁶, C. Schmidt ⁹⁶, M.O. Schmidt ³², M. Schmidt ⁹², N.V. Schmidt ⁸⁶, A.R. Schmier ¹²⁰, J. Schoengarth ⁶⁴, R. Schotter ¹⁰¹, A. Schröter ³⁸, J. Schukraft ³², K. Schweda ⁹⁶, G. Scioli ²⁵, E. Scomparin ⁵⁶, J.E. Seger ¹⁴, Y. Sekiguchi ¹²², D. Sekihata ¹²², M. Selina ⁸³, I. Selyuzhenkov ⁹⁶, S. Senyukov ¹²⁷, J.J. Seo ⁹³, D. Serebryakov ¹³⁹, L. Serkin ^{VII,65}, L. Šerkšnytė ⁹⁴, A. Sevcenco ⁶³, T.J. Shaba ⁶⁸, A. Shabetai ¹⁰², R. Shahoyan ³², A. Shangaraev ¹³⁹, B. Sharma ⁹⁰, D. Sharma ⁴⁷, H. Sharma ⁵⁴, M. Sharma ⁹⁰, S. Sharma ⁹⁰, U. Sharma ⁹⁰, A. Shatat ¹²⁹, O. Sheibani ^{135,114}, K. Shigaki ⁹¹, M. Shimomura ⁷⁶, S. Shirinkin ¹³⁹, Q. Shou ³⁹, Y. Sibiraki ¹³⁹, S. Siddhanta ⁵², T. Siemiarzczuk ⁷⁸, T.F. Silva ¹⁰⁹, D. Silvermyr ⁷⁴, T. Simantathammakul ¹⁰⁴, R. Simeonov ³⁵, B. Singh ⁹⁰, B. Singh ⁹⁴, K. Singh ⁴⁸, R. Singh ⁷⁹, R. Singh ^{54,96}, S. Singh ¹⁵, V.K. Singh ¹³³, V. Singhal ¹³³, T. Sinha ⁹⁸, B. Sitar ¹³, M. Sitta ^{131,56}, T.B. Skaali ¹⁹, G. Skorodumovs ⁹³, N. Smirnov ¹³⁶, R.J.M. Snellings ⁵⁹, E.H. Solheim ¹⁹, C. Sonnabend ^{32,96}, J.M. Sonneveld ⁸³, F. Soramel ²⁷, A.B. Soto-herandez ⁸⁷, R. Spijkers ⁸³, I. Sputowska ¹⁰⁶, J. Staa ⁷⁴, J. Stachel ⁹³, I. Stan ⁶³, P.J. Steffanic ¹²⁰, T. Stellhorn ¹²⁴, S.F. Stiefelmaier ⁹³, D. Stocco ¹⁰², I. Storehaug ¹⁹, N.J. Strangmann ⁶⁴, P. Stratmann ¹²⁴, S. Strazzi ²⁵, A. Sturniolo ^{30,53}, C.P. Stylianidis ⁸³, A.A.P. Suaide ¹⁰⁹, C. Suire ¹²⁹, A. Suiu ^{32,112}, M. Sukhanov ¹³⁹, M. Suljic ³², R. Sultanov ¹³⁹, V. Sumberia ⁹⁰, S. Sumowidagdo ⁸¹, L.H. Tabares ⁷, S.F. Taghavi ⁹⁴, J. Takahashi ¹¹⁰, G.J. Tambave ⁷⁹, S. Tang ⁶, Z. Tang ¹¹⁸, J.D. Tapia Takaki ¹¹⁶, N. Tapus ¹¹², L.A. Tarasovicova ³⁶, M.G. Tarzila ⁴⁵, A. Tauro ³², A. Tavira García ¹²⁹, G. Tejeda Muñoz ⁴⁴, L. Terlizzi ²⁴, C. Terrevoli ⁵⁰, D. Thakur ²⁴, S. Thakur ⁴, M. Thogersen ¹⁹, D. Thomas ¹⁰⁷, A. Tikhonov ¹³⁹, N. Tiltmann ^{32,124}, A.R. Timmins ¹¹⁴, M. Tkacik ¹⁰⁵, A. Toia ⁶⁴, R. Tokumoto ⁹¹, S. Tomassini ²⁵, K. Tomohiro ⁹¹, N. Topilskaya ¹³⁹, M. Toppi ⁴⁹, V.V. Torres ¹⁰², A. Trifiró ^{30,53}, T. Triloki ⁹⁵, A.S. Triolo ^{32,30,53}, S. Tripathy ³², T. Tripathy ^{125,47}, S. Trogolo ²⁴, V. Trubnikov ³, W.H. Trzaska ¹¹⁵, T.P. Trzcinski ¹³⁴, C. Tsolanta ¹⁹, R. Tu ³⁹, A. Tumkin ¹³⁹, R. Turrisi ⁵⁴, T.S. Tveter ¹⁹, K. Ullaland ²⁰, B. Ulukutlu ⁹⁴, S. Upadhyaya ¹⁰⁶, A. Uras ¹²⁶, M. Urioni ²³, G.L. Usai ²², M. Vaid ⁹⁰, M. Vala ³⁶, N. Valle ⁵⁵, L.V.R. van Doremalen ⁵⁹, M. van Leeuwen ⁸³, C.A. van Veen ⁹³, R.J.G. van Weelden ⁸³, D. Varga ⁴⁶, Z. Varga ^{136,46}, P. Vargas Torres ⁶⁵, M. Vasileiou ⁷⁷, A. Vasiliev ^{I,139}, O. Vázquez Doce ⁴⁹, O. Vazquez Rueda ¹¹⁴, V. Vechernin ¹³⁹, P. Veen ¹²⁸, E. Vercellin ²⁴, R. Verma ⁴⁷, R. Vértesi ⁴⁶, M. Verweij ⁵⁹, L. Vickovic ³³, Z. Vilakazi ¹²¹, O. Villalobos Baillie ⁹⁹, A. Villani ²³, A. Vinogradov ¹³⁹, T. Virgili ²⁸, M.M.O. Virta ¹¹⁵, A. Vodopyanov ¹⁴⁰, B. Volkel ³², M.A. Völkl ⁹⁹, S.A. Voloshin ¹³⁵, G. Volpe ³¹,

B. von Haller³², I. Vorobyev³², N. Vozniuk¹³⁹, J. Vrláková³⁶, J. Wan³⁹, C. Wang³⁹, D. Wang³⁹, Y. Wang³⁹, Y. Wang⁶, Z. Wang³⁹, A. Wegrzynek³², F.T. Weiglhofer³⁸, S.C. Wenzel³², J.P. Wessels¹²⁴, P.K. Wiacek², J. Wiechula⁶⁴, J. Wikne¹⁹, G. Wilk⁷⁸, J. Wilkinson⁹⁶, G.A. Willems¹²⁴, B. Windelband⁹³, M. Winn¹²⁸, J.R. Wright¹⁰⁷, W. Wu³⁹, Y. Wu¹¹⁸, K. Xiong³⁹, Z. Xiong¹¹⁸, R. Xu⁶, A. Yadav⁴², A.K. Yadav¹³³, Y. Yamaguchi⁹¹, S. Yang²⁰, S. Yano⁹¹, E.R. Yeats¹⁸, J. Yi⁶, Z. Yin⁶, I.-K. Yoo¹⁶, J.H. Yoon⁵⁸, H. Yu¹², S. Yuan²⁰, A. Yuncu⁹³, V. Zaccolo²³, C. Zampolli³², F. Zanone⁹³, N. Zardoshti³², A. Zarochentsev¹³⁹, P. Závada⁶², M. Zhalov¹³⁹, B. Zhang⁹³, C. Zhang¹²⁸, L. Zhang³⁹, M. Zhang^{125,6}, M. Zhang^{27,6}, S. Zhang³⁹, X. Zhang⁶, Y. Zhang¹¹⁸, Y. Zhang¹¹⁸, Z. Zhang⁶, M. Zhao¹⁰, V. Zhrebchevskii¹³⁹, Y. Zhi¹⁰, D. Zhou⁶, Y. Zhou⁸², J. Zhu^{54,6}, S. Zhu^{96,118}, Y. Zhu⁶, S.C. Zugravel⁵⁶, N. Zurlo^{132,55}

Affiliation Notes

^I Deceased

^{II} Also at: Max-Planck-Institut für Physik, Munich, Germany

^{III} Also at: Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Bologna, Italy

^{IV} Also at: Dipartimento DET del Politecnico di Torino, Turin, Italy

^V Also at: Department of Applied Physics, Aligarh Muslim University, Aligarh, India

^{VI} Also at: Institute of Theoretical Physics, University of Wrocław, Poland

^{VII} Also at: Facultad de Ciencias, Universidad Nacional Autónoma de México, Mexico City, Mexico

Collaboration Institutes

¹ A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia

² AGH University of Krakow, Cracow, Poland

³ Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kiev, Ukraine

⁴ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India

⁵ California Polytechnic State University, San Luis Obispo, California, United States

⁶ Central China Normal University, Wuhan, China

⁷ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba

⁸ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico

⁹ Chicago State University, Chicago, Illinois, United States

¹⁰ China Nuclear Data Center, China Institute of Atomic Energy, Beijing, China

¹¹ China University of Geosciences, Wuhan, China

¹² Chungbuk National University, Cheongju, Republic of Korea

¹³ Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovak Republic

¹⁴ Creighton University, Omaha, Nebraska, United States

¹⁵ Department of Physics, Aligarh Muslim University, Aligarh, India

¹⁶ Department of Physics, Pusan National University, Pusan, Republic of Korea

¹⁷ Department of Physics, Sejong University, Seoul, Republic of Korea

¹⁸ Department of Physics, University of California, Berkeley, California, United States

¹⁹ Department of Physics, University of Oslo, Oslo, Norway

²⁰ Department of Physics and Technology, University of Bergen, Bergen, Norway

²¹ Dipartimento di Fisica, Università di Pavia, Pavia, Italy

²² Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy

²³ Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy

²⁴ Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy

²⁵ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy

²⁶ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy

²⁷ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy

²⁸ Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy

²⁹ Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy

³⁰ Dipartimento di Scienze MIFT, Università di Messina, Messina, Italy

³¹ Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy

- ³² European Organization for Nuclear Research (CERN), Geneva, Switzerland
- ³³ Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia
- ³⁴ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic
- ³⁵ Faculty of Physics, Sofia University, Sofia, Bulgaria
- ³⁶ Faculty of Science, P.J. Šafárik University, Košice, Slovak Republic
- ³⁷ Faculty of Technology, Environmental and Social Sciences, Bergen, Norway
- ³⁸ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- ³⁹ Fudan University, Shanghai, China
- ⁴⁰ Gangneung-Wonju National University, Gangneung, Republic of Korea
- ⁴¹ Gauhati University, Department of Physics, Guwahati, India
- ⁴² Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany
- ⁴³ Helsinki Institute of Physics (HIP), Helsinki, Finland
- ⁴⁴ High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico
- ⁴⁵ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania
- ⁴⁶ HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- ⁴⁷ Indian Institute of Technology Bombay (IIT), Mumbai, India
- ⁴⁸ Indian Institute of Technology Indore, Indore, India
- ⁴⁹ INFN, Laboratori Nazionali di Frascati, Frascati, Italy
- ⁵⁰ INFN, Sezione di Bari, Bari, Italy
- ⁵¹ INFN, Sezione di Bologna, Bologna, Italy
- ⁵² INFN, Sezione di Cagliari, Cagliari, Italy
- ⁵³ INFN, Sezione di Catania, Catania, Italy
- ⁵⁴ INFN, Sezione di Padova, Padova, Italy
- ⁵⁵ INFN, Sezione di Pavia, Pavia, Italy
- ⁵⁶ INFN, Sezione di Torino, Turin, Italy
- ⁵⁷ INFN, Sezione di Trieste, Trieste, Italy
- ⁵⁸ Inha University, Incheon, Republic of Korea
- ⁵⁹ Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University/Nikhef, Utrecht, Netherlands
- ⁶⁰ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovak Republic
- ⁶¹ Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India
- ⁶² Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic
- ⁶³ Institute of Space Science (ISS), Bucharest, Romania
- ⁶⁴ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- ⁶⁵ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁶⁶ Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil
- ⁶⁷ Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁶⁸ iThemba LABS, National Research Foundation, Somerset West, South Africa
- ⁶⁹ Jeonbuk National University, Jeonju, Republic of Korea
- ⁷⁰ Johann-Wolfgang-Goethe Universität Frankfurt Institut für Informatik, Fachbereich Informatik und Mathematik, Frankfurt, Germany
- ⁷¹ Korea Institute of Science and Technology Information, Daejeon, Republic of Korea
- ⁷² Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France
- ⁷³ Lawrence Berkeley National Laboratory, Berkeley, California, United States
- ⁷⁴ Lund University Department of Physics, Division of Particle Physics, Lund, Sweden
- ⁷⁵ Nagasaki Institute of Applied Science, Nagasaki, Japan
- ⁷⁶ Nara Women's University (NWU), Nara, Japan
- ⁷⁷ National and Kapodistrian University of Athens, School of Science, Department of Physics, Athens, Greece
- ⁷⁸ National Centre for Nuclear Research, Warsaw, Poland
- ⁷⁹ National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India
- ⁸⁰ National Nuclear Research Center, Baku, Azerbaijan
- ⁸¹ National Research and Innovation Agency - BRIN, Jakarta, Indonesia
- ⁸² Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark

- 83 Nikhef, National institute for subatomic physics, Amsterdam, Netherlands
- 84 Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
- 85 Nuclear Physics Institute of the Czech Academy of Sciences, Husinec-Řež, Czech Republic
- 86 Oak Ridge National Laboratory, Oak Ridge, Tennessee, United States
- 87 Ohio State University, Columbus, Ohio, United States
- 88 Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia
- 89 Physics Department, Panjab University, Chandigarh, India
- 90 Physics Department, University of Jammu, Jammu, India
- 91 Physics Program and International Institute for Sustainability with Knotted Chiral Meta Matter (WPI-SKCM²), Hiroshima University, Hiroshima, Japan
- 92 Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany
- 93 Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- 94 Physik Department, Technische Universität München, Munich, Germany
- 95 Politecnico di Bari and Sezione INFN, Bari, Italy
- 96 Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
- 97 Saga University, Saga, Japan
- 98 Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India
- 99 School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
- 100 Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
- 101 Stefan Meyer Institut für Subatomare Physik (SMI), Vienna, Austria
- 102 SUBATECH, IMT Atlantique, Nantes Université, CNRS-IN2P3, Nantes, France
- 103 Sungkyunkwan University, Suwon City, Republic of Korea
- 104 Suranaree University of Technology, Nakhon Ratchasima, Thailand
- 105 Technical University of Košice, Košice, Slovak Republic
- 106 The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
- 107 The University of Texas at Austin, Austin, Texas, United States
- 108 Universidad Autónoma de Sinaloa, Culiacán, Mexico
- 109 Universidade de São Paulo (USP), São Paulo, Brazil
- 110 Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
- 111 Universidade Federal do ABC, Santo Andre, Brazil
- 112 Universitatea Nationala de Stiinta si Tehnologie Politehnica Bucuresti, Bucharest, Romania
- 113 University of Derby, Derby, United Kingdom
- 114 University of Houston, Houston, Texas, United States
- 115 University of Jyväskylä, Jyväskylä, Finland
- 116 University of Kansas, Lawrence, Kansas, United States
- 117 University of Liverpool, Liverpool, United Kingdom
- 118 University of Science and Technology of China, Hefei, China
- 119 University of South-Eastern Norway, Kongsberg, Norway
- 120 University of Tennessee, Knoxville, Tennessee, United States
- 121 University of the Witwatersrand, Johannesburg, South Africa
- 122 University of Tokyo, Tokyo, Japan
- 123 University of Tsukuba, Tsukuba, Japan
- 124 Universität Münster, Institut für Kernphysik, Münster, Germany
- 125 Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France
- 126 Université de Lyon, CNRS/IN2P3, Institut de Physique des 2 Infinis de Lyon, Lyon, France
- 127 Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France, Strasbourg, France
- 128 Université Paris-Saclay, Centre d'Etudes de Saclay (CEA), IRFU, Département de Physique Nucléaire (DPhN), Saclay, France
- 129 Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France
- 130 Università degli Studi di Foggia, Foggia, Italy
- 131 Università del Piemonte Orientale, Vercelli, Italy
- 132 Università di Brescia, Brescia, Italy
- 133 Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India
- 134 Warsaw University of Technology, Warsaw, Poland
- 135 Wayne State University, Detroit, Michigan, United States

¹³⁶ Yale University, New Haven, Connecticut, United States

¹³⁷ Yildiz Technical University, Istanbul, Turkey

¹³⁸ Yonsei University, Seoul, Republic of Korea

¹³⁹ Affiliated with an institute formerly covered by a cooperation agreement with CERN

¹⁴⁰ Affiliated with an international laboratory covered by a cooperation agreement with CERN.