

BB

E1-98-319

FRAGMENTATION  
OF TENSOR POLARIZED DEUTERONS  
INTO CUMULATIVE PIONS

Submitted to «Physics Letters B»

SCAN-9902073



CERN LIBRARIES, GENEVA

529908

1998

S.Afanasiev, V.Arhipov, V.Bondarev, M.Ehara<sup>1</sup>, G.Filipov<sup>2</sup>, S.Fukui<sup>1</sup>,  
S.Hasegawa<sup>1</sup>, T.Hasegawa<sup>3</sup>, N.Horikawa<sup>1</sup>, A.Isupov, T.Iwata<sup>1</sup>, V.Kashirin,  
M.Kawano<sup>3</sup>, T.Kageya<sup>1</sup>, A.Khrenov, V.Kolesnikov, V.Ladygin,  
A.Litvinenko, A.Malakhov, T.Matsuda<sup>3</sup>, H.Nakayama<sup>1</sup>, A.Nikiforov,  
E.Osada<sup>3</sup>, Yu.Pilipenko, S.Reznikov, P.Rukoyatkin, A.Semenov, I.Semenova,  
N.Takabayashi<sup>1</sup>, A.Wakai<sup>1</sup>, L.Zolin<sup>4</sup>

---

<sup>1</sup>Nagoya University, Nagoya 464-8062, Japan

<sup>2</sup>INPNE, 1784 Sofia, Bulgaria

<sup>3</sup>Miyazaki University, Miyazaki 889-21, Japan

<sup>4</sup>E-mail address: zolin@moonhe.jinr.ru

To study the deuteron core structure, a number of experiments in a GeV region including the polarization ones, have been performed in the last two decades. The nucleon momentum distributions in a deuteron, extracted from the deuteron breakup [1] and  $ed$ -inelastic scattering [2], are in good agreement with each other confirming that both hadron and electron probes give reliable information on the deuteron structure. These data were reproduced rather well by the relativistic impulse approximation (IA) calculations using conventional deuteron wave functions (DWF) [3, 4] at internal momenta up to  $k = 0.25$  GeV/c [5]. In the higher momentum region, however, a marked deviation from the calculations was observed. Taking into account additional mechanisms to IA [6, 7] gives a satisfactory description of the momentum distribution data.

A deviation from IA was also observed in the data on the tensor analyzing power  $T_{20}$  for deuteron inclusive breakup with the emission of the proton at a zero angle [8, 9, 10] and for  $dp$ -backward elastic scattering [11, 12] as shown in Fig.1. The same values of  $T_{20}$  are assumed in both reactions since one nucleon exchange mechanism is to be dominant in both of them. However, the data for these reactions disagree in a high momentum region. Furthermore, the  $dp$ -backward data demonstrate a strong oscillation structure in the region  $k > 0.3$  GeV/c. The experimental data over all internal momentum region cannot be described by the calculations taking account of additional mechanisms to IA [6, 13] or considering additional components to the DWF due to relativistic effects [14, 15, 16, 17]. On the other hand, the behavior of  $T_{20}$  at high nucleon momenta is explained by the models incorporating the non-nucleonic degrees of freedom in the deuteron [18, 19]. These theoretical investigations give different explanations to the data on different reactions, in particular in a high momentum region, based on different pictures of the deuteron core structure and nucleon correlations at short distances. It is desirable to have a global picture which could explain the data.

The deuteron fragmentation into cumulative hadrons is one of the suitable reactions to study the deuteron core structure. Cumulative hadrons are those produced in a nuclear reaction beyond the kinematic limit of free nucleon-nucleon collisions. [20] [21]. The simplest model used to describe their production is a direct mechanism, where a nucleon with a high momentum in the projectile, for instance in the deuteron, collides against a nucleon in the target nucleus and produces a cumulative hadron (e.g.  $NN \rightarrow NN\pi$ ) [22]. The fea-




Fig.1. The experimental data for the reactions induced by tensor polarized deuterons in GeV region: the  $T_{20}$  data for deuteron breakup  $dA \rightarrow pX$  on the hydrogen and carbon targets [8, 9, 10]; and the  $T_{20}$  data for backward elastic  $dp \rightarrow pd$  [11, 12]. The solid curve is the IA calculation with the Paris DWF [3] for both breakup and backward elastic scattering, and the dashed curve is the calculation [6] taking account of final state interactions for breakup at 9GeV/c. The direct mechanism IA-prediction for  $dC \rightarrow \pi(0^+)X$  [23] is also shown (dot-dashed curve).

tures of cumulative reactions reflect the nuclear structure at small distances where short range nucleon correlations take place. In case of the deuteron, spin observables in the fragmentation of deuterons into cumulative hadrons allow one to extract information on the deuteron core spin structure.

In order to compare the data on cumulative hadrons obtained under different kinematic conditions, the cumulative variable  $x_c$  ("cumulative number") [20] [23] is often used in the literature. For the reaction of cumulative pion production,  $A + B \rightarrow \pi + X$ , it is expressed as

$$x_c = \frac{(p_B p_\pi) - (m_\pi^2/2)}{(p_A p_B) - m_N^2 - (p_A p_\pi)}, \quad (1)$$

where  $p_A, p_B$  are the four-momenta per nucleon for projectile and target nuclei, respectively,  $p_h$  is the four-momentum of the produced pion,  $m_N$  and  $m_\pi$  are

the nucleon and pion masses. In its meaning, the cumulative number,  $x_c$ , is a minimum target mass (measured in nucleon mass units) in the rest frame of a fragmenting nucleus needed to obey the 4-momentum conservation law for particle production in nucleon-nucleus collision. Hence, its range is up to  $A$ , the mass number of the fragmenting nucleus. When the hadron with  $x_c$  larger than 1 is produced, it is assigned to a cumulative hadron. The backward elastic scattering  $dp \rightarrow pd$  is the special case of cumulative reactions when  $x_c = 2$ . In the framework of the direct mechanism, the cumulative hadron is produced by a nucleon with a high internal momentum  $k$  in the fragmenting nucleus. Hence, the minimum momentum of the nucleon,  $k_{min}$ , which is wanted for cumulative hadron production can be related to  $x_c$ :  $k_{min}$  increases from 0 as  $x_c$  increases from 1. In this paper, the data on the tensor analyzing power  $T_{20}$  at the fragmentation of deuterons into cumulative pions are presented vs the variables  $x_c$  and  $k_{min}$ .

The first measurement of  $T_{20}$  in cumulative pion production in the forward direction

$$\vec{d} + A \rightarrow \pi^\pm(0^\circ) + X \quad (2)$$

has been performed in our previous experiment [23] using a carbon target. The values of  $|T_{20}|$  were found to be about one order smaller than those predicted by the IA calculations. However, it might be due to the nuclear effect of carbon target. Moreover, it was difficult to achieve definite conclusion concerning the sign of  $T_{20}$  due to large errors.

Here, we report new data on  $T_{20}$  obtained in the fragmentation of deuterons into negative pions on hydrogen, beryllium and carbon targets ( $A = 1, 9, 12$ ) at a zero emission angle over the range of the cumulative variable  $x_c$  from 1.08 to 1.76. The values of  $T_{20}$  were determined by measuring the ratio of pion yields ( $N^+/N^-$ ) emitted at forward angles ( $\theta_\pi \leq 15mrad$ ) at different tensor polarizations ( $p_{zz}^\pm$ ) of the deuteron beam:

$$T_{20} = \frac{2\sqrt{2}(\frac{N^+}{N^-} - 1)}{\frac{N^+}{N^-}p_{zz}^- - p_{zz}^+}. \quad (3)$$

The measurements were carried out with a 9 GeV/c polarized deuteron beam slowly extracted from the Dubna 10 GeV accelerator (Synchrophasotron). The ion source POLARIS [24], operating in the tensor polarization mode, provided three tensor polarization states "+, -, 0" ("0" means the absence of polarization). A series of polarization states was cyclically repeated every three spills. The beam polarization was measured with the two-arm polarimeter ALPHA [25] by detecting  $dp$ - elastic scattering at a deuteron momentum of




Fig.2. The layout of the setup of the 4V beam line in the Synchrotron experimental hall.  $M_i$ ; bending magnets,  $L_i$ ; quadrupole magnets,  $F_i, C_i$ ; scintillation and Cherenkov counters,  $HT, HOXY, HOUV$ ; scintillation hodoscopes,  $T$ ; target.  $IC$ : ionization chamber.

3 GeV/c [26]. The values of tensor polarization for the "+" and "-" states were  $p_{zz}^+ = 0.643 \pm 0.033(stat) \pm 0.026(syst)$  and  $p_{zz}^- = -0.729 \pm 0.024(stat) \pm 0.029(syst)$ . To study beam polarization stability for a 5-day run, the beam line polarity was switched periodically to register protons and the  $T_{20}^p$  of the deuteron breakup reaction  $\vec{d} + C \rightarrow p(0^\circ) + X$  was measured (24 times in all). The value of  $T_{20}^p$  at  $k \sim 0.3 GeV/c$  averaged over the run is shown by a star in Fig.1. It is consistent with the data obtained in other experiments [8] [9]. The analysis of  $T_{20}^p$  fluctuations during the run showed that the values of polarization remained stable within  $\pm 2\%$  [27].

The deuteron beam intensity was  $1.5$  to  $2.5 \cdot 10^9 \vec{d}/spill$ . The duration of the beam spill was 0.5 s. The deuterons with a fixed momentum of 9 GeV/c were led to the target  $T$  located at focus  $F_5$ , the starting point of secondary beam line 4V as shown in Fig.2. The position of the beam and its profile ( $\sigma_x = 4$  mm,  $\sigma_y = 9$  mm) at the target point were monitored with the multiwire chambers. The beam intensity was monitored with the ionization chamber  $IC$  placed 1 m upstream the target. The magnetic optics of line 4V, consisting of four magnetic dipoles  $M_0$ - $M_3$  and three quadrupole doublets  $L_1$ - $L_3$ , provided a momentum resolution of  $\sigma_p/p = \pm 2.3\%$  and an angular acceptance of  $\Delta\Omega = 2.1 \cdot 10^{-4}$  sr. The particles produced in the target were

led to final focus F6 located 42 m downstream the target. The scintillation hodoscopes H0XY and H0UV were used to control the beam size at F6.

The detecting system consists of two counter sets as shown in Fig.2. One set, a scintillation hodoscope HT and counters  $F_{561-4}$ , was placed behind the collimator at the intermediate beam focus, and the other, counters  $F_{61-3}$ , was 27 m downstream at focus F6. Pions were selected by time of flight (TOF) between the intermediate focus and the focus F6 with a time resolution of  $\sigma_{TOF}=0.2\text{ns}$ . To reject background due to the pile-up effect in  $F_{561-4}$ , a correlation of two TOF pairs was required. Although events corresponding to pions are only seen in the TOF spectrum, muons from the pions decay should contribute to the peak. Their number was estimated to be no more than 8% [23]. It was confirmed by GEANT simulation [28] that they originate mainly from pions decaying in the final section of beam line 4V. For this reason we did not exclude muon events when we calculate the  $N^+/N^-$ -ratio to extract  $T_{20}$  according to (3).

The pion momentum  $p_\pi$  varied from 3.5 to 5.3 GeV/c at a fixed deuteron beam momentum of  $p_d=9.0$  GeV/c to range  $x_c$  from 1.08 to 1.76. The measurements at  $p_\pi=3.5, 4.0, 4.5$  and  $5.0$  GeV/c were carried out with the hydrogen target ( $7\text{ g/cm}^2$ ). To study the target dependence, the measurement at  $p_\pi=5.0$  GeV/c was performed also with the beryllium target ( $36\text{ g/cm}^2$ ), and the highest  $x_c$  point,  $x_c=1.76$ , was taken with the carbon target ( $55\text{ g/cm}^2$ ). Such relatively thick targets do not affect on the  $T_{20}$ . According to (3) the  $T_{20}$  depends on the  $N^+/N^-$ -ratio only at fixed  $p_{\perp z}^\pm$ . It means that the results of  $T_{20}$ -measurements are not affected by the distortion of the cumulative pion spectra due to secondary interactions in a target and beamline materials as the distortion factor is the same for both polarization states of the primary deuteron beam. Monte Carlo simulation shows that pion rescattering in the carbon target ( $55\text{ g/cm}^2$ ) increases the angular acceptance by 10% with respect to a thin target; an effective value of  $x_c$  increases by  $\simeq 2\%$  due to energy losses and  $\pi A$ -interactions in the target. Experience in studying  $T_{20}$  in deuteron breakup confirmed its independence of target thickness [9] [10]. The background pions not from the target, were measured to be about 3% for a  $36\text{ g/cm}^2$  target and 8% for a  $7\text{ g/cm}^2$  target. They cannot change radically the  $N^+/N^-$ -ratio. To estimate an effect of pions originated not from the reaction (2) which can affect on  $N^+/N^-$ , we estimated a contribution from the two step process where high momentum nucleons from a deuteron breakup produce high momentum pions in secondary collisions in a target:  $dA \rightarrow NX, NA \rightarrow \pi X$ . Our numerical estimations for a thick target ( $55\text{ g/cm}^2$ ), based on the known approximation of the cross sections near the kinematic limit for  $NA \rightarrow \pi X$  [29], showed that the contamination of pions from this process is no more than 1%.

Table 1 presents values on  $T_{20}$  obtained in this experiment for  $\pi^-$  production as a function of  $x_c$  and  $k_{min}$ . At Fig.3 the new  $T_{20}$  data are plotted together with the  $\pi^\pm$  data obtained in the previous experiment on carbon [23].

Table 1

$T_{20}$  as a function of  $x_c$  and  $k_{min}$ ;  $\delta x_c$  is the width of  $x_c$ -bin ( $1\sigma$ ) due to the beam momentum resolution and energy losses in the target with the indicated thickness  $\Delta_T$  ( $g/cm^2$ ). All the momenta ( $p_\pi, k_{min}$ ) are in GeV/c. The invariant cross section  $\sigma_{inv} = E_\pi d\sigma/d^3p_\pi$  is in  $\mu b \cdot GeV^{-2} \cdot c^3 \cdot sr^{-1}/nucleus$ , the uncertainty of  $\sigma_{inv}$  is estimated to be equal to 20%. The systematic error of  $T_{20}$  is caused by the uncertainty of beam polarization.

| $p_\pi$ | Target | $\Delta_T$ | $x_c$ | $\delta x_c$ | $k_{min}$ | $\sigma_{inv}(x_c)$ | $T_{20} \pm stat. \pm sys.$ |
|---------|--------|------------|-------|--------------|-----------|---------------------|-----------------------------|
| 3.5     | H      | 7.1        | 1.08  | $\pm 0.03$   | 0.051     | 23.5                | $0.064 \pm 0.059 \pm 0.004$ |
| 4.0     | H      | 7.1        | 1.25  | $\pm 0.03$   | 0.155     | 19.6                | $0.048 \pm 0.047 \pm 0.003$ |
| 4.5     | H      | 7.1        | 1.43  | $\pm 0.04$   | 0.270     | 3.53                | $0.120 \pm 0.058 \pm 0.008$ |
| 5.0     | H      | 7.1        | 1.61  | $\pm 0.04$   | 0.392     | 0.75                | $0.158 \pm 0.098 \pm 0.010$ |
| 5.0     | Be     | 36         | 1.63  | $\pm 0.04$   | 0.396     | 6.59                | $0.128 \pm 0.056 \pm 0.008$ |
| 5.3     | C      | 55         | 1.76  | $\pm 0.05$   | 0.520     | 1.81                | $0.162 \pm 0.118 \pm 0.010$ |

Comparing the values of  $T_{20}$  obtained on hydrogen, beryllium and carbon targets,  $T_{20}$  in pion production was found to be independent of the mass number of the target. Although a weak target dependence was seen in the inclusive cross sections for cumulative hadron production [30] [31], it is evident that  $T_{20}$  is predominantly ruled by the internal structure of the fragmenting nucleus, in this case the deuteron.

In the framework of the direct mechanism, the increasing  $x_c$  corresponds to the increasing  $k_{min}$ , thus decreasing internucleonic distances  $r_{NX}$ . Using the momentum-space transformation,  $r_{NX}$  is evaluated to be  $r_{NX} \simeq 0.4$  fm at  $x_c=1.7$ . This internucleonic distance corresponds to a large D-wave contribution to the DWF, and therefore a large value of the tensor analyzing power could be expected. The results of the calculations [23] performed in the framework of the direct mechanism using the Paris DWF [3] are shown in Fig.3. It is notable that the measured values of  $T_{20}$  have a positive sign opposite to the sign predicted by the calculations. The value of  $T_{20}$  is nearly zero at  $x_c=1.0$  and increases up to  $\simeq 0.15$  at  $x_c = 1.7$  not showing an extremum at  $k \simeq 0.3$  GeV/c peculiar to IA predictions (Fig.1). Taking account of relativistic effects






Fig.3.  $T_{20}$  vs  $x_c$  and  $k_{min}$  (MeV/c) for the reaction  $d + A \rightarrow \pi(0^\circ) + X$  at the fragmentation of 9 GeV deuterons on the hydrogen, beryllium and carbon targets obtained in this experiment along with the previous data [23] obtained with the carbon target for  $\pi^-$  (open squares) and  $\pi^+$  (open cross). The dashed line is a linear fit of the data ( $\chi^2/d.o.f.=0.5$ ). The solid curve is our calculation based on the Paris DWF [23].

[32] or processes additional to the direct mechanism like  $\pi$ -rescattering [33] cannot explain such a behaviour of  $T_{20}(k)$ .

The deuteron spin structure is also studied in the experiments with electron probes [35] [36] although the data are limited within a lower internal momentum region. Based on the hypothesis of NN-core in the deuteron, oscillating behavior of the tensor analyzing power due to variation of the D-wave contribution was predicted for  $ed$ - elastic scattering and also for cumulative pion production [34]. The available  $T_{20}$  data for  $ed$ - elastic scattering [35] [36] confirmed this prediction. Hence, we can conclude that IA calculations are successful in all previously studied reactions induced by deuterons (the deuteron breakup and dp-,ed-scattering) at low internal momenta; on the contrary, a marked deviation from IA is observed at higher  $k$  ( $> 0.3$  GeV/c). It is different in form in different reactions as shown in Fig.1. The reasons of these deviations were discussed by many authors: in particular, final state interactions

and non-nucleonic contributions ( $NN^*$ ,  $\Delta\Delta$  or  $6q$ - configurations) in the DWF were considered.

In the case of cumulative pion production, we met a more serious problem: the sign of  $T_{20}(k)$  was found to be opposite to the IA calculation over the whole region of internal momenta from 0 to 0.5 GeV/c studied in the experiment. This contradiction seems not to be solved by any current models, based on the traditional picture of the deuteron as a bound state of a neutron and a proton. It may be more successfully the  $T_{20}$  pion data can be explain by introducing quark scenarios for cumulative meson production [37] [38], where the deuteron core is considered as a multiquark configuration and a quark-spectator with a high momentum is softly hadronized into a cumulative particle. To explain the behaviour of spin observables in cumulative hadron production the similar quark approaches introducing in consideration of spin degrees of freedom in nuclei should be developed.

The results of the present work are the following:

- The tensor analyzing power  $T_{20}$  of the reaction  $d+H(Be, C) \rightarrow \pi^-(0^\circ)+X$  at a deuteron momentum of 9 GeV/c and pion momenta from 3.0 to 5.3 GeV/c has been measured . In the framework of the direct mechanism, this kinematic region corresponds to internal momenta of the nucleon in the deuteron up to  $k \simeq 500$  MeV/c.

- It has been observed that the behavior of  $T_{20}$  is independent of the atomic number of the target. This fact and a weak A-dependence of the cross section shape of deuteron fragmentation into cumulative pions [30] [31] evidently confirm that  $T_{20}$  is predominantly determined by the internal structure of the deuteron.

- The sign of the measured  $T_{20}$  is positive and opposite to that obtained by the calculations using standard DWFs assuming the direct mechanism. The  $T_{20}$  data cannot be reproduced by relativistic corrections or taking pion rescattering into account either. Attention should be given to more sophisticated approaches taking account of non-nucleonic degrees of freedom in nuclei.

The authors are grateful to the LHE accelerator staff and to the cryogenic division staff supporting POLARIS and the liquid hydrogen target operation, especially to A.Kovalenko, A.Kirillov, L.Golovanov, V.Finushkin, V.Ershov, Yu.Borzunov, A.Tsvinev. We would like to thank L.Strunov, V.Sharov, and A.Nomofilov for the primary deuteron polarization measurements by the ALPHA polarimeter. .

This research was supported in part by Russian Foundation for Fundamental Research under grant No. 96-02-17208 and by Japan Society for the

Promotion of Science(JSPS), Daikou Foundation and Toyota Foundation.

## References

- [1] V.G.Ableev et al., Nucl. Phys. A 393 (1983) 491, and A 411 (1983) 541E.
- [2] P.Bosted et al., Phys. Rev. Lett. 49 (1982) 1380.
- [3] M.Lacombe et al., Phys. Lett. B 101 (1981) 139.
- [4] R.V. Reid, Ann. Phys. 50 (1968) 411.
- [5] At high energy deuteron fragmentation it is preferable to use the definition of the internal momenta in the infinite momentum frame (IMF)  
 $k = \sqrt{(m_N^2 + p_{\perp}^2)/(4\alpha(1-\alpha)) - m_N^2}$ , where  $\alpha = (E_N + p_{N\parallel})/(E_d + p_d) = (E_q + q_{\parallel})/m_d$  is the light cone variable;  $E_N, p_{N\parallel}$  and  $E_q, q_{\parallel}$  are the nucleon energies and momenta parallel to the incident deuteron momentum  $p_d$  in the laboratory and in the deuteron frames, respectively;  $p_{\perp}$  is the transverse component of the nucleon momentum (for refs. see L.L.Frankfurt and M.Strikman, Phys. Rep. 76 (1981) 215).
- [6] G.I. Lykasov, Phys. Part. Nucl. 24 (1993) 59.
- [7] C.Ciofi degli Atti et al., Phys.Rev. C36 (1987) 1208.
- [8] C.F.Perdrisat et al., Phys. Rev. Lett. 59 (1987) 2840;  
V.Punjabi et al., Phys. Rev. C 39 (1989) 608.
- [9] T.Aono et al., Phys. Rev. Lett. 74 (1995) 4997.
- [10] L.S.Azhgirey et al., Phys. Lett. B 387 (1996) 37.
- [11] V.Punjabi et al., Phys. Lett. B 350 (1995) 178.
- [12] L.S.Azhgirey et al., Phys. Lett. B 391 (1997) 22.
- [13] A.Nakanura and L.Satta, Nucl.Phys. A445 (1985) 706.
- [14] W.W.Buck and F.Gross, Phys.Rev. D20 (1979) 2361.
- [15] M.A.Braun and M.V.Tokarev, Part. and Nucl. 22 (1991) 1237.
- [16] L.Kaptari et al., Phys.Lett. B351 (1995) 400.

- [17] B.D.Keister and J.A.Tjon, Phys.Rev.C26 (1982) 578;  
L.Kaptari et al., Phys.Rev. C57 (1998) 1097.
- [18] A.P.Kobushkin, J. Phys. G: Nucl. Phys. 19 (1993) 1993.
- [19] A.P.Kobushkin, Phys. Lett. B 421 (1998) 53.
- [20] A.M.Baldin, Nucl. Phys. A 434 (1985) 695c.
- [21] V.B.Kopeliovich, Phys. Rep., vol. 139, No. 2 (1986) 52-157.
- [22] I.A.Schmidt and R. Blankenbecler, Phys.Rev. D15 (1977) 3321;  
Ch.-Y. Wong and R. Blankenbecler, Phys.Rev. C22 (1980) 2433.
- [23] S.Afanasiev et al., Nucl. Phys. A 625 (1997) 817.
- [24] V.P.Ershov et al., in Proc. Intern. Workshop on Polarized Beams and  
Polarized Gas Targets, Cologne, Germany, June 6-9, 1995, p.193.
- [25] V.G.Ableev et al., Nucl. Instr. and Meth. A 306 (1991) 73.
- [26] V.Ghazikhanian et al., Phys.Rev. C 43 (1991) 1532.
- [27] L.S.Zolin et al., JINR Rapid Comm. 2[88]-98 (1998) 27.
- [28] R.Brun et al., GEANT 3, CERN report DD/EE/84-1,1989.
- [29] V.S.Barashenkov, N.V.Slavin, Phys. Part. Nucl. 15 (1984) 997.
- [30] E.Moeller et al., Phys. Rev. C28 (1983) 1246.
- [31] Yu.S.Anisimov et al., Sov. J. Nucl. Phys. 60 (1997) 1070.
- [32] M.V.Tokarev, Proc. Int. Workshop "Dubna Deuteron-91", Dubna, 1991,  
(JINR, E2-92-25, Dubna, 1992) p.84.
- [33] A.Wakai, Talk at Intern. Symposium "Dubna Deuteron-97", Dubna, July  
2-5, 1997 (to be published in Proc. of the Symposium).
- [34] L.L.Frankfurt and M.Strikman, Nucl. Phys. A 405 (1983) 557.
- [35] C.E. Jones et al., in Proc. of the 14th Intern. European Conference on  
Few Body Problems in Physics, Amsterdam, Aug. 23-27, 1993, p.112.
- [36] M.Garcon et al., Phys. Rev. C 49 (1994) 2516.
- [37] V.V.Burov et al., Phys. Lett. B 67 (1977) 46.
- [38] V.K.Lukyanov, A.I.Titov, Phys. Part. Nucl. 10 (1979) 815.

**Received by Publishing Department  
on November 12, 1998.**

**The Publishing Department  
of the Joint Institute for Nuclear Research  
offers you to acquire the following books:**

| Index         | Title   |
|---------------|---|
| 94-55         | Proceedings of the International Bogoliubov Memorial Meeting. Dubna, 1993 (216 p. in Russian and English)   |
| D3,14-95-323  | VII School on Neutron Physics. Lectures. Vol.1. Dubna, 1995 (356 p. in Russian and English)   |
| E10,11-95-387 | Proceedings of the ESONE International Conference 'RTD'94 on REAL TIME DATA 1994 with Emphasis on Distributed Front-End Processing. Dubna, 1994 (358 p. in English)                           |
| D15-96-18     | Proceedings of the International Workshop Charge and Nucleon Radii of Exotic Nuclei. Poznan, 1995 (172 p. in Russian and English)   |
| E9-96-21      | Proceedings of VII ICFA Beam Dynamics Workshop on «Beam Issues for Multibunch, High Luminosity Circular Colliders». Dubna, 1995 (198 p. in English)   |
| E2-96-100     | Proceedings of the 3rd International Symposium «Dubna Deuteron-95». Dubna, 1995 (374 p. in English)   |
| E2-96-224     | Proceedings of the VII International Conference «Symmetry Methods in Physics». Dubna, 1996 (2 volumes, 630 p., in English)  |
| E-96-321      | Proceedings of the International Conference «Path Integrals: Dubna'96». Dubna, 1996 (392 p. in English)   |
| E3-96-336     | Proceedings of the IV International Seminar on Interaction of Neutrons with Nuclei. Dubna, 1996 (396 p. in English)   |
| E3-96-369     | Proceedings of the X International Conference «Problems of Quantum Field Theory». Dubna, 1996 (437 p. in English)   |
| E3-96-507     | Proceedings of the International Workshop «Polarized Neutrons for Condensed Matter Investigations». Dubna, 1996 (154 p. in English)   |
| D1,2-97-6     | Proceedings of the International Workshop «Relativistic Nuclear Physics: from MeV to TeV». Dubna, 1996 (2 volumes 418 p. and 412 p. in English and Russian)                                   |
| E7-97-49      | Proceedings of the 3rd International Conference «Dynamical Aspects of Nuclear Fission». Slovakia, 1996 (426 p. in English)  |
| E1,2-97-79    | Proceedings of the XIII International Seminar on High Energy Physics Problems. Relativistic Nuclear Physics and Quantum Chromodynamics. Dubna, 1996 (2 volumes, 364 p. and 370 p. in English) |
| D5,11-97-112  | Proceedings of the 9th International Conference «Computational Modelling and Computing in Physics». Dubna, 1996 (378 p. in English)   |

| Index       | Title   |
|-------------|---|
| E2-97-213   | Proceedings of the V International Seminar on Interaction of Neutron with Nuclei «Neutron Spectroscopy, Nuclear Structure, Related Topics». Dubna, 1997 (446 p. in English) |
| E2,4-97-263 | Proceedings of the Third International Conference «Renormalization Group'96». Dubna, 1996 (436 p. in English)   |
| E10-97-272  | Proceedings of the Data Acquisition Systems of Neutron Experimental Facilities (DANEF'97). Dubna, 1997 (325 p. in English)  |
| D19-97-284  | Proceedings of the International Symposium «Problems of Biochemistry. Radiation and Space Biology». Dubna, 1997 (2 volumes 284 p. and 405 p. in Russian and English)        |
| E2-97-413   | Proceedings of the VII Workshop on High Energy Spin Physics (SPIN'97). Dubna, 1997 (398 p. in English)  |

Please apply to the Publishing Department of the Joint Institute for Nuclear Research for extra information. Our address is:

Publishing Department  
 Joint Institute for Nuclear Research  
 Dubna, Moscow Region  
 141980 Russia  
 E-mail: [publish@pds.jinr.dubna.su](mailto:publish@pds.jinr.dubna.su).

Афанасьев С. и др.

E1-98-319

Фрагментация тензорно-поляризованных дейтронов  
в кумулятивные пионы

Измерена тензорная анализирующая способность  $T_{20}$  реакции  $\vec{d} + A \rightarrow \pi^- (0^\circ) + X$  при фрагментации тензорно-поляризованных дейтронов с энергией 9 ГэВ в пионы с импульсом от 3,5 до 5,3 ГэВ/с на водородной, бериллиевой и углеродной мишенях. Данный кинематический интервал соответствует области кумулятивного рождения адронов с кумулятивной переменной  $x_c$  от 1,08 до 1,76. Величины  $T_{20}$  оказались небольшими и положительными по знаку, что противоречит предсказаниям, основанным на прямом механизме, исходящем из  $NN$ -взаимодействия высокоимпульсного нуклона в дейтроне с нуклоном мишени ( $NN \rightarrow NN\pi$ ).

Работа выполнена в Лаборатории высоких энергий ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 1998

Afanasiev S. et al.

E1-98-319

Fragmentation of Tensor Polarized Deuterons into Cumulative Pions

The tensor analyzing power  $T_{20}$  of the reaction  $\vec{d} + A \rightarrow \pi^- (0^\circ) + X$  has been measured in the fragmentation of 9 GeV tensor polarized deuterons into pions with momenta from 3.5 to 5.3 GeV/c on hydrogen, beryllium and carbon targets. This kinematic range corresponds to the region of cumulative hadron production with the cumulative variable  $x_c$  from 1.08 to 1.76. The values of  $T_{20}$  have been found to be small and consistent with positive values. This contradicts the predictions based on a direct mechanism assuming  $NN$  collision between a high momentum nucleon in the deuteron and a target nucleon ( $NN \rightarrow NN\pi$ ).

The investigation has been performed at the Laboratory of High Energies, JINR.

Preprint of the Joint Institute for Nuclear Research, Dubna, 1998

Макет Т.Е.Попеко

Подписано в печать 26.11.98  
Формат 60 × 90/16. Офсетная печать. Уч.-изд. листов 1,35  
Тираж 430. Заказ 51032. Цена 1 р. 62 к.

Издательский отдел Объединенного института ядерных исследований  
Дубна Московской области