

CERN-TH/2000-114
DFTT 15/2000
SPIN-2000/13
UB-ECM-PF-00/06
ITFA-2000-08
hep-th/0004080

On the Stringy Nature of Winding Modes in Noncommutative Thermal Field Theories

G. Arcioni^{a,b}, J.L.F. Barbon^{c,1}, Joaquim Gomis^{d,e} and M.A. Vazquez-Mozo^{e,b}

^a Dipartimento di Fisica Teorica, Università di Torino, via P. Giuria 1
I-10125 Torino, Italy and INFN Sezione di Torino
arcioni@to.infn.it

^b Spinoza Instituut, Universiteit Utrecht, Leuvenlaan 4, 3584 CE Utrecht, The Netherlands
g.arcioni@phys.uu.nl, M.Vazquez-Mozo@phys.uu.nl

^c Theory Division, CERN, CH-1211 Geneva 23, Switzerland
barbon@mail.cern.ch, gomis@mail.cern.ch

^d Departament ECM, Facultat de Física, Universitat de Barcelona
and Institut de Física d'Altes Energies, Diagonal 647, E-08028 Barcelona, Spain
gomis@ecm.ub.es

^e Instituut voor Theoretische Fysica, Universiteit van Amsterdam
Valckenierstraat 65, 1018 XE Amsterdam, The Netherlands
vazquez@wins.uva.nl

CERN-TH/2000-114

¹ On leave from : Departament de Física de Partícules, Universidad de Santiago de Compostela, Spain.

A B S T R A C T

We show that thermal noncommutative field theories admit a version of 'channel duality' reminiscent of open/closed string duality, where non-planar thermal loops can be replaced by an infinite tower of tree-level exchanges of effective fields. These effective fields resemble closed strings in three aspects: their mass spectrum is that of closed-string winding modes, their interaction vertices contain extra moduli, and they can be regarded as propagating in a higher-dimensional 'bulk' space-time. In noncommutative models that can be embedded in a D-brane, we show the precise relation between the effective 'winding fields' and closed strings propagating on the D-brane. The winding fields represent the coherent coupling of the infinite tower of closed-string oscillator states. We derive a sum rule that expresses this effective coupling in terms of the elementary couplings of closed strings to the D-brane. We furthermore clarify the relation between the effective propagating dimension of the winding fields and the true codimension of the D-brane.

1. Introduction

It has been realized that noncommutative field theories (NCFT) emerge as effective field theories of string/M-theory compactifications in the presence of constant antisymmetric tensor fields [1][2][3]. This result has triggered a renewed interest in the study of both perturbative [4][5][6] and nonperturbative [7] aspects of noncommutative field theories. The stringy connection of NCFT opens up the interesting possibility of trying to understand some of their physical features by embedding them in string theory. In particular a number of NCFT have been obtained as a low-energy limit of open-string theories in B-field backgrounds [8][9][10][11][12][13]. As a matter of example, one can try to understand the nonlocality inherent in these quantum field theories in terms of string theory after an appropriate low-energy limit is taken [3].

Among the most intriguing features of NCFT is a peculiar mixing between infrared and ultraviolet scales [5][6]. On physical grounds it can be understood as the result of the uncertainty principle between two noncommuting spatial dimensions, since probing ultraviolet physics in one direction leads to infrared effects in the other. At a more technical level, this mixing reflects itself in the appearances of extra poles at zero momentum in some amplitudes in the limit where the ultraviolet cutoff is sent to infinity. The authors of Refs. [5][6] interpreted these poles as resulting from the interchange of a new field with kinetic kernel $\partial^2 \delta^d(x)$. It would be very interesting to see if there is a stringy interpretation for these particles.

One of the obvious ways of spotting stringy behavior in NCFT would be to look at situations where the presence of extended objects is manifest, as for example studying these theories in spaces with nontrivial topology or at finite temperature [14][15][16]. In Refs. [17][18] it was pointed out that the two loop thermal partition function of some NCFT can be cast in a way that indicates the presence of states whose energy scales with the inverse temperature as $j' j$, with j' some integer number. This would suggest that NCFT contains certain extended degrees of freedom that are able to wrap around the euclidean time.

In this paper we will try to understand whether some kind of winding modes can be identified in noncommutative thermal perturbation theory, extending on the work of [17]. Actually, we shall see that the winding modes formally identified in thermodynamical quantities can be associated to effective fields with special propagators, much in the same fashion as the α' -fields of Refs. [5][6]. In fact, the UV/IR interpretation of these propagators is the same once we realize that the temperature acts as an ultraviolet cutoff in the field theory.

Therefore, this raises the question of whether the 'winding fields' could be interpreted

as ‘D-brane’ closed-string modes that survive the Seiberg-Witten (SW) decoupling limit. We find that this expectation is not fulfilled, at least in a literal sense. In particular, any closed-string picture amounts to the exchange of the infinite tower of closed-string excitations in the bulk, and therefore it is not a very transparent way of describing the dynamics. Instead, each winding field describes a sort of coherent exchange of an infinite number of closed-string modes. One of our results is the derivation of a sum rule for the effective coupling of the winding fields, in terms of the elementary couplings of closed strings to a D-brane. In fact, the interactions of these winding fields are not specified solely in terms of standard interaction vertices, except in very special kinematical situations. Generically, the vertices contain additional modular parameters that must be integrated over.

The paper is organized as follows. In Secs. 2 and 3 we extend the analysis of Refs. [17][18] to more general diagrams in NCFT at finite temperature and try to cast the loop amplitudes in a ‘dual channel’ picture, in terms of tree-level exchanges. In Sec. 4 we will obtain these amplitudes by studying the low energy SW limit from string theory in order to identify the low-energy winding modes with undecoupled winding strings. Finally in Sec. 5 we will summarize our conclusions.

2. Winding modes in noncommutative quantum field theory: an elementary example

The simplest situation where one can formally identify ‘winding modes’ is that of the two-loop contribution to the free energy in a $d=4$ theory. The planar diagram is independent of the deformation parameter θ , but a non-trivial phase $(p;q) = p \cdot q$ enters in the loop integral in the non-planar case²

$$F_{NP} = g^2 \int_{p,q} \frac{e^{i(p;q)}}{(p^2 + M^2)(q^2 + M^2)}; \quad (2.1)$$

where we have used the notation

$$p^2 = p^2 + \frac{4\theta^2 n^2}{2}; \quad \int_{p,q} = \int_{n \in \mathbb{Z}^d} \frac{1}{(2\pi)^d} \frac{dp}{(2\pi)^{d-1}};$$

The ultraviolet divergences of this integral can be appropriately eliminated by renormalization of the $T=0$ limit, as usual in thermal field theory [19]. In fact, the ultraviolet structure of this diagram is milder than that of the planar counterpart, because the divergence contributed by one of the loops is effectively cut-off by the noncommutative phase

² We will assume throughout that $\theta^i = 0$.




Fig. 1: ‘Channel duality’ in the nonplanar self-energy diagram in $d = 4$ theory.

provided $(p^2 - (p')^2)$ is non-vanishing. This is an example of the UV/IR mixing of [5], namely this divergence will reappear disguised as an infrared effect as $p \rightarrow 0$.

In Ref. [17] it was pointed out that one could ‘integrate out’ one of the loops and replace it by a statistical sum over objects living at the formally T-dual temperature $1 = (T)$, thus representing analogues of winding modes. The essential phenomenon can be understood by simply looking at the one-loop self-energy tadpole diagram (Fig. 1A)

$$(\Pi(p))_{NP} = g^2 \int \frac{d^d q}{(2\pi)^d} \frac{e^{i(p \cdot q)}}{q^2 + M^2} \quad (2.2)$$

Introducing a Schwinger-parameter representation of the propagator,

$$(\Pi(p))_{NP} = g^2 \int_0^\infty dt \int \frac{d^d q}{(2\pi)^d} e^{-t(q^2 + M^2 - i\frac{(p \cdot q)}{t})} \quad (2.3)$$

we can perform the gaussian integral over q . After a further Poisson resummation in the thermal frequency running in the loop we obtain

$$(\Pi(p))_{NP} = \frac{g^2}{4^{\frac{d}{2}}} \int_0^\infty ds s^{\frac{d-4}{2}} \sum_{n \in \mathbb{Z}} e^{-s[2n^2 + (p')^2] \frac{M^2}{4s}} \quad (2.4)$$

where we have changed variables to the ‘dual’ Schwinger parameter $s = 1=4t$. This form is very convenient to perform the subtraction of the $T = 0$ self-energy, since we simply have to restrict the integer sum to $n \neq 0$.

For $d < 4$, the explicit power of $s^{\frac{d-4}{2}}$ in the proper time integral can be ‘integrated in’ into the exponent by introducing $4 - d$ extra gaussian variables z_γ and we can write the full non-planar loop in the following suggestive form

$$(\Pi(p))_{NP} = \sum_{j \in \mathbb{Z}} \int \frac{d^d z_\gamma}{(2\pi)^d} \frac{j^2 (p \cdot z_\gamma)^2}{2n^2 + (p')^2 + z_\gamma^2} \quad (2.5)$$

That is, we have written the original loop diagram in a ‘dual channel’ in terms of an infinite number of tree-level exchanges of particles j , with momenta p , mass proportional to j^2

and extra momentum variables in $d^2 = 4 - d$ transverse dimensions (Fig. 1B). This complete expression renormalizes the mass of the particle running in the second loop³.

The mass of the ψ -elds, scaling as integer multiples of the thermal length, is characteristic of winding modes of closed strings. The effective coupling squared of these particles to the ϕ -elds in external legs is given by

$$\langle \phi(p; z_1) \phi(p; z_2) \rangle = \frac{g^2}{4^{2-d}} \int_0^1 ds e^{-s \frac{1}{4s} M^2 [z_1^2 + (p^2 + z_2^2)]}. \quad (2.6)$$

Thus, if the original ϕ -eld was massive, the coupling to the ψ -eld is suppressed at high values of momentum and winding number j , i.e. only ψ -elds with winding numbers $|j| < (M)^{1/2}$ contribute significantly to the tree-level exchange. The most interesting case is that of a massless ϕ -eld theory. In this case the effective coupling is constant and weights all winding numbers democratically, with the coupling strength $g = g(2-d)$. Finally, if the ϕ -eld is tachyonic, the whole expression is meaningless, since it diverges at the $s = 0$ end. For this matter this ‘channel duality’ in NCFT is reminiscent of open/closed-string channel duality in string theory. Since NCFT can be obtained in many cases as low-energy limits of open-string theories, we find it natural that the ‘dual channel’, obtained through a modular transformation $t = 1/4s$, exhibits the open-string tachyon as an ultraviolet divergence.

It is most interesting to compare the winding ψ -elds we have defined with the ϕ -particles of [5][6]. The structure of the propagator shows that these ψ -elds are formally similar

$$\langle \phi(p; z_1) \phi(p; z_2) \rangle = \frac{1}{2^{2-d} + (p^2 + z_2^2)^2}; \quad (2.7)$$

namely, they have a ‘static’ kinetic term with the kernel $\mathcal{G} = (p^2)^{-2}$ for a ϕ -eld non-canonical dimension. Furthermore, at least as long as $d \leq 4$, the non-standard power of the propagator can be understood in terms of a free propagation in a $4 - d$ dimensional ‘transverse bulk’. The effective mass $|j|$ plays also the role of the inverse ultraviolet cut-off

¹ in the treatment of [5] and, in the absence of the explicit ultraviolet cut-off, the original ultraviolet divergence is back as an infrared divergence at $p^2 = 0$. In our expression, this shows up as a pole in the zero-winding sector. It is precisely this contribution that is subtracted when renormalizing the self-energy by the zero-temperature one⁴. Therefore,

³ One could proceed in the standard way and perform a resummation of ring diagrams.

⁴ Notice that this procedure is different from the one followed in [17] where the authors worked with the two-loop free energy for $d = 4$ NCFT before subtracting the zero-temperature counterterms. In that case the ultraviolet divergence in one of the original loops partially transforms into an infrared one after Poisson resummation and integration over the loop momenta. This is just a consequence of UV/IR mixing.

we confirm that j plays the role of a regulator.

One important difference between our tree-level exchange interactions and the α' -elds of [5][6] is that our ultraviolet cutoff, Λ , has a physical interpretation, and we are free from the arbitrariness of the choice of Wilsonian cutoffs. In particular we can integrate out the complete non-planar loop in terms of the infinite tower of tree exchanges of α' particles. The manipulation is not a priori restricted to the extreme ultraviolet part.

One interesting aspect of the tree-exchange 'dual' representation (2.5) is that it admits an interpretation for the planar diagram too. The only difference in the planar case comes from setting $p = 0$. Therefore, the planar thermal loop can be replaced in this case by

$$(i; p)_P \quad (1; i; p)_P = \sum_{l \neq 0} \int dz_l \frac{\mathcal{Y}(i; p = 0; z_l)^2}{2l^2 + z_l^2} : \quad (2.8)$$

Now we must work with the fully renormalized quantity ($l \neq 0$ in the winding sum) and the propagator of the α' particles is inserted formally at zero noncommutative momentum, i.e. we have a sum over zero-momentum tadpoles of the α' -elds. This is also reminiscent of the closed-string interpretation, because closed strings have tree-level tadpoles on D-branes. Now we would be inclined to interpret the residue of the propagator poles as the product of the couplings g_{vac} (Fig. 2).




Fig. 2: Dual channel interpretation of the thermal loop in the planar contribution to the two-point function in α'^4 noncommutative eld theory.

3. Integrating out a general loop

In the above example we have seen how the effect of a thermal tadpole loop in a noncommutative α'^4 theory admits a 'dual channel' interpretation in terms of a tree-level interchange of some 'winding' α' -eld with inverse propagator $2l^2 + (p)^2$ which mixes with the fundamental α' -quantum. It would be interesting to decide to what extent this duality between thermal loops and tree-level α' -exchanges is a general feature of noncommutative eld theory at finite temperature, or just a property of a particular class of diagrams and theories.

3.1. Generic one-loop diagram in noncommutative \mathbb{R}^n theory

The first case we can consider is the generalization of the example studied in the previous section, a generic one-loop diagram with $N = N_+ + N_-$ vertices in the noncommutative version of \mathbb{R}^n field theory in d dimensions (Fig. 3), where we will take N_- vertices as ‘twisted’, so the amplitude will be nonplanar whenever $N_- \neq 0$. Thus, the fully amputated amplitude can be written as

$$A(p_1; \dots; p_N) = g^{2N} \prod_{a=1}^Z P_a^N \int \frac{e^{i \sum_{a=1}^Z p_a \cdot (q + Q_a)}}{(q + Q_a)^2 + M^2} (Q_N);$$

where $Q_a = \sum_{i=1}^{P_a} p_i$ and $\epsilon_a = \pm 1$ depending on whether the insertion is twisted or not; p_a indicates the total momentum entering in the loop through the a -th insertion. Using Feynman and Schwinger parameters we can write

$$A(p_1; \dots; p_N) = g^{2N} (N-1)! e^{i \sum_{a=1}^Z p_a \cdot Q_a} \int dt t^{N-1} e^{-tM^2} \int [dx] e^{i \sum_{a=1}^Z p_a \cdot x_a Q_a^2} \int \frac{1}{n^2 Z} \int e^{-t \left(\frac{4-2n^2}{2} + \frac{4-n}{2} \sum_{a=1}^Z p_a \cdot x_a Q_a^0 \right)} \int \frac{dq}{(2)^{d-1}} e^{-t(q^2 + 2q \cdot \sum_{a=1}^Z p_a \cdot x_a Q_a)} e^{i p_{np} \cdot q}; \quad (3.1)$$

the integration measure $[dx]$ over the Feynman parameters x_a ($a = 1; \dots; N$) is given by

$$[dx] = \prod_{a=1}^N x_a \prod_{a=1}^{N-1} dx_a$$

and p_{np} denotes the total nonplanar spatial momentum entering in the loop through the N_- ‘twisted’ insertions, $p_{np} = \frac{1}{2} \sum_{a=1}^{N_-} p_a$.

By integrating the loop spatial momentum and performing a Poisson resummation the total amplitude can be recast in terms of the dual Schwinger parameters $s = 1/(4t)$ in the form

$$A(p_1; \dots; p_N) = g^{2N} \frac{(N-1)!}{2^{2N} \frac{d}{2}} W_{NC} \int \frac{1}{ds} s^{\frac{d-2N-2}{2}} \int e^{-s \left[\frac{d-2}{2} + (p_{np})^2 \right]} F(s; p_1; \dots; p_N); \quad (3.2)$$

where W_{NC} is the overall noncommutative phase of the diagram

$$W_{NC} = e^{i \sum_{a=1}^Z p_a \cdot Q_a} \quad (3.3)$$

and the function $F(s; \{p_a\})$ is expressed in terms of an integral over the x^a as

$$F(s; \{p_a\}) = e^{-\frac{1}{4s} M^2} \int_0^1 [dx] e^{-\frac{1}{4s} \sum_a x_a Q_a^2} \left(\sum_a x_a Q_a \right)^2 e^{i \sum_a x_a (Q_a^0)^2} e^{i \sum_a x_a Q_a \cdot (P_P)}; \quad (3.4)$$

As in the simpler case of the tadpole of the $\mathcal{N}=4$ theory, whenever $d < 2N + 2$ we can replace the factor $s^{-\frac{d-2N-2}{2}}$ by an integral over $2 + 2N - d$ extra variables, so we can finally write the diagram in the form of a tree-level exchange of effective fields propagating in $d^2 = 2N + 2 - d$ additional 'bulk' dimensions,

$$A(p_1; \dots; p_N) = \int_{\mathbb{R}^{2Z}} dz_? \frac{f(\{p_a; z_?\})}{s^2 + (P_{np})^2 + z_?^2} \quad (3.5)$$

where the function $f(\{p_a; z_?\})$ is given by

$$f(\{p_a; z_?\}) = g^{2N} \frac{(N-1)!}{2^{2N-\frac{d}{2}}} W_{NC} \int_0^1 ds F(s; \{p_a; z_?\})$$




Fig. 3: Channel duality for a nonplanar thermal loop in $\mathcal{N}=n$ noncommutative field theory.

In the same spirit of the $\mathcal{N}=4$ tadpole one would like to interpret the amplitude (3.5) as a 'dual channel' representation of the original loop diagram in terms of a tree-level exchange of χ -particles with propagators (2.7), so the function $f(\{p_a; z_?\})$ would be interpreted as the product of the couplings in Fig. 3B

$$g_{(nN+)} g_{(nN)} f(\{p_a; z_?\}); \quad (3.6)$$

However, such an identification is rather problematic. Unlike the case of the $\mathcal{N}=4$ tadpole, there seems to be no unambiguous way to define the individual couplings $g_{(nN+)}$ and

$g_{(nN)}$, since their product (3.6) is expressed in terms of a function which does not factorize into the contributions of the two vertices. Moreover, because of the integration over the Feynman parameters in Eq. (3.4), the interaction on the two vertices cannot be disentangled, even for massless elds. It is only in the tadpole case ($N = 1$) that the integration over Feynman parameters disappears and the whole loop can be understood as resulting from the mixing of the eld with an effective ϕ -eld, thus generalizing the result of the previous section to n .

Therefore, even if we can formally replace the generic thermal loop by the exchange of an effective ϕ -particle, we cannot assign ordinary Feynman rules to this eld, since the total amplitude is expressed as a convolution of the two interaction vertices, and not just a product as it is the case of ordinary (and noncommutative) quantum eld theory. We summarize this state of affairs by saying that the vertices of the ϕ -elds have relative moduli that must be integrated over.

In principle, the ϕ -elds introduced here could become bona fide elds, with standard Feynman rules, when considering only the behaviour of the diagram at singularities of the integral over Feynman parameters. We suspect that this is the precise link between the ϕ -elds defined here and the ϕ -elds of ref. [5][6].

The appearance of moduli in the ‘dual channel vertices’ will find a string-theory explanation in the next section. First, we shall discuss some special instances in which the formalism simplifies.

3.2. Some special cases at two loops

The previous example seems to indicate that, although in general we can replace nonplanar loops in NCFT by tree-level exchanges of an infinite tower of some effective ϕ -elds, in a generic situation the nonlocal character of this eld makes the effective description not very transparent. Here we will further comment on two examples where this effective description is useful.

Let us first consider noncommutative super Yang-Mills (NCSYM) theories at finite temperature. The two loop free energy density can be written for $U(N)$ NCSYM_d as [16]

$$F(\beta; \theta) = F(\beta; \theta = 0) + C_{sc} g^2 N \int \frac{dp}{(2\pi)^{d-1}} \left[\frac{n_b(p)}{!_p} + \frac{n_f(p)}{!_p} \right]^2 + \int \frac{dp}{(2\pi)^{d-1}} \int \frac{dq}{(2\pi)^{d-1}} \frac{n_b(p)}{!_p} + \frac{n_f(p)}{!_p} \frac{n_b(q)}{!_q} + \frac{n_f(q)}{!_q} e^{i(p \cdot q)}; \quad (3:7)$$

where $C_{sc} = 16; 4; 1$ for theories with 16, 8 and 4 supercharges respectively [20], $!_p = \prod_j p_j$ and $n_{b(f)}(p) = (e^{-\beta p_j} - 1)^{-1}$ are the Bose-Einstein and Fermi-Dirac distribution functions.

It is interesting to notice how, for NC SYM theories, the ‘nonplanar’ part of the two-loop free energy [the last term in (3.7)] factorizes into the product of two independent loop contributions only linked through the noncommutative phase, much in the same fashion of $d = 4$ NCFT. Following Ref. [17] we can now integrate one of these loops to try to spot winding states (Fig. 4). When $d < 4$ again we can introduce d extra variables z_i to write

$$\begin{aligned} \int \frac{dq}{(2\pi)^{d-1}} \frac{n_b(q)}{i_q} + \frac{n_f(q)}{i_q} e^{i(p \cdot \pi)} &= \frac{1}{(2\pi)^2} \int \frac{dz_i}{(2\pi)^2 + (p^2 + z_i^2)} \\ &= \frac{1}{2^{d-2}} \int \frac{dz_i}{(2^{d-1} + 1)^2 + (p^2 + z_i^2)} \end{aligned} \quad (3.8)$$




Fig. 4: Dual channel description of the nonplanar thermal loop in $d = 3$ -like two-loop nonplanar vacuum diagram.

That is, we find a standard tower of d particles, restricted to odd winding numbers. From this expression we learn that, in general, the fermion loops will give rise to d particles with negative norm for even d . For the supersymmetric case, there is a cancellation with the tower coming from the bosonic loop and we find the projection onto odd winding numbers. We shall give a string theory explanation of this phenomenon in the next section.

Notice that the factorization of the two-loop free energy into contribution of two independent loops is not a general property of any quantum field theory. In particular, for $d = 3$ NCFT the integrand of $F(\pi; \pi)$ does not have this property even in the massless case. What is special about NC SYM is the fact that many of the individual diagrams contributing to (3.7) have loops with two external insertions, so one would need at least one Feynman parameter in order to formally ‘integrate out’ the loop, along the lines of the general discussion above. Yet, the complete two-loop diagram shows factorized form and one can introduce effective d particles with standard Feynman rules (apart from the negative-norm feature in the fermionic case).

The reason behind this simplification is two-fold. First, gauge symmetry relates the $d = 3$ -like diagrams to the $d = 4$ -like diagrams. Second, the theory is massless, so that the effective coupling (2.6) is a momentum-independent pure number, and thus both $d = 4$ -like loops

are completely disentangled from the kinematical point of view. Therefore, this factorization is, in principle, specific of two-loop diagrams in massless theories whose symmetries can relate all diagrams, contributing to a given physical quantity, to $d=4$ -like ones. Another example, considered in [17], is the massless $\mathcal{N}=4$ Wess-Zumino model, where supersymmetry plays the relevant role. In the massless limit the $\mathcal{N}=4$ Wess-Zumino model reduces itself to a supersymmetric version of $d=4$ NCFT. Thus the factorization of the two-loop free energy follows from the factorization of the corresponding diagram in $d=4$ theory under the substitution $n_b(p) \rightarrow n_b(p) + n_f(p)$. In the NC SYM case it is not supersymmetry, but rather gauge symmetry, the one playing the simplifying role, because the two-loop factorization is true already for non-supersymmetric noncommutative Yang-Mills theories [16].

4. Windings and closed strings

4.1. Heuristic considerations

Given that many NCFT derive from open-string theory in background B -fields in the SW limit, it is natural to associate the winding modes of the previous representations to closed strings in intermediate states. Namely, the structure of (2.5) is reminiscent of a closed-string tree-level propagator between boundary states of a D_{d-1} brane (Fig. 5). Heuristically, we expect

$$\langle j|p\rangle_{NP} = \sum_{\text{closed strings}} \int dz \frac{\langle j|p;z\rangle^2}{2\alpha' + (p^2 + z^2)} \xrightarrow{\text{SW}} \sum_l D_{d-1}|V_p\rangle \frac{1}{c_l} \langle V_p|j\rangle_{D_{d-1}} \quad (4.1)$$

whereas the planar diagram would be a low-energy limit of $D_{d-1}|V_p\rangle \langle V_p|j\rangle_{D_{d-1}}$, for suitably defined boundary states.




Fig. 5: Seiberg-Witten limit of the nonplanar two-point function in the closed-string channel.

There are various pieces of the previous tentative equation that fit nicely. First, the closed-string inverse propagator is

$$c_l = \frac{0}{2} g_{pp} + \frac{2\alpha'}{4\alpha'^2} + M_{cl}^2 \quad (4.2)$$

where g is the closed-string or sigma-model metric, to be distinguished from G or open-string metric. The precise relation is defined in [3]:

$$G = \frac{1}{g + 2^{-1} B_S} ; \quad = 2^{-1} \frac{1}{g + 2^{-1} B_A} \quad (4:3)$$

where by the subscripts S and A we indicate the symmetric and antisymmetric part respectively. We can take $g =$ in commutative directions, including the $d_7 = D - d$ Dirichlet/Dirichlet directions transverse to the D_{d-1} brane. On the other hand, in the noncommutative directions the SW scaling assigns

$$g \rightarrow \frac{1}{4^{-2} \alpha'} (\alpha')^2 \quad (4:4)$$

as $\alpha' \rightarrow 0$, with $G =$ and fixed. Therefore, the inverse propagator scales

$$c_1 = \frac{1}{8^{-2} \alpha'} (\alpha')^2 + (p)^2 + \alpha' p_7^2 + \alpha' M_{cl}^2 \quad (4:5)$$

and we see that our familiar combination $(\alpha')^2 + (p)^2$ scales together and dominates over the other terms in the SW limit, since $\alpha' M_{cl}^2 \rightarrow 0_{osc} \neq 0$. This is the main evidence for the stringy origin of the winding modes. Indeed, we have Neumann boundary conditions in the thermal circle, and therefore the closed-string cylinder can wind in this direction. On the other hand, there cannot be momentum flow through the Neumann directions unless it is explicitly inserted via the open-string vertices into the boundary states, but there is an arbitrary flow of momentum in Dirichlet/Dirichlet directions. In the noncommutative directions, having a nonzero B-field, one could have momentum flow induced just by the B-field. The boundary conditions set to zero only a linear combination of momentum and winding numbers. However, since we are assuming a noncompact D-brane in spatial directions, there are no winding modes in spatial directions and thus no extra momentum flow induced by the B-field.

If we are willing to naively neglect the nominally subleading terms in (4.5) we can almost get (4.1) with the coupling of the ψ -elds defined through the 'sum rule' over all closed-string elds j_i (the oscillator excitations)

$$\langle j_i j_j \rangle = \sum^X \langle h_{D_{d-1}}; V_{p_j} | h_{D_{d-1}}; V_{p_i} \rangle \quad (4:6)$$

in the SW limit, perhaps with appropriate powers of α' in front. According to this picture, the low-energy ψ -elds are not some low-lying closed-string modes that fail to decouple. In fact, the whole infinite tower of closed-string modes fails to decouple, but the interaction with the boundary states defines an effective coupling for the ψ -eld, which represents

the coherent exchange of an infinite number of closed-string excitations. Formally, the SW limit squeezes the complete tower of string excited states into an approximately continuous band, as compared to the gap of the winding modes

$$\frac{\text{Oscillator Gap}}{\text{Winding Gap}} \sim \frac{0}{2} \neq 0: \quad (4:7)$$

This is an interesting compromise between the general lore that the closed-string channel should be intractable whenever the open-string channel is simple [11], and the factual existence of the dual channel representation in terms of the fields.

Actually, the sum rule (4.6) is too naive. The first indication that something is missing in (4.6) is the fact that a naive attempt to associate the z_2 degrees of freedom with Dirichlet/Dirichlet momenta p_2 in the D-brane codimension fails quantitatively, because in general $d_2 \notin d^2 = 2 + 2N - d$ for a D_{d-1} brane. The resolution of the puzzle amounts to recognize that one cannot simply neglect $\alpha^2 M_{cl}^2$ in the closed-string propagator, as compared to α'^2 , even in the low-energy SW limit, because there are an infinite number of states contributing to the sum. In other words, the truncated sum rule (4.6) is not convergent in general.

The second reason for concern lies in the definition of the low-energy effective coupling g as a proper effective vertex. In the full string theory diagram, the boundary states with open-string insertions have moduli (the Kobayashi-Nielsen parameters) that must be integrated over. Therefore, the stringy diagram does not have in general the structure of an ordinary tree-level exchange, since both vertices are convoluted in an integral over Kobayashi-Nielsen parameters. Only at the boundaries of the moduli space, when the string diagram degenerates into proper field theory diagrams, one finds standard Feynman rules. This means that the correct sum rule replacing (4.6) must hold for the integrand over moduli space (including the Kobayashi-Nielsen moduli).

4.2. Open-string channel

In the remainder of this section we obtain the correct sum rule by careful consideration of the general one-loop open-string diagram. This is a weighted sum over spin structures f, g , each one given by a path integral on the annulus with arbitrary vertex operator insertions on both the inner () and outer (+) boundaries

$$\begin{aligned} A &= \int_C \int_C \int_0^1 \frac{dZ}{2} \int_0^1 [dy] \int_{Y^*} V(p; y) \\ &= (\alpha')^{N_0} G_s^{N-2} \int_C \int_C \int_0^1 \frac{dZ}{2} \int_0^1 [dy] V(p; y;); \end{aligned} \quad (4:8)$$

where r is the modulus of the annulus, y_i are the Kobayashi-Nielsen parameters in each boundary, $V(p; y; \epsilon)$ is the normalized correlator of vertex operators in the spin structure ϵ , and $Z(\epsilon)$ is the normalization, i.e. the path integral without vertex insertions. The power of G_s comes from the normalization of the vertex operators with the open-string coupling defined in [3]

$$G_s = g_s \frac{\det G}{\det(g + 2\alpha' B)}^{\frac{1}{2}} = \frac{(\alpha')^{\frac{4-d}{2}}}{(2\pi\alpha')^{\frac{d-3}{2}}} g_{YM}^2 \quad (4.9)$$

The complete amplitude has an appropriate power of the Regge slope α' so that the SW limit with fixed g_{YM} leads to the field-theoretical expression of the amplitude. This depends on the overall dimension of the amplitude and the number of insertions.

The function $V(p; y; \epsilon)$ is a polynomial in external field polarizations, the fermionic Green function G_F and derivatives of the bosonic Green function G_B on the annulus with appropriate B -dependent boundary conditions, times the contraction of the tachyonic part of the vertices

$$V(p; y; \epsilon) = P(p; G_F; G_B) e^{P \cdot Q} \quad (4.10)$$

This Green function can be parametrized completely by the open-string metric G and noncommutativity parameters θ , except for a single constant term from the purely bosonic component [21], which contributes to nonplanar diagrams and depends explicitly on the sigma model metric g . If we separate this contribution from (4.10) we can write

$$V(p; y; \epsilon) = \bar{V}(p; y; \epsilon) \exp \left[-\frac{1}{2} p^\mu p^\nu \theta_{\mu\nu} \right] \quad (4.11)$$

where $p_{\text{cyl}}^\mu = \sum P p^\mu = \sum P p^\mu$ is the total momentum circulating in the closed-string channel. We recognize this term as the standard kinetic term in (4.2).

If we assume, for simplicity, that the external insertions are space-time bosons with no thermal frequency, i.e. we have a purely static bosonic correlator, then the world-sheet partition sum can be written directly in operator form as

$$Z(\epsilon) = \frac{1}{\text{Vol}_g} \text{Tr}_{\text{open}} S e^{-\int \mathcal{H}} \quad ; \quad (4.12)$$

where S is a piece of the GSO projector, $\frac{1}{2}$ or $\frac{1}{2}(-1)^F$ depending on the spin structure, \mathcal{H} is the open-string world-sheet hamiltonian, and we have normalized by the volume in the open-string metric G . The temperature-dependent part of (4.12) is unaffected by the B -field, as long as we keep $B_{0i} = 0$. Therefore, it has the form

$$Z(\epsilon) = \int d^d p (p^0)^2 = \frac{1}{2} \int d^d p p^2 \quad ; \quad (4.13)$$

with n integer in those spin structures running space-time bosons in the loop, and half-integer in those running space-time fermions. From here we can read off the relation between the annulus modular parameter and the low-energy Schwinger parameter of the field theory expressions in eq. (3.1) of the previous section; it is

$$t = \tau^2 : \tag{4.14}$$

For the comparison with the low-energy expression, it is useful to work with Koba-Nielsen parameters normalized to unity, $y = x$, so that a further factor of $N = (t = \tau^2)^N$ appears for a total of N insertions. Consistency of the SW low-energy limit requires that, for an appropriate choice of N_0 , the field-theoretical amplitude is obtained as

$$A_{\text{NCFT}} = \lim_{\text{SW}} (t = \tau^2)^{N_0 + \frac{4-d}{4}N} g_{\text{YM}}^N \int_0^1 dt t^{N-1} \int_0^1 dx \int_{\mathbb{Z}} \frac{t}{\tau} \mathcal{V}_p(x; \frac{t}{\tau}) : \tag{4.15}$$

The precise details of this limit in various examples of the bosonic theory at zero temperature can be found in recent papers [8][9][10][11][12][13]. The important feature is that the SW limit is dominated by massless open strings (in the bosonic examples one is forced to discard the open-string tachyon by hand). Thus, in comparing with (3.1), we must set $M = 0$ and interpret the normalized Koba-Nielsen parameters $x = y =$ as Feynman parameters of the field theory diagram.

On general grounds, the massless open-string dominance means that we do not expect the closed-string channel expression to be simple, in the sense of being saturated by a finite number of closed-string fields.

4.3. The closed-string channel sum rule

Ideally, we would like to specify explicitly, in the closed-string Fock space, the boundary states appearing in eq. (4.1). This is a very complicated task in general, and can be carried out in detail only for the 'vacuum' boundary states $\mathcal{D}_{\text{diag}}$ without open-string vertex insertions. On the other hand, we can obtain explicitly the overlap in (4.1), by direct modular transformation of the open-string channel expression (4.8).

In order to keep track of the right normalization of winding modes, we perform a Poisson resummation of the discrete frequency sums

$$\frac{1}{n} \sum_n e^{i\pi n^2 \tau} = (4\tau)^{-1/2} \sum_{n \in \mathbb{Z}} e^{-\frac{2\pi n^2}{4\tau}} (1)^{F'} \tag{4.16}$$

where F' is the space-time fermion number in the open-string channel. It is correlated with the closed-string sector in the cylinder channel in such a way that $F = 0$ corresponds

in terms of which, we have

$$A_{\text{NCFT}} = \lim_{\text{SW}} \int_{\mathcal{X}} g_{\text{YM}}^N \int_{\mathcal{C}} (0)^{N_0 + \frac{4-d}{4}N} \int_0^1 ds s^{d-2} \int_0^1 [dx] \int_{\mathcal{D}} e^{s[2,2+(p)^2]} U,; \int_{\mathcal{D}_d} V_{p^+}; \mathbf{x}^+ e^{s(2^{0M_{cl}})^2} \int_{\mathcal{D}_d} V_p; \mathbf{x} \quad (4.19)$$

with $U,;$ is the Atiyah-Witten phase. We can recast this expression in the form of eq. (3.2)

$$A_{\text{NCFT}} = \lim_{\text{SW}} W_{\text{NC}} \int_{\mathcal{X}} g_{\text{YM}}^N \int_0^1 ds s^{N + \frac{d-2}{2}} \int_{\mathcal{D}} e^{s[2,2+(p)^2]} \int_0^1 [dx] F(s;p;\mathbf{x}) \quad (4.20)$$

where W_{NC} is the global noncommutative phase of the diagram given by Eq. (3.3), up to numerical constants. This expression implies a ‘sum rule’ for the function $F(s;p;\mathbf{x})$:

$$W_{\text{NC}} F(s;p;\mathbf{x}) = \lim_{\text{SW}} (0)^{N_0 + \frac{4-d}{4}N} s^{\frac{2-d}{2}+N} \int_{\mathcal{D}} e^{s(2^{0M})^2} \int_{\mathcal{D}_d} V_{p^+}; \mathbf{x}^+ \int_{\mathcal{D}} e^{s(2^{0M})^2} \int_{\mathcal{D}_d} V_p; \mathbf{x} \quad (4.21)$$

where the sum over closed-string states j_i runs over all oscillator degrees of freedom of the closed strings in the bulk. In comparing (4.21) with the field-theoretical expression in (3.4), we must take into account that $M = 0$ and (3.4) was derived for a purely bosonic loop, hence there is no non-trivial Atiyah-Witten phase in (3.4). Furthermore, (4.21) was derived under the assumption that external states were bosonic and static, i.e. external momenta have vanishing time components. That explains the absence of the phase $\exp[i \int_{\mathcal{P}} x_a (Q_a^0)^2]$ in (4.21).

We conclude this subsection with some observations on the interpretation of (4.21):

i) In terms of the dimensionless closed-string modulus, the SW limit in (4.21) takes $\alpha' \rightarrow 0$. In this region of moduli space, the infinite tower of closed-string fields contributes to the sum rule, which is by no means saturated by a few closed-string fields. This was already obvious from the fact that the open-string channel expression was saturated by massless open strings.

ii) The sum rule (4.21) replaces the naive one in (4.6). One of the defects of (4.6), the mismatch between the powers of the Schwinger parameter and the true number of transverse dimensions of the brane, is resolved by noticing that the sum rule includes a non-trivial power of $s^{\frac{d-2}{2}}$, together with an explicit exponential kernel, which gives back the field-theoretical measure in (3.2) in the SW limit, and ensures the convergence of the sum over closed-string states. In fact, since this limit takes $\alpha' \rightarrow 0$, the way to

evaluate the infinite sum over states in (4.21) is to perform a modular transformation back to the open-string variables. In this process we get the appropriate powers of the Schwinger parameter from the modular anomaly of the oscillator traces (Jacobi's theta functions).

This discussion makes also manifest the formal character of the extra 'bulk dimensions' $d^2 = 2N + 2$ of the winding fields, something already clear from the fact that d^2 depends on the number of insertions in the loop. We see that, in those models with a string-theory embedding, there is a 'bulk' codimension $d_2 = D - d$, but its relation with d^2 is rather indirect.

iii) Another deficiency of (4.6), the absence of Koba-Nielsen parameters, is remedied in (4.21). The interpretation of the full diagram in the NCFT as a tree-level exchange of fields was all right provided we make a further convolution of the vertices with Feynman parameters. We now understand this feature as a residue of the full string picture, since Koba-Nielsen parameters map consistently to Feynman parameters in the SW limit. Therefore, the field picture of the NCFT mimics closely the structure of the closed-string channel in the full string theory.

iv) The sum rule (4.21) holds for the integrand of the moduli-space integral. Therefore, it holds independently of the possible occurrence of open- or closed-channel tachyons in the full string theory. This is in contrast with (4.6), which would be invalidated by open-string tachyons, and perhaps also by closed-string tachyons. It would be very interesting to study particular examples in detail to see the interplay between the various open/closed tachyons that could appear, including the finite temperature Hagedorn tachyon.

v) Our discussion is tailored to the case of thermal amplitudes. However, it is clear that the general features generalize to other toroidal compactifications with various degrees of supersymmetry.

4.4. An illustrative example

Unlike Eq. (4.6), the sum rule (4.21) is valid point by point in the $(s; x)$ moduli space. As a consequence, it is well defined even for tachyonic theories for which the integrated expressions would diverge due to the contribution coming from the moduli space boundaries. This being so, we can illustrate our sum rule (4.21) by considering the simplest possible example and take the two-point function of open-string tachyons on a D_{d-1} brane of the $D = 26$ critical bosonic string theory. In order to avoid unnecessary complications we will consider the static amplitude where incoming states do not carry time-components of the momenta. Thus, the amplitude in the open-string channel can be

written as [9][10][13]

$$A(p; p)_{\text{tachyon}} = G_s \int_0^1 \frac{dx}{2} (4 - x^2)^{\frac{d}{2}} \frac{i}{2} \int_0^1 \frac{dx}{2} e^{-\frac{1}{2} x_{12}^2} e^{-\frac{i}{2} x_{12}} e^{-\frac{0}{2} p \cdot G} e^{-\frac{0}{2} p \cdot G}$$

where we have defined $x_{12} = x^+ - x^-$. Comparing this expression with Eqs. (4.8) and (4.10) we can read both $Z(\tau)^{\text{osc}}$ and $\bar{V}(p; x; \tau)$, in terms of which the overlap (4.17) is expressed,

$$Z(\tau)^{\text{osc}} = \frac{i}{2} \int_0^1 \frac{dx}{2} e^{-\frac{1}{2} x_{12}^2} e^{-\frac{i}{2} x_{12}} e^{-\frac{0}{2} p \cdot G} e^{-\frac{0}{2} p \cdot G}$$

Switching from τ to the closed-string modular parameter $\tau_2 = 2\tau = \tau_2$ and performing the inversion on the modular functions we find for the partition function of the oscillators

$$Z\left(\frac{2}{\tau_2}\right)^{\text{osc}} = \frac{i}{2} \int_0^1 \frac{dx}{2} e^{-\frac{1}{2} x_{12}^2} e^{-\frac{i}{2} x_{12}} e^{-\frac{0}{2} p \cdot G} e^{-\frac{0}{2} p \cdot G}$$

whereas the function $\bar{V}(p; x; \tau)$ is written

$$\bar{V}\left(p; x; \frac{2}{\tau_2}\right) = \frac{i}{2} \int_0^1 \frac{dx}{2} e^{-\frac{1}{2} x_{12}^2} e^{-\frac{i}{2} x_{12}} e^{-\frac{0}{2} p \cdot G} e^{-\frac{0}{2} p \cdot G}$$

Using Eq. (4.17) we can now obtain the expression for the overlap, namely

$$D_{d-1}(V_{p^+}; x^+) e^{-\frac{0}{2} p \cdot G} D_{d-1}(V_p; x) = e^{-\frac{1}{2} x_{12}^2} \frac{i}{2} \int_0^1 \frac{dx}{2} e^{-\frac{i}{2} x_{12}} e^{-\frac{0}{2} p \cdot G} e^{-\frac{0}{2} p \cdot G}$$

where we have used the on-shell condition for the external tachyons, $p \cdot G = 1 = 0$ and also the relation $\int_0^1 dx (0j) = 2^{-3}(\dots)$. Actually, the modular functions can be rewritten using their product representations. Expanding the resulting infinite products in power series of $e^{-2\tau_2}$ we finally arrive at

$$\begin{aligned} & D_{d-1}(V_{p^+}; x^+) e^{-\frac{0}{2} p \cdot G} D_{d-1}(V_p; x) = e^{-\frac{1}{2} x_{12}^2} \frac{i}{2} \int_0^1 \frac{dx}{2} e^{-\frac{i}{2} x_{12}} e^{-\frac{0}{2} p \cdot G} e^{-\frac{0}{2} p \cdot G} \\ & = e^{-\frac{1}{2} x_{12}^2} \prod_{k=1}^{\infty} (1 - e^{-2k\tau_2})^{-20} \prod_{k=1}^{\infty} (1 - e^{-(2k-1)\tau_2})^{-2} e^{2ix^+} e^{-2ix^-} \\ & = \sum_{n=0}^{\infty} D(n) C_n^+(x^+) e^{-\frac{1}{2} x_{12}^2} \sum_{n=0}^{\infty} C_n(x^-) : \end{aligned} \tag{4.22}$$

where $M_n^2 = \frac{4}{\alpha'}(n-1)$ is the mass of the level- n oscillator states, and $\rho_D(n)$ is the level-density of those states with non-vanishing coupling to the D_{d-1} brane. Notice that, because of the structure of the product representation for the modular functions, the coefficient of $e^{-2(n-1)\tau}$ in the series always factorizes into contributions from the two different boundaries, $C_n(x)$, weighted by the level-density, $\rho_D(n)$. Comparing this expression with the sum rule (4.21) we read-off the couplings of a level- n closed-string state $|j_n\rangle$ to the boundary state with an external open-string tachyon insertion

$$\langle D_{d-1}; V_{p^+}; x_n | E = C_n(x) : \quad (4.23)$$

In this example we explicitly see how the coupling between $|j_n\rangle$ and the boundary state with a tachyon insertion is in general nonvanishing for all values of n and thus all closed-string oscillator levels run in the cylinder. As a consequence, the effective ϕ -elds cannot be seen as some undecoupled winding string state in the SW limit, but rather as a superposition of all massive closed-string modes with coherent couplings to the elementary quanta of the NCFT.

5. Concluding remarks

In the present paper we have tried to identify the stringy connection of the recently conjectured winding modes emerging in NCFT [17]. We have seen how a thermal loop in NCFT can be represented in a ‘dual channel’ picture as an infinite tower of tree-level exchanges of some effective ϕ -elds ϕ , with masses proportional to j^2 ($j \in \mathbb{Z}$) and kinetic term $\partial^2 \phi$ in the effective action. The scaling of the masses of these ϕ -elds with the length of the euclidean time suggests a winding mode interpretation for them.

In many respects these ϕ -elds are similar to the ϕ -elds introduced in Refs. [5][6]. It is important to notice however that there are a number of differences. First of all, in the thermal case we have not just one, but an infinite tower of effective ϕ -elds replacing the thermal loop in nonplanar amplitudes. As a consequence, one is able to replace the whole thermal loop by a tree-level exchange of these ϕ -elds and not just the high energy part as in [5]. Most importantly, the ϕ -elds have nonstandard Feynman rules. If the nonlocality in NCFT is just encoded in a nonpolynomial dependence of the interaction vertices on the incoming momenta, in the perturbation theory for the effective ϕ -elds the interaction vertices are convoluted in finite dimensional integrals over the Feynman parameters of the original diagram.

Actually, both features, winding-like masses and integration over the relative moduli of the interaction vertices, strongly suggest a stringy interpretation. We have found that such interpretation exists in those models which can be obtained from a D-brane theory

in the presence of a constant B - field in the SW limit⁵. In this case, we found that the scaling of the masses of the α' -fields with the length of the thermal circle is a residue of winding closed-string states, whereas the convolution over the vertex moduli descend from the integration over the Kobayashi-Nielsen moduli of the D-brane boundary states.

It is however important to notice that the 'winding states' identified in [17] have no simple interpretation in terms of individual string states that fail to decouple in the SW limit. On the contrary, the α' -fields have to be considered formal devices to represent a coherent coupling of an infinite number of closed-string states. These fields have effective couplings to the ordinary fields that can be derived from the elementary coupling of closed strings to D-brane boundaries via 'sum rules' involving the full tower of closed-string oscillator modes in the bulk. This is a rather unusual picture, and essentially is telling us that the SW limit is not an ordinary low-energy limit in the closed-string channel since all massive states are squeezed below the gap of the winding modes. As a consequence, the resulting NCFT present a degenerate version of the open/closed string duality of the original string theory: the ordinary 'open' representation of the Feynman diagram in NCFT and the 'closed' dual channel in terms of the winding α' -fields.

Acknowledgements

It is a pleasure to thank Luis Alvarez-Gaume, Jan de Boer, Karl Landsteiner, Esperanza Lopez, Niels Obers and Eliezer Rabinovici for useful discussions. G.A. warmly thanks the Spinoza Instituut and specially Gerard 't Hooft for hospitality. J.G. is partially supported by CERN and by Spanish Science Ministry Grant AEN98-043 and Catalan Science Foundation Grant GC 1998 SGR. The work of M.A.V.M. has been supported by FOM (Fundamenteel Onderzoek van de Materie) and University of the Basque Country Grants UPV 063.310-EB187/98 and UPV 172.310-G02/99 and Spanish Science Ministry Grant AEN99-0315. M.A.V.M. wishes also to thank CERN Theory Division for hospitality during the completion of this work.

⁵ The phenomenon of UV/IR mixing is also present in nonrelativistic noncommutative field theories for which no obvious embedding into a string theory seems to exist [24].

References

- [1] M. R. Douglas and C. M. Hull, *JHEP* 02 (1998) 008. (hep-th/9711165)
- [2] A. Connes, M. Douglas and A. Schwarz, *JHEP* 02 (1998) 003. (hep-th/9711162)
- [3] N. Seiberg and E. Witten, *JHEP* 09 (1999) 032. (hep-th/9908142)
- [4] T. Filk, *Phys. Lett. B* 376 (1996) 53;
 J.C. Varilly and J.M. Gracia-Bondía, *Int. J. Mod. Phys. A* 14 (1999) 1305 (hep-th/9804001);
 C.P. Martin and D. Sanchez-Ruiz, *Phys. Rev. Lett.* 83 (1999) 476 (hep-th/9903077);
 M.M. Sheikh-Jabbari, *JHEP* 06 (1999) 015 (hep-th/9903107);
 T. Krajewski and R. Wulkenhaar, *Perturbative quantum gauge fields on the noncommutative torus*, hep-th/9903187;
 D. Bigatti and L. Susskind, *Magnetic fields, branes and noncommutative geometry*, hep-th/9908056;
 I.Ya. Aref'eva, D.M. Belov and A.S. Koshelev, *Phys. Lett. B* 476 (2000) 431 (hep-th/9912075);
 M. Hayakawa, *Perturbative analysis on infrared aspects of noncommutative QED on R^4* , hep-th/9912094; *Perturbative analysis on infrared and ultraviolet aspects of noncommutative QED on R^4* , hep-th/9912167;
 I.Ya. Aref'eva, D.M. Belov and A.S. Koshelev, *A note on UV/IR for noncommutative complex scalar field*, hep-th/0001215;
 A. Matulis, L. Susskind and N. Toumbas, *The IR/UV connection in noncommutative gauge theories*, hep-th/0002075.
- [5] S. Minwalla, M. Van Raamsdonk and N. Seiberg, *Noncommutative perturbative dynamics*, (hep-th/9912072).
- [6] M. Van Raamsdonk and N. Seiberg, *JHEP* 03 (2000) 035. (hep-th/0002186)
- [7] A. Hashimoto and N. Itzaki, *Phys. Lett. B* 465 (1999) 142 (hep-th/9907166);
 J.M. Maldacena and J.G. Russo, *JHEP* 09 (1999) 025 (hep-th/9908134);
 M. Alishahiha, Y. Oz and M.M. Sheikh-Jabbari, *JHEP* 11 (1999) 007 (hep-th/9909215);
 A. Hashimoto and N. Itzaki, *JHEP* 12 (1999) 007 (hep-th/9911057);
 T. Harmark and N.A. Obers, *JHEP* 03 (2000) 024. (hep-th/9911169)
- [8] O. Andreev and H. Dom, *Diagrams of Noncommutative 3 Theory from String Theory*, hep-th/0003113.
- [9] Y. Kim and S. Lee, *UV/IR mixing in noncommutative field theory via open string loops*, hep-th/0003145

- [10] A .B ilal, C .S .C hu and R .R usso, String theory and noncom m utative eld theories at one loop, hep-th/0003180.
- [11] J .G om is, M .K leban, T .M ehen, M .R angam ani and S . Shenker, Noncom m utative gauge dynam ics from the string worldsheet, hep-th/0003215.
- [12] A .R a jaram an and M .R ozali, Noncom m utative gauge theory, divergences and closed strings, hep-th/0003227.
- [13] H .L iu and J .M ichelson, Stretched strings in noncom m utative eld theory, hep-th/0004013.
- [14] J .L .F .B arbon and E .R abinovici, JHEP 04 (1999) 015. (hep-th/9910019)
- [15] R .G .C ai and N .O hta, On the therm odynam ics of large-N noncom m utative super Yang-M ills theory, hep-th/9910092.
- [16] G .A rcioni and M .A .V azquez-M ozo, JHEP 01 (2000) 028. (hep-th/9912140)
- [17] W .F ischler, J .G om is, E .G orbatov, A .K ashani-Poor, S .Paban and P .Pouliot, Evidence for winding states in noncom m utative quantum eld theory, hep-th/0002067.
- [18] W .F ischler, E .G orbatov, A .K ashani-Poor, R .M cNees, S .Paban and P .Pouliot, The interplay between and T , hep-th/0003216.
- [19] J .I .K apusta, Therm alF ield Theory, C am bridge 1989.
- [20] M .A .V azquez-M ozo, Phys. Rev. D 60 (1999) 106010. (hep-th/9905030)
- [21] A .A bouelsaood, C .G .C allan, C .R .N appi and S .A .Y ost, Nucl. Phys. B 280 (1987) 599.
- [22] M .A .V azquez-M ozo, Phys. Lett. B 388 (1996) 494. (hep-th/9607052)
- [23] J .J .A tick and E .W itten, Nucl. Phys. B 310 (1988) 291.
- [24] Joaquin G om is, K .L andsteiner and E .Lopez, work in progress.