

STATUS OF THEORETICAL $\bar{B} \rightarrow X_s \gamma$ AND $\bar{B} \rightarrow X_s L^+ L^-$ ANALYSES

MIKOŁAJ MISIAK

Theory Division, CERN, CH-1211 Geneva 23, Switzerland

Status of the theoretical $\bar{B} \rightarrow X_s \gamma$ and $\bar{B} \rightarrow X_s l^+ l^-$ analyses is reviewed. Recently completed perturbative calculations are mentioned. The level at which non-perturbative effects are controlled is discussed.

The present talk will be devoted to discussion of the SM predictions only. Let us begin with $\bar{B} \rightarrow X_s \gamma$. Since the completion of NLO QCD calculations¹ 4 years ago, many new analyses have been performed. They include evaluation of non-perturbative Λ^2/m_c^2 corrections² and the leading electroweak corrections.^{3,4} None of these results exceeds half of the overall $\sim 10\%$ uncertainty, and there are cancellations among them. In consequence, the prediction for $BR[\bar{B} \rightarrow X_s \gamma]$ remains almost unchanged: $(3.29 \pm 0.33) \times 10^{-4}$. This prediction agrees very well with the measurements of CLEO⁵, ALEPH⁶ and BELLE⁷, whose combined result is $(3.21 \pm 0.40) \times 10^{-4}$.

The dominant contribution to the perturbative $b \rightarrow s \gamma$ amplitude originates from charm-quark loops. After including QCD corrections, the top-quark contribution is less than half of the charm-quark one, and it comes with an opposite sign. This fact should be remembered when one attempts to extract $|V_{ts}|$ from $b \rightarrow s \gamma$. The u -quark contribution is suppressed with respect to the charm one by $|V_{ub}V_{us}|/|V_{cb}V_{cs}| \simeq 2\%$.

The results of CLEO, ALEPH and BELLE have to be understood as the ones with subtracted intermediate ψ background, i.e. the background from $\bar{B} \rightarrow \psi X_s$ followed by $\psi \rightarrow X' \gamma$. This background gives more than 4×10^{-4} in the “total” BR, but gets suppressed when only high-energy photons are counted. A rough estimate⁸ of the effect of the photon energy cutoff on this background can be made when X_s in $\bar{B} \rightarrow \psi X_s$ is assumed to be massless, and the non-zero spin

of ψ is ignored. Then,^a the intermediate ψ background is less than 5%, for the present experimental cutoff $E_\gamma > 2.1$ GeV in the \bar{B} -meson rest frame.^b However, the background grows fast when the cutoff goes down.

The photon energy cutoff *will* have to go down by at least 200 or 300 MeV in the future. With the present one, non-perturbative effects related to the unknown \bar{B} -meson shape function⁴ considerably weaken the power of $b \rightarrow s \gamma$ for testing new physics. For the same reason, future measurements of $\bar{B} \rightarrow X_s \gamma$ should rely as little as possible on theoretical predictions for the precise shape of the photon spectrum above $E_\gamma \sim 2$ GeV.

A systematic analysis of non-perturbative effects in $\bar{B} \rightarrow X_s \gamma$ at order $\mathcal{O}(\alpha_s(m_b))$ is missing. There is no straightforward method to perform such an analysis, because there is no obvious operator product expansion for the matrix elements of the 4-quark operators, in the presence of one or more hard gluons (i.e. the gluons with momenta of order m_b). At present, we have only intuitive arguments to convince ourselves that such non-perturbative effects are probably significantly smaller than the overall $\sim 10\%$ theoretical uncertainty in $BR[\bar{B} \rightarrow X_s \gamma]$, when the energy cutoff is between 1 and 2 GeV, and when the intermediate $\psi^{(\prime)}$ contribution(s) are subtracted.

^a The $\psi \rightarrow X \gamma$ spectrum is available from the ancient MARK II data⁹. New results are expected soon from the BES experiment in Beijing.

^b A further suppression (to less than 1%) is found when X_s is not treated as massless but the measured¹⁰ mass spectrum is used.

As far as the decay $\bar{B} \rightarrow X_s l^+ l^-$ is concerned (for $l = e$ or μ), the best control over non-perturbative effects can be achieved in the region of low dilepton invariant mass ($\hat{s} \equiv m_{l^+ l^-}^2 / m_b^2 \in [0.05, 0.25]$). The present prediction¹¹ for the branching ratio integrated over this domain is $(1.46 \pm 0.19) \times 10^{-6}$. The quoted uncertainty is only the perturbative one. The non-perturbative Λ^2 / m_c^2 and Λ^2 / m_b^2 contributions¹² have been included in the central value. They are around 2% and 5%, respectively.

A calculation of $\mathcal{O}(\alpha_s)$ terms in all the relevant Wilson coefficients $C_i(m_b)$ has been recently completed¹¹, up to small effects originating from 3-loop RGE evolution of C_9 . However, the perturbative uncertainty in the above-mentioned prediction remains close to $\sim 13\%$, because 2-loop matrix elements of the 4-quark operators are unknown.

The low- \hat{s} branching ratio is as sensitive to new physics as the forward-backward or energy asymmetries, i.e. $\sim 100\%$ effects are observed when $C_7(m_b)$ changes sign.

The background from $\bar{B} \rightarrow \psi X_s$ followed by $\psi \rightarrow l^+ l^-$ is removed by the cutoff $\hat{s} < 0.25$. Analogous contributions from virtual $c\bar{c}$ states are, in principle, included in the calculated Λ^2 / m_c^2 correction. An independent verification of this fact can be performed with help of dispersion relations and the factorization approximation.¹³ Indeed, for $\hat{s} < 0.25$, the difference between results obtained with help of the two methods is quite small, and can be attributed to higher-order perturbative effects.

On the other hand, the background from $\bar{B} \rightarrow \psi X_s$ followed by $\psi \rightarrow X' l^+ l^-$ has never been studied. Most probably, for $\hat{s} < 0.25$, it is less important than the analogous background in the case of $\bar{B} \rightarrow X_s \gamma$. Experiment-based calculations of these backgrounds are awaited, because they are essential for performing theoretical estimates of similar non-perturbative contributions from other $c\bar{c}$ states.

References

1. K. Chetyrkin, M. Misiak and M. Münz, *Phys. Lett. B* **400**, 206 (1997); C. Greub, T. Hurth and D. Wyler, *Phys. Rev. D* **54**, 3350 (1996); K. Adel and Y.P. Yao, *Phys. Rev. D* **49**, 4945 (1994); A. Ali and C. Greub, *Phys. Lett. B* **361**, 146 (1995).
2. G. Buchalla, G. Isidori and S.J. Rey, *Nucl. Phys. B* **511**, 594 (1998); Z. Ligeti, L. Randall and M.B. Wise, *Phys. Lett. B* **402**, 178 (1997); A.K. Grant et al., *Phys. Rev. D* **56**, 3151 (1997); A. Khodjamirian et al., *Phys. Lett. B* **402**, 167 (1997); M.B. Voloshin, *Phys. Lett. B* **397**, 275 (1997).
3. P. Gambino and U. Haisch, hep-ph/0007259; K. Baranowski and M. Misiak, *Phys. Lett. B* **483**, 410 (2000); A. Strumia, *Nucl. Phys. B* **532**, 28 (1998); A. Czarnecki and W. Marciano, *Phys. Rev. Lett.* **81**, 277 (1998).
4. A.L. Kagan and M. Neubert, *Eur. Phys. J. C* **7**, 5 (1999).
5. S. Ahmed et al., hep-ex/9908022.
6. R. Barate et al., *Phys. Lett. B* **429**, 169 (1998).
7. M. Nakao, this conference; A. Abashian et al., preprint BELLE-CONF-0003.
8. A. Khodjamirian and M. Misiak, in preparation.
9. D.L. Scharre et al., *Phys. Rev. D* **23**, 43 (1981).
10. R. Balest et al., *Phys. Rev. D* **52**, 2661 (1995).
11. C. Bobeth, M. Misiak and J. Urban, *Nucl. Phys. B* **574**, 291 (2000).
12. G. Buchalla, G. Isidori and S.-J. Rey, *Nucl. Phys. B* **511**, 594 (1998); G. Buchalla and G. Isidori, *Nucl. Phys. B* **525**, 333 (1998); A.F. Falk, M. Luke and M.J. Savage, *Phys. Rev. D* **49**, 3367 (1994).
13. F. Krüger and L.M. Sehgal, *Phys. Lett. B* **380**, 199 (1996).