

## NNLO THRESHOLD RESUMMATION IN HEAVY FLAVOUR DECAYS

**U. Aglietti**<sup>\*)</sup>

CERN-TH, Geneva, Switzerland  
e-mail address: ugo.aglietti@cern.ch.

**G. Ricciardi**

Dipartimento di Scienze Fisiche,  
Università di Napoli “Federico II”  
and I.N.F.N., Sezione di Napoli, Italy  
e-mail address: giulia.ricciardi@na.infn.it

### Abstract

We present a NNLO evaluation of the QCD form factor resumming large logarithmic perturbative contributions in semi-inclusive heavy flavour decays.

<sup>\*)</sup> On leave of absence from Dipartimento di Fisica, Università di Roma I, Piazzale Aldo Moro 2, 00185 Roma, Italy.

# 1 Introduction

In this note we present a next-to-next-to-leading order (NNLO) evaluation of the QCD form factor resumming threshold logarithmic contributions in semi-inclusive heavy flavour decays.

In semi-inclusive processes, final gluon radiation is strongly inhibited in the phase space regions where the observed final state obtains its maximum energy, therefore opening the way to soft and collinear singularities. The perturbative calculation of the differential cross section is plagued, in that limit, by large logarithms. In order to improve the reliability of the perturbative calculation, these large logarithms need to be resummed.

Let us consider the rate of semi-inclusive heavy flavour decays. The Mellin  $N$ -moments of the rate contain double logarithmic contributions and have an expansion of the form:

$$\begin{aligned} \frac{1}{\Gamma_B} \Gamma_N(\alpha_S) &\equiv \int_0^1 dx x^{N-1} \frac{1}{\Gamma_B} \frac{d\Gamma}{dx}(x; \alpha_S) & (1) \\ &= 1 + \sum_{n=1}^{\infty} \sum_{m=0}^{2n} G_{nm} \alpha_S^n L^m + \sum_{l=1}^{\infty} \alpha_S^l R_N^{(l)} \\ &= 1 + G_{12} \alpha_S L^2 + G_{11} \alpha_S L + G_{10} \alpha_S + G_{24} \alpha_S^2 L^4 + G_{23} \alpha_S^2 L^3 + G_{22} \alpha_S^2 L^2 + \dots \\ &\quad + \alpha_S R_N^{(1)} + \alpha_S^2 R_N^{(2)} + \alpha_S^3 R_N^{(3)} + \dots, & (2) \end{aligned}$$

where

$$L \equiv \ln N. \quad (3)$$

$R_N(\alpha_S)$  is a remainder function, which does not contain large logarithms and has a perturbative expansion of the form:

$$R_N(\alpha_S) \equiv \sum_{l=1}^{\infty} \alpha_S^l R_N^{(l)}. \quad (4)$$

The running coupling is evaluated at a general renormalization scale  $\mu \neq Q$ , where  $Q$  is the hard scale:

$$\alpha_S = \alpha_S(\mu^2). \quad (5)$$

The logarithmic terms have an exponential structure, so one can write [1, 2]:

$$\frac{1}{\Gamma_B} \Gamma_N(\alpha_S) = C(\alpha_S) f_N(\alpha_S) \quad (6)$$

where the form factor  $f_N(\alpha_S)$  reads:

$$\begin{aligned} f_N(\alpha_S) &= \exp \left[ \sum_{n=1}^{\infty} \sum_{k=1}^{n+1} c_{nm} \alpha_S^n L^k \right] \\ &= \exp [c_{12} \alpha_S L^2 + c_{11} \alpha_S L + c_{23} \alpha_S^2 L^3 + c_{22} \alpha_S^2 L^2 + c_{21} \alpha_S^2 L + c_{34} \alpha_S^3 L^4 + \dots]. & (7) \end{aligned}$$

Note that the exponent contains only the first term  $\alpha_S L^2$  of the double-logarithmic series  $(\alpha_S L^2)^n$ . The advantage of the exponentiation is therefore that we can predict  $\ln f_N$  reliably for  $\alpha_S L \ll 1$ , that is, for a larger region than  $\alpha_S L^2 \ll 1$ , where the perturbative expansion of  $f_N$  holds. In fact, the other terms of  $(\alpha_S L^2)^n$  come purely from the expansion of the exponential function, as can be seen in formula (2).

The prefactor in (6) is the coefficient function, having an expansion in powers of  $\alpha_S$ :

$$C(\alpha_S) = 1 + \sum_{n=1}^{\infty} C_n \alpha_S^n = 1 + C_1 \alpha_S + C_2 \alpha_S^2 + \dots \quad (8)$$

The double sum in the exponent is usually organized as a series of functions, which resum “strips” in the  $(n, k)$  plane:

$$\begin{aligned} f_N(\alpha_S) &= \exp \left[ L g_1(\beta_0 \alpha_S L) + \sum_{n=2}^{\infty} \alpha_S^{n-2} g_n(\beta_0 \alpha_S L) \right] \\ &= \exp \left[ L g_1(\beta_0 \alpha_S L) + g_2(\beta_0 \alpha_S L) + \alpha_S g_3(\beta_0 \alpha_S L) + \alpha_S^2 g_4(\beta_0 \alpha_S L) + \dots \right]. \end{aligned} \quad (9)$$

The functions  $g_i(\lambda)$  have a power-series expansion:

$$g_i(\lambda) = \sum_{n=1}^{\infty} g_{in} \lambda^n. \quad (10)$$

where  $\lambda = \beta_0 \alpha_S L$ . They are all homogeneous functions:  $g_i(0) = 0$ . This property insures the normalization of the form factor:  $f_{N=1} = 1$ . The resummation at LO, NLO and NNLO is referred, respectively, to series in  $L(\alpha_s L)^n$ ,  $(\alpha_s L)^n$  and  $\alpha_s(\alpha_s L)^n$ .

We have computed the function  $g_3(\lambda)$ , which is necessary for the NNLO resummation. Not all the quantities determining  $g_3(\lambda)$  are known exactly.  $A_3$  is only known numerically from a fit of the three-loop Altarelli-Parisi (AP) splitting function to the known moments.  $B_2$  is unknown and we approximate it with the infrared-regular part of the two-loop AP splitting function. The coefficient functions and the remainder functions for radiative and semileptonic  $B$  decays have been computed to NLO in ref. [3]. A complete NNLO computation of the distributions requires also the knowledge of the two-loop coefficient function, which at present is unknown for any distribution. After the resummation of the threshold logarithms in the  $N$ -space, we have returned to the form factor in the  $x$ -space, by inverting  $f_N(\alpha_s)$  with analytic and numerical procedures.

## 2 QCD form factor at NNLO

Let us briefly describe the derivation of the NNLO form factor in the  $N$ -space. By NNLO accuracy, we mean the resummation of all the infrared logarithms up to and including

$$NNLO: \quad \alpha_S^n L^{n-1}. \quad (11)$$

The general expression of the form factor is:

$$\log f_N(\alpha_S) = \int_0^1 dz \frac{z^{N-1} - 1}{1-z} \left\{ \int_{Q^2(1-z)^2}^{Q^2(1-z)} \frac{dk^2}{k^2} A[\alpha_S(k^2)] + D[\alpha_S(Q^2(1-z)^2)] + B[\alpha_S(Q^2(1-z))] \right\}. \quad (12)$$

Let us remind one difference between annihilation processes, as Drell-Yan, and other processes, like the present one or DIS. In the annihilation processes, the initial partons reduce their momenta by irradiation, before actually annihilating; when  $\tau = Q^2/s \rightarrow 1$ , only the emission of soft partons is allowed. Likewise, in the other processes, as f.i. DIS, and in the limit  $x = Q^2/2Q \cdot p \rightarrow 1$ , only soft emission is allowed, before the scattering; however, after the scattering, the parton fragments with the only kinematical constraint of having a low virtuality and collinear emission is no longer forbidden.

The functions  $A(\alpha_S)$ ,  $D(\alpha_S)$  and  $B(\alpha_S)$  have a perturbative expansion in powers of  $\alpha_S$ :

$$\begin{aligned} A(\alpha_S) &= \sum_{n=1}^{\infty} A_n \alpha_S^n = A_1 \alpha_S + A_2 \alpha_S^2 + A_3 \alpha_S^3 + \dots; \\ D(\alpha_S) &= \sum_{n=1}^{\infty} D_n \alpha_S^n = D_1 \alpha_S + D_2 \alpha_S^2 + \dots; \\ B(\alpha_S) &= \sum_{n=1}^{\infty} B_n \alpha_S^n = B_1 \alpha_S + B_2 \alpha_S^2 + \dots. \end{aligned} \quad (13)$$

Let us observe, that in general the transverse momentum rule is a guess, as it has not been proven to such an accuracy. The universality of  $B_2$ , in general, is a debated problem. If we neglect the variation of the coupling with the scale (frozen coupling), we find logarithmic terms of the form:

$$A_1\alpha_S L^2, \quad A_2\alpha_S^2 L^2, \quad A_3\alpha_S^3 L^2, \dots \quad D_1\alpha_S L, \quad D_2\alpha_S^2 L, \dots \quad B_1\alpha_S L, \quad B_2\alpha_S^2 L, \dots \quad (14)$$

Then, to NNLO accuracy, one needs the first three terms in the expansion of  $A(\alpha_S)$  and the first two terms of the functions  $D(\alpha_S)$  and  $B(\alpha_S)$ .

The three-loop coupling, according to the definition given in the PDG [4], reads:

$$\alpha_s(\mu^2) = \frac{1}{\beta_0 \log \mu^2 / \Lambda^2} - \frac{\beta_1}{\beta_0^3} \frac{\log \log \mu^2 / \Lambda^2}{\log^2 \mu^2 / \Lambda^2} + \frac{\beta_1^2}{\beta_0^5} \frac{\log^2 \log \mu^2 / \Lambda^2 - \log \log \mu^2 / \Lambda^2 - 1}{\log^3 \mu^2 / \Lambda^2} + \frac{\beta_2}{\beta_0^4} \frac{1}{\log^3 \mu^2 / \Lambda^2}. \quad (15)$$

The asymptotic expansion of the coupling is basically an expansion in inverse powers of  $\log \mu^2 / \Lambda^2$ . The first three coefficients of the  $\beta$ -function, defined as

$$\frac{d\alpha_s}{d \log \mu^2} = -\beta_0 \alpha_S^2 - \beta_1 \alpha_S^3 - \beta_2 \alpha_S^4 - \dots, \quad (16)$$

read:

$$\begin{aligned} \beta_0 &= \frac{11C_A - 2n_F}{12\pi} = \frac{33 - 2n_F}{12\pi} = 0.87535 - 0.05305 n_F, \\ \beta_1 &= \frac{17C_A^2 - 5C_A n_F - 3C_F n_F}{24\pi^2} = \frac{153 - 19n_F}{24\pi^2} = 0.64592 - 0.08021 n_F, \\ \beta_2 &= \frac{2857 - 5033/9 n_F + 325/27 n_F^2}{128\pi^3} = 0.71986 - 0.140904 n_F + 0.003032 n_F^2. \end{aligned} \quad (17)$$

Let us note that  $\beta_0$  and  $\beta_1$  are renormalization-scheme independent, while  $\beta_2$  is not and we have given its value in the  $\overline{MS}$  scheme [5].

Integrating the  $\beta$ -function equation (16) on both sides, one obtains:

$$\alpha_S(k^2) = \alpha_S(Q^2) - \beta_0 \alpha_S^2(Q^2) \log \frac{k^2}{Q^2} - \beta_1 \alpha_S^3(Q^2) \log \frac{k^2}{Q^2} - \beta_2 \alpha_S^4(Q^2) \log \frac{k^2}{Q^2} + (\text{iterations}). \quad (18)$$

Substituting the above expression into (12), one sees that a  $\beta_0$ -insertion corresponds to the additional factor  $\alpha_S L$ , that of  $\beta_1$  to  $\alpha_S^2 L$  and that of  $\beta_2$  to  $\alpha_S^3 L$ :

$$\beta_0 : \alpha_S L, \quad \beta_1 : \alpha_S^2 L, \quad \beta_2 : \alpha_S^3 L. \quad (19)$$

Therefore, in the terms containing NNLO coefficients, the coupling can be replaced with the one-loop one and in the NLO terms the coupling can be replaced with the two-loop one, so that one has:

$$\begin{aligned} A(\alpha_S) &= A_1\alpha_{S,3L} + A_2\alpha_{S,2L}^2 + A_3\alpha_{S,1L}^3; \\ D(\alpha_S) &= D_1\alpha_{S,2L} + D_2\alpha_{S,1L}^2; \\ B(\alpha_S) &= B_1\alpha_{S,2L} + B_2\alpha_{S,1L}^2. \end{aligned} \quad (20)$$

Furthermore, the  $\beta_1^2$  term in  $A_2\alpha_{S,2L}^2$  can be neglected, as it is a N<sup>3</sup>LO contribution. After replacing eqs. (20) into eq. (12), one performs a straightforward integration over  $k^2$ , the gluon transverse momentum. The integration over  $z$ , the longitudinal gluon momentum, is easily done using the approximation [1]:

$$z^{N-1} - 1 \simeq -\theta \left( 1 - z - \frac{1}{n} \right). \quad (21)$$

This approximation misses the term proportional to  $A_1\zeta_2$ , where  $\zeta_2 = \pi^2/6$ , which can be obtained in the following way. Using the large- $N$  approximation derived in [6], one obtains, in the  $n$  variable:

$$\begin{aligned} \int_0^1 dz \frac{z^{N-1} - 1}{1-z} \int_{Q^2(1-z)^2}^{Q^2(1-z)} \frac{dk^2}{k^2} A_1 \alpha_S(k^2) &= \sum_{k=1}^{\infty} c_k \int_0^1 dz \frac{z^{N-1} - 1}{1-z} \log^k(1-z) \\ &\simeq \sum_{k=1}^{\infty} c_k \frac{(-1)^{k+1}}{k+1} \left[ \log^{k+1} n + \frac{\zeta_2}{2} k(k+1) \log^{k-1} n \right] \\ &= [\text{lo}] + [\text{mlo}]. \end{aligned} \quad (22)$$

Therefore:

$$[\text{mlo}] = \frac{\zeta_2}{2} \frac{\partial^2}{\partial (\log n)^2} [\text{lo}]. \quad (23)$$

We have used the variable  $n = N/N_0$  instead of  $N$ , where  $N_0 \equiv e^{-\gamma_E} = 0.561459\dots$ . Within this alternative representation, the terms proportional to  $\gamma_E$  and to  $\gamma_E^2$  disappear. This scheme is probably more accurate as Feynman diagram computation directly in  $N$ -space brings factors containing  $\Gamma(1-\epsilon)$  with  $D = 4 - 2\epsilon$  the space-time dimension.

The advantage of the variable  $N$  is that the total rate is directly reproduced by setting  $N = 1$ , while in the variable  $n$  it is given by  $f_{n=1/N_0}$ . These two variables differ by terms of higher order in  $\gamma_E$ .

The lowest-order term, computed within the approximation (21), reads:

$$[\text{lo}] = -\frac{A_1}{2\beta_0} \left[ \log \frac{s}{n^2} \log \log \frac{s}{n^2} + \log s \log \log s - 2 \log \frac{s}{n} \log \log \frac{s}{n} \right]. \quad (24)$$

One then obtains:

$$\begin{aligned} \log f_N(\alpha_S) &= -\frac{A_1}{2\beta_0} \left[ \log s/n^2 \log \log s/n^2 + \log s \log \log s - 2 \log s/n \log \log s/n \right] + \\ &+ \frac{\beta_0 A_2 - \beta_1 A_1}{2\beta_0^3} \left[ \log \log s - 2 \log \log s/n + \log \log s/n^2 \right] + \\ &- \frac{\beta_1 A_1}{4\beta_0^3} \left[ \log^2 \log s/n^2 - 2 \log^2 \log s/n + \log^2 \log s \right] + \\ &+ \frac{D_1}{2\beta_0} \left[ \log \log s/n^2 - \log \log s \right] + \frac{B_1}{\beta_0} \left[ \log \log s/n - \log \log s \right] \\ &- \frac{A_3}{4\beta_0^3} \left[ \frac{1}{\log s/n^2} - \frac{2}{\log s/n} + \frac{1}{\log s} \right] - \frac{A_1 \zeta_2}{2\beta_0} \left[ \frac{2}{\log s/n^2} - \frac{1}{\log s/n} - \frac{1}{\log s} \right] + \\ &- \frac{A_1 \beta_2}{8\beta_0^4} \left[ \frac{1}{\log s/n^2} - \frac{2}{\log s/n} + \frac{1}{\log s} \right] + \\ &+ \frac{A_2 \beta_1}{4\beta_0^4} \left[ 2 \frac{\log \log s/n^2}{\log s/n^2} - 4 \frac{\log \log s/n}{\log s/n} + 2 \frac{\log \log s}{\log s} + \frac{3}{\log s/n^2} - \frac{6}{\log s/n} + \frac{3}{\log s} \right] + \\ &- \frac{A_1 \beta_1^2}{4\beta_0^5} \left[ \frac{\log^2 \log s/n^2}{\log s/n^2} - 2 \frac{\log^2 \log s/n}{\log s/n} + \frac{\log^2 \log s}{\log s} + 2 \frac{\log \log s/n^2}{\log s/n^2} + \right. \\ &\quad \left. - 4 \frac{\log \log s/n}{\log s/n} + 2 \frac{\log \log s}{\log s} + \frac{1}{\log s/n^2} - \frac{2}{\log s/n} + \frac{1}{\log s} \right] + \\ &+ \frac{D_1 \beta_1}{2\beta_0^3} \left[ \frac{\log \log s/n^2}{\log s/n^2} - \frac{\log \log s}{\log s} + \frac{1}{\log s/n^2} - \frac{1}{\log s} \right] - \frac{D_2}{2\beta_0^2} \left[ \frac{1}{\log s/n^2} - \frac{1}{\log s} \right] + \\ &+ \frac{B_1 \beta_1}{\beta_0^3} \left[ \frac{\log \log s/n}{\log s/n} - \frac{\log \log s}{\log s} + \frac{1}{\log s/n} - \frac{1}{\log s} \right] - \frac{B_2}{\beta_0^2} \left[ \frac{1}{\log s/n} - \frac{1}{\log s} \right]. \end{aligned} \quad (25)$$

The next step is to write the above result as a function of the three-loop coupling. The inverse of (15) reads:

$$\log s = \frac{1}{\beta_0 \alpha_S} + \frac{\beta_1}{\beta_0^2} \log(\beta_0 \alpha_S) - \left( \frac{\beta_1^2}{\beta_0^3} - \frac{\beta_2}{\beta_0^2} \right) \alpha_S + O(\alpha_S^2), \quad (26)$$

where  $s$  is the square of the hard scale in unit of the QCD scale,

$$s \equiv \frac{Q^2}{\Lambda^2}. \quad (27)$$

After the replacement (26), terms of higher order with respect to NNLO are generated, which must be discarded. One finally makes the expansion in  $\gamma_E$  up to second order to obtain eq. (40).

The quantities  $A_1$  and  $A_2$  are known analytically [1, 7]:

$$A_1 = \frac{C_F}{\pi} = 0.424413, \quad A_2 = \frac{C_F}{\pi^2} \left[ C_A \left( \frac{67}{36} - \frac{\pi^2}{12} \right) - \frac{5}{9} n_F T_R \right] = 0.42095 - 0.03753 n_F, \quad (28)$$

where  $C_A = N_c = 3$ ,  $T_R = 1/2$  and  $n_F = 3$  is the number of active quark flavours in  $b$  decay. The value given for  $A_2$  in the  $\overline{MS}$  scheme for the coupling constant. At present, only a numerical estimate of  $A_3$  is available:

$$A_3 = 0.59413 - 0.09272 n_F - 0.00040 n_F^2. \quad (29)$$

The soft quantities  $D_1$  and  $D_2$  [8] are known analytically:

$$\begin{aligned} D_1 &= -\frac{C_F}{\pi} = -0.424413, \\ D_2 &= -\frac{C_F}{\pi^2} \left[ \left( \frac{37}{108} + \frac{7}{18} \pi^2 - \frac{9}{4} \zeta(3) \right) C_A + \left( \frac{1}{27} - \frac{1}{9} \pi^2 \right) T_R n_F \right] = -0.59826 - 0.07157 n_F. \end{aligned} \quad (30)$$

Numerically,  $\zeta(3) \cong 1.20206$ . The constant  $D_1$  is renormalization-scheme independent, while  $D_2$  is not and we have given its value in the  $\overline{MS}$  scheme. The latter quantity turns out to be the most important one to determine the NNLO effects.

The collinear quantity  $B_1$  is known analytically,

$$B_1 = -\frac{3 C_F}{4 \pi} = -0.31831. \quad (31)$$

The two-loop quantity  $B_2$  is unknown and we approximate it with the infrared-regular part in the two-loop Altarelli-Parisi splitting function  $P_{qq}(z)$ . In general, the latter is naturally decomposed in the soft limit as:

$$P_{qq}(z) = \left[ \frac{A(\alpha_S(Q^2(1-z)))}{1-z} \right]_+ - K(z; \alpha_S) + K(\alpha_S) \delta(1-z). \quad (32)$$

The functions  $K(z; \alpha_S)$  and  $K(\alpha_S)$  have an expansion in powers of  $\alpha_S$ :

$$\begin{aligned} K(\alpha_S) &= K_1 \alpha_S + K_2 \alpha_S^2 + \dots, \\ K(z; \alpha_S) &= K_1(z) \alpha_S + K_2(z) \alpha_S^2 + \dots. \end{aligned} \quad (33)$$

Since the splitting function has vanishing first moment:

$$K(\alpha_S) = \int_0^1 K(z; \alpha_S) dz, \quad (34)$$

it holds that

$$K_1 = \int_0^1 K_1(z) dz \quad \text{and} \quad K_2 = \int_0^1 K_2(z) dz. \quad (35)$$

In leading order, it holds  $B_1 = K_1$ . Our approximation then is:

$$B_2 \approx K_2 \tag{36}$$

where, in the  $\overline{MS}$  scheme [9],

$$\begin{aligned} K_2 &= -\frac{C_F}{\pi^2} \left[ C_F \left( \frac{3}{32} - \frac{\pi^2}{8} + \frac{3}{2} \zeta(3) \right) + C_A \left( \frac{17}{96} + \frac{11}{72} \pi^2 - \frac{3}{4} \zeta(3) \right) - \left( \frac{1}{24} + \frac{\pi^2}{18} \right) T_R n_F \right] \\ &= -0.43695 + 0.03985 n_F. \end{aligned} \tag{37}$$

## 2.1 Results

The functions  $g_1$  and  $g_2$  have the following expressions [10, 11]:

$$g_1 \left( \lambda; \frac{\mu^2}{Q^2} \right) = -\frac{A_1}{2\beta_0} \frac{1}{\lambda} [(1-2\lambda) \log(1-2\lambda) - 2(1-\lambda) \log(1-\lambda)]; \tag{38}$$

$$\begin{aligned} g_2 \left( \lambda; \frac{\mu^2}{Q^2} \right) &= +\frac{A_2}{2\beta_0^2} [\log(1-2\lambda) - 2\log(1-\lambda)] + \frac{A_1 \gamma_E}{\beta_0} [\log(1-2\lambda) - \log(1-\lambda)] + \\ &\quad -\frac{\beta_1 A_1}{4\beta_0^3} [\log^2(1-2\lambda) - 2\log^2(1-\lambda) + 2\log(1-2\lambda) - 4\log(1-\lambda)] + \\ &\quad +\frac{D_1}{2\beta_0} \log(1-2\lambda) + \frac{B_1}{\beta_0} \log(1-\lambda) + \frac{A_1}{2\beta_0} [\log(1-2\lambda) - 2\log(1-\lambda)] \log \frac{\mu^2}{Q^2}. \end{aligned} \tag{39}$$

Our result for the NNLO function  $g_3$  reads:

$$\begin{aligned}
g_3 \left( \lambda; \frac{\mu^2}{Q^2} \right) = & -\frac{A_3}{2\beta_0^2} \left[ \frac{\lambda}{1-2\lambda} - \frac{\lambda}{1-\lambda} \right] - \frac{A_1\zeta_2}{2} \left[ \frac{4\lambda}{1-2\lambda} - \frac{\lambda}{1-\lambda} \right] + \\
& -\frac{A_1\beta_2}{4\beta_0^3} \left[ \frac{2\lambda}{1-2\lambda} - \frac{2\lambda}{1-\lambda} + \log(1-2\lambda) - 2\log(1-\lambda) \right] + \\
& +\frac{A_2\beta_1}{2\beta_0^3} \left[ \frac{\log(1-2\lambda)}{1-2\lambda} - \frac{2\log(1-\lambda)}{1-\lambda} + \frac{3\lambda}{1-2\lambda} - \frac{3\lambda}{1-\lambda} \right] + \\
& -\frac{A_1\beta_1^2}{2\beta_0^4} \left[ \frac{1}{2} \frac{\log^2(1-2\lambda)}{1-2\lambda} - \frac{\log^2(1-\lambda)}{1-\lambda} + \frac{\log(1-2\lambda)}{1-2\lambda} + \right. \\
& \left. -\frac{2\log(1-\lambda)}{1-\lambda} + \frac{\lambda}{1-2\lambda} - \frac{\lambda}{1-\lambda} - \log(1-2\lambda) + 2\log(1-\lambda) \right] + \\
& +\frac{D_1\beta_1}{2\beta_0^2} \left[ \frac{\log(1-2\lambda)}{1-2\lambda} + \frac{2\lambda}{1-2\lambda} \right] + \frac{B_1\beta_1}{\beta_0^2} \left[ \frac{\log(1-\lambda)}{1-\lambda} + \frac{\lambda}{1-\lambda} \right] + \\
& -\frac{D_2}{\beta_0} \frac{\lambda}{1-2\lambda} - \frac{B_2}{\beta_0} \frac{\lambda}{1-\lambda} - \frac{A_1\gamma_E^2}{2} \left[ \frac{4\lambda}{1-2\lambda} - \frac{\lambda}{1-\lambda} \right] + \\
& +\frac{A_1\beta_1\gamma_E}{\beta_0^2} \left[ \frac{\log(1-2\lambda)}{1-2\lambda} - \frac{\log(1-\lambda)}{1-\lambda} + \frac{1}{1-2\lambda} - \frac{1}{1-\lambda} \right] + \\
& -\frac{A_2\gamma_E}{\beta_0} \left[ \frac{1}{1-2\lambda} - \frac{1}{1-\lambda} \right] - \frac{D_1\gamma_E 2\lambda}{1-2\lambda} - \frac{B_1\gamma_E \lambda}{1-\lambda} + \\
& -\frac{A_1}{2\beta_0} \left[ \frac{2\lambda^2}{1-2\lambda} - \frac{\lambda^2}{1-\lambda} \right] \log^2 \frac{\mu^2}{Q^2} - \frac{A_2}{\beta_0^2} \left[ \frac{\lambda}{1-2\lambda} - \frac{\lambda}{1-\lambda} \right] \log \frac{\mu^2}{Q^2} + \\
& -\frac{A_1\gamma_E}{\beta_0} \left[ \frac{2\lambda}{1-2\lambda} - \frac{\lambda}{1-\lambda} \right] \log \frac{\mu^2}{Q^2} - \frac{D_1}{\beta_0} \frac{\lambda}{1-2\lambda} \log \frac{\mu^2}{Q^2} - \frac{B_1}{\beta_0} \frac{\lambda}{1-\lambda} \log \frac{\mu^2}{Q^2} + \\
& +\frac{A_1\beta_1}{\beta_0^3} \left[ \frac{\lambda \log(1-2\lambda)}{1-2\lambda} - \frac{\lambda \log(1-\lambda)}{1-\lambda} + \frac{\lambda}{1-2\lambda} + \right. \\
& \left. -\frac{\lambda}{1-\lambda} + \frac{1}{2} \log(1-2\lambda) - \log(1-\lambda) \right] \log \frac{\mu^2}{Q^2}. \tag{40}
\end{aligned}$$

Arbitrary constants have been added to the function  $g_3$  in order to make it homogenous. The quantity  $\gamma_E = 0.577216\dots$  is the Euler constant and  $\zeta(n)$  is the Riemann zeta function,

$$\zeta(n) \equiv \sum_{k=1}^{\infty} \frac{1}{k^n}. \tag{41}$$

$\zeta(2) = \pi^2/6 = 1.64493$ . The functions  $g_2$  and  $g_3$  depend on the renormalization scale  $\mu$ , while  $g_1$  does not.

The expansion up to order  $\alpha_S^3$  reads:

$$\begin{aligned}
\log f_N = & -\frac{1}{2}A_1\alpha_S L^2 - D_1\alpha_S L - B_1\alpha_S L + \\
& -\frac{1}{2}A_1\beta_0\alpha_S^2 L^3 - \left( \frac{1}{2}A_2 + \beta_0 D_1 + \frac{1}{2}\beta_0 B_1 \right) \alpha_S^2 L^2 - \left( D_2 + B_2 + \frac{3}{2}A_1\zeta_2\beta_0 \right) \alpha_S^2 L + \\
& -\frac{7}{12}A_1\beta_0^2\alpha_S^3 L^4 - \left( A_2\beta_0 + \frac{1}{2}A_1\beta_1 + \frac{4}{3}D_1\beta_0^2 + \frac{1}{3}B_1\beta_0^2 \right) \alpha_S^3 L^3 + \\
& - \left( \frac{1}{2}A_3 + \frac{7}{2}A_1\zeta_2\beta_0^2 + 2D_2\beta_0 + B_2\beta_0 + D_1\beta_1 + \frac{1}{2}B_1\beta_1 \right) \alpha_S^3 L^2 + O(\alpha_S^4). \tag{42}
\end{aligned}$$

Let us note that  $\beta_2$  appears only at order  $O(\alpha_S^4)$ .



The functions  $g_i$  become singular when

$$\lambda \rightarrow \frac{1}{2}^- . \quad (43)$$

Since

$$\lambda = \beta_0 \alpha_S (\mu^2) L, \quad (44)$$

this means that a singularity in  $N$ -space occurs when:

$$N \rightarrow \exp \left[ \frac{1}{2\beta_0 \alpha_S (\mu^2)} \right] \approx \frac{\mu}{\Lambda}, \quad (45)$$

where  $\Lambda$  is the QCD scale. Let us observe that, in general, this singularity signals non-perturbative effects but its precise position is completely unphysical, as we can move it with a change of renormalization scale.

## 2.2 Renormalization-scale dependence

In this section we consider renormalization-scale dependence. In principle, such scale  $\mu$  should not appear in the cross sections, as it does not correspond to any fundamental constant or kinematical scale in the problem. The completely resummed perturbative expansion of an observable<sup>1</sup> is indeed formally independent on  $\mu$ . In practise, truncated perturbative expansions exhibit a residual scale dependence, because of neglected higher orders.

We start with the form factor as a function of  $\alpha_S (Q^2)$  and we derive its expression as a function of  $\alpha_S (\mu^2)$  and  $\mu^2/Q^2$ .

Since

$$a(Q) = a(\mu) + c a^2(\mu) + c' a^3(\mu) + O(a^4), \quad (46)$$

where  $a \equiv \beta_0 \alpha_S$  and

$$c = \log \frac{\mu^2}{Q^2}, \quad c' = \log^2 \frac{\mu^2}{Q^2} + \frac{\beta_1}{\beta_0^2} \log \frac{\mu^2}{Q^2}, \quad (47)$$

one has, where now  $\lambda = \lambda(\mu)$ :

$$\begin{aligned} L g_1 [\lambda + ac\lambda + a^2 c' \lambda] &= L g_1 [\lambda] + c \lambda^2 g_1' [\lambda] + ac' \lambda^2 g_1' [\lambda] + a \frac{1}{2} c^2 \lambda^3 g_1'' [\lambda] + \dots \\ g_2 [\lambda + ac\lambda + a^2 c' \lambda] &= g_2 [\lambda] + a c \lambda g_2' [\lambda] + \dots \end{aligned} \quad (48)$$

The additional terms in the functions  $g_i$ , to (partially) compensate for the scale change  $Q^2 \rightarrow \mu^2$  therefore read:

$$\begin{aligned} \delta g_1 \left[ \lambda, \frac{\mu^2}{Q^2} \right] &= 0 \\ \delta g_2 \left[ \lambda, \frac{\mu^2}{Q^2} \right] &= \lambda^2 g_1' [\lambda] \log \frac{\mu^2}{Q^2} \\ \delta g_3 \left[ \lambda, \frac{\mu^2}{Q^2} \right] &= \frac{1}{2} \lambda^3 g_1'' [\lambda] \log^2 \frac{\mu^2}{Q^2} + \lambda^2 g_1' [\lambda] \left( \log^2 \frac{\mu^2}{Q^2} + \frac{\beta_1}{\beta_0^2} \log \frac{\mu^2}{Q^2} \right) + \lambda g_2' [\lambda] \log \frac{\mu^2}{Q^2} \\ &= \frac{1}{2} \lambda \frac{d}{d\lambda} (\lambda^2 g_1' [\lambda]) \log^2 \frac{\mu^2}{Q^2} + \left( \frac{\beta_1}{\beta_0^2} \lambda^2 g_1' [\lambda] + \lambda g_2' [\lambda] \right) \log \frac{\mu^2}{Q^2}. \end{aligned} \quad (49)$$

The leading function  $g_1$  is therefore renormalization-scale independent, analogously to the coefficient of the leading term in the expansion of inclusive observables (such as the  $\alpha_S/\pi$  term in the total  $e^+e^-$  hadronic cross-section). The higher-order functions  $g_{i>1}$  instead explicitly depend on  $\mu$ .

<sup>1</sup>We do not mean here the resummation of towers of logarithmic contributions, but the resummation of the whole series in  $\alpha_S$ .

### 3 Analytic inverse Mellin transform

The  $N$ -moments are physical quantities, but in practice a measure of the moments for large  $N$  is difficult. It is therefore convenient to perform the inverse transform back to momentum space.

Let us derive an analytical expression of the inverse Mellin transform at NNLO. We start by introducing the inverse Mellin transform of  $f_N(\alpha_s)/N$ :

$$G(\alpha_s; x) = \frac{1}{2\pi i} \int_{C-i\infty}^{C+i\infty} \frac{dN}{N} x^{-N} f_N(\alpha_s, L) \quad (50)$$

The inverse Mellin transform of  $f_N(\alpha_s)$  is the logarithmic derivative of  $G(\alpha_s, L; x)$ :

$$f(x) = -x \frac{d}{dx} G(x). \quad (51)$$

After the substitutions  $x \equiv e^{-s}$  and  $u \equiv Ns$ , we have:

$$\begin{aligned} G(\alpha_s; x) &= \frac{1}{2\pi i} \int_{C'-i\infty}^{C'+i\infty} \frac{du}{u} e^u f_N(\alpha_s, L) \\ &= \frac{1}{2\pi i} \int_{C'-i\infty}^{C'+i\infty} \frac{du}{u} e^u \exp \left[ L g_1(\beta_0 \alpha_s L) + \sum_{n=2}^{\infty} \alpha_s^{n-2} g_n(\beta_0 \alpha_s L) \right] \end{aligned} \quad (52)$$

We Taylor expand the exponent with respect to  $L$  around  $L = l = \ln 1/s$  [12]. In the  $x$ -variable,  $l \equiv -\ln(-\ln x)$ . Note that  $l \rightarrow -\ln(1-x)$  when  $x \rightarrow 1$ .

We have, at NNLO:

$$G(\alpha_s; x) = \frac{1}{2\pi i} \int_{C'-i\infty}^{C'+i\infty} \frac{du}{u} e^{u+F_0(l)+F_1(l) \ln u + \frac{1}{2} F_2(l) \ln^2 u + \dots} \quad (53)$$

where, at scale  $Q^2$  and at the same NNLO:

$$\begin{aligned} F_0(l) &= l g_1(\beta_0 \alpha_s l) + g_2(\beta_0 \alpha_s l) + \alpha_s g_3(\beta_0 \alpha_s l), \\ F_1(l) &= g_1(\beta_0 \alpha_s l) + \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l) + \beta_0 \alpha_s g_2'(\beta_0 \alpha_s l), \\ F_2(l) &= 2\beta_0 \alpha_s g_1'(\beta_0 \alpha_s l) + \beta_0^2 \alpha_s^2 l g_1''(\beta_0 \alpha_s l). \end{aligned}$$

By keeping in the exponent terms up to NLO, while expanding the NNLO terms, we obtain:

$$G(\alpha_s; x) = \frac{e^{F_0(l)}}{2\pi i} \int_{C'-i\infty}^{C'+i\infty} du e^{u-[1-F_1^{NL}(l)] \ln u} \left[ 1 + F_1^{N^2L}(l) \ln u + \frac{1}{2} F_2(l) \ln^2 u + \dots \right] \quad (54)$$

where:

$$\begin{aligned} F_1(l) &\equiv F_1^{NL}(l) + F_1^{N^2L}(l) \\ F_1^{NL}(l) &\equiv g_1(\beta_0 \alpha_s l) + \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l) \\ F_1^{N^2L}(l) &\equiv \beta_0 \alpha_s g_2'(\beta_0 \alpha_s l). \end{aligned} \quad (55)$$

By using the result:

$$\frac{1}{2\pi i} \int_C du \log^k u e^{u-[1-F_1^{NL}(l)] \log u} = \frac{d^k}{dF_1^{NLk}} \frac{1}{\Gamma(1-F_1^{NL})}, \quad (56)$$

where  $\Gamma$  is the Euler Gamma function, we obtain, after the integration:

$$\begin{aligned}
G(\alpha_s; x) &= \frac{e^{F_0(l)}}{\Gamma(1 - F_1^{NL})} \left[ 1 + F_1^{N^2 L} \psi(1 - F_1^{NL}) + \frac{1}{2} F_2(l) (\psi^2(1 - F_1^{NL}) - \psi'(1 - F_1^{NL})) \right] \\
&= \frac{e^{l g_1(\beta_0 \alpha_s l) + g_2(\beta_0 \alpha_s l) + \alpha_s g_3(\beta_0 \alpha_s l)}}{\Gamma(1 - g_1(\beta_0 \alpha_s l) - \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l))} \left[ 1 + \beta_0 \alpha_s g_2'(\beta_0 \alpha_s l) \psi(1 - g_1(\beta_0 \alpha_s l) - \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l)) \right. \\
&\quad \left. + \frac{1}{2} F_2(l) (\psi^2(1 - g_1(\beta_0 \alpha_s l) - \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l)) - \psi'(1 - g_1(\beta_0 \alpha_s l) - \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l))) \right] \quad (57)
\end{aligned}$$

with  $\psi(x) = d \log \Gamma(x) / dx$ , the digamma function.

At the end, we find the explicit analytic formula for the inverse Mellin transform at NNLO:

$$\begin{aligned}
f(x) &= -x \frac{d}{dx} G(\alpha_s; x) \\
&= -x \frac{d}{dx} \left\{ \frac{e^{l g_1(\beta_0 \alpha_s l) + g_2(\beta_0 \alpha_s l) + \alpha_s g_3(\beta_0 \alpha_s l)}}{\Gamma(1 - g_1(\beta_0 \alpha_s l) - \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l))} \left[ 1 + \beta_0 \alpha_s g_2'(\beta_0 \alpha_s l) \psi(1 - g_1(\beta_0 \alpha_s l) - \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l)) \right. \right. \\
&\quad \left. \left. + \frac{1}{2} F_2(l) (\psi^2(1 - g_1(\beta_0 \alpha_s l) - \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l)) - \psi'(1 - g_1(\beta_0 \alpha_s l) - \beta_0 \alpha_s l g_1'(\beta_0 \alpha_s l))) \right] \right\}. \quad (58)
\end{aligned}$$

### 3.1 Discussion and conclusions

There are two physical effects involved in the inverse Mellin transform:

1. *exact longitudinal momentum conservation*; the terms which enforce longitudinal momentum conservation are formally subleading in the infrared logarithm counting;
2. *infrared pole in the coupling*; the inverse transform involves an integration over all moments and arbitrarily large values of  $|N|$  enter. This implies that a prescription for the infrared pole in the coupling has to be given. In general, the transformation mixes all the momentum scales in the problem.

One can study one problem at a time; for example, one can study the problem 1. only, by considering the frozen coupling approximation for the distributions in  $N$ -space. In one-loop, this is equivalent to consider the QED case.

The frozen coupling approximation means neglecting the variation of  $\alpha_s$  with the scale. Let us start from formula (12) at NNLO:

$$\begin{aligned}
\log f_N(\alpha_S) &= \int_0^1 dz \frac{z^{N-1} - 1}{1 - z} \left\{ \int_{Q^2(1-z)^2}^{Q^2(1-z)} \frac{dk^2}{k^2} [A_1 \alpha_S + A_2 \alpha_S^2 + A_3 \alpha_S^3 + \dots] + \right. \\
&\quad \left. + B_1 \alpha_S + B_2 \alpha_S^2 + \dots + D_1 \alpha_S + D_2 \alpha_S^2 + \dots \right\} \\
&\simeq \int_0^1 dz \frac{z^{N-1} - 1}{1 - z} \left\{ (A_1 \alpha_S + A_2 \alpha_S^2 + A_3 \alpha_S^3) \ln \frac{1}{1 - z} + (B_1 + D_1) \alpha_S + (B_2 + D_2) \alpha_S^2 \right\} \quad (59)
\end{aligned}$$

After integration in  $z$ , at the lowest order in  $\lambda = \beta_0 \alpha_2 \ln N$ , we have:

$$g_1 = -\frac{A_1}{2\beta_0} \lambda \quad (60)$$

$$g_2 = \left( -\frac{B_1}{\beta_0} - \frac{D_1}{\beta_0} - \frac{A_1 \gamma_E}{\beta_0} \right) \lambda \quad (61)$$

$$g_3 = \left( -\frac{B_2}{\beta_0} - \frac{D_2}{\beta_0} - \frac{A_2 \gamma_E}{\beta_0} \right) \lambda. \quad (62)$$

We have two ways, analytical and numerical, to compute the inverse Mellin transform of  $f_N(\alpha_s)$ :

$$f(\alpha_s; x) = \frac{1}{2\pi i} \int_{C-i\infty}^{C+i\infty} dN x^{-N} f_N(\alpha_s, L). \quad (63)$$

In the case of frozen coupling, the  $g_i$  are linear in  $\lambda$  and therefore the numerical path does not include the Landau pole; the numerical integration is therefore exact. We have compared the inverse Mellin transform calculated numerically, directly from the definition (63), with the analytical result (58). We find a very good agreement, up to energies around the charm scale, provided that we go to NNLO. The exact longitudinal momentum conservation does not seem to spoil the reliability of the perturbative series.

In the real world, we cannot use linearized  $g_i$ , but we have to use the  $g_i$  computed in section 2; therefore, problems due to infrared poles come into play.

Let us first make an important observation: the degree of singularity of the functions  $g_i$  for  $\lambda \rightarrow 1/2$ , and therefore also of the form factor, increases with the order of the function, i.e. with  $i$  [13].

At LO,  $g_1$  has at most a logarithmic singularity of the form

$$g_1 : (1 - 2\lambda) \log(1 - 2\lambda); \quad (64)$$

at NLO,  $g_2$  has at most a singularity of the form

$$g_2 : \log^2(1 - 2\lambda). \quad (65)$$

The prefactor  $1 - 2\lambda$  in front of the logarithm, vanishing for  $\lambda = 1/2$ , is now absent, but the singularity is still logarithmic. At NNLO,  $g_3$  has at most a singularity of the form

$$g_3 : \frac{\log^2(1 - 2\lambda)}{1 - 2\lambda}. \quad (66)$$

The latter is basically a pole singularity, no more a logarithmic singularity anymore, i.e. a much stronger singularity. Note that the above term is proportional to  $A_1\beta_1^2$  and so it is scheme-independent.

Another observation concerns the size of the two-loop soft term. For three active flavours, the size of the two-loop correction with respect to the one-loop one is rather large,

$$\frac{D_2}{D_1} \alpha_S \approx 2 \alpha_S \approx 40\%, \quad (67)$$

since  $\alpha_S(m_B) \simeq 0.21$ . The inclusion of  $D_2$  is therefore important. The size of the two-loop term compared to the one-loop term is expected on general ground to be of order:

$$\frac{D_2}{D_1} \approx \frac{C_A}{C_F} = 2.25, \quad (68)$$

since in two-loop order gluons start to be radiated by gluons instead of quarks, the former having a larger colour charge  $C_A$  instead of  $C_F$ .

Let us now consider the inverse Mellin transform (63), releasing the frozen coupling approximation; in formula (12) the coupling runs over the whole integration range.

The form factor in  $N$ -space is computed in such a way that infrared-pole effects appear in a sharp way for  $N > N_c$ , where it acquires an unphysical (imaginary) part. In other terms, the numerical distribution is not real for any value of  $N$  because of the integration over the Landau pole. An exact numerical evaluation of the inverse transform then requires a prescription for the pole. An alternative strategy is to give a prescription for the infrared pole directly in  $N$ -space, in such a way that the form factors are well-defined for any  $N$ . In general, this results into a softening of the form factor for large  $N$ . It is then not necessary to give a prescription for the pole in the inverse transform.

We have used the minimal prescription (mp) [14], over two different paths (with the same results); the first path was made by two straight lines parallel to the negative real axes, closed by a half-circle centered around the origin and crossing the positive axes between the origin and the first Landau pole; the second path was composed by two lines almost vertical, meeting on the positive real axes between the origin and the first Landau pole. The precise crossing point is irrelevant, as far as it is before the Landau pole.

The inverse Mellin transform can be also derived analytically, as seen in section 3. We compare the analytic (NLO and NNLO) and numerical (NNLO) plots in fig. 1 (at  $\alpha_s = 0.21$ ). We conclude that, at such high energies, the perturbative effects dominate also for  $x$  very close to 1, the first Landau pole (see eq. (43)) occurring at large values of  $N$  ( $N = e^{1/2\beta_0\alpha_s} \simeq 3000 - 4000$ ). The perturbative resummation keeps under control the infrared divergences and cancel the oscillatory behaviour. We are in a trustable perturbative region for  $f(x)$ ; the observed final state can reach very high values of energy and we can be confident that soft gluon effects are consistently resummed by the perturbative form factor.

#### Acknowledgements

We wish to thank S. Catani for discussions. G.R. thanks the CERN theory department for hospitality during part of this work.

## References

- [1] S. Catani and L. Trentadue, Nucl. Phys. B 327, 323 (1989).
- [2] S. Catani and L. Trentadue, Nucl. Phys. B 353, 183 (1991).
- [3] U. Aglietti, Nucl. Phys. B 610, 293 (2001).
- [4] Particle Data Book coll., Eur. Phys. J. C 15, 1 (2000).
- [5] O. Tarasov, A. Vladimirov and A. Zharkov, Phys. Lett. B 93, 429 (1980).
- [6] A. Vogt, Phys. Lett. B 497, 228 (2001).
- [7] J. Kodaira and L. Trentadue, preprint SLAC-PUB-2934 (1982); Phys. Lett. B 112, 66 (1982); S. Catani, E. D’Emilio and L. Trentadue, Phys. Lett. B 211, 335 (1988).
- [8] G.P. Korchemsky and G. Marchesini, Nucl. Phys. B 406, 225 (1993).
- [9] G.P. Korchemsky, Mod. Phys. Lett. A 4, 1257 (1989).
- [10] R. Akhouri and I. Rothstein, Phys. Rev. D 54, 2349 (1996); G. Korchemsky and G. Sterman, Phys. Lett. B 340, 96 (1994).
- [11] U. Aglietti, Phys. Lett. B 515, 308 (2001).
- [12] S. Catani, L. Trentadue, G. Turnock and B. Webber, Nucl. Phys. B 407, 3 (1993).
- [13] E. Gardi and J. Rathsman, Nucl. Phys. B 609, 123 (2001).
- [14] S. Catani, M. Mangano, L. Trentadue and P. Nason, Nucl. Phys. B 478, 273 (1996).




Figure 1: mp numerical distribution (solid line), NLO (dashed line) and NNLO (dot-dashed line) analytic distributions of  $f(x)$ , at  $\alpha_s = 0.1$ .