

## The universal freeze-out criterion at SPS and RHIC

Boris Tomášik and Urs Achim Wiedemann

CERN, Theory Division, CH-1211 Geneva 23, Switzerland

We formulate a freeze-out criterion for ultra-relativistic heavy ion collisions in terms of the pion escape probability from the collision region. We find that the increase in pion phase-space density from SPS to RHIC reported at this conference has a small influence on particle freeze-out because of the small  $\pi\pi$  cross-section. Our treatment takes into account dynamical expansion, chemical composition and momentum of the escaping particle for particle freeze-out. It supports a freeze-out at rather low temperature—below 100 MeV—and earlier decoupling of high  $p_{\perp}$  particles.

A precise formulation of the condition under which particles decouple (“freeze-out”) from a heavy ion collision is important for understanding the dynamical origin of the final state. The observation of STAR [1] that the pion phase-space density increases at RHIC indicates that freeze-out is not characterised solely by a universal value of *pion phase-space density*, in contrast to a recent suggestion [2]. Also, freeze-out is not characterised solely by a universal value of the *total particle density* since the volume from which particles decouple shows a non-monotonous dependence on collision energy [3].

It has been argued [4] and demonstrated [3] that the chemical composition of the system must be taken into account. For pion freeze-out, the nucleon and anti-nucleon density is more important than that of pions since the pion-nucleon cross-section is larger than the pion-pion one. With increasing collision energy the pion abundance and their contribution to scattering grows, but it is argued that the *momentum-averaged* mean free path of a pion at freeze-out stays constant [3].

This argumentation still ignores the influence of global dynamics on the freeze-out criterion. A strong expansion makes the particle density decrease fast and if the characteristic time scale for density decrease  $\tau_{\text{exp}} = (\partial_{\mu} u^{\mu})^{-1}$  is smaller than the mean free time between collisions  $\tau_{\text{scatt}}$  a particle is likely to seize interacting, i.e., to freeze-out [5].

In our work we study—in addition to these two aspects—how the probability for a particle to decouple depends on its momentum [6]. We formulate freeze-out as condition for an individual particle: a *particle freezes-out* at the time of its last scattering. This is in contrast to the “fireball freeze-out” which is understood as a boundary in space-time between interacting matter and free-streaming particles. In the extreme case of a hydrodynamic model the “fireball freeze-out” is usually treated by the Cooper-Frye prescription [7] in which *all* particles decouple along the same three-dimensional hypersurface when *the medium* fulfils certain condition. Our treatment goes beyond Cooper-Frye: particle freeze-out depends on its *type, position, and momentum* in addition to the characteristics of the medium. This corresponds to freeze-out as described by a cascade

event generator.

The freeze-out criterion which we explore is based on the *escape probability* [8]

$$\mathcal{P}(x, p, \tau) = \exp \left( - \int_{\tau}^{\infty} d\bar{\tau} \mathcal{R}(x + v\bar{\tau}, p) \right), \quad (1)$$

which determines the probability that a pion emitted with momentum  $p$  from the position  $(x, \tau)$  escapes from the medium without further interaction. Here, the scattering rate  $\mathcal{R}(x, p)$  is integrated along the trajectory of the pion. Freeze-out is assumed to occur when  $\mathcal{P}(x, p, \tau)$  reaches a *universal* value of order one.

The role of collective expansion: To illustrate how collective expansion affects particle freeze-out, we consider the case of particles with vanishing momentum in the centre of a fireball with longitudinally boost-invariant and transversely linear expansion velocity profile of the form

$$u^{\mu} = (\cosh \eta \cosh \eta_t, \cos \phi \sinh \eta_t, \sin \phi \sinh \eta_t, \sinh \eta \cosh \eta_t). \quad (2)$$

Here,  $\eta = \text{Arctanh}(z/t)$  and  $\eta_t = \xi r$  where  $\xi = 0.08 \text{ fm}^{-1}$  is a typical value for the transverse expansion gradient [9]. At the time  $\tau_{\text{em}}$  at which the pion is emitted, the density of scattering centres decreases by the rate

$$-\frac{1}{\rho} \frac{\partial \rho}{\partial \tau} \Big|_{\tau=\tau_{\text{em}}} = \partial_{\mu} u^{\mu} = \frac{1}{\tau_{\text{em}}} + 2\xi. \quad (3)$$

A power-law decrease of density with time is consistent with (3) if

$$\rho(\tau) = \rho_{\text{em}} \left( \frac{\tau_{\text{em}}}{\tau} \right)^{\alpha}, \quad \alpha = 1 + 2\xi \tau_{\text{em}}. \quad (4)$$

Under the assumption  $\mathcal{R} \propto \rho$  we obtain for the opacity integral of a particle of zero momentum

$$\int_{\tau_{\text{em}}}^{\infty} d\tau \mathcal{R}(\tau) = \frac{\mathcal{R}_{\text{em}}}{\alpha - 1} \tau_{\text{em}} = \frac{\mathcal{R}_{\text{em}}}{2\xi}, \quad (5)$$

where  $\mathcal{R}_{\text{em}}$  is the scattering rate at  $\tau_{\text{em}}$ . Technically, the calculation of this opacity integral for particles of non-zero momentum  $p$  is more complicated, since one has to follow the propagation of the particle through layers of different density. For a quantitative statement, this requires a model of the space-time evolution of the fireball. Qualitatively, however, the general result will be consistent with the main feature of the case  $p = 0$  considered here: as seen from (5), freeze-out from a denser fireball (larger  $\mathcal{R}_{\text{em}}$ ) is possible if it is compensated by a stronger expansion gradient such that the value of opacity integral stays constant.

The role of chemical composition: To determine the value of  $\mathcal{R}_{\text{em}}$  for collisions at  $\sqrt{s} = 17 \text{ AGeV}$  (SPS) and  $\sqrt{s} = 130 \text{ AGeV}$  (RHIC), we calculate the scattering rate

$$\mathcal{R}(p) = \sum_i \int d^3k \rho_i(k) \sigma'_i(s) |v_{\pi} - v_i|, \quad (6)$$

where  $\sigma'_i$  is the cross-section for pion scattering on species  $i$ . For the collinear cross-section we use the parametrisation

$$\sigma_i(\sqrt{s}) = \sum_r \langle j_i, m_i, j_{\pi}, m_{\pi} || J_r, M_r \rangle \frac{2S_r + 1}{(2S_i + 1)(2S_{\pi} + 1)} \frac{\pi}{p_{\text{CMS}}^2} \frac{\Gamma_{r \rightarrow \pi i} \Gamma_{i, \text{tot}}}{(M_r - \sqrt{s})^2 + \Gamma_{i, \text{tot}}^2/4}, \quad (7)$$




Figure 1. The pion scattering rate as a function of pion momentum with respect to medium calculated at  $T = 100$  MeV and the highest estimates of chemical potentials allowed by data from SPS (left column) and RHIC (right column). Contributions to the total scattering rate from scattering on nucleons, anti-nucleons and pions are indicated. The lower row shows the baryonic and mesonic relative contributions.

where the sum runs over all resonances in the  $\pi i$  scattering. Particle densities  $\rho_i(k)$  are assumed to be thermal. For realistic temperatures, only low-lying resonances are relevant and we include  $i = \pi, N, \bar{N}, K, \rho, \Delta, \bar{\Delta}$  when summing in (6) over scattering partners. Since the freeze-out temperature is not known a priori, we scan three values:  $T = 90, 100, 120$  MeV. Chemical potentials for pions are estimated from data on phase-space density, those for other species from measured ratios of  $dN/dy$  at mid-rapidity (see [6] for more details on the estimates).

Figure 1 shows the scattering rates as a function of the pion momentum *relative to the heat bath* calculated for SPS and RHIC, as well as the most important contributions to  $\mathcal{R}$ . In spite of the strong increase of the pion phase-space density at RHIC [1], the relative meson contribution to the total scattering rate does not grow proportionally since the  $\pi\pi$  cross-section is much smaller than the one for  $\pi N$ . The total baryonic contribution changes very little since the smaller amount of baryons at RHIC is roughly compensated by anti-baryons. In summary, there is about 10% of the relative contribution shifted from baryons to mesons when going from SPS to RHIC.

As seen in Figure 2, the scattering rate depends strongly on the assumed freeze-out temperature since the pion-nucleon scattering is typically dominated by total CMS energy below the  $\Delta$  resonance peak. At higher temperature, average CMS energies move closer to the peak value thus leading to increased scattering cross-sections. Estimating the pion escape probability (1) from eq. (5) and  $\xi = 0.08 \text{ fm}^{-1}$ , one finds that a pion only has a




Figure 2. The pion scattering rate as a function of pion momentum with respect to the medium calculated for different temperatures and a range of chemical potentials allowed by data (see [6]). New results of STAR collaboration [1] correspond rather to the “high  $\mu$ 's” estimates.

reasonable chance to escape (of order 50%) when the temperature is 100 MeV at most.

We finally point out that the scattering rates in Figure 2 generically decreases with increasing pion momentum. This leads us to conjecture that high  $p_{\perp}$  pions may decouple earlier, from hotter and smaller system. This may provide a novel as yet unexplored contribution to the observed  $M_{\perp}$  dependence of HBT radii.

The freeze-out criterion studied here differs from the Cooper-Frye formalism typically employed in hydrodynamical model studies. It will be interesting to explore to what extent this affects the predictions of hydrodynamical simulations for the collision evolution.

## REFERENCES

1. L. Ray for the STAR collaboration, these proceedings.
2. D. Ferenc *et al.*, Phys. Lett. B **457** (1999) 347.
3. D. Adamová *et al.* [CERES collab.], arXiv:nucl-ex/0207008 and these proceedings.
4. S. Nagamiya, Phys. Rev. Lett. **49** (1982) 1383.
5. J. P. Bondorf, S. I. Garpman and J. Zimányi, Nucl. Phys. A **296** (1978) 320.
6. B. Tomášik and U. A. Wiedemann, arXiv:nucl-th/0207074.
7. F. Cooper and G. Frye, Phys. Rev. D **10** (1974) 186.
8. Y. M. Sinyukov, S. V. Akkelin and Y. Hama, Phys. Rev. Lett. **89** (2002) 052301.
9. B. Tomášik, U. A. Wiedemann and U. W. Heinz, arXiv:nucl-th/9907096.