

Towards the NNLL precision in the decay $\bar{B} \rightarrow X_s \gamma$

Hrachia M. Asatrian, Artyom Hovhannisyan, and Vahagn Poghosyan

Yerevan Physics Institute, 375036, Yerevan, Armenia

Christoph Greub

Institute for Theoretical Physics, University Berne, CH-3012 Berne, Switzerland

Tobias Hurth^{*†}

CERN, Dept. of Physics, Theory Unit, CH-1211 Geneva 23, Switzerland

SLAC, Stanford University, Stanford, CA 94309, USA

The present NLL prediction for the decay rate of the rare inclusive process $\bar{B} \rightarrow X_s \gamma$ has a large uncertainty due to the charm mass renormalization scheme ambiguity. We estimate that this uncertainty will be reduced by a factor of 2 at the NNLL level. This is a strong motivation for the on-going NNLL calculation, which will thus significantly increase the sensitivity of the observable $\bar{B} \rightarrow X_s \gamma$ to possible new degrees of freedom beyond the SM. We also give a brief status report of the NNLL calculation.

HEP2005 International Europhysics Conference on High Energy Physics

21–27 July 2005

Lisbon, Portugal

CERN-PH-TH/2005-233, SLAC-PUB-11586

^{*}Speaker.

[†]Heisenberg Fellow

The inclusive decay $\bar{B} \rightarrow X_s \gamma$ is well known as one of the most important flavour observables within the indirect search for new physics [1]. The present experimental accuracy already reached the 10% level, as reflected in the world average of the present measurements [2]:

$$\text{BR}[\bar{B} \rightarrow X_s \gamma] = (3.39_{-0.27}^{+0.30}) \times 10^{-4}. \quad (1)$$

In the near future, more precise data on this mode are expected from the B -factories. Thus, it is mandatory to reduce the present theoretical uncertainty accordingly. Non-perturbative effects are naturally small within inclusive modes [1]; also additional non-perturbative corrections due to necessary cuts in the photon energy spectrum are under control (see [3]). As was first noticed in [4], there exists a large uncertainty in the theoretical NLL prediction related to the renormalization scheme of the charm-quark mass on which we focus in this article. The reason is that the matrix elements $\langle s \gamma | O_{1,2} | b \rangle$, through which the charm-quark mass dependence dominantly enters, vanish at the lowest order (LL) and, as a consequence, the charm-quark mass does not get renormalized in a NLL calculation, which means that the symbol m_c can be identified with $m_{c,\text{pole}}$ or with the $\overline{\text{MS}}$ mass $\bar{m}_c(\mu_c)$ at some scale μ_c or with some other definition of m_c . In a recent theoretical update of the NLL prediction of this branching fraction the ratio m_c/m_b was varied in the conservative range $0.18 \leq m_c/m_b \leq 0.31$ that covers both the pole mass value (with its numerical error) and the running mass $\bar{m}_c(\mu_c)$ value (with $\mu_c \in [m_c, m_b]$), leading to [5]:

$$\text{BR}[\bar{B} \rightarrow X_s \gamma] = (3.70 \pm 0.35|_{m_c/m_b} \pm 0.02|_{\text{CKM}} \pm 0.25|_{\text{param.}} \pm 0.15|_{\text{scale}}) \times 10^{-4}. \quad (2)$$

The only way to resolve this scheme ambiguity in a satisfactory way is to perform a systematic NNLL calculation. Working to next-to-next-to-leading-log (NNLL) precision means that one is resumming all the terms of the form

$$(\alpha_s(m_b))^p \alpha_s^n(m_b) \log^n(m_b/M), \quad (p = 0, 1, 2). \quad (3)$$

where $M = m_t$ or $M = m_W$, $n = 0, 1, 2, \dots$. Such a calculation is most suitably done in the framework of an effective low-energy theory. The effective interaction Hamiltonian can be written as

$$H_{\text{eff}} = -4G_F/\sqrt{2} \ V_{tb}V_{ts}^* \sum C_i(\mu, M) \ O_i(\mu), \quad (4)$$

where $O_i(\mu)$ are the relevant dimension 6 operators and $C_i(\mu, M)$ are the Wilson coefficients.

Parts of the three principal calculational steps leading to the NNLL result within the effective field theory approach are already done: **(a)** The full SM theory has to be matched with the effective theory at the scale $\mu = \mu_W$, where μ_W denotes a scale of order m_W or m_t . The Wilson coefficients $C_i(\mu_W)$ only pick up small QCD corrections, which can be calculated in fixed-order perturbation theory. In the NNLL program, the matching has to be worked out at the order α_s^2 . The matching calculation to this precision is already finished, including the most difficult piece, the three-loop matching of the operators $O_{7,8}$ [6]. **(b)** The evolution of these Wilson coefficients from $\mu = \mu_W$ down to $\mu = \mu_b$ then has to be performed with the help of the renormalization group, where μ_b is of the order of m_b . As the matrix elements of the operators evaluated at the low scale μ_b are free of large logarithms, the latter are contained in resummed form in the Wilson coefficients. For the NNLL calculation, this RGE step has to be done using the anomalous–dimension matrix up to

order α_s^3 . While the three-loop mixing among the four-quark operators O_i ($i = 1, \dots, 6$) [7] and among the dipole operators $O_{7,8}$ [8] are already available, the four-loop mixing of the four-quark into the dipole operators is still an open issue. (c) To achieve NNLL precision, the matrix elements $\langle X_s \gamma | O_i(\mu_b) | b \rangle$ have to be calculated to order α_s^2 precision. This includes also bremsstrahlung corrections. In 2003, the $(\alpha_s^2 n_f)$ corrections to the matrix elements of the operators O_1, O_2, O_7, O_8 were calculated [9]. Complete order α_s^2 results are available to the (O_7, O_7) contribution to the decay width [10]. Recently, also order α_s^2 terms to the photon energy spectrum (away from the endpoint E_γ^{\max}) were worked out for the operator O_7 [11].

In ref. [12] a strong motivation for this complicated NNLL effort was given by calculating those NNLL terms that are induced by renormalizing the charm-quark mass in the NLL expressions, i.e. those terms that are sensitive to the definition of the charm-quark mass. These terms correspond to δm_c insertions in the diagrams related to the NLL order matrix elements $M_{1,2}^{\text{virt}}(m_c) = \langle s \gamma | O_{1,2}(\mu_b) | b \rangle$ and $M_{1,2}^{\text{brems}}(m_c) = \langle s \gamma g | O_{1,2}(\mu_b) | b \rangle$ (for an example, see the left diagram in Fig. 1).




Figure 1: Left: Typical δm_c insertion diagram. Right: Typical diagram with a self energy insertion.

The sum $\delta M_{1,2}^{\text{virt}(\epsilon)}(m_c) \cdot \delta m_c$ of all these insertions can be obtained by replacing m_c by $m_c + \delta m_c$ in the $O(\alpha_s^1)$ results $M_{1,2}^{\text{virt}(\epsilon)}(m_c)$, followed by expanding in δm_c up to linear order:

$$M_{1,2}^{\text{virt}(\epsilon)}(m_c + \delta m_c) = M_{1,2}^{\text{virt}(\epsilon)}(m_c) + \delta M_{1,2}^{\text{virt}(\epsilon)}(m_c) \cdot \delta m_c + O((\delta m_c)^2). \quad (5)$$

As δm_c is ultraviolet-divergent, the matrix elements $M_{1,2}^{\text{virt}(\epsilon)}(m_c)$ are needed in our application up to order ϵ^1 , as indicated by the notation in eq. (5). In [12] the explicit analytical results for these matrix elements are given in such a way that they can be used in a future complete NNLL calculation. The explicit shift δm_c depends of course on the renormalization scheme. When aiming at expressing the results for $M_{1,2}^{\text{virt}(\epsilon)}(m_c)$ in terms of $\bar{m}_c(\mu_b)$ or $m_{c,\text{pole}}$, the shift reads ($C_F = 4/3$)

$$\delta \bar{m}_c(\mu_b) = -\frac{\alpha_s(\mu_b)}{4\pi} C_F \frac{3}{\epsilon} \bar{m}_c(\mu_b) \quad \text{or} \quad \delta m_{c,\text{pole}} = -\frac{\alpha_s(\mu_b)}{4\pi} C_F \left(\frac{3}{\epsilon} + 3 \ln \frac{\mu_b^2}{m_c^2} + 4 \right) m_{c,\text{pole}}.$$

The infinities induced by the $1/\epsilon$ terms in δm_c get cancelled in a full NNLL calculation, in particular by self-energy diagrams depicted in the right diagram in Fig. 1. When implementing these self-energy insertions, we only took into account the $\Sigma_1(p^2 = m_c^2)$ piece, i.e. that part of the one-loop self-energy which only gets renormalized by the mass parameter. When used at the fixed momentum $p^2 = m_c^2$, this piece is gauge-independent.

Our final estimates are given in Fig. 2 for three different values of μ_b , where μ_b represents the usual renormalization scale of the effective field theory. Within each vertical string, the solid dot represents the branching ratio using the pole mass $m_{c,\text{pole}}$, while the open symbols correspond to the $\overline{\text{MS}}$ mass $\bar{m}_c(\mu_c)$ for $\mu_c = 1.25$ GeV (triangle), $\mu_c = 2.5$ GeV (quadrangle) and $\mu_c = 5.0$ GeV (pentagon). For each μ_b the left string shows the value of the branching ratio at the NLL level, while the right string shows the corresponding value where, in addition, δm_c mass insertions and $\Sigma_1(p^2 = m_c^2)$ insertions were taken into account. Because the combination of these insertions is zero by construction for the pole scheme, the solid dots are at the same place in the left and the right string for a given value of μ_b . We stress that all the statements made in the following are independent of this absolute normalization introduced by the additional $\Sigma_1(p^2 = m_c^2)$ insertions, because we refer to the *reduction of the error* only. From Fig. 2 we see that the error related to the charm-quark mass definition is significantly reduced when the NNLL terms connected with mass insertions are taken into account. Taking as an example the results for $\mu_b = 5$ GeV, we find that at the NLL level the branching ratio evaluated for $\bar{m}_c(2.5$ GeV) is 12.6% higher than the one based on $m_{c,\text{pole}}$. Including the new contributions, these 12.6% get reduced to 5.1%. One also can read off an analogous significant reduction within the $\overline{\text{MS}}$ scheme itself. However, to obtain a NNLL prediction for the central value of the branching ratio, it is of course necessary to calculate all NNLL terms.




Figure 2: $\text{BR}(b \rightarrow X_s \gamma)$ for three values of μ_b (see text for more details).

References

- [1] T. Hurth, Rev. Mod. Phys. **75** (2003) 1159 [arXiv:hep-ph/0212304].
- [2] Heavy Flavor Averaging Group (HFAG), <http://www.slac.stanford.edu/xorg/hfag/>
- [3] M. Neubert, Eur. Phys. J. C **40** (2005) 165 [arXiv:hep-ph/0408179].
- [4] P. Gambino and M. Misiak, Nucl. Phys. B **611** (2001) 338 [arXiv:hep-ph/0104034].
- [5] T. Hurth, E. Lunghi and W. Porod, Nucl. Phys. B **704** (2005) 56 [arXiv:hep-ph/0312260].
- [6] M. Misiak and M. Steinhauser, Nucl. Phys. B **683** (2004) 277 [arXiv:hep-ph/0401041].
- [7] M. Gorbahn and U. Haisch, Nucl. Phys. B **713** (2005) 291 [arXiv:hep-ph/0411071].
- [8] M. Gorbahn, U. Haisch and M. Misiak, Phys. Rev. Lett. **95** (2005) 102004 [arXiv:hep-ph/0504194].
- [9] K. Bieri, C. Greub and M. Steinhauser, Phys. Rev. D **67** (2003) 114019 [arXiv:hep-ph/0302051].
- [10] I. Blokland, A. Czarnecki, M. Misiak, M. Slusarczyk and F. Tkachov, Phys. Rev. D **72** (2005) 033014 [arXiv:hep-ph/0506055].
- [11] K. Melnikov and A. Mitov, Phys. Lett. B **620** (2005) 69 [arXiv:hep-ph/0505097].
- [12] H. M. Asatrian, C. Greub, A. Hovhannisyanyan, T. Hurth and V. Poghosyan, Phys. Lett. B **619** (2005) 322 [arXiv:hep-ph/0505068].