

THE SEARCH FOR SOLAR AXIONS IN THE CAST EXPERIMENT

DONGHWA KANG^{10, a}, S. ANDRIAMONJE², V. ARSOV¹³, S. AUNE², D. AUTIERO¹,
 F. T. AVIGNONE³, K. BARTH¹, A. BELOV¹¹, B. BELTRÁN⁶, H. BRÄUNINGER⁵,
 J. M. CARMONA⁶, S. CEBRIÁN⁶, E. CHESI¹, J. I. COLLAR⁷, R. CRESWICK³, T. DAFNI⁴,
 M. DAVENPORT¹, L. Di LELLA¹, C. ELEFThERIADIS⁸, J. ENGLHAUSER⁵, G. FANOURLAKIS⁹,
 H. FARACH³, E. FERRER², H. FISCHER¹⁰, J. FRANZ¹⁰, P. FRIEDRICH⁵, T. GERALIS⁹,
 I. GIOMATARIS², S. GNINENKO¹¹, N. GOLOUBEV¹¹, M. D. HASINOFF¹², F. H. HEINSIUS¹⁰,
 D. H. H. HOFFMANN⁴, I. G. IRASTORZA², J. JACOBY¹³, K. KÖNIGSMANN¹⁰, R. KOTTHAUS¹⁴,
 M. KRČMAR¹⁵, K. KOUSOURIS⁹, M. KUSTER⁵, B. LAKIĆ¹⁵, C. LASSEUR¹, A. LIOLIOS⁸,
 A. LJUBIČIĆ¹⁵, G. LUTZ¹⁴, G. LUZÓN⁶, D. W. MILLER⁷, A. MORALES⁶, J. MORALES⁶,
 M. MUTTERER⁴, A. NIKOLAIDIS⁸, A. ORTIZ⁶, T. PAPAEVANGELOU¹, A. PLACCI¹,
 G. RAFFELT¹⁴, J. RUZ⁶, H. RIEGE⁴, M. L. SARSA⁶, I. SAVVIDIS⁸, P. SERPICO¹⁴,
 Y. SEMERTZIDIS⁴, L. STEWART¹, J. D. VIEIRA⁷, J. VILLAR⁶, J. VOGEL¹⁰, L. WALCKIERS¹,
 K. ZACHARIADOU⁹ and K. ZIOUTAS¹⁶

(CAST Collaboration)

¹European Organization for Nuclear Research (CERN), Genève, Switzerland

²DAPNIA, Centre d'Études Nucléaires de Saclay (CEA-Saclay), Gif-sur-Yvette, France

³Department of Physics and Astronomy, University of South Carolina, Columbia, South Carolina, USA

⁴GSI-Darmstadt and Institut für Kernphysik, Technische Universität Darmstadt, Darmstadt, Germany

⁵Max-Planck-Institut für Extraterrestrische Physik, Garching, Germany

⁶Instituto de Física Nuclear y Altas Energías, Universidad de Zaragoza, Zaragoza, Spain

⁷Enrico Fermi Institute and KICP, University of Chicago, Chicago, Illinois, USA

⁸Aristotle University of Thessaloniki, Thessaloniki, Greece

⁹National Center for Scientific Research "Demokritos", Athens, Greece

¹⁰Albert-Ludwigs-Universität Freiburg, Freiburg, Germany

¹¹Institute for Nuclear Research (INR), Russian Academy of Sciences, Moscow, Russia

¹²Department of Physics and Astronomy, University of British Columbia, Vancouver, Canada

¹³Johann Wolfgang Goethe-Universität, Institut für Angewandte Physik, Frankfurt am Main, Germany

¹⁴Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Munich, Germany

¹⁵Rudjer Bošković Institute, Zagreb, Croatia

¹⁶Department of Physics, University of Patras, Patras, Greece

The CAST (CERN Axion Solar Telescope) experiment at CERN searches for solar axions with energies in the keV range. It is possible that axions are produced in the core of the sun by the interaction of thermal photons with virtual photons of strong electromagnetic fields. In this experiment, the solar axions can be reconverted to photons in the transversal field of a 9 Tesla superconducting magnet. At both ends of the 10m-long dipole magnet three different X-ray detectors were installed, which are sensitive in the interesting photon energy range. Preliminary results from the analysis of the 2004 data are presented: $g_{a\gamma} < 0.9 \times 10^{-10} \text{ GeV}^{-1}$ at 95% C.L. for axion masses $m_a < 0.02 \text{ eV}$. At the end of 2005, data started to be taken with a buffer gas in the magnet pipes in order to extend the sensitivity to axion masses up to 0.8 eV.

^aattending speaker, e-mail: donghwa.kang@cern.ch

1 Introduction

In the standard model the QCD Lagrangian¹ can be written with a gluon interaction term violating charge conjugation times parity (CP) and time reversal (T)

$$L_{QCD} = L_{pert} + \bar{\theta} \frac{g^2}{32\pi^2} G_a^{\mu\nu} \tilde{G}_{\mu\nu}^a, \quad (1)$$

where $G_a^{\mu\nu}$ is the gluon field strength tensor, $\tilde{G}_{\mu\nu}^a$ is the corresponding dual tensor, and the parameter $\bar{\theta}$ represents the effective QCD vacuum² in the basis where all quark masses are real. $\bar{\theta}$ can attain any value between 0 and 2π and a nonzero $\bar{\theta}$ value would imply CP violation. An observable evidence of CP violation would be an electric dipole moment of neutrons, which has theoretically been estimated³ to $d_n \approx 5 \times 10^{-16} \bar{\theta}$ e cm. The present experimental upper limit⁴, however, is smaller than $d_n < 3 \times 10^{-26}$ e cm. Consequently, the arbitrary parameter $\bar{\theta}$ should be smaller than 10^{-10} . Are there physical reasons for $\bar{\theta}$ being fixed so close to zero? This open question is referred to as the strong CP problem.

An elegant solution of the strong CP problem has been proposed by Peccei and Quinn⁵ in 1977 by introducing an additional global chiral symmetry $U(1)_{PQ}$, which is known as PQ-symmetry. This symmetry is spontaneously broken at an unknown scale f_a . A new, light, and neutral pseudoscalar particle could thus arise, namely an axion⁶ associated with the spontaneous symmetry breaking. The axion field gives an additional term L_a in the QCD Lagrangian, and Eq. (1) reads

$$L_{QCD} = L_{pert} + \bar{\theta} \frac{g^2}{32\pi^2} G_a^{\mu\nu} \tilde{G}_{\mu\nu}^a + L_a \quad (2)$$

$$L_a = -\frac{1}{2}(\partial_\mu a)^2 + C_a \frac{a}{f_a} \frac{g^2}{32\pi^2} G_a^{\mu\nu} \tilde{G}_{\mu\nu}^a, \quad (3)$$

where a is the axion field, f_a is the scale of the spontaneous PQ symmetry breaking and C_a is a model dependent constant. The presence of the second term in Eq. (3) provides an effective potential for the axion field. At the potential minimum $\langle a \rangle = -(f_a/C_a)\bar{\theta}$ of this Lagrangian, the CP violating term is eliminated, so that the strong CP problem is solved in the presence of an axion.

All important axion properties depend on the PQ symmetry breaking scale. The axion coupling to ordinary matter is proportional to the axion mass m_a and, equivalently, to the inverse of the PQ scale $1/f_a$. Axions are generated without mass, however, they get an effective mass by their interaction with gluons. The axion-gluon vertex induces an axion to quark-antiquark transition, therefore axions and neutral pions are mixing with each other. The axion mass has been estimated by Bardeen and Tye⁷ using the current algebra method and is given by

$$m_a = \frac{f_\pi m_{\pi^0}}{f_a} \frac{\sqrt{z}}{1+z} = 0.6 \text{eV} \frac{10^7 \text{GeV}}{f_a}, \quad (4)$$

where $m_{\pi^0} = 135$ MeV, $f_\pi = 93$ MeV, and $z = m_u/m_d = 0.56$ are the pion mass, the pion decay constant, and the mass ratio of up and down quarks, respectively.

Most experimental searches for axions focus on the axion interaction with two photons which is described by

$$L_{int} = -\frac{1}{4} g_{a\gamma} F_{\mu\nu} \tilde{F}^{\mu\nu} a = g_{a\gamma} \mathbf{E} \cdot \mathbf{B} a, \quad (5)$$

where F is the electromagnetic field strength tensor, \tilde{F} its dual, \mathbf{E} and \mathbf{B} the electric and magnetic field, and a the axion field. The axion photon coupling constant is

$$g_{a\gamma} = \frac{\alpha}{2\pi f_a} \left[\frac{E}{N} - \frac{2(4+z)}{3(1+z)} \right], \quad (6)$$

where E/N is a model dependent parameter⁸. This coupling is usually bound by the energy loss arguments of stars⁹ and a constraint from globular cluster stars¹⁰ is $g_{a\gamma} \leq 10^{-10} \text{GeV}^{-1}$. CAST is sensitive up to this range. Axions decay to two photons with a mean lifetime $\tau_{a \rightarrow 2\gamma} \sim \tau_{\pi^0 \rightarrow 2\gamma} (m_\pi/m_a)^5 \approx 10^{24}$ s for 1 eV axion mass. Hence, axions are quite stable; they live longer than the present estimated age of the universe of 10^{17} s. Therefore, axions would be excellent candidates for cold dark matter and possibly constitute some of cosmic dark matter¹¹.

2 The CAST experiment

Axions could be produced at the core of the sun and other stars by interacting with thermal photons in the Coulomb field of electric charges, namely by Primakoff¹² conversion. The expected solar axion flux on earth is $\Phi_a = 3.67 \times 10^{11} \text{cm}^{-2} \text{s}^{-1} (g_{a\gamma}/10^{-10} \text{GeV}^{-1})^2$ with an approximate spectrum¹³

$$\frac{d\Phi_a}{dE_a} = 3.821 \times 10^{10} \left(\frac{g_{a\gamma}}{10^{-10} \text{GeV}^{-1}} \right)^2 \frac{(E_a/\text{keV})^3}{e^{E_a/1.103\text{keV}} - 1} \text{cm}^{-2} \text{s}^{-1} \text{keV}^{-1} \quad (7)$$

and an average energy of 4.2 keV. In a transverse magnetic field, the axions can be reconverted to X-ray photons which have the energy and momentum of the original axions. The conversion probability¹⁴ of an axion to a photon in a strong magnetic field is

$$P_{a \rightarrow \gamma} = 1.7 \times 10^{-17} \left(\frac{\mathbf{B} \cdot \mathbf{L}}{9.0\text{T} \cdot 9.26\text{m}} \right)^2 \left(\frac{g_{a\gamma}}{10^{-10} \text{GeV}^{-1}} \right)^2 \frac{\sin^2(qL)}{(qL)^2}, \quad (8)$$

where B and L are the magnetic field strength and magnet length, respectively, $q = m_a^2/2E_a$ is the momentum difference between the axion and the photon in vacuum. Full coherence over the length of the magnet requires $qL \ll 1$. This constrains the range of axion masses detectable in the CAST experiment to be $m_a < 0.02$ eV. This is valid for the evacuated superconductive magnet with parameter L being 9.26 m. The expected number of photons reaching the detector is $N_\gamma = \Phi_a \cdot A \cdot P_{a \rightarrow \gamma} \approx 7$ events per day for $g_{a\gamma} = 10^{-10} \text{GeV}^{-1}$ and the magnet bore area $A = 14.5 \text{cm}^2$. By filling the beam pipe inside the magnet with a buffer gas, the X-ray photon acquires an effective mass and the coherence can be restored for a narrow mass range. The experimental sensitivity can thus be extended to higher axion masses of about 1 eV, proper gas pressure settings assumed.

The first experiment based on the axion helioscope principle proposed by Sikivie¹⁵ was performed at BNL. Later on, the Tokyo axion helioscope¹⁶ with $L = 2.3$ m and $B = 3.9$ T obtained $g_{a\gamma} < 6.0 \times 10^{-10} \text{GeV}^{-1}$ at 95% C.L. for $m_a < 0.03$ eV in vacuum and $g_{a\gamma} < 6.8 - 10.9 \times 10^{-10} \text{GeV}^{-1}$ at 95% C.L. for $m_a < 0.3$ eV using a buffer gas with different pressures. Limits from crystal detectors are much less restrictive¹⁷⁻¹⁹.

The CAST experiment searches for solar axions of energies of a few keV. The axions from the core of the sun could convert back into photons in the 9 Tesla Large Hadron Collider (LHC) prototype superconducting magnet²⁰, where the two beam pipes of length 9.26 m inside the magnet are straight with an effective cross sectional area $2 \times 14.5 \text{cm}^2$. Each aperture of the bores fully covers the potentially axion-emitting solar core (about 1/10th of the solar radius). A full

cryogenic system²¹ is used to cool the superconducting magnet down to 9 T. The magnet can be moved automatically to follow the sun track. The data were taken every morning and evening for approximately 1.5 hours by three different kinds of detectors installed at both ends of the dipole magnet, which are sensitive in the energy range up to 10 keV. The time where the magnet is not aligned with the sun is dedicated to background measurements. The CAST tracking system has been accurately calibrated by geometric survey measurements; the pointing precision is better than 0.01° . A plexiglas Time Projection Chamber (TPC) covering both beam bores looks for photons coming from the magnet after axion-to-photon conversion during sunset, while during sunrise, the photons are detected with a MICROMesh Gaseous Structure (MicroMegas)²² and a position-sensitive Charge Coupled Device (CCD)²³, installed at the other end of the magnet. The reason for using different detectors based on complementary techniques is the possibility of cross check and their systematic effects. An X-ray focusing mirror telescope designed for the German X-ray satellite mission ABRIXAS is mounted between the magnet and the CCD. It consists of 27 gold-coated mirror shells with a focal length of 1.6 m and the total efficiency is approximately 35% in the energy range of 1 to 7 keV. The X-ray mirror telescope focuses the photons from the magnet bore of 14.5 cm^2 aperture to a spot size of about 6 mm^2 on the CCD, and thus improving the signal to background ratio by more than 2 orders of magnitude. First results were obtained from the 2003 data taking¹³. Here we present new results from 2004.

3 Detector improvements for 2004 operation

All CAST detectors were operated almost throughout the 2004 data collection periods with improved conditions. The differences between 2003 and 2004 measurements were mainly due to an upgraded experimental setup and additional shielding for detectors. Here the detector improvements for the 2004 operation are described in detail. The technical aspects of the detectors can be found in Refs.^{24 25}.

TPC : In 2004 the TPC improvements focused mainly on two points: a differential pumping system for the TPC X-ray windows as well as passive and active detector shielding. The aim of the differential pumping system is to effectively decrease the gas leaks toward the magnet and to reduce the damage to the thin windows due to unexpected pressure changes or breakdown. By the continuous pumping of an intermediate volume, the leak rate has been reduced, and thus the total time for data taking during 2004 is five times more than 2003. A passive shielding for the TPC was installed in 2004. It consists of an innermost copper cage of thickness 5 mm, surrounded by a layer of 2.5 cm ancient lead, a layer of 1 mm cadmium, and finally a wall of 22.5 cm polyethylene. A plastic box covering the full setting is slightly pressurized by injection of clean nitrogen to reduce the radon contamination from the air in the space close to the detector. Moreover, an active shield has been installed on top of the TPC to clearly identify and reject the muons producing background. The full detector shielding of the TPC has reduced the average background level by a factor 2.4 as compared to 2003. In 2004 the background counting rate was about $4 \times 10^{-5} \text{ counts cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$ in the energy range of the solar axion spectrum.

MicroMegas : A new version of the MicroMegas detector and new electronics have been installed for the 2004 data taking period. The new MicroMegas was designed to reduce the cross talk effect of the read out strips and thus, the quality of the data is improved. In addition, a faster VME Digitizing Board, which records the time structure of the mesh signal, was installed and has improved the data transfer, reducing the detector dead time. An automatic controller has been installed in order to control the calibration source directly from the data acquisition system to avoid manual operations. The improved MicroMegas detector reduces the background level to about $5 \times 10^{-5} \text{ counts cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$ in the energy range of 1 to 8.5 keV.

CCD : In 2004 the CCD detector was operated with an extra shield inside the detector chamber. The detector shield was modified by adding a 17 to 25 mm thick layer of low activity ancient lead outside of the copper shielding. The lead shield inside the CCD chamber is encapsulated in sealed boxes made of high purity copper. The additional shielding has reduced the background level by a factor of 1.5 as compared to the previous setup. The differential mean background rate is about 7.5×10^{-5} counts $\text{cm}^{-2} \text{s}^{-1} \text{keV}^{-1}$ with a flat distribution over the 1 to 7 keV energy range. Very important for the CCD setup is the monitoring of the position of the solar spot on the CCD. An X-ray source based on a pyroelectric crystal producing a peak activity of 100 MBq is mounted on a manipulator at the TPC end of the CAST magnet more than 10 m away from the telescope focal plane. The image of this source allows one to monitor the stability of the alignment of the X-ray telescope and correspondingly to monitor the position of the solar spot on the CCD. In addition, using a parallel laser beam which is focused by the X-ray telescope on to the CCD, the location of the expected solar axion signal on the CCD can be determined. From the regularly performed laser and X-ray measurements, the center of the spot was determined and it was shown to be stable within one pixel $(150\mu\text{m})^2$ diameter of the CCD.

4 Data analysis and results

All detectors were operated with improved conditions for the whole 2004 data collection period. Each detector acquired data during 1.5 h per day when the magnet was pointing to the sun. Background measurements were taken during non alignment periods and have about a 10 fold exposure time. The data analysis procedures are along the guideline given in Ref. ¹³. The theoretically expected spectrum for axions in the allowed mass range (< 0.02 eV) has been estimated and detector and software efficiencies have been applied. For the TPC and the MicroMegas these spectra which are proportional to $g_{a\gamma}^4$ are directly used as fit functions to the experimental spectra after background subtraction. For the CCD data collected in 2004 the analysis could be restricted to a small 6 mm^2 spot on the CCD since the telescope stability was continuously monitored. Due to the low counting statistics on the spot a likelihood function for the expected signal plus background was required instead of using a χ^2 analysis. All the data sets analyzed for the different detectors are consistent with the absence of a signal and therefore 95% confidence limits on $g_{a\gamma}$ for each of the data sets have been extracted. The preliminary results can be combined by multiplying the Bayesian probability distribution and integrating them over the physically allowed region, i.e. positive signals, to find the combined preliminary result for the 2004 CAST data:

$$g_{a\gamma} < 0.9 \times 10^{-10} \text{ GeV}^{-1} \text{ (95\% C.L.)}.$$

This result is limited to the axion mass range $m_a < 0.02$ eV where the expected signal is independent of the mass since the axion photon oscillation length is larger than the length of the magnet. For higher masses, the number of expected signal counts decreases rapidly and the shape of the spectral curve differs. This analysis procedure was repeated for different values of axion masses to derive the whole exclusion line for 95% C.L. as shown in Figure 1. The systematic uncertainties are estimated to have an effect of less than about 10% on the final results.

5 Summary and outlook

The CAST experiment has been operated successfully since May 2003 and during 2004 all three detectors were taking data in upgraded versions with higher sensitivity. No signal above




Figure 1: Exclusion limit at 95% C.L. from the 2004 CAST data compared with constraints from other experiments. The shaded region represents the theoretical models. The future CAST sensitivity is shown as prospects.

background was observed in any of the data taken so far. First results of the 2003 data have been published¹³. The preliminary CAST results of the 2004 data yield a lower limit on $g_{a\gamma} < 0.9 \times 10^{-10} \text{ GeV}^{-1}$ at 95% C.L. for axion masses $m_a < 0.02 \text{ eV}$. This result improves the previous constraints given by other experiments by a factor of 7. It is the first time that an experimental limit falls below the astrophysical limit of globular clusters. So far the CAST experiment has been operated with an evacuated magnet bore and thus was restricted to axion masses below 0.02 eV. During 2005 a major modification to the magnet pipe system was undertaken. End of 2005 CAST started with measurements for its second phase. The magnet pipes were filled with a low Z buffer gas (starting with ^4He and in the future ^3He) of various pressures in order to restore coherence for higher axion masses $m_a > 0.02 \text{ eV}$. The extended sensitivity will allow CAST to reach masses up to 0.8 eV and thus reach into the region of theoretical axion models.

References

1. C. Quigg, *Gauge Theories of the Strong, Weak and Electromagnetic Interactions* (Addison Wesley, 1982).
2. C. Callan, R. Dashen and D. Gross, *Phys. Lett. B* **63**, 334 (1976).
3. V. Baluni, *Phys. Rev. D* **19**, 2227 (1979).
4. W. B. Dress *et al.*, *Phys. Rev. D* **15**, 9 (1977); C. A. Baker *et al.*, hep-ex/0602020.
5. R. D. Peccei and H. R. Quinn, *Phys. Rev. D* **38**, 1440 (1977).
6. S. Weinberg, *Phys. Rev. Lett.* **40**, 223 (1978).
7. W. A. Bardeen and S.-H. H. Tye, *Phys. Lett. B* **74**, 229 (1978).
8. J. E. Kim, *Phys. Rep.* 150 (1987).
9. G. G. Raffelt, *Stars as Laboratories for Fundamental Physics* (Univ. of Chicago Press, 1996).
10. G. G. Raffelt, *Phys. Rep.* 198 (1990).

11. E. W. Kolb and M. S. Turner, *The Early Universe* (Addison Wesley, 1990).
12. H. Primakoff, *Phys. Rev.* **81**, 899 (1951).
13. K. Zioutas *et al.*, *Phys. Rev. Lett.* **94**, 121301 (2005).
14. K. van Bibber *et al.*, *Phys. Rev. Lett.* **59**, 759 (1987).
15. P. Sikivie, *Phys. Rev. Lett.* **51**, 1415 (1983).
16. S. Moriyama *et al.*, *Phys. Lett. B* **434**, 147 (1998).
17. F. T. Avignone *et al.*, *Phys. Rev. Lett.* **81**, 5068 (1998).
18. A. Morales *et al.*, *Astropart. Phys.* **16** (2002) 325.
19. R. Bernabei *et al.*, *Phys. Lett. B* **515**, 6 (2001).
20. K. Zioutas *et al.*, *Nucl. Instrum. Methods A* **425**, 480 (1999).
21. K. Barth *et al.*, Divisional Report, CERN LHC/2002-17 (ACR/ECR/MTA).
22. Y. Giomataris *et al.*, *Nucl. Instrum. Methods A* **376**, 29 (1996).
23. L. Strüder, *Nucl. Instrum. Methods A* **454**, 73 (2000).
24. G. Lutz *et al.*, *Nucl. Instrum. Methods A* **518**, 252 (2004); M. Kuster *et al.*, Proceedings of SPIE, vol. 5500 (2004) 139; R. Kotthaus *et al.*, astro-ph/0511390.
25. S. Andriamonje *et al.*, *Nucl. Instrum. Methods A* **518**, 252 (2004).