

L3 Results up to 208 GeV

Maria Novella Kienzle-Focacci


Université de Genève

On behalf of the L3 Collaboration


October 10th, 2000

Most of the results are preliminary.

In parallel with the main physics stream.




Differential momentum spectrum for vertical muons




Coincidence with shower array

Final result: 10^{10} events, $20 \text{ GeV} \leq p_\mu \leq 2 \text{ TeV}$,
 $<1\%$ stat and 2.5% syst errors.



Accurate measurements necessary for atmospheric ν .



Fermion Pair Production - Leptons



Hadronic $\gamma-Z$ interference term $j_{\text{had}}^{\text{tot}}$




Data	$j_{\text{had}}^{\text{tot}}$	M_Z [MeV]
Z pole	0.44 ± 0.59	91185.2 ± 10.3
$\sqrt{s} \rightarrow 208$ GeV	0.29 ± 0.12	91188.4 ± 3.7
Standard Analysis	0.22 fixed	91189.5 ± 3.1



Contact interactions:



$$\mathcal{L} = \frac{1}{1+\delta_{ef}} \sum_{i,j=L,R} \eta_{ij} \frac{g^2}{\Lambda_{ij}^2} (\bar{e}_i \gamma^\mu e_i) (\bar{f}_j \gamma_\mu f_j)$$

L3 $e^+e^- \rightarrow l^+l^-$



$e^+e^- \rightarrow ff$





From single photon:


$N_\nu = 2.86 \pm 0.13$ (LEP 2)

$N_\nu = 2.98 \pm 0.10$ (LEP 1)

From Z width:

$N_\nu = 2.978 \pm 0.014$

Search for $e^+e^- \rightarrow \tilde{G}\tilde{\chi}_1^0 \rightarrow \tilde{G}\tilde{G}\gamma$ No signal observed





Limits on deviation from QED :

$$\Lambda > 1536 \text{ GeV} \quad m_{e^*} > 319 \text{ GeV}$$




Limits on graviton exchange:

$$M_S(\lambda = +1) > 0.84 \text{ TeV}$$


$$M_S(\lambda = -1) > 0.95 \text{ TeV}$$



W Gauge Couplings



Single W:
130 - 202 GeV





1-parameter fit:



$$g_1^z = +0.93 \pm 0.06$$

$$\kappa_\gamma = +0.96 \pm 0.12$$

$$\lambda_\gamma = -0.08 \pm 0.06$$




CP-violating [1/GeV²]:
 $-0.41 < a_n/\Lambda^2 < 0.37$




CP-conserving [1/GeV²]:
 $-0.008 < a_0/\Lambda^2 < 0.005$
 $-0.006 < a_c/\Lambda^2 < 0.012$

$WW \rightarrow q\bar{q}\ell\nu$



$WW \rightarrow q\bar{q}q\bar{q}$



172 GeV – 202 GeV data:

$q\bar{q}q\bar{q}$

$80.46 \pm 0.08 \pm 0.07$ GeV

$q\bar{q}\ell\nu$

$80.27 \pm 0.09 \pm 0.05$ GeV


Mass difference

$0.19 \pm 0.12 \pm 0.05$ GeV

calculated without FSI systematics


All combined

$80.375 \pm 0.058 \pm 0.051$ GeV




Cross talk in $W^+W^- \rightarrow q\bar{q}q\bar{q}$?

Distribution of identical pions close in phase space:



$Q = 4$ -momentum difference of identical pions.




$$D'(Q) = (1 + \epsilon Q) (1 + \Lambda \exp(-k^2 Q^2))$$

$$\Lambda = 0.013 \pm 0.023$$

$$\Delta M_W \leq 20 \text{ MeV}$$


Fit of the Standard Model to all L3 electro-weak data:



$$M_W = 80.52 \pm 0.08 \text{ GeV}$$

$$M_{\text{top}} = 195^{+31}_{-17} \text{ GeV}$$

within 1σ of measured values




$$M_H = 58^{+73}_{-33} \text{ GeV}$$

Higgs must be around the corner !
or "New Physics" !

Discriminant variable combining:
b-tag, neural network, recoil mass to Z.


Total luminosity 170.3 pb^{-1}
 82.9 pb^{-1} for $\sqrt{s} \leq 206.5 \text{ GeV}$
 87.4 pb^{-1} for $\sqrt{s} > 206.5 \text{ GeV}$

For qqqq selection:




Final Discriminant

Combining $qqqq$, $qq\nu\nu$ and all $q\bar{q}ll$ channels




$$Q = \frac{L(s+b)}{L(b)}$$




$M_H > 113.2$ GeV
 at 95 % CL
 expected limit : 111.2 GeV

Combining $qqqq$, $qq\nu\nu$ and all $qql\bar{l}$ channels
 Use mass-independent cuts
 S/B defined for recoil mass > 109 GeV

L3 preliminary ($\sqrt{s} = 200-209$ GeV)






$\sqrt{s} = 206.7 \text{ GeV}$, S/B = 2 , btag = 3.2

5-C fit $e^+e^- \rightarrow HZ$: $M_H = 109.9 \text{ GeV}$



5-C fit with equal masses : 89.3 GeV




$\sqrt{s} = 205.4 \text{ GeV}, S/B = 2$

5-C fit $e^+e^- \rightarrow HZ$: $M_H = 105.5 \text{ GeV}$
4-C fit: $M_H = 108.0 \text{ GeV}$

Preliminary analysis looking for the channels:






Run 868610, Event 378, $\sqrt{s} = 206.8$ GeV

$$\sqrt{s} = 183 - 209 \text{ GeV}$$


Require genuine $e^+e^- \rightarrow \text{qqqq}$ events :
reduce QCD (qqgg) and W-pair background
5-C fit with equal mass (M) constraint



Excess of events at $\simeq 68$ GeV.

Expectation calculated for the reaction $e^+e^- \rightarrow H^+H^-$ with the HZHA Monte Carlo.

$$\sqrt{s} = 183 - 209 \text{ GeV}$$




The effect
 (2.7σ above background in 183-202 GeV)
 increases to 3.6σ

$$\Delta M = M_{\tilde{\chi}_1^\pm} - M_{\tilde{\chi}_1^0} > 3 \text{ GeV}$$

Topologies: $l^+l^- + \cancel{E}$, $ljj + \cancel{E}$, $jjjj + \cancel{E}$

ΔM (GeV)	3–10		20–40		> 50		.OR.	
\sqrt{s} (GeV)	data	exp	data	exp	data	exp	data	exp
208	0	1.6	2	1.2	0	0.9	2	3.7
205–208	40	35.8	26	27.8	17	19.9	83	83.6

Gaugino like and $M_{\tilde{\nu}} > 300$ GeV :
 Kinematic Limit reached




$$\Delta M < 3 \text{ GeV}$$


Tag chargino decays with ISR photons

ΔM	$\sim 3 \text{ GeV}$		$\sim 1 \text{ GeV}$		$\sim 0.3 \text{ GeV}$.OR.	
$\sqrt{s} \text{ (GeV)}$	data	exp	data	exp	data	exp	data	exp
205-208	6	5.6	2	2.0	3	1.2	8	6.8

Higgsino Like (CMSSM): Mass limit independant of ΔM



For $\mu = -200 \text{ GeV}$, $\tan \beta = 2$, $M_{\tilde{\chi}_1^0} > 15 \text{ GeV}$, $\Delta M > 10 \text{ GeV}$




Indirect limits from slepton, chargino, neutralino:

$M_{\tilde{\chi}_1^0} > 39.4 \text{ GeV}$

Search for $\tilde{\chi}_1^\pm$, $\tilde{\chi}_{J=1,4}^0$ and $\tilde{\ell}$


$\tilde{\chi}_1^0$ is no more stable \rightarrow final states with:
many leptons (λ), many jets (λ'') or both (λ').

Including 2000 data: (λ'' only)





Limits on $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^0$ similar to the one obtained assuming R–Parity conservation

Running α_S from 4 event shape variables.
 QCD fits to $\mathcal{O}(\alpha_S^2)$ with resummed LO and NLO terms.



$$\alpha_S(M_Z) = 0.1218 \pm 0.0012(\text{exp}) \pm 0.0061(\text{th})$$


Number of active flavours: $5.0 \pm 1.3(\text{exp}) \pm 2.0(\text{th})$




DLA : $\chi^2 = 57$ for 16 points

MLLA : $\chi^2 = 16$ for 16 points


Evidence of gluon coherence




VDM



Direct




Single Resolved



BFKL


$$e^+ e^- \rightarrow e^+ e^- c\bar{c}, b\bar{b} X$$

NLO QCD: $\gamma\gamma \rightarrow q\bar{q}$ (direct)
 $g\gamma \rightarrow q\bar{q}$ (single resolved)




$$\sigma(e^+ e^- \rightarrow e^+ e^- b\bar{b} X) = 14 \pm 3 \text{ pb}$$

(Expected 4.6 pb)




$$\langle Q_1^2 \rangle = \langle Q_2^2 \rangle = 15 \text{ GeV}^2$$




Test of BFKL models

$$\gamma\gamma \rightarrow \Lambda\bar{\Lambda}$$



Test of diquark models for baryon production.



The Standard Model is a wonderful construction...

The San Petronio church in Bologna

... let's complete it!

Long life to LHC and to Linear Colliders