



$$I(J^P) = \frac{1}{2}(0^-)$$

## $D^\pm$ MASS

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1869.61 ± 0.10 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>1869.5 ± 0.4 OUR AVERAGE</b>				
1869.53 ± 0.49 ± 0.20	110 ± 15	ANASHIN	10A KEDR	$e^+e^-$ at $\psi(3770)$
1870.0 ± 0.5 ± 1.0	317	BARLAG	90C ACCM	$\pi^-$ Cu 230 GeV
1869.4 ± 0.6		<sup>1</sup> TRILLING	81 RVUE	$e^+e^-$ 3.77 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1875 ± 10	9	ADAMOVICH	87 EMUL	Photoproduction
1860 ± 16	6	ADAMOVICH	84 EMUL	Photoproduction
1863 ± 4		DERRICK	84 HRS	$e^+e^-$ 29 GeV
1868.4 ± 0.5		<sup>1</sup> SCHINDLER	81 MRK2	$e^+e^-$ 3.77 GeV
1874 ± 5		GOLDHABER	77 MRK1	$D^0$ , $D^+$ recoil spectra
1868.3 ± 0.9		<sup>1</sup> PERUZZI	77 LGW	$e^+e^-$ 3.77 GeV
1874 ± 11		PICCOLO	77 MRK1	$e^+e^-$ 4.03, 4.41 GeV
1876 ± 15	50	PERUZZI	76 MRK1	$K^\mp \pi^\pm \pi^\pm$

<sup>1</sup>PERUZZI 77 and SCHINDLER 81 errors do not include the 0.13% uncertainty in the absolute SPEAR energy calibration. TRILLING 81 uses the high precision  $J/\psi(1S)$  and  $\psi(2S)$  measurements of ZHOLENTZ 80 to determine this uncertainty and combines the PERUZZI 77 and SCHINDLER 81 results to obtain the value quoted.

## $D^\pm$ MEAN LIFE

Measurements with an error  $> 100 \times 10^{-15}$  s have been omitted from the Listings.

VALUE ( $10^{-15}$ s)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1040 ± 7 OUR AVERAGE</b>				
1039.4 ± 4.3 ± 7.0	110k	LINK	02F FOCS	$\gamma$ nucleus, $\approx 180$ GeV
1033.6 ± 22.1 <sup>+9.9</sup> <sub>-12.7</sub>	3777	BONVICINI	99 CLEO	$e^+e^- \approx \Upsilon(4S)$
1048 ± 15 ± 11	9k	FRABETTI	94D E687	$D^+ \rightarrow K^- \pi^+ \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1075 ± 40 ± 18	2455	FRABETTI	91 E687	$\gamma$ Be, $D^+ \rightarrow K^- \pi^+ \pi^+$
1030 ± 80 ± 60	200	ALVAREZ	90 NA14	$\gamma$ , $D^+ \rightarrow K^- \pi^+ \pi^+$
1050 <sup>+77</sup> <sub>-72</sub>	317	<sup>1</sup> BARLAG	90C ACCM	$\pi^-$ Cu 230 GeV
1050 ± 80 ± 70	363	ALBRECHT	88i ARG	$e^+e^-$ 10 GeV
1090 ± 30 ± 25	2992	RAAB	88 E691	Photoproduction

<sup>1</sup>BARLAG 90C estimates the systematic error to be negligible.

**$D^+$  DECAY MODES**

Most decay modes (other than the semileptonic modes) that involve a neutral  $K$  meson are now given as  $K_S^0$  modes, not as  $\bar{K}^0$  modes. Nearly always it is a  $K_S^0$  that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that  $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$ .

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
<b>Inclusive modes</b>		
$\Gamma_1$ $D^+ \rightarrow e^+$ semileptonic	$(16.07 \pm 0.30) \%$	
$\Gamma_2$ $D^+ \rightarrow \mu^+$ anything	$(17.6 \pm 3.2) \%$	
$\Gamma_3$ $D^+ \rightarrow K^-$ anything	$(25.7 \pm 1.4) \%$	
$\Gamma_4$ $D^+ \rightarrow \bar{K}^0$ anything + $K^0$ anything	$(61 \pm 5) \%$	
$\Gamma_5$ $D^+ \rightarrow K^+$ anything	$(5.9 \pm 0.8) \%$	
$\Gamma_6$ $D^+ \rightarrow K^*(892)^-$ anything	$(6 \pm 5) \%$	
$\Gamma_7$ $D^+ \rightarrow \bar{K}^*(892)^0$ anything	$(23 \pm 5) \%$	
$\Gamma_8$ $D^+ \rightarrow K^*(892)^0$ anything	$< 6.6 \%$	CL=90%
$\Gamma_9$ $D^+ \rightarrow \eta$ anything	$(6.3 \pm 0.7) \%$	
$\Gamma_{10}$ $D^+ \rightarrow \eta'$ anything	$(1.04 \pm 0.18) \%$	
$\Gamma_{11}$ $D^+ \rightarrow \phi$ anything	$(1.03 \pm 0.12) \%$	
<b>Leptonic and semileptonic modes</b>		
$\Gamma_{12}$ $D^+ \rightarrow e^+ \nu_e$	$< 8.8 \times 10^{-6}$	CL=90%
$\Gamma_{13}$ $D^+ \rightarrow \mu^+ \nu_\mu$	$(3.82 \pm 0.33) \times 10^{-4}$	
$\Gamma_{14}$ $D^+ \rightarrow \tau^+ \nu_\tau$	$< 1.2 \times 10^{-3}$	CL=90%
$\Gamma_{15}$ $D^+ \rightarrow \bar{K}^0 e^+ \nu_e$	$(8.83 \pm 0.22) \%$	
$\Gamma_{16}$ $D^+ \rightarrow \bar{K}^0 \mu^+ \nu_\mu$	$(9.2 \pm 0.6) \%$	
$\Gamma_{17}$ $D^+ \rightarrow K^- \pi^+ e^+ \nu_e$	$(4.00 \pm 0.10) \%$	
$\Gamma_{18}$ $D^+ \rightarrow \bar{K}^*(892)^0 e^+ \nu_e,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(3.68 \pm 0.10) \%$	
$\Gamma_{19}$ $D^+ \rightarrow (K^- \pi^+)_{S\text{-wave}} e^+ \nu_e$	$(2.32 \pm 0.10) \times 10^{-3}$	
$\Gamma_{20}$ $D^+ \rightarrow \bar{K}^*(1410)^0 e^+ \nu_e,$ $\bar{K}^*(1410)^0 \rightarrow K^- \pi^+$	$< 6 \times 10^{-3}$	CL=90%
$\Gamma_{21}$ $D^+ \rightarrow \bar{K}_2^*(1430)^0 e^+ \nu_e,$ $\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{22}$ $D^+ \rightarrow K^- \pi^+ e^+ \nu_e$ nonresonant	$< 7 \times 10^{-3}$	CL=90%
$\Gamma_{23}$ $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$	$(3.8 \pm 0.4) \%$	
$\Gamma_{24}$ $D^+ \rightarrow \bar{K}^*(892)^0 \mu^+ \nu_\mu,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(3.52 \pm 0.10) \%$	
$\Gamma_{25}$ $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	$(2.0 \pm 0.5) \times 10^{-3}$	

$\Gamma_{26}$	$D^+ \rightarrow K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	$< 1.6 \times 10^{-3}$	CL=90%
$\Gamma_{27}$	$D^+ \rightarrow \pi^0 e^+ \nu_e$	$(4.05 \pm 0.18) \times 10^{-3}$	
$\Gamma_{28}$	$D^+ \rightarrow \eta e^+ \nu_e$	$(1.14 \pm 0.10) \times 10^{-3}$	
$\Gamma_{29}$	$D^+ \rightarrow \rho^0 e^+ \nu_e$	$(2.18^{+0.17}_{-0.25}) \times 10^{-3}$	
$\Gamma_{30}$	$D^+ \rightarrow \rho^0 \mu^+ \nu_\mu$	$(2.4 \pm 0.4) \times 10^{-3}$	
$\Gamma_{31}$	$D^+ \rightarrow \omega e^+ \nu_e$	$(1.82 \pm 0.19) \times 10^{-3}$	
$\Gamma_{32}$	$D^+ \rightarrow \eta'(958) e^+ \nu_e$	$(2.2 \pm 0.5) \times 10^{-4}$	
$\Gamma_{33}$	$D^+ \rightarrow \phi e^+ \nu_e$	$< 9 \times 10^{-5}$	CL=90%

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\Gamma_{34}$	$D^+ \rightarrow \bar{K}^*(892)^0 e^+ \nu_e$	$(5.52 \pm 0.15) \%$	
$\Gamma_{35}$	$D^+ \rightarrow \bar{K}^*(892)^0 \mu^+ \nu_\mu$	$(5.28 \pm 0.15) \%$	
$\Gamma_{36}$	$D^+ \rightarrow \bar{K}_0^*(1430)^0 \mu^+ \nu_\mu$	$< 2.4 \times 10^{-4}$	CL=90%
$\Gamma_{37}$	$D^+ \rightarrow \bar{K}^*(1680)^0 \mu^+ \nu_\mu$	$< 1.5 \times 10^{-3}$	CL=90%

### Hadronic modes with a $\bar{K}$ or $\bar{K}K\bar{K}$

$\Gamma_{38}$	$D^+ \rightarrow K_S^0 \pi^+$	$(1.47 \pm 0.07) \%$	S=2.0
$\Gamma_{39}$	$D^+ \rightarrow K_L^0 \pi^+$	$(1.46 \pm 0.05) \%$	
$\Gamma_{40}$	$D^+ \rightarrow K^- 2\pi^+$	[a] $(9.13 \pm 0.19) \%$	
$\Gamma_{41}$	$D^+ \rightarrow (K^- \pi^+)_{S\text{-wave}} \pi^+$	$(7.32 \pm 0.19) \%$	
$\Gamma_{42}$	$D^+ \rightarrow \bar{K}_0^*(800)^0 \pi^+$ , $\bar{K}_0^*(800) \rightarrow K^- \pi^+$		
$\Gamma_{43}$	$D^+ \rightarrow \bar{K}_0^*(1430)^0 \pi^+$ , $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	[b] $(1.21 \pm 0.06) \%$	
$\Gamma_{44}$	$D^+ \rightarrow \bar{K}^*(892)^0 \pi^+$ , $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(1.01 \pm 0.11) \%$	
$\Gamma_{45}$	$D^+ \rightarrow \bar{K}^*(1410)^0 \pi^+$ , $\bar{K}^{*0} \rightarrow K^- \pi^+$	not seen	
$\Gamma_{46}$	$D^+ \rightarrow \bar{K}_2^*(1430)^0 \pi^+$ , $\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	[b] $(2.2 \pm 0.7) \times 10^{-4}$	
$\Gamma_{47}$	$D^+ \rightarrow \bar{K}^*(1680)^0 \pi^+$ , $\bar{K}^*(1680)^0 \rightarrow K^- \pi^+$	[b] $(2.1 \pm 1.1) \times 10^{-4}$	
$\Gamma_{48}$	$D^+ \rightarrow K^- (2\pi^+)_{I=2}$	$(1.41 \pm 0.26) \%$	
$\Gamma_{49}$	$D^+ \rightarrow K^- 2\pi^+$ nonresonant		
$\Gamma_{50}$	$D^+ \rightarrow K_S^0 \pi^+ \pi^0$	[a] $(6.99 \pm 0.27) \%$	
$\Gamma_{51}$	$D^+ \rightarrow K_S^0 \rho^+$	$(4.8 \pm 1.0) \%$	
$\Gamma_{52}$	$D^+ \rightarrow \bar{K}^*(892)^0 \pi^+$ , $\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	$(1.3 \pm 0.6) \%$	
$\Gamma_{53}$	$D^+ \rightarrow K_S^0 \pi^+ \pi^0$ nonresonant	$(9 \pm 7) \times 10^{-3}$	
$\Gamma_{54}$	$D^+ \rightarrow K^- 2\pi^+ \pi^0$	[c] $(5.99 \pm 0.18) \%$	

$\Gamma_{55}$	$D^+ \rightarrow K_S^0 2\pi^+ \pi^-$	[c]	$(3.12 \pm 0.11) \%$	
$\Gamma_{56}$	$D^+ \rightarrow K^- 3\pi^+ \pi^-$	[a]	$(5.6 \pm 0.5) \times 10^{-3}$	S=1.1
$\Gamma_{57}$	$D^+ \rightarrow \bar{K}^*(892)^0 2\pi^+ \pi^-$ , $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		$(1.2 \pm 0.4) \times 10^{-3}$	
$\Gamma_{58}$	$D^+ \rightarrow \bar{K}^*(892)^0 \rho^0 \pi^+$ , $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		$(2.2 \pm 0.4) \times 10^{-3}$	
$\Gamma_{59}$	$D^+ \rightarrow \bar{K}^*(892)^0 a_1(1260)^+$	[d]	$(9.0 \pm 1.8) \times 10^{-3}$	
$\Gamma_{60}$	$D^+ \rightarrow \bar{K}^*(892)^0 2\pi^+ \pi^-$ no- $\rho$ , $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$\Gamma_{61}$	$D^+ \rightarrow K^- \rho^0 2\pi^+$		$(1.68 \pm 0.27) \times 10^{-3}$	
$\Gamma_{62}$	$D^+ \rightarrow K^- 3\pi^+ \pi^-$ nonresonant		$(3.9 \pm 2.9) \times 10^{-4}$	
$\Gamma_{63}$	$D^+ \rightarrow K^+ 2K_S^0$		$(4.5 \pm 2.0) \times 10^{-3}$	
$\Gamma_{64}$	$D^+ \rightarrow K^+ K^- K_S^0 \pi^+$		$(2.4 \pm 0.6) \times 10^{-4}$	

### Pionic modes

$\Gamma_{65}$	$D^+ \rightarrow \pi^+ \pi^0$		$(1.19 \pm 0.06) \times 10^{-3}$	
$\Gamma_{66}$	$D^+ \rightarrow 2\pi^+ \pi^-$		$(3.18 \pm 0.18) \times 10^{-3}$	
$\Gamma_{67}$	$D^+ \rightarrow \rho^0 \pi^+$		$(8.1 \pm 1.5) \times 10^{-4}$	
$\Gamma_{68}$	$D^+ \rightarrow \pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$		$(1.78 \pm 0.16) \times 10^{-3}$	
$\Gamma_{69}$	$D^+ \rightarrow \sigma \pi^+$ , $\sigma \rightarrow \pi^+ \pi^-$		$(1.34 \pm 0.12) \times 10^{-3}$	
$\Gamma_{70}$	$D^+ \rightarrow f_0(980) \pi^+$ , $f_0(980) \rightarrow \pi^+ \pi^-$		$(1.52 \pm 0.33) \times 10^{-4}$	
$\Gamma_{71}$	$D^+ \rightarrow f_0(1370) \pi^+$ , $f_0(1370) \rightarrow \pi^+ \pi^-$		$(8 \pm 4) \times 10^{-5}$	
$\Gamma_{72}$	$D^+ \rightarrow f_2(1270) \pi^+$ , $f_2(1270) \rightarrow \pi^+ \pi^-$		$(4.9 \pm 0.9) \times 10^{-4}$	
$\Gamma_{73}$	$D^+ \rightarrow \rho(1450)^0 \pi^+$ , $\rho(1450)^0 \rightarrow \pi^+ \pi^-$	$< 8$	$\times 10^{-5}$	CL=95%
$\Gamma_{74}$	$D^+ \rightarrow f_0(1500) \pi^+$ , $f_0(1500) \rightarrow \pi^+ \pi^-$		$(1.1 \pm 0.4) \times 10^{-4}$	
$\Gamma_{75}$	$D^+ \rightarrow f_0(1710) \pi^+$ , $f_0(1710) \rightarrow \pi^+ \pi^-$	$< 5$	$\times 10^{-5}$	CL=95%
$\Gamma_{76}$	$D^+ \rightarrow f_0(1790) \pi^+$ , $f_0(1790) \rightarrow \pi^+ \pi^-$	$< 6$	$\times 10^{-5}$	CL=95%
$\Gamma_{77}$	$D^+ \rightarrow (\pi^+ \pi^+)_{S\text{-wave}} \pi^-$	$< 1.2$	$\times 10^{-4}$	CL=95%
$\Gamma_{78}$	$D^+ \rightarrow 2\pi^+ \pi^-$ nonresonant	$< 1.1$	$\times 10^{-4}$	CL=95%
$\Gamma_{79}$	$D^+ \rightarrow \pi^+ 2\pi^0$		$(4.6 \pm 0.4) \times 10^{-3}$	
$\Gamma_{80}$	$D^+ \rightarrow 2\pi^+ \pi^- \pi^0$		$(1.13 \pm 0.08) \%$	
$\Gamma_{81}$	$D^+ \rightarrow \eta \pi^+$ , $\eta \rightarrow \pi^+ \pi^- \pi^0$		$(8.0 \pm 0.5) \times 10^{-4}$	
$\Gamma_{82}$	$D^+ \rightarrow \omega \pi^+$ , $\omega \rightarrow \pi^+ \pi^- \pi^0$	$< 3$	$\times 10^{-4}$	CL=90%
$\Gamma_{83}$	$D^+ \rightarrow 3\pi^+ 2\pi^-$		$(1.61 \pm 0.16) \times 10^{-3}$	

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\Gamma_{84}$	$D^+ \rightarrow \eta \pi^+$	$(3.53 \pm 0.21) \times 10^{-3}$	
$\Gamma_{85}$	$D^+ \rightarrow \eta \pi^+ \pi^0$	$(1.38 \pm 0.35) \times 10^{-3}$	
$\Gamma_{86}$	$D^+ \rightarrow \omega \pi^+$	$< 3.4 \times 10^{-4}$	CL=90%
$\Gamma_{87}$	$D^+ \rightarrow \eta'(958) \pi^+$	$(4.67 \pm 0.29) \times 10^{-3}$	
$\Gamma_{88}$	$D^+ \rightarrow \eta'(958) \pi^+ \pi^0$	$(1.6 \pm 0.5) \times 10^{-3}$	

### Hadronic modes with a $K\bar{K}$ pair

$\Gamma_{89}$	$D^+ \rightarrow K^+ K_S^0$	$(2.83 \pm 0.16) \times 10^{-3}$	S=2.2
$\Gamma_{90}$	$D^+ \rightarrow K^+ K^- \pi^+$	[a] $(9.54 \pm 0.26) \times 10^{-3}$	S=1.1
$\Gamma_{91}$	$D^+ \rightarrow \phi \pi^+, \phi \rightarrow K^+ K^-$	$(2.65^{+0.08}_{-0.09}) \times 10^{-3}$	
$\Gamma_{92}$	$D^+ \rightarrow K^+ \bar{K}^*(892)^0,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(2.45^{+0.09}_{-0.14}) \times 10^{-3}$	
$\Gamma_{93}$	$D^+ \rightarrow K^+ \bar{K}_0^*(1430)^0,$ $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	$(1.79 \pm 0.34) \times 10^{-3}$	
$\Gamma_{94}$	$D^+ \rightarrow K^+ \bar{K}_2^*(1430)^0,$ $\bar{K}_2^* \rightarrow K^- \pi^+$	$(1.6^{+1.2}_{-0.8}) \times 10^{-4}$	
$\Gamma_{95}$	$D^+ \rightarrow K^+ \bar{K}_0^*(800), \bar{K}_0^* \rightarrow$ $K^- \pi^+$	$(6.7^{+3.4}_{-2.1}) \times 10^{-4}$	
$\Gamma_{96}$	$D^+ \rightarrow a_0(1450)^0 \pi^+, a_0^0 \rightarrow$ $K^+ K^-$	$(4.4^{+7.0}_{-1.8}) \times 10^{-4}$	
$\Gamma_{97}$	$D^+ \rightarrow \phi(1680) \pi^+, \phi \rightarrow$ $K^+ K^-$	$(4.9^{+4.0}_{-1.9}) \times 10^{-5}$	
$\Gamma_{98}$	$D^+ \rightarrow$ $K^+ K^- \pi^+$ nonresonant	not seen	
$\Gamma_{99}$	$D^+ \rightarrow K^+ K_S^0 \pi^+ \pi^-$	$(1.75 \pm 0.18) \times 10^{-3}$	
$\Gamma_{100}$	$D^+ \rightarrow K_S^0 K^- 2\pi^+$	$(2.40 \pm 0.18) \times 10^{-3}$	
$\Gamma_{101}$	$D^+ \rightarrow K^+ K^- 2\pi^+ \pi^-$	$(2.2 \pm 1.2) \times 10^{-4}$	

A few poorly measured branching fractions:

$\Gamma_{102}$	$D^+ \rightarrow \phi \pi^+ \pi^0$	$(2.3 \pm 1.0) \%$	
$\Gamma_{103}$	$D^+ \rightarrow \phi \rho^+$	$< 1.5 \%$	CL=90%
$\Gamma_{104}$	$D^+ \rightarrow K^+ K^- \pi^+ \pi^0$ non- $\phi$	$(1.5^{+0.7}_{-0.6}) \%$	
$\Gamma_{105}$	$D^+ \rightarrow K^*(892)^+ K_S^0$	$(1.6 \pm 0.7) \%$	

### Doubly Cabibbo-suppressed modes

$\Gamma_{106}$	$D^+ \rightarrow K^+ \pi^0$	$(1.83 \pm 0.26) \times 10^{-4}$	S=1.4
$\Gamma_{107}$	$D^+ \rightarrow K^+ \eta$	$(1.08 \pm 0.17) \times 10^{-4}$	
$\Gamma_{108}$	$D^+ \rightarrow K^+ \eta'(958)$	$(1.76 \pm 0.22) \times 10^{-4}$	
$\Gamma_{109}$	$D^+ \rightarrow K^+ \pi^+ \pi^-$	$(5.27 \pm 0.23) \times 10^{-4}$	
$\Gamma_{110}$	$D^+ \rightarrow K^+ \rho^0$	$(2.0 \pm 0.5) \times 10^{-4}$	

$\Gamma_{111}$	$D^+ \rightarrow K^*(892)^0 \pi^+,$ $K^*(892)^0 \rightarrow K^+ \pi^-$	$( 2.5 \pm 0.4 ) \times 10^{-4}$
$\Gamma_{112}$	$D^+ \rightarrow K^+ f_0(980),$ $f_0(980) \rightarrow \pi^+ \pi^-$	$( 4.7 \pm 2.8 ) \times 10^{-5}$
$\Gamma_{113}$	$D^+ \rightarrow K_2^*(1430)^0 \pi^+,$ $K_2^*(1430)^0 \rightarrow K^+ \pi^-$	$( 4.2 \pm 2.9 ) \times 10^{-5}$
$\Gamma_{114}$	$D^+ \rightarrow K^+ \pi^+ \pi^-$ nonreso-	not seen
	nant	
$\Gamma_{115}$	$D^+ \rightarrow 2K^+ K^-$	$( 8.7 \pm 2.0 ) \times 10^{-5}$

**$\Delta C = 1$  weak neutral current (C1) modes, or  
Lepton Family number (LF) or Lepton number (L) violating modes**

$\Gamma_{116}$	$D^+ \rightarrow \pi^+ e^+ e^-$	C1	$< 1.1$	$\times 10^{-6}$	CL=90%
$\Gamma_{117}$	$D^+ \rightarrow \pi^+ \phi, \phi \rightarrow e^+ e^-$	[e]	$( 1.7^{+1.4}_{-0.9} )$	$\times 10^{-6}$	
$\Gamma_{118}$	$D^+ \rightarrow \pi^+ \mu^+ \mu^-$	C1	$< 7.3$	$\times 10^{-8}$	CL=90%
$\Gamma_{119}$	$D^+ \rightarrow \pi^+ \phi, \phi \rightarrow$ $\mu^+ \mu^-$	[e]	$( 1.8 \pm 0.8 )$	$\times 10^{-6}$	
$\Gamma_{120}$	$D^+ \rightarrow \rho^+ \mu^+ \mu^-$	C1	$< 5.6$	$\times 10^{-4}$	CL=90%
$\Gamma_{121}$	$D^+ \rightarrow K^+ e^+ e^-$	[f]	$< 1.0$	$\times 10^{-6}$	CL=90%
$\Gamma_{122}$	$D^+ \rightarrow K^+ \mu^+ \mu^-$	[f]	$< 4.3$	$\times 10^{-6}$	CL=90%
$\Gamma_{123}$	$D^+ \rightarrow \pi^+ e^+ \mu^-$	LF	$< 2.9$	$\times 10^{-6}$	CL=90%
$\Gamma_{124}$	$D^+ \rightarrow \pi^+ e^- \mu^+$	LF	$< 3.6$	$\times 10^{-6}$	CL=90%
$\Gamma_{125}$	$D^+ \rightarrow K^+ e^+ \mu^-$	LF	$< 1.2$	$\times 10^{-6}$	CL=90%
$\Gamma_{126}$	$D^+ \rightarrow K^+ e^- \mu^+$	LF	$< 2.8$	$\times 10^{-6}$	CL=90%
$\Gamma_{127}$	$D^+ \rightarrow \pi^- 2e^+$	L	$< 1.1$	$\times 10^{-6}$	CL=90%
$\Gamma_{128}$	$D^+ \rightarrow \pi^- 2\mu^+$	L	$< 2.2$	$\times 10^{-8}$	CL=90%
$\Gamma_{129}$	$D^+ \rightarrow \pi^- e^+ \mu^+$	L	$< 2.0$	$\times 10^{-6}$	CL=90%
$\Gamma_{130}$	$D^+ \rightarrow \rho^- 2\mu^+$	L	$< 5.6$	$\times 10^{-4}$	CL=90%
$\Gamma_{131}$	$D^+ \rightarrow K^- 2e^+$	L	$< 9$	$\times 10^{-7}$	CL=90%
$\Gamma_{132}$	$D^+ \rightarrow K^- 2\mu^+$	L	$< 1.0$	$\times 10^{-5}$	CL=90%
$\Gamma_{133}$	$D^+ \rightarrow K^- e^+ \mu^+$	L	$< 1.9$	$\times 10^{-6}$	CL=90%
$\Gamma_{134}$	$D^+ \rightarrow K^*(892)^- 2\mu^+$	L	$< 8.5$	$\times 10^{-4}$	CL=90%
$\Gamma_{135}$	Unaccounted decay modes		$( 51.2 \pm 1.0 )$	%	

[a] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.

[b] These subfractions of the  $K^- 2\pi^+$  mode are uncertain: see the Particle Listings.

[c] Submodes of the  $D^+ \rightarrow K^- 2\pi^+ \pi^0$  and  $K_S^0 2\pi^+ \pi^-$  modes were studied by ANJOS 92C and COFFMAN 92B, but with at most 142 events for the first mode and 229 for the second – not enough for precise results. With

nothing new for 18 years, we refer to our 2008 edition, *Physics Letters B* **667** 1 (2008), for those results.

- [d] The unseen decay modes of the resonances are included.
- [e] This is *not* a test for the  $\Delta C=1$  weak neutral current, but leads to the  $\pi^+ \ell^+ \ell^-$  final state.
- [f] This mode is not a useful test for a  $\Delta C=1$  weak neutral current because both quarks must change flavor in this decay.

### CONSTRAINED FIT INFORMATION

An overall fit to 22 branching ratios uses 31 measurements and one constraint to determine 15 parameters. The overall fit has a  $\chi^2 = 32.0$  for 17 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_{29}$	0										
$x_{34}$	0	0									
$x_{35}$	22	0	0								
$x_{38}$	6	0	0	1							
$x_{40}$	15	0	0	3	44						
$x_{50}$	5	0	0	1	14	31					
$x_{54}$	6	0	0	1	18	40	56				
$x_{55}$	7	0	0	2	22	50	50	0			
$x_{56}$	3	0	0	1	10	24	7	10	12		
$x_{83}$	3	0	0	1	10	22	7	9	11	76	
$x_{89}$	6	0	0	1	75	38	12	15	19	9	
$x_{90}$	10	0	0	2	29	66	24	38	36	16	
$x_{106}$	2	0	0	0	6	13	4	5	6	3	
$x_{135}$	-75	-2	-15	-32	-32	-58	-54	-48	-42	-20	
	$x_{16}$	$x_{29}$	$x_{34}$	$x_{35}$	$x_{38}$	$x_{40}$	$x_{50}$	$x_{54}$	$x_{55}$	$x_{56}$	
$x_{89}$	8										
$x_{90}$	14	25									
$x_{106}$	3	5	9								
$x_{135}$	-18	-27	-43	-8							
	$x_{83}$	$x_{89}$	$x_{90}$	$x_{106}$							

## $D^+$ BRANCHING RATIOS

Some now-obsolete measurements have been omitted from these Listings.

### ———— c-quark decays ————

#### $\Gamma(c \rightarrow e^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$

For the Summary Table, we only use the average of  $e^+$  and  $\mu^+$  measurements from  $Z^0 \rightarrow c\bar{c}$  decays; see the second data block below.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.103 \pm 0.009</math></b> $_{-0.008}^{+0.009}$	378	<sup>1</sup> ABBIENDI	99K OPAL	$Z^0 \rightarrow c\bar{c}$

<sup>1</sup> ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed  $D^*(2010)^+ \rightarrow D^0\pi^+$  decays in  $Z^0 \rightarrow c\bar{c}$ .

#### $\Gamma(c \rightarrow \mu^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$

For the Summary Table, we only use the average of  $e^+$  and  $\mu^+$  measurements from  $Z^0 \rightarrow c\bar{c}$  decays; see the next data block.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.082 \pm 0.005</math></b> <b>OUR AVERAGE</b>				
$0.073 \pm 0.008 \pm 0.002$	73	KAYIS-TOPAK.05	CHRS	$\nu_\mu$ emulsion
$0.095 \pm 0.007$ $_{-0.013}^{+0.014}$	2829	ASTIER	00D NOMD	$\nu_\mu \text{ Fe} \rightarrow \mu^- \mu^+ X$
$0.090 \pm 0.007$ $_{-0.006}^{+0.007}$	476	<sup>1</sup> ABBIENDI	99K OPAL	$Z^0 \rightarrow c\bar{c}$
$0.086 \pm 0.017$ $_{-0.007}^{+0.008}$	69	<sup>2</sup> ALBRECHT	92F ARG	$e^+e^- \approx 10$ GeV
$0.078 \pm 0.009 \pm 0.012$		ONG	88 MRK2	$e^+e^- 29$ GeV
$0.078 \pm 0.015 \pm 0.02$		BARTEL	87 JADE	$e^+e^- 34.6$ GeV
$0.082 \pm 0.012$ $_{-0.01}^{+0.02}$		ALTHOFF	84G TASS	$e^+e^- 34.5$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.093 \pm 0.009 \pm 0.009$	88	KAYIS-TOPAK.02	CHRS	See KAYIS-TOPAKSU 05
$0.089 \pm 0.018 \pm 0.025$		BARTEL	85J JADE	See BARTEL 87

<sup>1</sup> ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed  $D^*(2010)^+ \rightarrow D^0\pi^+$  decays in  $Z^0 \rightarrow c\bar{c}$ .

<sup>2</sup> ALBRECHT 92F uses the excess of right-sign over wrong-sign leptons in a sample of events tagged by fully reconstructed  $D^*(2010)^+ \rightarrow D^0\pi^+$  decays.

#### $\Gamma(c \rightarrow \ell^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$

This is an average (not a sum) of  $e^+$  and  $\mu^+$  measurements.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.096 \pm 0.004</math></b> <b>OUR AVERAGE</b>				
$0.0958 \pm 0.0042 \pm 0.0028$	1828	<sup>1</sup> ABREU	000 DLPH	$Z^0 \rightarrow c\bar{c}$
$0.095 \pm 0.006$ $_{-0.006}^{+0.007}$	854	<sup>2</sup> ABBIENDI	99K OPAL	$Z^0 \rightarrow c\bar{c}$

<sup>1</sup> ABREU 000 uses leptons opposite fully reconstructed  $D^*(2010)^+$ ,  $D^+$ , or  $D^0$  mesons.

<sup>2</sup> ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed  $D^*(2010)^+ \rightarrow D^0\pi^+$  decays in  $Z^0 \rightarrow c\bar{c}$ .

### $\Gamma(c \rightarrow D^*(2010)^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.255 ± 0.015 ± 0.008</b>	2371	<sup>1</sup> ABREU	000	DLPH $Z^0 \rightarrow c\bar{c}$

<sup>1</sup> ABREU 000 uses slow pions opposite fully reconstructed  $D^*(2010)^+$ ,  $D^+$ , or  $D^0$  mesons as a signal of  $D^*(2010)^-$  production.

### ———— Inclusive modes ————

### $\Gamma(e^+ \text{ semileptonic}) / \Gamma_{\text{total}}$

$\Gamma_1 / \Gamma$

The sum of our  $\bar{K}^0 e^+ \nu_e$ ,  $\bar{K}^*(892)^0 e^+ \nu_e$ ,  $\pi^0 e^+ \nu_e$ ,  $\eta e^+ \nu_e$ ,  $\rho^0 e^+ \nu_e$ , and  $\omega e^+ \nu_e$  branching fractions is  $15.3 \pm 0.4\%$ .

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

#### **16.07 ± 0.30 OUR AVERAGE**

16.13 ± 0.10 ± 0.29	26.2 ± 0.2k	<sup>1</sup> ASNER	10	CLEO $e^+ e^-$ at 3774 MeV
15.2 ± 0.9 ± 0.8	521 ± 32	ABLIKIM	07G	BES2 $e^+ e^- \approx \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

16.13 ± 0.20 ± 0.33	8798 ± 105	<sup>2</sup> ADAM	06A	CLEO See ASNER 10
17.0 ± 1.9 ± 0.7	158	BALTRUSAIT ..85B	MRK3	$e^+ e^-$ 3.77 GeV

<sup>1</sup> Using the  $D^+$  and  $D^0$  lifetimes, ASNER 10 finds that the ratio of the  $D^+$  and  $D^0$  semileptonic widths is  $0.985 \pm 0.015 \pm 0.024$ .

<sup>2</sup> Using the  $D^+$  and  $D^0$  lifetimes, ADAM 06A finds that the ratio of the  $D^+$  and  $D^0$  inclusive  $e^+$  widths is  $0.985 \pm 0.028 \pm 0.015$ , consistent with the isospin-invariance prediction of 1.

### $\Gamma(\mu^+ \text{ anything}) / \Gamma_{\text{total}}$

$\Gamma_2 / \Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

<b>17.6 ± 2.7 ± 1.8</b>	100 ± 12	<sup>1</sup> ABLIKIM	08L	BES2 $e^+ e^- \approx \psi(3772)$
-------------------------	----------	----------------------	-----	-----------------------------------

<sup>1</sup> ABLIKIM 08L finds the ratio of  $D^+ \rightarrow \mu^+ X$  and  $D^0 \rightarrow \mu^+ X$  branching fractions to be  $2.59 \pm 0.70 \pm 0.25$ , in accord with the ratio of  $D^+$  and  $D^0$  lifetimes,  $2.54 \pm 0.02$ .

### $\Gamma(K^- \text{ anything}) / \Gamma_{\text{total}}$

$\Gamma_3 / \Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

#### **25.7 ± 1.4 OUR AVERAGE**

24.7 ± 1.3 ± 1.2	631 ± 33	ABLIKIM	07G	BES2 $e^+ e^- \approx \psi(3770)$
27.8 <sup>+3.6</sup> <sub>-3.1</sub>		BARLAG	92C	ACCM $\pi^-$ Cu 230 GeV
27.1 ± 2.3 ± 2.4		COFFMAN	91	MRK3 $e^+ e^-$ 3.77 GeV

### $[\Gamma(\bar{K}^0 \text{ anything}) + \Gamma(K^0 \text{ anything})] / \Gamma_{\text{total}}$

$\Gamma_4 / \Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

#### **61 ± 5 OUR AVERAGE**

60.5 ± 5.5 ± 3.3	244 ± 22	ABLIKIM	06U	BES2 $e^+ e^-$ at 3773 MeV
61.2 ± 6.5 ± 4.3		COFFMAN	91	MRK3 $e^+ e^-$ 3.77 GeV

### $\Gamma(K^+ \text{ anything}) / \Gamma_{\text{total}}$

$\Gamma_5 / \Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

#### **5.9 ± 0.8 OUR AVERAGE**

6.1 ± 0.9 ± 0.4	189 ± 27	ABLIKIM	07G	BES2 $e^+ e^- \approx \psi(3770)$
5.5 ± 1.3 ± 0.9		COFFMAN	91	MRK3 $e^+ e^-$ 3.77 GeV

$\Gamma(K^*(892)^- \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$5.7 \pm 5.2 \pm 0.7$	$7.2 \pm 6.5$	ABLIKIM	06U BES2	$e^+ e^-$ at 3773 MeV

$\Gamma(\bar{K}^*(892)^0 \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$23.2 \pm 4.5 \pm 3.0$	$189 \pm 36$	ABLIKIM	05P BES	$e^+ e^- \approx 3773$ MeV

$\Gamma(K^*(892)^0 \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.6$	90	ABLIKIM	05P BES	$e^+ e^- \approx 3773$ MeV

$\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$

This ratio includes  $\eta$  particles from  $\eta'$  decays.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$6.3 \pm 0.5 \pm 0.5$	$1972 \pm 142$	HUANG	06B CLEO	$e^+ e^-$ at $\psi(3770)$

$\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.04 \pm 0.16 \pm 0.09$	$82 \pm 13$	HUANG	06B CLEO	$e^+ e^-$ at $\psi(3770)$

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.03 \pm 0.10 \pm 0.07$	$248 \pm 21$	HUANG	06B CLEO	$e^+ e^-$ at $\psi(3770)$

————— Leptonic and semileptonic modes —————

$\Gamma(e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8.8 \times 10^{-6}$	90	EISENSTEIN	08 CLEO	$e^+ e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.4 \times 10^{-5}$	90	ARTUSO	05A CLEO	See EISENSTEIN 08
------------------------	----	--------	----------	-------------------

$\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$

See the note on "Decay Constants of Charged Pseudoscalar Mesons" in the  $D_s^+$  Listings.

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.82 \pm 0.32 \pm 0.09$	$150 \pm 12$	<sup>1</sup> EISENSTEIN	08 CLEO	$e^+ e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$12.2 \begin{smallmatrix} +11.1 \\ -5.3 \end{smallmatrix} \pm 1.0$	3	<sup>2</sup> ABLIKIM	05D BES	$e^+ e^- \approx 3.773$ GeV
$4.40 \pm 0.66 \begin{smallmatrix} +0.09 \\ -0.12 \end{smallmatrix}$	$47 \pm 7$	<sup>3</sup> ARTUSO	05A CLEO	See EISENSTEIN 08
$3.5 \pm 1.4 \pm 0.6$	7	<sup>4</sup> BONVICINI	04A CLEO	Incl. in ARTUSO 05A
$8 \begin{smallmatrix} +16 \\ -5 \end{smallmatrix} \begin{smallmatrix} +5 \\ -2 \end{smallmatrix}$	1	<sup>5</sup> BAI	98B BES	$e^+ e^- \rightarrow D^{*+} D^-$

- <sup>1</sup> EISENSTEIN 08, using the  $D^+$  lifetime and assuming  $|V_{cd}| = |V_{us}|$ , gets  $f_{D^+} = (205.8 \pm 8.5 \pm 2.5)$  MeV from this measurement.
- <sup>2</sup> ABLIKIM 05D finds a background-subtracted  $2.67 \pm 1.74$   $D^+ \rightarrow \mu^+ \nu_\mu$  events, and from this obtains  $f_{D^+} = 371^{+129}_{-119} \pm 25$  MeV.
- <sup>3</sup> ARTUSO 05A obtains  $f_{D^+} = 222.6 \pm 16.7^{+2.8}_{-3.4}$  MeV from this measurement.
- <sup>4</sup> BONVICINI 04A finds eight events with an estimated background of one, and from the branching fraction obtains  $f_{D^+} = 202 \pm 41 \pm 17$  MeV.
- <sup>5</sup> BAI 98B obtains  $f_{D^+} = (300^{+180+80}_{-150-40})$  MeV from this measurement.

$\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$					$\Gamma_{14}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b><math>&lt;1.2 \times 10^{-3}</math></b>	90	EISENSTEIN 08	CLEO	$e^+ e^-$ at $\psi(3770)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.1 \times 10^{-3}$	90	RUBIN 06A	CLEO	See EISENSTEIN 08	

$\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma_{\text{total}}$					$\Gamma_{15}/\Gamma$
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>8.83±0.22 OUR AVERAGE</b>					
$8.83 \pm 0.10 \pm 0.20$	8467	<sup>1</sup> BESSON 09	CLEO	$e^+ e^-$ at $\psi(3770)$	
$8.95 \pm 1.59 \pm 0.67$	$34 \pm 6$	<sup>2</sup> ABLIKIM 05A	BES	$e^+ e^-$ at $\psi(3770)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$8.53 \pm 0.13 \pm 0.23$		<sup>3</sup> DOBBS 08	CLEO	See BESSON 09	
$8.71 \pm 0.38 \pm 0.37$	$545 \pm 24$	HUANG 05B	CLEO	See DOBBS 08	

- <sup>1</sup> See the form-factor parameters near the end of this  $D^+$  Listing.
- <sup>2</sup> The ABLIKIM 05A result together with the  $D^0 \rightarrow K^- e^+ \nu_e$  branching fraction of ABLIKIM 04C and Particle Data Group lifetimes gives  $\Gamma(D^0 \rightarrow K^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^0 e^+ \nu_e) = 1.08 \pm 0.22 \pm 0.07$ ; isospin invariance predicts the ratio is 1.0.
- <sup>3</sup> DOBBS 08 establishes  $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}| = 0.188 \pm 0.008 \pm 0.002$  from the  $D^+$  and  $D^0$  decays to  $\bar{K} e^+ \nu_e$  and  $\pi e^+ \nu_e$ . It also finds  $\Gamma(D^0 \rightarrow K^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^0 e^+ \nu_e) = 1.06 \pm 0.02 \pm 0.03$ ; isospin invariance predicts the ratio is 1.0.

$\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$					$\Gamma_{16}/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.092±0.006 OUR FIT</b>					
<b>0.103±0.023±0.008</b>	$29 \pm 6$	ABLIKIM 07	BES2	$e^+ e^-$ at 3773 MeV	

$\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma(K^- 2\pi^+)$					$\Gamma_{16}/\Gamma_{40}$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.00 ±0.07 OUR FIT</b>					
<b>1.019±0.076±0.065</b>	$555 \pm 39$	LINK 04E	FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV	

$\Gamma(K^- \pi^+ e^+ \nu_e)/\Gamma_{\text{total}}$					$\Gamma_{17}/\Gamma$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$3.50 \pm 0.75 \pm 0.27$	$29 \pm 6$	ABLIKIM 060	BES2	$e^+ e^-$ at 3773 MeV	
$3.5^{+1.2}_{-0.7} \pm 0.4$	14	BAI 91	MRK3	$e^+ e^- \approx 3.77$ GeV	

$\Gamma(K^- \pi^+ e^+ \nu_e) / \Gamma(K^- 2\pi^+)$   $\Gamma_{17} / \Gamma_{40}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.4380 ± 0.0036 ± 0.0042</b>	70k ± 363	DEL-AMO-SA..11I	BABR	$e^+ e^- \approx 10.6$ GeV

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{34} / \Gamma$

Unseen decay modes of  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**5.52 ± 0.15 OUR FIT**

**5.52 ± 0.07 ± 0.13**  $\approx 5k$  BRIERE 10 CLEO  $e^+ e^-$  at  $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.06 ± 1.21 ± 0.40 28 ± 7 ABLIKIM 06O BES2  $e^+ e^-$  at 3773 MeV

5.56 ± 0.27 ± 0.23 422 ± 21 <sup>1</sup>HUANG 05B CLEO  $e^+ e^-$  at  $\psi(3770)$

<sup>1</sup>HUANG 05B finds  $\Gamma(D^0 \rightarrow K^{*-} e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e) = 0.98 \pm 0.08 \pm 0.04$ ; isospin invariance predicts the ratio is 1.0.

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e) / \Gamma(K^- 2\pi^+)$   $\Gamma_{34} / \Gamma_{40}$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.74 ± 0.04 ± 0.05 BRANDENB... 02 CLEO  $e^+ e^- \approx \Upsilon(4S)$

0.62 ± 0.15 ± 0.09 35 ADAMOVICH 91 OMEG  $\pi^-$  340 GeV

0.55 ± 0.08 ± 0.10 880 ALBRECHT 91 ARG  $e^+ e^- \approx 10.4$  GeV

0.49 ± 0.04 ± 0.05 ANJOS 89B E691 Photoproduction

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow K^- \pi^+) / \Gamma(K^- \pi^+ e^+ \nu_e)$   $\Gamma_{18} / \Gamma_{17}$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

**94.11 ± 0.74 ± 0.75** DEL-AMO-SA..11I BABR  $e^+ e^- \approx 10.6$  GeV

$\Gamma((K^- \pi^+)_{S\text{-wave}} e^+ \nu_e) / \Gamma(K^- \pi^+ e^+ \nu_e)$   $\Gamma_{19} / \Gamma_{17}$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

**5.79 ± 0.16 ± 0.15** DEL-AMO-SA..11I BABR  $e^+ e^- \approx 10.6$  GeV

$\Gamma(\bar{K}^*(1410)^0 e^+ \nu_e, \bar{K}^*(1410)^0 \rightarrow K^- \pi^+) / \Gamma_{\text{total}}$   $\Gamma_{20} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**< 6 × 10<sup>-3</sup>** 90 DEL-AMO-SA..11I BABR  $e^+ e^- \approx 10.6$  GeV

$\Gamma(\bar{K}_2^*(1430)^0 e^+ \nu_e, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+) / \Gamma_{\text{total}}$   $\Gamma_{21} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**< 5 × 10<sup>-4</sup>** 90 DEL-AMO-SA..11I BABR  $e^+ e^- \approx 10.6$  GeV

$\Gamma(K^- \pi^+ e^+ \nu_e \text{ nonresonant}) / \Gamma_{\text{total}}$   $\Gamma_{22} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**< 0.007** 90 ANJOS 89B E691 Photoproduction

$\Gamma(K^- \pi^+ \mu^+ \nu_\mu) / \Gamma(\bar{K}^0 \mu^+ \nu_\mu)$   $\Gamma_{23} / \Gamma_{16}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.417 ± 0.030 ± 0.023</b>	555 ± 39	LINK	04E	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma_{\text{total}}$   $\Gamma_{35} / \Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.28 ± 0.15 OUR FIT</b>				
<b>5.27 ± 0.07 ± 0.14</b>	$\approx 5k$	BRIERE	10	CLEO $e^+ e^-$ at $\psi(3770)$

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma(\bar{K}^0 \mu^+ \nu_\mu)$   $\Gamma_{35} / \Gamma_{16}$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.58 ± 0.04 OUR FIT</b>				
<b>0.594 ± 0.043 ± 0.033</b>	555 ± 39	LINK	04E	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma(K^- 2\pi^+)$   $\Gamma_{35} / \Gamma_{40}$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.578 ± 0.021 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>0.57 ± 0.06 OUR AVERAGE</b>	Error includes scale factor of 1.2.			
0.72 ± 0.10 ± 0.05		BRANDENB...	02	CLEO $e^+ e^- \approx \gamma(4S)$
0.56 ± 0.04 ± 0.06	875	FRABETTI	93E	E687 $\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV
0.46 ± 0.07 ± 0.08	224	KODAMA	92C	E653 $\pi^-$ emulsion 600 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.602 ± 0.010 ± 0.021	12k	<sup>1</sup> LINK	02J	FOCS $\gamma$ nucleus, $\approx 180$ GeV

<sup>1</sup>This LINK 02J result includes the effects of an interference of a small  $S$ -wave  $K^- \pi^+$  amplitude with the dominant  $\bar{K}^{*0}$  amplitude. (The interference effect is reported in LINK 02E.) This result is redundant with results of LINK 04E elsewhere in these Listings.

$\Gamma(K^- \pi^+ \mu^+ \nu_\mu \text{ nonresonant}) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$   $\Gamma_{25} / \Gamma_{23}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0530 ± 0.0074<sup>+0.0099</sup><sub>-0.0096</sub></b>	14k	LINK	05I	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^- \pi^+ \pi^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$   $\Gamma_{26} / \Gamma_{23}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.042</b>	90	FRABETTI	93E	E687 $\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV

$\Gamma(\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$   $\Gamma_{36} / \Gamma_{23}$

Unseen decay modes of the  $\bar{K}_0^*(1430)^0$  are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0064</b>	90	LINK	05I	FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\bar{K}^*(1680)^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$   $\Gamma_{37} / \Gamma_{23}$

Unseen decay modes of the  $\bar{K}^*(1680)^0$  are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	LINK	05I	FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\pi^0 e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{27} / \Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.405 ± 0.016 ± 0.009</b>	838	<sup>1</sup> BESSON	09	CLEO $e^+ e^-$ at $\psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.373 ± 0.022 ± 0.013		<sup>2</sup> DOBBS	08	CLEO See BESSON 09
0.44 ± 0.06 ± 0.03	63 ± 9	HUANG	05B	CLEO See DOBBS 08

<sup>1</sup> See the form-factor parameters near the end of this  $D^+$  Listing.

<sup>2</sup> DOBBS 08 establishes  $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}| = 0.188 \pm 0.008 \pm 0.002$  from the  $D^+$  and  $D^0$  decays to  $\bar{K} e^+ \nu_e$  and  $\pi e^+ \nu_e$ . It finds  $\Gamma(D^0 \rightarrow \pi^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \pi^0 e^+ \nu_e) = 2.03 \pm 0.14 \pm 0.08$ ; isospin invariance predicts the ratio is 2.0.

$\Gamma(\eta e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{28} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.4 ± 0.9 ± 0.4</b>		YELTON	11	CLEO $e^+ e^-$ at $\psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
13.3 ± 2.0 ± 0.6	46 ± 8	MITCHELL	09B	CLEO See YELTON 11

$\Gamma(\rho^0 e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{29} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.18<sup>+0.17</sup><sub>-0.25</sub> OUR FIT</b>				

<b>2.17 ± 0.12<sup>+0.12</sup><sub>-0.22</sub></b>	447 ± 25	<sup>1</sup> DOBBS	13	CLEO $e^+ e^-$ at $\psi(3770)$
--	----------	--------------------	----	--------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.1 ± 0.4 ± 0.1	27 ± 6	<sup>2</sup> HUANG	05B	CLEO See DOBBS 13
-----------------	--------	--------------------	-----	-------------------

<sup>1</sup> DOBBS 13 finds  $\Gamma(D^0 \rightarrow \rho^- e^+ \nu_e) / 2 \Gamma(D^+ \rightarrow \rho^0 e^+ \nu_e) = 1.03 \pm 0.09^{+0.08}_{-0.02}$ ; isospin invariance predicts the ratio is 1.0.

<sup>2</sup> HUANG 05B finds  $\Gamma(D^0 \rightarrow \rho^- e^+ \nu_e) / 2 \Gamma(D^+ \rightarrow \rho^0 e^+ \nu_e) = 1.2^{+0.4}_{-0.3} \pm 0.1$ ; isospin invariance predicts the ratio is 1.0.

$\Gamma(\rho^0 e^+ \nu_e) / \Gamma(\bar{K}^*(892)^0 e^+ \nu_e)$   $\Gamma_{29} / \Gamma_{34}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0396<sup>+0.0033</sup><sub>-0.0050</sub> OUR FIT</b>				

<b>0.045 ± 0.014 ± 0.009</b>	49	<sup>1</sup> AITALA	97	E791 $\pi^-$ nucleus, 500 GeV
------------------------------	----	---------------------	----	-------------------------------

<sup>1</sup> AITALA 97 explicitly subtracts  $D^+ \rightarrow \eta' e^+ \nu_e$  and other backgrounds to get this result.

$\Gamma(\rho^0 \mu^+ \nu_\mu) / \Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$   $\Gamma_{30} / \Gamma_{35}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.045 ± 0.007 OUR AVERAGE</b>				Error includes scale factor of 1.1.
0.041 ± 0.006 ± 0.004	320 ± 44	LINK	06B FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
0.051 ± 0.015 ± 0.009	54	<sup>1</sup> AITALA	97 E791	$\pi^-$ nucleus, 500 GeV
0.079 ± 0.019 ± 0.013	39	<sup>2</sup> FRABETTI	97 E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV

<sup>1</sup> AITALA 97 explicitly subtracts  $D^+ \rightarrow \eta' \mu^+ \nu_\mu$  and other backgrounds to get this result.

<sup>2</sup> Because the reconstruction efficiency for photons is low, this FRABETTI 97 result also includes any  $D^+ \rightarrow \eta' \mu^+ \nu_\mu \rightarrow \gamma \rho^0 \mu^+ \nu_\mu$  events in the numerator.

$\Gamma(\omega e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{31} / \Gamma$

<u>VALUE (units 10<sup>-3</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.82 ± 0.18 ± 0.07</b>	129 ± 13	DOBBS	13 CLEO	$e^+ e^-$ at $\psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.6 <sup>+0.7</sup> <sub>-0.6</sub> ± 0.1	7.6 <sup>+3.3</sup> <sub>-2.7</sub>	HUANG	05B CLEO	See DOBBS 13

$\Gamma(\eta'(958) e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{32} / \Gamma$

<u>VALUE (units 10<sup>-4</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.16 ± 0.53 ± 0.07</b>		YELTON	11 CLEO	$e^+ e^-$ at $\psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<3.5	90	MITCHELL	09B CLEO	See YELTON 11

$\Gamma(\phi e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{33} / \Gamma$

Unseen decay modes of the  $\phi$  are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.9 × 10<sup>-4</sup></b>	90	YELTON	11 CLEO	$e^+ e^-$ at $\psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.6 × 10 <sup>-4</sup>	90	MITCHELL	09B CLEO	See YELTON 11
<0.0201	90	ABLIKIM	06P BES2	$e^+ e^-$ at 3773 MeV
<0.0209	90	BAI	91 MRK3	$e^+ e^- \approx 3.77$ GeV

———— Hadronic modes with a  $\bar{K}$  or  $\bar{K}K\bar{K}$  ————

$\Gamma(K_S^0 \pi^+) / \Gamma_{\text{total}}$   $\Gamma_{38} / \Gamma$

<u>VALUE (units 10<sup>-2</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.526 ± 0.022 ± 0.038		<sup>1</sup> DOBBS	07 CLEO	See MENDEZ 10
1.55 ± 0.05 ± 0.06	2230 ± 60	<sup>1</sup> HE	05 CLEO	See DOBBS 07
1.6 ± 0.3 ± 0.1	161	ADLER	88C MRK3	$e^+ e^-$ 3.77 GeV

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

$\Gamma(K_S^0 \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{38}/\Gamma_{40}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.161 ± 0.007 OUR FIT</b>				Error includes scale factor of 3.4.
<b>0.158 ± 0.007 OUR AVERAGE</b>				Error includes scale factor of 3.2.
0.1682 ± 0.0012 ± 0.0037	30k	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV
0.1530 ± 0.0023 ± 0.0016	10.6k	LINK	02B	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.174 ± 0.012 ± 0.011	473	<sup>1</sup> BISHAI	97	CLEO $e^+ e^- \approx \Upsilon(4S)$
0.137 ± 0.015 ± 0.016	264	ANJOS	90C	E691 Photoproduction

<sup>1</sup> See BISHAI 97 for an isospin analysis of  $D^+ \rightarrow \bar{K} \pi$  amplitudes.

$\Gamma(K_L^0 \pi^+)/\Gamma_{total}$   $\Gamma_{39}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.460 ± 0.040 ± 0.035</b>	2023 ± 54	<sup>1</sup> HE	08	CLEO $e^+ e^-$ at $\psi(3770)$

<sup>1</sup> The difference of CLEO  $D^+ \rightarrow K_S^0 \pi^+$  and  $K_L^0 \pi^+$  branching fractions over the sum (DOBBS 07 and HE 08) is  $+0.022 \pm 0.016 \pm 0.018$ .

$\Gamma(K^- 2\pi^+)/\Gamma_{total}$   $\Gamma_{40}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.13 ± 0.19 OUR FIT</b>				
<b>9.14 ± 0.10 ± 0.17</b>		<sup>1</sup> DOBBS	07	CLEO $e^+ e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.5 ± 0.2 ± 0.3	15.1k ± 130	<sup>1</sup> HE	05	CLEO See DOBBS 07
9.3 ± 0.6 ± 0.8	1502	<sup>2</sup> BALEST	94	CLEO $e^+ e^- \approx \Upsilon(4S)$
6.4 <sup>+1.5</sup> <sub>-1.4</sub>		<sup>3</sup> BARLAG	92C	ACCM $\pi^-$ Cu 230 GeV
9.1 ± 1.3 ± 0.4	1164	ADLER	88C	MRK3 $e^+ e^-$ 3.77 GeV
9.1 ± 1.9	239	<sup>4</sup> SCHINDLER	81	MRK2 $e^+ e^-$ 3.771 GeV

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

<sup>2</sup> BALEST 94 measures the ratio of  $D^+ \rightarrow K^- \pi^+ \pi^+$  and  $D^0 \rightarrow K^- \pi^+$  branching fractions to be  $2.35 \pm 0.16 \pm 0.16$  and uses their absolute measurement of the  $D^0 \rightarrow K^- \pi^+$  fraction (AKERIB 93).

<sup>3</sup> BARLAG 92C computes the branching fraction by topological normalization.

<sup>4</sup> SCHINDLER 81 (MARK-2) measures  $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$  branching fraction to be  $0.38 \pm 0.05$  nb. We use the MARK-3 (ADLER 88C) value of  $\sigma = 4.2 \pm 0.6 \pm 0.3$  nb.

$\Gamma((K^- \pi^+)_{S\text{-wave}} \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{41}/\Gamma_{40}$

This is the "fit fraction" from the Dalitz-plot analysis. The  $K^- \pi^+$  S-wave includes a broad scalar  $\kappa$  ( $\bar{K}_0^*(800)$ ), the  $\bar{K}_0^*(1430)^0$ , and non-resonant background.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.801 ± 0.012 OUR AVERAGE</b>			
0.8024 ± 0.0138 ± 0.0043	<sup>1</sup> LINK	09	FOCS MIPWA fit, 53k evts
0.838 ± 0.038	<sup>2</sup> BONVICINI	08A	CLEO QMIPWA fit, 141k evts
0.786 ± 0.014 ± 0.018	AITALA	06	E791 Dalitz fit, 15.1k events
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.8323 ± 0.0150 ± 0.0008	<sup>3</sup> LINK	07B	FOCS See LINK 09

<sup>1</sup> This LINK 09 model-independent partial-wave analysis of the  $K^- \pi^+$   $S$ -wave slices the  $K^- \pi^+$  mass range into 39 bins.

<sup>2</sup> The BONVICINI 08A QMIPWA (quasi-model-independent partial-wave analysis) of the  $K^- \pi^+$   $S$ -wave amplitude slices the  $K^- \pi^+$  mass range into 26 bins but keeps the Breit-Wigner  $\bar{K}_0^*(1430)^0$ .

<sup>3</sup> This LINK 07B fit uses a K matrix. The  $K^- \pi^+$   $S$ -wave fit fraction given above breaks down into  $(207.3 \pm 25.5 \pm 12.4)\%$  isospin-1/2 and  $(40.5 \pm 9.6 \pm 3.2)\%$  isospin-3/2 — with large interference between the two. The isospin-1/2 component includes the  $\kappa$  (or  $\bar{K}_0^*(800)^0$ ) and  $\bar{K}_0^*(1430)^0$ .

$$\Gamma(\bar{K}_0^*(800)^0 \pi^+, \bar{K}_0^*(800)^0 \rightarrow K^- \pi^+) / \Gamma(K^- 2\pi^+) \quad \Gamma_{42} / \Gamma_{40}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.478 \pm 0.121 \pm 0.053$	AITALA	02	E791 See AITALA 06
-----------------------------	--------	----	--------------------

$$\Gamma(\bar{K}^*(892)^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+) / \Gamma(K^- 2\pi^+) \quad \Gamma_{44} / \Gamma_{40}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.111 ± 0.012 OUR AVERAGE** Error includes scale factor of 3.7.

$0.1236 \pm 0.0034 \pm 0.0034$	LINK	09	FOCS MIPWA fit, 53k evts
$0.0988 \pm 0.0046$	BONVICINI	08A	CLEO QMIPWA fit, 141k evts
$0.119 \pm 0.002 \pm 0.020$	AITALA	06	E791 Dalitz fit, 15.1k events

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.1361 \pm 0.0041 \pm 0.0030$	<sup>1</sup> LINK	07B	FOCS See LINK 09
$0.123 \pm 0.010 \pm 0.009$	AITALA	02	E791 See AITALA 06
$0.137 \pm 0.006 \pm 0.009$	FRABETTI	94G	E687 Dalitz fit, 8800 evts
$0.170 \pm 0.009 \pm 0.034$	ANJOS	93	E691 $\gamma$ Be 90–260 GeV
$0.14 \pm 0.04 \pm 0.04$	ALVAREZ	91B	NA14 Photoproduction
$0.13 \pm 0.01 \pm 0.07$	ADLER	87	MRK3 $e^+ e^-$ 3.77 GeV

<sup>1</sup> The statistical error on this LINK 07B value is corrected in LINK 09.

$$\Gamma(\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow K^- \pi^+) / \Gamma(K^- 2\pi^+) \quad \Gamma_{45} / \Gamma_{40}$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

<b>not seen</b>	LINK	09	FOCS MIPWA fit, 53k evts
<b>not seen</b>	BONVICINI	08A	CLEO QMIPWA fit, 141k evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.8 \pm 2.1 \pm 1.7$	LINK	07B	FOCS See LINK 09
-----------------------	------	-----	------------------

$$\Gamma(\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+) / \Gamma(K^- 2\pi^+) \quad \Gamma_{43} / \Gamma_{40}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.1330 ± 0.0062** BONVICINI 08A CLEO QMIPWA fit, 141k evts

• • • We do not use the following data for averages, fits, limits, etc. • • •


$0.125 \pm 0.014 \pm 0.005$	AITALA	02	E791 See AITALA 06
$0.284 \pm 0.022 \pm 0.059$	FRABETTI	94G	E687 Dalitz fit, 8800 evts
$0.248 \pm 0.019 \pm 0.017$	ANJOS	93	E691 $\gamma$ Be 90–260 GeV

$\Gamma(\bar{K}_2^*(1430)^0 \pi^+, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{46}/\Gamma_{40}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.24 ± 0.08 OUR AVERAGE</b>	Error includes scale factor of 2.2. See the ideogram below.		
0.58 ± 0.10 ± 0.06	LINK	09	FOCS MIPWA fit, 53k evts
0.204 ± 0.040	BONVICINI	08A	CLEO QMIPWA fit, 141k evts
0.2 ± 0.1 ± 0.1	AITALA	06	E791 Dalitz fit, 15.1k events
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.39 ± 0.09 ± 0.05	LINK	07B	FOCS See LINK 09
0.5 ± 0.1 ± 0.2	AITALA	02	E791 See AITALA 06

WEIGHTED AVERAGE  
0.24±0.08 (Error scaled by 2.2)



$\Gamma(\bar{K}_2^*(1430)^0 \pi^+, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{46}/\Gamma_{40}$   
(units  $10^{-2}$ )

$\Gamma(\bar{K}^*(1680)^0 \pi^+, \bar{K}^*(1680)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{47}/\Gamma_{40}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.23 ± 0.12 OUR AVERAGE</b>			
1.75 ± 0.62 ± 0.54	LINK	09	FOCS MIPWA fit, 53k evts
0.196 ± 0.118	BONVICINI	08A	CLEO QMIPWA fit, 141k evts
1.2 ± 0.6 ± 1.2	AITALA	06	E791 Dalitz fit, 15.1k events
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.90 ± 0.63 ± 0.43	LINK	07B	FOCS See LINK 09
2.5 ± 0.7 ± 0.3	AITALA	02	E791 See AITALA 06
4.7 ± 0.6 ± 0.7	FRABETTI	94G	E687 Dalitz fit, 8800 evts
3.0 ± 0.4 ± 1.3	ANJOS	93	E691 $\gamma$ Be 90–260 GeV

$\Gamma(K^-(2\pi^+)_{I=2})/\Gamma(K^-2\pi^+)$   $\Gamma_{48}/\Gamma_{40}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.155±0.028</b>	BONVICINI 08A	CLEO	QMIPWA fit, 141k evts

$\Gamma(K^-2\pi^+ \text{ nonresonant})/\Gamma(K^-2\pi^+)$   $\Gamma_{49}/\Gamma_{40}$

This is the "fit fraction" from the Dalitz-plot analysis. Later analyses find little need for this decay mode.

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.130±0.058±0.044	AITALA 02	E791	See AITALA 06
0.998±0.037±0.072	FRABETTI 94G	E687	Dalitz fit, 8800 evts
0.838±0.088±0.275	ANJOS 93	E691	$\gamma$ Be 90–260 GeV
0.79 ±0.07 ±0.15	ADLER 87	MRK3	$e^+e^-$ 3.77 GeV

$\Gamma(K_S^0\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.99±0.27 OUR FIT</b>				
<b>6.99±0.09±0.25</b>		<sup>1</sup> DOBBS	07	CLEO $e^+e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.2 ±0.2 ±0.4	5090 ±100	<sup>1</sup> HE	05	CLEO See DOBBS 07
5.1 ±1.3 ±0.8	159	ADLER	88C	MRK3 $e^+e^-$ 3.77 GeV
<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.				

$\Gamma(K_S^0\rho^+)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{51}/\Gamma_{50}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.68±0.08±0.12</b>	ADLER 87	MRK3	$e^+e^-$ 3.77 GeV

$\Gamma(\bar{K}^*(892)^0\pi^+, \bar{K}^*(892)^0 \rightarrow K_S^0\pi^0)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{52}/\Gamma_{50}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.19±0.06±0.06</b>	ADLER 87	MRK3	$e^+e^-$ 3.77 GeV

$\Gamma(K_S^0\pi^+\pi^0 \text{ nonresonant})/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{53}/\Gamma_{50}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.13±0.07±0.08</b>	ADLER 87	MRK3	$e^+e^-$ 3.77 GeV

$\Gamma(K^-2\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{54}/\Gamma$

See our 2008 Review (Physics Letters **B667** 1 (2008)) for measurements of submodes of this mode. There is nothing new since 1992, and the two papers, ANJOS 92C, with  $91 \pm 12$  events above background, and COFFMAN 92B, with  $142 \pm 20$  such events, could not determine submode fractions with much accuracy.

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.99±0.18 OUR FIT</b>				
<b>5.98±0.08±0.16</b>		<sup>1</sup> DOBBS	07	CLEO $e^+e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.0 ±0.2 ±0.2	4840 ±100	<sup>1</sup> HE	05	CLEO	See DOBBS 07
5.8 ±1.2 ±1.2	142	COFFMAN	92B	MRK3	e <sup>+</sup> e <sup>-</sup> 3.77 GeV
6.3 <sup>+1.4</sup> / <sub>-1.3</sub> ±1.2	175	BALTRUSAIT..86E	MRK3	See COFFMAN	92B

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

$\Gamma(K_S^0 2\pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{55} / \Gamma$

See our 2008 Review (Physics Letters **B667** 1 (2008)) for measurements of submodes of this mode. There is nothing new since 1992, and the two papers, ANJOS 92C, with 229 ± 17 events above background, and COFFMAN 92B, with 209 ± 20 such events, could not determine submode fractions with much accuracy.

VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

**3.12 ±0.11 OUR FIT**

**3.122 ±0.046 ±0.096** <sup>1</sup> DOBBS 07 CLEO e<sup>+</sup>e<sup>-</sup> at ψ(3770)

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.2 ±0.1 ±0.2	3210 ± 85	<sup>1</sup> HE	05	CLEO	See DOBBS 07
2.1 <sup>+1.0</sup> / <sub>-0.9</sub>		<sup>2</sup> BARLAG	92C	ACCM	π <sup>-</sup> Cu 230 GeV
3.3 ±0.8 ±0.2	168	ADLER	88C	MRK3	e <sup>+</sup> e <sup>-</sup> 3.77 GeV

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

<sup>2</sup> BARLAG 92C computes the branching fraction by topological normalization.

$\Gamma(K^- 3\pi^+ \pi^-) / \Gamma(K^- 2\pi^+)$   $\Gamma_{56} / \Gamma_{40}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.061 ±0.005 OUR FIT** Error includes scale factor of 1.1.

**0.062 ±0.008 OUR AVERAGE** Error includes scale factor of 1.3.

0.058 ±0.002 ±0.006	2923	LINK	03D	FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
0.077 ±0.008 ±0.010	239	FRABETTI	97C	E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.09 ±0.01 ±0.01	113	ANJOS	90D	E691	Photoproduction
------------------	-----	-------	-----	------	-----------------

$\Gamma(\bar{K}^*(892)^0 2\pi^+ \pi^-, \bar{K}^*(892)^0 \rightarrow K^- \pi^+) / \Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{57} / \Gamma_{56}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.21 ±0.04 ±0.06** LINK 03D FOCS γ A,  $\bar{E}_\gamma \approx 180$  GeV

$\Gamma(\bar{K}^*(892)^0 \rho^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+) / \Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{58} / \Gamma_{56}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.40 ±0.03 ±0.06** LINK 03D FOCS γ A,  $\bar{E}_\gamma \approx 180$  GeV

$\Gamma(\bar{K}^*(892)^0 \rho^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+) / \Gamma(K^- 2\pi^+)$   $\Gamma_{58} / \Gamma_{40}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.016 ±0.007 ±0.004	FRABETTI	97C	E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV
---------------------	----------	-----	------	--

$\Gamma(\bar{K}^*(892)^0 2\pi^+ \pi^- \text{no-}\rho, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{60}/\Gamma_{40}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.032 \pm 0.010 \pm 0.008$	FRABETTI	97C	E687 $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV
-----------------------------	----------	-----	--

$\Gamma(K^- \rho^0 2\pi^+)/\Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{61}/\Gamma_{56}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b><math>0.30 \pm 0.04 \pm 0.01</math></b>	LINK	03D	FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
--	------	-----	---

$\Gamma(K^- \rho^0 2\pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{61}/\Gamma_{40}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.034 \pm 0.009 \pm 0.005$	FRABETTI	97C	E687 $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV
-----------------------------	----------	-----	--

$\Gamma(\bar{K}^*(892)^0 a_1(1260)^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{59}/\Gamma_{40}$

Unseen decay modes of the  $\bar{K}^*(892)^0$  and  $a_1(1260)^+$  are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b><math>0.099 \pm 0.008 \pm 0.018</math></b>	LINK	03D	FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
---	------	-----	---

$\Gamma(K^- 3\pi^+ \pi^- \text{nonresonant})/\Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{62}/\Gamma_{56}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b><math>0.07 \pm 0.05 \pm 0.01</math></b>		LINK	03D	FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
--	--	------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.026$	90	FRABETTI	97C	E687 $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV
----------	----	----------	-----	--

$\Gamma(K^+ 2K_S^0)/\Gamma(K^- 2\pi^+)$   $\Gamma_{63}/\Gamma_{40}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**$0.049 \pm 0.022$  OUR AVERAGE** Error includes scale factor of 2.4.

$0.035 \pm 0.010 \pm 0.005$	$39 \pm 9$	ALBRECHT	94I	ARG $e^+ e^- \approx 10$ GeV
-----------------------------	------------	----------	-----	------------------------------

$0.085 \pm 0.018$	$70 \pm 12$	AMMAR	91	CLEO $e^+ e^- \approx 10.5$ GeV
-------------------	-------------	-------	----	---------------------------------

$\Gamma(K^+ K^- K_S^0 \pi^+)/\Gamma(K_S^0 2\pi^+ \pi^-)$   $\Gamma_{64}/\Gamma_{55}$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b><math>7.7 \pm 1.5 \pm 0.9</math></b>	$35 \pm 7$	LINK	01C	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
---	------------	------	-----	---

————— Pionic modes —————

$\Gamma(\pi^+ \pi^0)/\Gamma(K^- 2\pi^+)$   $\Gamma_{65}/\Gamma_{40}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**$1.31 \pm 0.06$  OUR AVERAGE**

$1.29 \pm 0.04 \pm 0.05$	$2649 \pm 76$	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV
--------------------------	---------------	--------	----	----------------------------

$1.33 \pm 0.11 \pm 0.09$	$1229 \pm 99$	AUBERT,B	06F	BABR $e^+ e^- \approx \Upsilon(4S)$
--------------------------	---------------	----------	-----	-------------------------------------

$1.44 \pm 0.19 \pm 0.10$	$171 \pm 22$	ARMS	04	CLEO $e^+ e^- \approx 10$ GeV
--------------------------	--------------	------	----	-------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •


$1.33 \pm 0.07 \pm 0.06$	$914 \pm 46$	RUBIN	06	CLEO See MENDEZ 10
--------------------------	--------------	-------	----	--------------------

$\Gamma(2\pi^+\pi^-)/\Gamma(K^-2\pi^+)$

$\Gamma_{66}/\Gamma_{40}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.48±0.19 OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		
3.52±0.11±0.12	3303 ± 95	RUBIN	06 CLEO	$e^+e^-$ at $\psi(3770)$
4.1 ±1.1 ±0.3	85 ± 22	ABLIKIM	05F BES	$e^+e^- \approx \psi(3770)$
3.11±0.18 <sup>+0.16</sup> <sub>-0.26</sub>	1172	AITALA	01B E791	$\pi^-$ nucleus, 500 GeV
4.3 ±0.3 ±0.3	236	FRABETTI	97D E687	$\gamma$ Be $\approx$ 200 GeV
3.5 ±0.7 ±0.3	83	ANJOS	89 E691	Photoproduction

WEIGHTED AVERAGE  
3.48±0.19 (Error scaled by 1.4)




$\Gamma(\rho^0\pi^+)/\Gamma(2\pi^+\pi^-)$

$\Gamma_{67}/\Gamma_{66}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.25 ±0.04 OUR AVERAGE</b>	Error includes scale factor of 2.4. See the ideogram below.		
0.200 ±0.023 ±0.009	BONVICINI	07 CLEO	Dalitz fit, $\approx$ 2240 evts
0.3082±0.0314±0.0230	LINK	04 FOCS	Dalitz fit, 1527 ± 51 evts
0.336 ±0.032 ±0.022	AITALA	01B E791	Dalitz fit, 1172 evts



**$\Gamma(\pi^+(\pi^+\pi^-)_{S\text{-wave}}) / \Gamma(2\pi^+\pi^-)$   $\Gamma_{68} / \Gamma_{66}$**

This is the "fit fraction" from the Dalitz-plot analysis. See also the next three data blocks.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.5600 \pm 0.0324 \pm 0.0214</math></b>	<sup>1</sup> LINK	04	FOCS Dalitz fit, $1527 \pm 51$ evts

<sup>1</sup> LINK 04 borrows a K-matrix parametrization from ANISOVICH 03 of the full  $\pi\text{-}\pi$  S-wave isoscalar scattering amplitude to describe the  $\pi^+\pi^-$  S-wave component of the  $\pi^+\pi^+\pi^-$  state. The fit fraction given above is a sum over five  $f_0$  mesons, the  $f_0(980)$ ,  $f_0(1300)$ ,  $f_0(1200\text{--}1600)$ ,  $f_0(1500)$ , and  $f_0(1750)$ . See LINK 04 for details and discussion.

**$\Gamma(\sigma\pi^+, \sigma \rightarrow \pi^+\pi^-) / \Gamma(2\pi^+\pi^-)$   $\Gamma_{69} / \Gamma_{66}$**

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.422 \pm 0.027</math> OUR AVERAGE</b>			
$0.418 \pm 0.014 \pm 0.025$	BONVICINI	07	CLEO Dalitz fit, $\approx 2240$ evts
$0.463 \pm 0.090 \pm 0.021$	AITALA	01B	E791 Dalitz fit, 1172 evts

**$\Gamma(f_0(980)\pi^+, f_0(980) \rightarrow \pi^+\pi^-) / \Gamma(2\pi^+\pi^-)$   $\Gamma_{70} / \Gamma_{66}$**

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.048 \pm 0.010</math> OUR AVERAGE</b>	Error includes scale factor of 1.3.		
$0.041 \pm 0.009 \pm 0.003$	BONVICINI	07	CLEO Dalitz fit, $\approx 2240$ evts
$0.062 \pm 0.013 \pm 0.004$	AITALA	01B	E791 Dalitz fit, 1172 evts

$\Gamma(f_0(1370)\pi^+, f_0(1370) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{71}/\Gamma_{66}$


This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.024 ± 0.013 OUR AVERAGE</b>			
0.026 ± 0.018 ± 0.006	BONVICINI 07	CLEO	Dalitz fit, ≈ 2240 evts
0.023 ± 0.015 ± 0.008	AITALA 01B	E791	Dalitz fit, 1172 evts

$\Gamma(f_2(1270)\pi^+, f_2(1270) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{72}/\Gamma_{66}$

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.154 ± 0.025 OUR AVERAGE</b>	Error includes scale factor of 1.9. See the ideogram below.		
0.182 ± 0.026 ± 0.007	BONVICINI 07	CLEO	Dalitz fit, ≈ 2240 evts
0.1174 ± 0.0190 ± 0.0029	LINK 04	FOCS	Dalitz fit, 1527 ± 51 evts
0.194 ± 0.025 ± 0.004	AITALA 01B	E791	Dalitz fit, 1172 evts



$\Gamma(\rho(1450)^0\pi^+, \rho(1450)^0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{73}/\Gamma_{66}$

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.024</b>	95	BONVICINI 07	CLEO	Dalitz fit, ≈ 2240 evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.007 ± 0.007 ± 0.003		AITALA 01B	E791	Dalitz fit, 1172 evts

$\Gamma(f_0(1500)\pi^+, f_0(1500) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{74}/\Gamma_{66}$

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.034 ± 0.010 ± 0.008</b>	BONVICINI 07	CLEO	Dalitz fit, ≈ 2240 evts

$\Gamma(f_0(1710)\pi^+, f_0(1710) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{75}/\Gamma_{66}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.016</b>	95	BONVICINI 07	CLEO	Dalitz fit, $\approx 2240$ evts

$\Gamma(f_0(1790)\pi^+, f_0(1790) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{76}/\Gamma_{66}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.02</b>	95	BONVICINI 07	CLEO	Dalitz fit, $\approx 2240$ evts

$\Gamma((\pi^+\pi^+)_{S\text{-wave}}\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{77}/\Gamma_{66}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.037</b>	95	BONVICINI 07	CLEO	Dalitz fit, $\approx 2240$ evts

$\Gamma(2\pi^+\pi^- \text{ nonresonant})/\Gamma(2\pi^+\pi^-)$   $\Gamma_{78}/\Gamma_{66}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.035</b>	95	BONVICINI 07	CLEO	Dalitz fit, $\approx 2240$ evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.078 $\pm$ 0.060 $\pm$ 0.027      AITALA      01B      E791      Dalitz fit, 1172 evts

$\Gamma(\pi^+2\pi^0)/\Gamma(K^-2\pi^+)$   $\Gamma_{79}/\Gamma_{40}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.0<math>\pm</math>0.3<math>\pm</math>0.3</b>	1535 $\pm$ 89	RUBIN 06	CLEO	$e^+e^-$ at $\psi(3770)$

$\Gamma(2\pi^+\pi^-\pi^0)/\Gamma(K^-2\pi^+)$   $\Gamma_{80}/\Gamma_{40}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.4<math>\pm</math>0.5<math>\pm</math>0.6</b>	5701 $\pm$ 205	RUBIN 06	CLEO	$e^+e^-$ at $\psi(3770)$

$\Gamma(\eta\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{84}/\Gamma$

Unseen decay modes of the  $\eta$  are included.

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>34.3<math>\pm</math>1.4<math>\pm</math>1.7</b>	1033 $\pm$ 42	ARTUSO 08	CLEO	See MENDEZ 10

$\Gamma(\eta\pi^+)/\Gamma(K^-2\pi^+)$   $\Gamma_{84}/\Gamma_{40}$

Unseen decay modes of the  $\eta$  are included.

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.87<math>\pm</math>0.09<math>\pm</math>0.19</b>	2940 $\pm$ 68	MENDEZ 10	CLEO	$e^+e^-$ at 3774 MeV
3.81 $\pm$ 0.26 $\pm$ 0.21	377 $\pm$ 26	RUBIN 06	CLEO	See ARTUSO 08

$\Gamma(\omega\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{86}/\Gamma$

Unseen decay modes of the  $\omega$  are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.4 <math>\times 10^{-4}</math></b>	90	RUBIN 06	CLEO	$e^+e^-$ at $\psi(3770)$

$\Gamma(3\pi^+2\pi^-)/\Gamma(K^-2\pi^+)$   $\Gamma_{83}/\Gamma_{40}$

VALUE (units  $10^{-2}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**1.77±0.17 OUR FIT**

**1.73±0.20±0.17**    732 ± 77    RUBIN    06    CLEO     $e^+e^-$  at  $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ± 0.4 ± 0.2    58    FRABETTI    97C E687     $\gamma$ Be,  $\bar{E}_\gamma \approx 200$  GeV

$\Gamma(3\pi^+2\pi^-)/\Gamma(K^-3\pi^+\pi^-)$   $\Gamma_{83}/\Gamma_{56}$

VALUE    EVTS    DOCUMENT ID    TECN    COMMENT

**0.289±0.019 OUR FIT**

**0.290±0.017±0.011**    835    LINK    03D FOCS     $\gamma$  A,  $\bar{E}_\gamma \approx 180$  GeV

$\Gamma(\eta\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{85}/\Gamma$

VALUE (units  $10^{-4}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**13.8±3.1±1.6**    149 ± 34    ARTUSO    08    CLEO     $e^+e^-$  at  $\psi(3770)$

$\Gamma(\eta'(958)\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{87}/\Gamma$

Unseen decay modes of the  $\eta'(958)$  are included.

VALUE (units  $10^{-4}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

44.2±2.5±2.9    352 ± 20    ARTUSO    08    CLEO    See MENDEZ 10

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^-2\pi^+)$   $\Gamma_{87}/\Gamma_{40}$

Unseen decay modes of the  $\eta'(958)$  are included.

VALUE (units  $10^{-2}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**5.12±0.17±0.25**    1037 ± 35    MENDEZ    10    CLEO     $e^+e^-$  at 3774 MeV

$\Gamma(\eta'(958)\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{88}/\Gamma$

Unseen decay modes of the  $\eta'(958)$  are included.

VALUE (units  $10^{-4}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**15.7±4.3±2.5**    33 ± 9    ARTUSO    08    CLEO     $e^+e^-$  at  $\psi(3770)$

————— **Hadronic modes with a  $K\bar{K}$  pair** —————

$\Gamma(K^+K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{89}/\Gamma$

VALUE (units  $10^{-3}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.14±0.09±0.08    1971 ± 51    BONVICINI    08    CLEO    See MENDEZ 10

$\Gamma(K^+K_S^0)/\Gamma(K_S^0\pi^+)$   $\Gamma_{89}/\Gamma_{38}$

VALUE    EVTS    DOCUMENT ID    TECN    COMMENT

**0.193 ± 0.007 OUR FIT**    Error includes scale factor of 3.2.

**0.1901±0.0024 OUR AVERAGE**

0.1899±0.0011±0.0022    101k±561    WON    09    BELL     $e^+e^-$  at  $\Upsilon(4S)$

0.1892±0.0155±0.0073    278 ± 21    ARMS    04    CLEO     $e^+e^- \approx 10$  GeV

0.1996±0.0119±0.0096    949    LINK    02B FOCS     $\gamma$  A,  $\bar{E}_\gamma \approx 180$  GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.222 ± 0.037 ± 0.013	63 ± 10	ABLIKIM	05F	BES	$e^+e^- \approx \psi(3770)$
0.222 ± 0.041 ± 0.019	70	BISHAI	97	CLEO	See ARMS 04
0.25 ± 0.04 ± 0.02	129	FRABETTI	95	E687	$\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV
0.271 ± 0.065 ± 0.039	69	ANJOS	90C	E691	$\gamma$ Be
0.317 ± 0.086 ± 0.048	31	BALTRUSAIT..85E	MRK3		$e^+e^-$ 3.77 GeV
0.25 ± 0.15	6	SCHINDLER	81	MRK2	$e^+e^-$ 3.771 GeV

### $\Gamma(K^+ K_S^0)/\Gamma(K^- 2\pi^+)$

$\Gamma_{89}/\Gamma_{40}$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**3.11 ± 0.16 OUR FIT** Error includes scale factor of 3.3.

**3.35 ± 0.06 ± 0.07** 5161 ± 86 MENDEZ 10 CLEO  $e^+e^-$  at 3774 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.02 ± 0.18 ± 0.15 949 <sup>1</sup> LINK 02B FOCS  $\gamma$  nucleus,  $\bar{E}_\gamma \approx 180$  GeV

<sup>1</sup> This LINK 02B result is redundant with a result in the previous datablock.

### $\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$

$\Gamma_{90}/\Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**0.954 ± 0.026 OUR FIT** Error includes scale factor of 1.1.

**0.935 ± 0.017 ± 0.024** <sup>1</sup> DOBBS 07 CLEO  $e^+e^-$  at  $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.97 ± 0.04 ± 0.04 1250 ± 40 <sup>1</sup> HE 05 CLEO See DOBBS 07

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

### $\Gamma(K^+ K^- \pi^+)/\Gamma(K^- 2\pi^+)$

$\Gamma_{90}/\Gamma_{40}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

**0.1045 ± 0.0022 OUR FIT** Error includes scale factor of 1.3.

**0.1058 ± 0.0029 OUR AVERAGE** Error includes scale factor of 1.4.

0.117 ± 0.013 ± 0.007 181 ± 20 ABLIKIM 05F BES  $e^+e^- \approx \psi(3770)$

0.107 ± 0.001 ± 0.002 43k AUBERT 05s BABR  $e^+e^- \approx \Upsilon(4S)$

0.093 ± 0.010  $\begin{smallmatrix} +0.008 \\ -0.006 \end{smallmatrix}$  JUN 00 SELX  $\Sigma^-$  nucleus, 600 GeV

0.0976 ± 0.0042 ± 0.0046 FRABETTI 95B E687  $\gamma$  Be,  $\bar{E}_\gamma \approx 200$  GeV

### $\Gamma(\phi\pi^+, \phi \rightarrow K^+ K^-)/\Gamma(K^+ K^- \pi^+)$

$\Gamma_{91}/\Gamma_{90}$

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
------------------	--------------------	-------------	----------------

**27.8 ± 0.4  $\begin{smallmatrix} +0.2 \\ -0.5 \end{smallmatrix}$**  RUBIN 08 CLEO Dalitz fit, 19,458 ± 163 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

29.2 ± 3.1 ± 3.0 FRABETTI 95B E687 Dalitz fit, 915 evts

### $\Gamma(K^+ \bar{K}^*(892)^0, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+)$

$\Gamma_{92}/\Gamma_{90}$

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
------------------	--------------------	-------------	----------------

**25.7 ± 0.5  $\begin{smallmatrix} +0.4 \\ -1.2 \end{smallmatrix}$**  RUBIN 08 CLEO Dalitz fit, 19,458 ± 163 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

30.1 ± 2.0 ± 2.5 FRABETTI 95B E687 Dalitz fit, 915 evts

$\Gamma(K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+) / \Gamma(K^+ K^- \pi^+)$   $\Gamma_{93} / \Gamma_{90}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$18.8 \pm 1.2^{+3.3}_{-3.4}$	RUBIN	08	CLEO Dalitz fit, 19,458 ± 163 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

$37.0 \pm 3.5 \pm 1.8$	FRABETTI	95B	E687 Dalitz fit, 915 evts
------------------------	----------	-----	---------------------------

$\Gamma(K^+ \bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow K^- \pi^+) / \Gamma(K^+ K^- \pi^+)$   $\Gamma_{94} / \Gamma_{90}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$1.7 \pm 0.4^{+1.2}_{-0.7}$	RUBIN	08	CLEO Dalitz fit, 19,458 ± 163 evts

$\Gamma(K^+ \bar{K}_0^*(800), \bar{K}_0^* \rightarrow K^- \pi^+) / \Gamma(K^+ K^- \pi^+)$   $\Gamma_{95} / \Gamma_{90}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$7.0 \pm 0.8^{+3.5}_{-2.0}$	RUBIN	08	CLEO Dalitz fit, 19,458 ± 163 evts

$\Gamma(a_0(1450)^0 \pi^+, a_0^0 \rightarrow K^+ K^-) / \Gamma(K^+ K^- \pi^+)$   $\Gamma_{96} / \Gamma_{90}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$4.6 \pm 0.6^{+7.2}_{-1.8}$	RUBIN	08	CLEO Dalitz fit, 19,458 ± 163 evts

$\Gamma(\phi(1680) \pi^+, \phi \rightarrow K^+ K^-) / \Gamma(K^+ K^- \pi^+)$   $\Gamma_{97} / \Gamma_{90}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$0.51 \pm 0.11^{+0.37}_{-0.16}$	RUBIN	08	CLEO Dalitz fit, 19,458 ± 163 evts

$\Gamma(K^*(892)^+ K_S^0) / \Gamma(K_S^0 \pi^+)$   $\Gamma_{105} / \Gamma_{38}$

Unseen decay modes of the  $K^*(892)^+$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.1 \pm 0.3 \pm 0.4$	67	FRABETTI	95	E687 $\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV

$\Gamma(\phi \pi^+ \pi^0) / \Gamma_{\text{total}}$   $\Gamma_{102} / \Gamma$

Unseen decay modes of the  $\phi$  are included.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.023 \pm 0.010$	<sup>1</sup> BARLAG	92C	ACCM $\pi^-$ Cu 230 GeV

<sup>1</sup> BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\phi \rho^+) / \Gamma(K^- 2\pi^+)$   $\Gamma_{103} / \Gamma_{40}$

Unseen decay modes of the  $\phi$  are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.16$	90	DAOUDI	92	CLEO $e^+ e^- \approx 10.5$ GeV

$\Gamma(K^+ K^- \pi^+ \pi^0 \text{ non-}\phi) / \Gamma_{\text{total}}$   $\Gamma_{104} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.015^{+0.007}_{-0.006}$	<sup>1</sup> BARLAG	92C	ACCM $\pi^-$ Cu 230 GeV

<sup>1</sup> BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^+ K^- \pi^+ \pi^0 \text{non-}\phi) / \Gamma(K^- 2\pi^+)$   $\Gamma_{104} / \Gamma_{40}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.25	90	ANJOS	89E E691	Photoproduction
-------	----	-------	----------	-----------------

$\Gamma(K^+ K_S^0 \pi^+ \pi^-) / \Gamma(K_S^0 2\pi^+ \pi^-)$   $\Gamma_{99} / \Gamma_{55}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>5.62 ± 0.39 ± 0.40</b>	469 ± 32	LINK	01C FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
---------------------------	----------	------	----------	--

$\Gamma(K_S^0 K^- 2\pi^+) / \Gamma(K_S^0 2\pi^+ \pi^-)$   $\Gamma_{100} / \Gamma_{55}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>7.68 ± 0.41 ± 0.32</b>	670 ± 35	LINK	01C FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
---------------------------	----------	------	----------	--

$\Gamma(K^+ K^- 2\pi^+ \pi^-) / \Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{101} / \Gamma_{56}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

<b>0.040 ± 0.009 ± 0.019</b>	38	LINK	03D FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
------------------------------	----	------	----------	--

———— Doubly Cabibbo-suppressed modes ————

$\Gamma(K^+ \pi^0) / \Gamma_{\text{total}}$   $\Gamma_{106} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>1.83 ± 0.26 OUR FIT</b>	Error includes scale factor of 1.4.			
----------------------------	-------------------------------------	--	--	--

<b>2.52 ± 0.47 ± 0.26</b>	189 ± 37	AUBERT,B	06F BABR	$e^+ e^- \approx \Upsilon(4S)$
---------------------------	----------	----------	----------	--------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.28 ± 0.36 ± 0.17	148 ± 23	DYTMAN	06 CLEO	See MENDEZ 10
--------------------	----------	--------	---------	---------------

$\Gamma(K^+ \pi^0) / \Gamma(K^- 2\pi^+)$   $\Gamma_{106} / \Gamma_{40}$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>2.01 ± 0.29 OUR FIT</b>	Error includes scale factor of 1.4.			
----------------------------	-------------------------------------	--	--	--

<b>1.9 ± 0.2 ± 0.1</b>	343 ± 37	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV
------------------------	----------	--------	---------	-----------------------

$\Gamma(K^+ \eta) / \Gamma(\eta \pi^+)$   $\Gamma_{107} / \Gamma_{84}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

<b>3.06 ± 0.43 ± 0.14</b>	166 ± 23	WON	11 BELL	$e^+ e^- \approx \Upsilon(4S)$
---------------------------	----------	-----	---------	--------------------------------

$\Gamma(K^+ \eta) / \Gamma(K^- 2\pi^+)$   $\Gamma_{107} / \Gamma_{40}$

Unseen decay modes of the  $\eta$  are included.

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.15	90	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV
-------	----	--------	---------	-----------------------

$\Gamma(K^+ \eta'(958)) / \Gamma(\eta'(958) \pi^+)$   $\Gamma_{108} / \Gamma_{87}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

<b>3.77 ± 0.39 ± 0.10</b>	180 ± 19	WON	11 BELL	$e^+ e^- \approx \Upsilon(4S)$
---------------------------	----------	-----	---------	--------------------------------

$\Gamma(K^+ \eta'(958))/\Gamma(K^- 2\pi^+)$   $\Gamma_{108}/\Gamma_{40}$ Unseen decay modes of the  $\eta'(958)$  are included.

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.20	90	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV
-------	----	--------	----	----------------------------

 $\Gamma(K^+ \pi^+ \pi^-)/\Gamma(K^- 2\pi^+)$   $\Gamma_{109}/\Gamma_{40}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**5.77 ± 0.22 OUR AVERAGE**

5.69 ± 0.18 ± 0.14	2638 ± 84	KO	09	BELL $e^+ e^-$ at $\Upsilon(4S)$
6.5 ± 0.8 ± 0.4	189 ± 24	LINK	04F	FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
7.7 ± 1.7 ± 0.8	59 ± 13	AITALA	97C	E791 $\pi^-$ A, 500 GeV
7.2 ± 2.3 ± 1.7	21	FRABETTI	95E	E687 $\gamma$ Be, $\bar{E}_\gamma = 220$ GeV

 $\Gamma(K^+ \rho^0)/\Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{110}/\Gamma_{109}$ 

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.39 ± 0.09 OUR AVERAGE**

0.3943 ± 0.0787 ± 0.0815	LINK	04F	FOCS Dalitz fit, 189 evts
0.37 ± 0.14 ± 0.07	AITALA	97C	E791 Dalitz fit, 59 evts

 $\Gamma(K^+ f_0(980), f_0(980) \rightarrow \pi^+ \pi^-)/\Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{112}/\Gamma_{109}$ 

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>0.0892 ± 0.0333 ± 0.0412</b>	LINK	04F	FOCS Dalitz fit, 189 evts
---------------------------------	------	-----	---------------------------

 $\Gamma(K^*(892)^0 \pi^+, K^*(892)^0 \rightarrow K^+ \pi^-)/\Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{111}/\Gamma_{109}$ 

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.47 ± 0.08 OUR AVERAGE**

0.5220 ± 0.0684 ± 0.0638	LINK	04F	FOCS Dalitz fit, 189 evts
0.35 ± 0.14 ± 0.01	AITALA	97C	E791 Dalitz fit, 59 evts

 $\Gamma(K_2^*(1430)^0 \pi^+, K_2^*(1430)^0 \rightarrow K^+ \pi^-)/\Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{113}/\Gamma_{109}$ 

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>0.0803 ± 0.0372 ± 0.0391</b>	LINK	04F	FOCS Dalitz fit, 189 evts
---------------------------------	------	-----	---------------------------

 $\Gamma(K^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{114}/\Gamma_{109}$ 

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.36 ± 0.14 ± 0.07	<sup>1</sup> AITALA	97C	E791 Dalitz fit, 59 evts
--------------------	---------------------	-----	--------------------------

<sup>1</sup> LINK 04F, with three times as many events, finds no need for a nonresonant amplitude. $\Gamma(2K^+ K^-)/\Gamma(K^- 2\pi^+)$   $\Gamma_{115}/\Gamma_{40}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>9.49 ± 2.17 ± 0.22</b>	65	<sup>1</sup> LINK	02I	FOCS $\gamma$ nucleus, $\approx 180$ GeV
---------------------------	----	-------------------	-----	--

<sup>1</sup> LINK 02I finds little evidence for  $\phi K^+$  or  $f_0(980) K^+$  submodes.

————— Rare or forbidden modes —————

$\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_{116}/\Gamma$

A test for the  $\Delta C = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<5.9 \times 10^{-6}$	90	<sup>1</sup> RUBIN	10 CLEO	$e^+ e^-$ at $\psi(3770)$
$<7.4 \times 10^{-6}$	90	HE	05A CLEO	See RUBIN 10
$<5.2 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<1.1 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<6.6 \times 10^{-5}$	90	AITALA	96 E791	$\pi^- N$ 500 GeV
$<2.5 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
$<2.6 \times 10^{-3}$	90	HAAS	88 CLEO	$e^+ e^-$ 10 GeV

<sup>1</sup>This RUBIN 10 limit is for the  $e^+ e^-$  mass in the continuum away from the  $\phi(1020)$ . See the next data block.

$\Gamma(\pi^+ \phi, \phi \rightarrow e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_{117}/\Gamma$

This is *not* a test for the  $\Delta C = 1$  weak neutral current, but leads to the  $\pi^+ e^+ e^-$  final state.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$(1.7^{+1.4}_{-0.9} \pm 0.1) \times 10^{-6}$	4	<sup>1</sup> RUBIN	10 CLEO	$e^+ e^-$ at $\psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$(2.7^{+3.6}_{-1.8} \pm 0.2) \times 10^{-6}$	2	HE	05A CLEO	See RUBIN 10

<sup>1</sup>This RUBIN 10 result is consistent with the known  $D^+ \rightarrow \phi \pi^+$  and  $\phi \rightarrow e^+ e^-$  fractions.

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$

$\Gamma_{118}/\Gamma$

A test for the  $\Delta C = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.3 \times 10^{-8}$	90	AAIJ	13AF LHCb	$p\bar{p}$ at 7 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<6.5 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$<3.9 \times 10^{-6}$	90	<sup>1</sup> ABAZOV	08D D0	$p\bar{p}$ , $E_{\text{cm}} = 1.96$ TeV
$<8.8 \times 10^{-6}$	90	LINK	03F FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
$<1.5 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<8.9 \times 10^{-5}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<1.8 \times 10^{-5}$	90	AITALA	96 E791	$\pi^- N$ 500 GeV
$<2.2 \times 10^{-4}$	90	KODAMA	95 E653	$\pi^-$ emulsion 600 GeV
$<5.9 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
$<2.9 \times 10^{-3}$	90	HAAS	88 CLEO	$e^+ e^-$ 10 GeV

<sup>1</sup>This ABAZOV 08D limit is for the  $\mu^+ \mu^-$  mass in the continuum away from the  $\phi(1020)$ . See the next data block.

$\Gamma(\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-) / \Gamma_{\text{total}}$   $\Gamma_{119} / \Gamma$

This is *not* a test for the  $\Delta C = 1$  weak neutral current, but leads to the  $\pi^+ \mu^+ \mu^-$  final state.

VALUE	DOCUMENT ID	TECN	COMMENT
$(1.8 \pm 0.5 \pm 0.6) \times 10^{-6}$	<sup>1</sup> ABAZOV 08D D0		$p\bar{p}$ , $E_{\text{cm}} = 1.96$ TeV

<sup>1</sup> This ABAZOV 08D value is consistent with the known  $D^+ \rightarrow \phi \pi^+$  and  $\phi \rightarrow \mu^+ \mu^-$  fractions.

$\Gamma(\rho^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$   $\Gamma_{120} / \Gamma$

A test for the  $\Delta C = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-4}$	90	KODAMA 95 E653		$\pi^-$ emulsion 600 GeV

$\Gamma(K^+ e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{121} / \Gamma$

Both quarks would have to change flavor for this decay to occur.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.0 \times 10^{-6}$	90	LEES 11G BABR		$e^+ e^- \approx \mathcal{T}(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.0 \times 10^{-6}$	90	RUBIN 10 CLEO		$e^+ e^-$ at $\psi(3770)$
$< 6.2 \times 10^{-6}$	90	HE 05A CLEO		See RUBIN 10
$< 2.0 \times 10^{-4}$	90	AITALA 99G E791		$\pi^- N$ 500 GeV
$< 2.0 \times 10^{-4}$	90	FRABETTI 97B E687		$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$< 4.8 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+ e^-$ 29 GeV

$\Gamma(K^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$   $\Gamma_{122} / \Gamma$

Both quarks would have to change flavor for this decay to occur.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.3 \times 10^{-6}$	90	LEES 11G BABR		$e^+ e^- \approx \mathcal{T}(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 9.2 \times 10^{-6}$	90	LINK 03F FOCS		$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
$< 4.4 \times 10^{-5}$	90	AITALA 99G E791		$\pi^- N$ 500 GeV
$< 9.7 \times 10^{-5}$	90	FRABETTI 97B E687		$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$< 3.2 \times 10^{-4}$	90	KODAMA 95 E653		$\pi^-$ emulsion 600 GeV
$< 9.2 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+ e^-$ 29 GeV

$\Gamma(\pi^+ e^+ \mu^-) / \Gamma_{\text{total}}$   $\Gamma_{123} / \Gamma$

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.9 \times 10^{-6}$	90	LEES 11G BABR		$e^+ e^- \approx \mathcal{T}(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.1 \times 10^{-4}$	90	FRABETTI 97B E687		$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$< 3.3 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+ e^-$ 29 GeV

**$\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$**   **$\Gamma_{124}/\Gamma$**

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;3.6 \times 10^{-6}</math></b>	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.3 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<3.3 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

**$\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$**   **$\Gamma_{125}/\Gamma$**

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.2 \times 10^{-6}</math></b>	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.3 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<3.4 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

**$\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$**   **$\Gamma_{126}/\Gamma$**

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;2.8 \times 10^{-6}</math></b>	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.2 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<3.4 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

**$\Gamma(\pi^- 2e^+)/\Gamma_{\text{total}}$**   **$\Gamma_{127}/\Gamma$**

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.1 \times 10^{-6}</math></b>	90	RUBIN	10 CLEO	$e^+ e^-$ at $\psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.9 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$<3.6 \times 10^{-6}$	90	HE	05A CLEO	See RUBIN 10
$<9.6 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<1.1 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<4.8 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

**$\Gamma(\pi^- 2\mu^+)/\Gamma_{\text{total}}$**   **$\Gamma_{128}/\Gamma$**

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;2.2 \times 10^{-8}</math></b>	90	AAIJ	13AF LHCB	$pp$ at 7 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<2.0 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$<4.8 \times 10^{-6}$	90	LINK	03F FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
$<1.7 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<8.7 \times 10^{-5}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<2.2 \times 10^{-4}$	90	KODAMA	95 E653	$\pi^-$ emulsion 600 GeV
$<6.8 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$   $\Gamma_{129}/\Gamma$

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<5.0 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<1.1 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<3.7 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

$\Gamma(\rho^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{130}/\Gamma$

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-4}$	90	KODAMA	95 E653	$\pi^-$ emulsion 600 GeV

$\Gamma(K^- 2e^+)/\Gamma_{\text{total}}$   $\Gamma_{131}/\Gamma$

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.9 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<3.5 \times 10^{-6}$	90	RUBIN	10 CLEO	$e^+ e^-$ at $\psi(3770)$
$<4.5 \times 10^{-6}$	90	HE	05A CLEO	See RUBIN 10
$<1.2 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<9.1 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

$\Gamma(K^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{132}/\Gamma$

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<10 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.3 \times 10^{-5}$	90	LINK	03F FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
$<1.2 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<3.2 \times 10^{-4}$	90	KODAMA	95 E653	$\pi^-$ emulsion 600 GeV
$<4.3 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$   $\Gamma_{133}/\Gamma$

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.9 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.3 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<4.0 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

$\Gamma(K^*(892)^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{134}/\Gamma$

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.5 \times 10^{-4}$	90	KODAMA	95 E653	$\pi^-$ emulsion 600 GeV

**$D^\pm$  CP-VIOLATING DECAY-RATE ASYMMETRIES**

This is the difference between  $D^+$  and  $D^-$  partial widths for these modes divided by the sum of the widths.

 **$A_{CP}(\mu^\pm \nu)$  in  $D^+ \rightarrow \mu^+ \nu_\mu$ ,  $D^- \rightarrow \mu^- \bar{\nu}_\mu$** 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>+8±8</b>	EISENSTEIN 08	CLEO	$e^+ e^-$ at $\psi(3770)$

 **$A_{CP}(K_S^0 \pi^\pm)$  in  $D^\pm \rightarrow K_S^0 \pi^\pm$** 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-0.41 ± 0.09 OUR AVERAGE</b>				
-0.363 ± 0.094 ± 0.067	1738k	<sup>1</sup> KO	12A BELL	$e^+ e^- \approx \Upsilon(nS)$
-0.44 ± 0.13 ± 0.10	807k	DEL-AMO-SA..11H	BABR	$e^+ e^- \approx \Upsilon(4S)$
-1.3 ± 0.7 ± 0.3	30k	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV
-1.6 ± 1.5 ± 0.9	10.6k	<sup>2</sup> LINK	02B FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.71 ± 0.19 ± 0.20		KO	10 BELL	See KO 12A
-0.6 ± 1.0 ± 0.3		DOBBS	07 CLEO	See MENDEZ 10

<sup>1</sup> KO 12A finds that after subtracting the contribution due to  $K^0 - \bar{K}^0$  mixing, the CP asymmetry due to the change of charm is  $(-0.024 \pm 0.094 \pm 0.067)\%$ , consistent with zero.

<sup>2</sup> LINK 02B measures  $N(D^+ \rightarrow K_S^0 \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

 **$A_{CP}(K^\mp 2\pi^\pm)$  in  $D^+ \rightarrow K^- 2\pi^+$ ,  $D^- \rightarrow K^+ 2\pi^-$** 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-0.1±0.4±0.9</b>	231k	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.5±0.4±0.9		DOBBS	07 CLEO	See MENDEZ 10
--------------	--	-------	---------	---------------

 **$A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0)$  in  $D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$ ,  $D^- \rightarrow K^+ \pi^- \pi^- \pi^0$** 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>+1.0±0.9±0.9</b>	DOBBS 07	CLEO	$e^+ e^-$ at $\psi(3770)$

 **$A_{CP}(K_S^0 \pi^\pm \pi^0)$  in  $D^+ \rightarrow K_S^0 \pi^+ \pi^0$ ,  $D^- \rightarrow K_S^0 \pi^- \pi^0$** 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>+0.3±0.9±0.3</b>	DOBBS 07	CLEO	$e^+ e^-$ at $\psi(3770)$

 **$A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-)$  in  $D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$ ,  $D^- \rightarrow K_S^0 \pi^- \pi^- \pi^+$** 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>+0.1±1.1±0.6</b>	DOBBS 07	CLEO	$e^+ e^-$ at $\psi(3770)$

 **$A_{CP}(\pi^\pm \pi^0)$  in  $D^\pm \rightarrow \pi^\pm \pi^0$** 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>+2.9±2.9±0.3</b>	2.6k	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV

### $A_{CP}(\pi^\pm \eta)$ in $D^\pm \rightarrow \pi^\pm \eta$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.0 \pm 1.5</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.4.		
$+1.74 \pm 1.13 \pm 0.19$		WON	11	BELL $e^+ e^- \approx \Upsilon(4S)$
$-2.0 \pm 2.3 \pm 0.3$	2.9k	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV

### $A_{CP}(\pi^\pm \eta'(958))$ in $D^\pm \rightarrow \pi^\pm \eta'(958)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.5 \pm 1.2</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.1.		
$-0.12 \pm 1.12 \pm 0.17$		WON	11	BELL $e^+ e^- \approx \Upsilon(4S)$
$-4.0 \pm 3.4 \pm 0.3$	1.0k	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV

### $A_{CP}(K_S^0 K^\pm)$ in $D^\pm \rightarrow K_S^0 K^\pm$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.11 \pm 0.25</math></b>	<b>OUR AVERAGE</b>			
$-0.25 \pm 0.28 \pm 0.14$	277k	<sup>1</sup> KO	13	BELL $e^+ e^-$ at $\Upsilon(nS)$
$0.13 \pm 0.36 \pm 0.25$	159k	<sup>2</sup> LEES	13E	BABR $e^+ e^-$ at $\Upsilon(4S)$
$-0.2 \pm 1.5 \pm 0.9$	5.2k	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV
$7.1 \pm 6.1 \pm 1.2$	949	<sup>3</sup> LINK	02B	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.16 \pm 0.58 \pm 0.25$		KO	10	BELL $e^+ e^- \approx \Upsilon(4S)$
$6.9 \pm 6.0 \pm 1.5$	949	<sup>4</sup> LINK	02B	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

<sup>1</sup> KO 13 finds that after subtracting the contribution due to  $K^0 - \bar{K}^0$  mixing, the  $CP$  asymmetry is  $(+0.08 \pm 0.28 \pm 0.14)\%$ .

<sup>2</sup> LEES 13E finds that after subtracting the contribution due to  $K^0 - \bar{K}^0$  mixing, the  $CP$  asymmetry is  $(+0.46 \pm 0.36 \pm 0.25)\%$ .

<sup>3</sup> LINK 02B measures  $N(D^+ \rightarrow K_S^0 K^+)/N(D^+ \rightarrow K_S^0 \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

<sup>4</sup> LINK 02B measures  $N(D^+ \rightarrow K_S^0 K^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

### $A_{CP}(K^+ K^- \pi^\pm)$ in $D^\pm \rightarrow K^+ K^- \pi^\pm$

See also AAIJ 11G for a search for  $CP$  asymmetry in the  $D^\pm \rightarrow K^+ K^- \pi^\pm$  Dalitz plots using 370k decays and four different binning schemes. No evidence for  $CP$  asymmetry was found.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.36 \pm 0.29</math></b>	<b>OUR AVERAGE</b>			
$0.37 \pm 0.30 \pm 0.15$	224k	<sup>1</sup> LEES	13F	BABR $e^+ e^-$ at $\Upsilon(4S)$
$-0.03 \pm 0.84 \pm 0.29$		RUBIN	08	CLEO $e^+ e^-$ at 3774 MeV
$-0.1 \pm 1.5 \pm 0.8$		DOBBS	07	CLEO $e^+ e^-$ at $\psi(3770)$
$+1.4 \pm 1.0 \pm 0.8$	43k	<sup>2</sup> AUBERT	05S	BABR $e^+ e^-$ at $\Upsilon(4S)$
$+0.6 \pm 1.1 \pm 0.5$	14k	<sup>3</sup> LINK	00B	FOCS
$-1.4 \pm 2.9$		<sup>3</sup> AITALA	97B	E791 $-0.062 < A_{CP} < +0.034$ (90% CL)
$-3.1 \pm 6.8$		<sup>3</sup> FRABETTI	94I	E687 $-0.14 < A_{CP} < +0.081$ (90% CL)

<sup>1</sup> This is the integrated  $CP$  asymmetry. LEES 13F also searches for  $CP$  asymmetries in four regions of the Dalitz plots (two of which are listed below); in comparisons of binned  $D^+$  and  $D^-$  Dalitz plots; in parametrized fits to those plots, including 2-body submodes; and in comparisons of Legendre-polynomial distributions for the  $K^+K^-$  and  $K^-\pi^+$  systems.

<sup>2</sup> AUBERT 05S measures  $N(D^+ \rightarrow K^+K^-\pi^+)/N(D_s^+ \rightarrow K^+K^-\pi^+)$ , the ratio of the numbers of events observed, and similarly for the  $D^-$ .

<sup>3</sup> FRABETTI 94I, AITALA 98C, and LINK 00B measure  $N(D^+ \rightarrow K^-K^+\pi^+)/N(D^+ \rightarrow K^-\pi^+\pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

### $A_{CP}(K^\pm K^{*0})$ in $D^+ \rightarrow K^+\bar{K}^{*0}$ , $D^- \rightarrow K^-K^{*0}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.3 \pm 0.4</math> OUR AVERAGE</b>				
$-0.3 \pm 0.4 \pm 0.2$	73k	<sup>1</sup> LEES	13F BABR	$e^+e^-$ at $\Upsilon(4S)$
$-0.4 \pm 2.0 \pm 0.6$		RUBIN	08 CLEO	Fit-fraction asymmetry
$+0.9 \pm 1.7 \pm 0.7$	11k	<sup>2</sup> AUBERT	05S BABR	$e^+e^-$ at $\Upsilon(4S)$
$-1.0 \pm 5.0$		<sup>3</sup> AITALA	97B E791	$-0.092 < A_{CP} < +0.072$ (90% CL)
$-12 \pm 13$		<sup>3</sup> FRABETTI	94I E687	$-0.33 < A_{CP} < +0.094$ (90% CL)

<sup>1</sup> This LEES 13F result is for the  $K^\mp\pi^\pm$  mass-squared between 0.4 and 1.0  $\text{GeV}^2$ , and does not actually separate out the  $K^*$ .

<sup>2</sup> AUBERT 05S measures  $N(D^+ \rightarrow K^+\bar{K}^{*0})/N(D_s^+ \rightarrow K^+K^-\pi^+)$ , the ratio of the numbers of events observed, and similarly for the  $D^-$ .

<sup>3</sup> FRABETTI 94I and AITALA 97B measure  $N(D^+ \rightarrow K^+\bar{K}^*(892)^0)/N(D^+ \rightarrow K^-\pi^+\pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

### $A_{CP}(\phi\pi^\pm)$ in $D^\pm \rightarrow \phi\pi^\pm$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.09 \pm 0.19</math> OUR AVERAGE</b>				
		Error includes scale factor of 1.2.		
$-0.04 \pm 0.14 \pm 0.14$	1.58M	AAIJ	13W LHCb	$pp$ at 7 TeV
$-0.3 \pm 0.3 \pm 0.5$	97k	<sup>1</sup> LEES	13F BABR	$e^+e^-$ at $\Upsilon(4S)$
$+0.51 \pm 0.28 \pm 0.05$	237k	STARIC	12 BELL	Mainly at $\Upsilon(4S)$
$-1.8 \pm 1.6 \begin{smallmatrix} +0.2 \\ -0.4 \end{smallmatrix}$		RUBIN	08 CLEO	Fit-fraction asymmetry
$+0.2 \pm 1.5 \pm 0.6$	10k	<sup>2</sup> AUBERT	05S BABR	$e^+e^-$ at $\Upsilon(4S)$
$-2.8 \pm 3.6$		<sup>3</sup> AITALA	97B E791	$-0.087 < A_{CP} < +0.031$ (90% CL)
$+6.6 \pm 8.6$		<sup>3</sup> FRABETTI	94I E687	$-0.075 < A_{CP} < +0.21$ (90% CL)

<sup>1</sup> This LEES 13F result is for the  $K^+K^-$  mass-squared less than 1.3  $\text{GeV}^2$  and the  $K^\mp\pi^\pm$  mass-squared above 1.0  $\text{GeV}^2$ , and does not actually separate out the  $\phi$ .

<sup>2</sup> AUBERT 05S measures  $N(D^+ \rightarrow \phi\pi^+)/N(D_s^+ \rightarrow K^+K^-\pi^+)$ , the ratio of the numbers of events observed, and similarly for the  $D^-$ .

<sup>3</sup> FRABETTI 94I and AITALA 97B measure  $N(D^+ \rightarrow \phi\pi^+)/N(D^+ \rightarrow K^-\pi^+\pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

### $A_{CP}(K^\pm K_0^*(1430)^0)$ in $D^+ \rightarrow K^+\bar{K}_0^*(1430)^0$ , $D^- \rightarrow K^-K_0^*(1430)^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b><math>+8 \pm 6 \begin{smallmatrix} +4 \\ -2 \end{smallmatrix}</math></b>	RUBIN	08 CLEO	Fit-fraction asymmetry

**$A_{CP}(K^\pm K_2^*(1430)^0)$  in  $D^+ \rightarrow K^+ \bar{K}_2^*(1430)^0$ ,  $D^- \rightarrow K^- K_2^*(1430)^0$**

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$+43 \pm 19^{+5}_{-18}$	RUBIN	08	CLEO Fit-fraction asymmetry

**$A_{CP}(K^\pm K_0^*(800))$  in  $D^+ \rightarrow K^+ \bar{K}_0^*(800)$ ,  $D^- \rightarrow K^- K_0^*(800)$**

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-12 \pm 11^{+14}_{-6}$	RUBIN	08	CLEO Fit-fraction asymmetry

**$A_{CP}(a_0(1450)^0 \pi^\pm)$  in  $D^\pm \rightarrow a_0(1450)^0 \pi^\pm$**

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-19 \pm 12^{+8}_{-11}$	RUBIN	08	CLEO Fit-fraction asymmetry

**$A_{CP}(\phi(1680) \pi^\pm)$  in  $D^\pm \rightarrow \phi(1680) \pi^\pm$**

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-9 \pm 22 \pm 14$	RUBIN	08	CLEO Fit-fraction asymmetry

**$A_{CP}(\pi^+ \pi^- \pi^\pm)$  in  $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm$**

See also AAIJ 14C for a search for  $CP$  violation in  $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm$  Dalitz plots using model-independent binned and unbinned methods. No evidence was found.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-1.7 \pm 4.2$	<sup>1</sup> AITALA	97B E791	$-0.086 < A_{CP} < +0.052$ (90% CL)

<sup>1</sup> AITALA 97B measure  $N(D^+ \rightarrow \pi^+ \pi^- \pi^+) / N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

**$A_{CP}(K_S^0 K^\pm \pi^+ \pi^-)$  in  $D^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$-4.2 \pm 6.4 \pm 2.2$	$523 \pm 32$	LINK	05E FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV

**$A_{CP}(K^\pm \pi^0)$  in  $D^\pm \rightarrow K^\pm \pi^0$**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$-3.5 \pm 10.7 \pm 0.9$	$343 \pm 37$	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV

**$D^+-D^-$  T-VIOLATING DECAY-RATE ASYMMETRIES**

**$A_{Tviol}(K_S^0 K^\pm \pi^+ \pi^-)$  in  $D^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$**

$C_T \equiv \vec{p}_{K^+} \cdot (\vec{p}_{\pi^+} \times \vec{p}_{\pi^-})$  is a parity-odd correlation of the  $K^+$ ,  $\pi^+$ , and  $\pi^-$  momenta for the  $D^+$ .  $\bar{C}_T \equiv \vec{p}_{K^-} \cdot (\vec{p}_{\pi^-} \times \vec{p}_{\pi^+})$  is the corresponding quantity for the  $D^-$ . Then

$A_T \equiv [\Gamma(C_T > 0) - \Gamma(C_T < 0)] / [\Gamma(C_T > 0) + \Gamma(C_T < 0)]$ , and

$\bar{A}_T \equiv [\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)] / [\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)]$ , and

$A_{Tviol} \equiv \frac{1}{2}(A_T - \bar{A}_T)$ .  $C_T$  and  $\bar{C}_T$  are commonly referred to as  $T$ -odd moments, because they are odd under  $T$  reversal. However, the  $T$ -conjugate process  $K_S^0 K^\pm \pi^+ \pi^- \rightarrow D^\pm$  is not accessible, while the  $P$ -conjugate process is.

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$-12.0 \pm 10.0 \pm 4.6$	$21.2 \pm 0.4k$	LEES	11E BABR	$e^+ e^- \approx \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

**$D^+ \rightarrow (\bar{K}^0/\pi^0/\eta/\rho^0/\bar{K}^{*0})\ell^+\nu_\ell$  FORM FACTORS** **$f_+(0)|V_{cs}|$  in  $D^+ \rightarrow \bar{K}^0\ell^+\nu_\ell$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.707 \pm 0.010 \pm 0.009</math></b>	BESSION 09	CLEO	$\bar{K}^0 e^+ \nu_e$ 3-parameter fit

 **$r_1 \equiv a_1/a_0$  in  $D^+ \rightarrow \bar{K}^0\ell^+\nu_\ell$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>-1.66 \pm 0.44 \pm 0.10</math></b>	BESSION 09	CLEO	$\bar{K}^0 e^+ \nu_e$ 3-parameter fit

 **$r_2 \equiv a_2/a_0$  in  $D^+ \rightarrow \bar{K}^0\ell^+\nu_\ell$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>-14 \pm 11 \pm 1</math></b>	BESSION 09	CLEO	$\bar{K}^0 e^+ \nu_e$ 3-parameter fit

 **$f_+(0)|V_{cd}|$  in  $D^+ \rightarrow \pi^0\ell^+\nu_\ell$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.146 \pm 0.007 \pm 0.002</math></b>	BESSION 09	CLEO	$\pi^0 e^+ \nu_e$ 3-parameter fit

 **$r_1 \equiv a_1/a_0$  in  $D^+ \rightarrow \pi^0\ell^+\nu_\ell$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>-1.37 \pm 0.88 \pm 0.24</math></b>	BESSION 09	CLEO	$\pi^0 e^+ \nu_e$ 3-parameter fit

 **$r_2 \equiv a_2/a_0$  in  $D^+ \rightarrow \pi^0\ell^+\nu_\ell$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>-4 \pm 5 \pm 1</math></b>	BESSION 09	CLEO	$\pi^0 e^+ \nu_e$ 3-parameter fit

 **$f_+(0)|V_{cd}|$  in  $D^+ \rightarrow \eta e^+\nu_e$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.086 \pm 0.006 \pm 0.001</math></b>	YELTON 11	CLEO	z expansion

 **$r_1 \equiv a_1/a_0$  in  $D^+ \rightarrow \eta e^+\nu_e$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>-1.83 \pm 2.23 \pm 0.28</math></b>	YELTON 11	CLEO	z expansion

 **$r_\nu \equiv V(0)/A_1(0)$  in  $D^+, D^0 \rightarrow \rho e^+\nu_e$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>1.48 \pm 0.15 \pm 0.05</math></b>	<sup>1</sup> DOBBS 13	CLEO	$e^+ e^-$ at $\psi(3770)$

<sup>1</sup> Uses both  $D^+$  and  $D^0$  events. Using PDG 10 values of  $V_{cd}$  and lifetimes, DOBBS 13 gets  $A_1(0) = 0.56 \pm 0.01^{+0.02}_{-0.03}$ ,  $A_2(0) = 0.47 \pm 0.06 \pm 0.04$ , and  $V(0) = 0.84 \pm 0.09^{+0.05}_{-0.06}$ .

$r_2 \equiv A_2(0)/A_1(0)$  in  $D^+, D^0 \rightarrow \rho e^+ \nu_e$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.83±0.11±0.04</b>	<sup>1</sup> DOBBS	13	CLEO $e^+ e^-$ at $\psi(3770)$

<sup>1</sup> Uses both  $D^+$  and  $D^0$  events. Using PDG 10 values of  $V_{cd}$  and lifetimes, DOBBS 13 gets  $A_1(0) = 0.56 \pm 0.01^{+0.02}_{-0.03}$ ,  $A_2(0) = 0.47 \pm 0.06 \pm 0.04$ , and  $V(0) = 0.84 \pm 0.09^{+0.05}_{-0.06}$ .

$r_V \equiv V(0)/A_1(0)$  in  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.51 ± 0.07 OUR AVERAGE</b>				Error includes scale factor of 2.2. See the ideogram below.

1.463±0.017±0.031		<sup>1</sup> DEL-AMO-SA...11I	BABR	
1.504±0.057±0.039	15k	<sup>2</sup> LINK	02L	FOCS $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.45 ± 0.23 ± 0.07	763	ADAMOVICH	99	BEAT $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.90 ± 0.11 ± 0.09	3000	<sup>3</sup> AITALA	98B	E791 $\bar{K}^*(892)^0 e^+ \nu_e$
1.84 ± 0.11 ± 0.09	3034	AITALA	98F	E791 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.74 ± 0.27 ± 0.28	874	FRABETTI	93E	E687 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
2.00 <sup>+0.34</sup> <sub>-0.32</sub> ± 0.16	305	KODAMA	92	E653 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$

• • • We do not use the following data for averages, fits, limits, etc. • • •


2.0 ± 0.6 ± 0.3	183	ANJOS	90E	E691 $\bar{K}^*(892)^0 e^+ \nu_e$
-----------------	-----	-------	-----	-----------------------------------

<sup>1</sup> DEL-AMO-SANCHEZ 11I finds the pole mass  $m_A = (2.63 \pm 0.10 \pm 0.13)$  GeV ( $m_V$  is fixed at 2 GeV).

<sup>2</sup> LINK 02L includes the effects of interference with an  $S$ -wave background. This much improves the goodness of fit, but does not much shift the values of the form factors.

<sup>3</sup> This is slightly different from the AITALA 98B value: see ref. [5] in AITALA 98F.

WEIGHTED AVERAGE  
1.51±0.07 (Error scaled by 2.2)



$r_V \equiv V(0)/A_1(0)$  in  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

### $r_2 \equiv A_2(0)/A_1(0)$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.807±0.025 OUR AVERAGE</b>				
0.801±0.020±0.020		<sup>1</sup> DEL-AMO-SA..11i	BABR	
0.875±0.049±0.064	15k	<sup>2</sup> LINK	02L	FOCS $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.00 ±0.15 ±0.03	763	ADAMOVICH	99	BEAT $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.71 ±0.08 ±0.09	3000	AITALA	98B	E791 $\bar{K}^*(892)^0 e^+ \nu_e$
0.75 ±0.08 ±0.09	3034	AITALA	98F	E791 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.78 ±0.18 ±0.10	874	FRABETTI	93E	E687 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.82 <sup>+0.22</sup> <sub>-0.23</sub> ±0.11	305	KODAMA	92	E653 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0 ±0.5 ±0.2	183	ANJOS	90E	E691 $\bar{K}^*(892)^0 e^+ \nu_e$
---------------	-----	-------	-----	-----------------------------------

<sup>1</sup> DEL-AMO-SANCHEZ 11i finds the pole mass  $m_A = (2.63 \pm 0.10 \pm 0.13)$  GeV ( $m_V$  is fixed at 2 GeV).

<sup>2</sup> LINK 02L includes the effects of interference with an  $S$ -wave background. This much improves the goodness of fit, but does not much shift the values of the form factors.

### $r_3 \equiv A_3(0)/A_1(0)$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.04±0.33±0.29</b>				
	3034	AITALA	98F	E791 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$

### $\Gamma_L/\Gamma_T$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.13±0.08 OUR AVERAGE</b>				
1.09±0.10±0.02	763	ADAMOVICH	99	BEAT $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.20±0.13±0.13	874	FRABETTI	93E	E687 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.18±0.18±0.08	305	KODAMA	92	E653 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 <sup>+0.6</sup> <sub>-0.4</sub> ±0.3	183	ANJOS	90E	E691 $\bar{K}^*(892)^0 e^+ \nu_e$
--	-----	-------	-----	-----------------------------------

### $\Gamma_+/\Gamma_-$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.22±0.06 OUR AVERAGE</b> Error includes scale factor of 1.6.				
0.28±0.05±0.02	763	ADAMOVICH	99	BEAT $\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.16±0.05±0.02	305	KODAMA	92	E653 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.15 <sup>+0.07</sup> <sub>-0.05</sub> ±0.03	183	ANJOS	90E	E691 $\bar{K}^*(892)^0 e^+ \nu_e$
--	-----	-------	-----	-----------------------------------

**$D^\pm$  REFERENCES**

AAIJ	14C	PL B728 585	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AF	PL B724 203	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13W	JHEP 1306 112	R. Aaij <i>et al.</i>	(LHCb Collab.)
DOBBS	13	PRL 110 131802	S. Dobbs <i>et al.</i>	(CLEO Collab.)
KO	13	JHEP 1302 098	B.R. Ko <i>et al.</i>	(BELLE Collab.)
LEES	13E	PR D87 052012	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
KO	12A	PRL 109 119903 (errata)	B.R. Ko <i>et al.</i>	(BELLE Collab.)
Also		PRL 109 021601	B.R. Ko <i>et al.</i>	(BELLE Collab.)
STARIC	12	PRL 108 071801	M. Staric <i>et al.</i>	(BELLE Collab.)
AAIJ	11G	PR D84 112008	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA...	11H	PR D83 071103	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
DEL-AMO-SA...	11I	PR D83 072001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
LEES	11E	PR D84 031103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11G	PR D84 072006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WON	11	PRL 107 221801	E. Won <i>et al.</i>	(BELLE Collab.)
YELTON	11	PR D84 032001	J. Yelton <i>et al.</i>	(CLEO Collab.)
ANASHIN	10A	PL B686 84	V.V. Anashin <i>et al.</i>	(VEPP-4M KEDR Collab.)
ASNER	10	PR D81 052007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	10	PR D81 112001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
KO	10	PRL 104 181602	B.R. Ko <i>et al.</i>	(BELLE Collab.)
MENDEZ	10	PR D81 052013	H. Mendez <i>et al.</i>	(CLEO Collab.)
PDG	10	JP G37 075021	K. Nakamura <i>et al.</i>	(PDG Collab.)
RUBIN	10	PR D82 092007	P. Rubin <i>et al.</i>	(CLEO Collab.)
BESSION	09	PR D80 032005	D. Besson <i>et al.</i>	(CLEO Collab.)
Also		PR D79 052010	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
KO	09	PRL 102 221802	B.R. Ko <i>et al.</i>	(BELLE Collab.)
LINK	09	PL B681 14	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
MITCHELL	09B	PRL 102 081801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
WON	09	PR D80 111101	E. Won <i>et al.</i>	(BELLE Collab.)
ABAZOV	08D	PRL 100 101801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABLIKIM	08L	PL B665 16	M. Ablikim <i>et al.</i>	(BES Collab.)
ARTUSO	08	PR D77 092003	M. Artuso <i>et al.</i>	(CLEO Collab.)
BONVICINI	08	PR D77 091106	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
DOBBS	08	PR D77 112005	S. Dobbs <i>et al.</i>	(CLEO Collab.)
Also		PRL 100 251802	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
EISENSTEIN	08	PR D78 052003	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)
HE	08	PRL 100 091801	Q. He <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
RUBIN	08	PR D78 072003	P. Rubin <i>et al.</i>	(CLEO Collab.)
ABLIKIM	07	PL B644 20	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07G	PL B658 1	M. Ablikim <i>et al.</i>	(BES Collab.)
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
DOBBS	07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)
LINK	07B	PL B653 1	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABLIKIM	06O	EPJ C47 31	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06P	EPJ C47 39	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06U	PL B643 246	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	06A	PRL 97 251801	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
Also		PR D74 059901 (errata)	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AUBERT,B	06F	PR D74 011107	B. Aubert <i>et al.</i>	(BABAR Collab.)
DYTMAN	06	PR D74 071102	S.A. Dytman <i>et al.</i>	(CLEO Collab.)
HUANG	06B	PR D74 112005	G.S. Huang <i>et al.</i>	(CLEO Collab.)
LINK	06B	PL B637 32	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
RUBIN	06	PRL 96 081802	P. Rubin <i>et al.</i>	(CLEO Collab.)
RUBIN	06A	PR D73 112005	P. Rubin <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05A	PL B608 24	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05D	PL B610 183	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05F	PL B622 6	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05P	PL B625 196	M. Ablikim <i>et al.</i>	(BES Collab.)
ARTUSO	05A	PRL 95 251801	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT	05S	PR D71 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	05	PRL 95 121801	Q. He <i>et al.</i>	(CLEO Collab.)
Also		PRL 96 199903 (errata)	Q. He <i>et al.</i>	(CLEO Collab.)
HE	05A	PRL 95 221802	Q. He <i>et al.</i>	(CLEO Collab.)
HUANG	05B	PRL 95 181801	G.S. Huang <i>et al.</i>	(CLEO Collab.)
KAYIS-TOPAK...	05	PL B626 24	A. Kayis-Topaksu <i>et al.</i>	(CERN CHORUS Collab.)

LINK	05E	PL B622 239	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05I	PL B621 72	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABLIKIM	04C	PL B597 39	M. Ablikim <i>et al.</i>	(BEPC BES Collab.)
ARMS	04	PR D69 071102	K. Arms <i>et al.</i>	(CLEO Collab.)
BONVICINI	04A	PR D70 112004	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LINK	04	PL B585 200	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04E	PL B598 33	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04F	PL B601 10	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
LINK	03D	PL B561 225	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	03F	PL B572 21	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BRANDENB...	02	PRL 89 222001	G. Brandenburg <i>et al.</i>	(CLEO Collab.)
KAYIS-TOPAK...	02	PL B549 48	A. Kayis-Topaksu <i>et al.</i>	(CERN CHORUS Collab.)
LINK	02B	PRL 88 041602	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
Also		PRL 88 159903 (errata)	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02E	PL B535 43	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02F	PL B537 192	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02I	PL B541 227	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02J	PL B541 243	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02L	PL B544 89	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
LINK	01C	PRL 87 162001	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABREU	00O	EPJ C12 209	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ASTIER	00D	PL B486 35	P. Astier <i>et al.</i>	(CERN NOMAD Collab.)
JUN	00	PRL 84 1857	S.Y. Jun <i>et al.</i>	(FNAL SELEX Collab.)
LINK	00B	PL B491 232	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
Also		PL B495 443 (errata)	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABBIENDI	99K	EPJ C8 573	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ADAMOVICH	99	EPJ C6 35	M. Adamovich <i>et al.</i>	(CERN BEATRICE Collab.)
AITALA	99G	PL B462 401	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BONVICINI	99	PRL 82 4586	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AITALA	98B	PRL 80 1393	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	98C	PL B421 405	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	98F	PL B440 435	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BAI	98B	PL B429 188	J.Z. Bai <i>et al.</i>	(BEPC BES Collab.)
AITALA	97	PL B397 325	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	97B	PL B403 377	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	97C	PL B404 187	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BISHAI	97	PRL 78 3261	M. Bishai <i>et al.</i>	(CLEO Collab.)
FRABETTI	97	PL B391 235	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	97B	PL B398 239	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	97C	PL B401 131	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AITALA	96	PRL 76 364	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
FRABETTI	95	PL B346 199	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	95B	PL B351 591	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	95E	PL B359 403	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	94I	ZPHY C64 375	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	94	PRL 72 2328	R. Balest <i>et al.</i>	(CLEO Collab.)
FRABETTI	94D	PL B323 459	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	94G	PL B331 217	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	94I	PR D50 R2953	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AKERIB	93	PRL 71 3070	D.S. Akerib <i>et al.</i>	(CLEO Collab.)
ANJOS	93	PR D48 56	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
FRABETTI	93E	PL B307 262	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT	92F	PL B278 202	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	92C	PR D46 1941	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
BARLAG	92C	ZPHY C55 383	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
Also		ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
COFFMAN	92B	PR D45 2196	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
DAOUDI	92	PR D45 3965	M. Daoudi <i>et al.</i>	(CLEO Collab.)
KODAMA	92	PL B274 246	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
KODAMA	92C	PL B286 187	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ADAMOVICH	91	PL B268 142	M.I. Adamovich <i>et al.</i>	(WA82 Collab.)
ALBRECHT	91	PL B255 634	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALVAREZ	91B	ZPHY C50 11	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
AMMAR	91	PR D44 3383	R. Ammar <i>et al.</i>	(CLEO Collab.)
BAI	91	PRL 66 1011	Z. Bai <i>et al.</i>	(Mark III Collab.)
COFFMAN	91	PL B263 135	D.M. Coffman <i>et al.</i>	(Mark III Collab.)

FRABETTI	91	PL B263 584	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALVAREZ	90	ZPHY C47 539	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ANJOS	90C	PR D41 2705	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	90D	PR D42 2414	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	90E	PRL 65 2630	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
BARLAG	90C	ZPHY C46 563	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
WEIR	90B	PR D41 1384	A.J. Weir <i>et al.</i>	(Mark II Collab.)
ANJOS	89	PRL 62 125	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	89B	PRL 62 722	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	89E	PL B223 267	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ADLER	88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)
ALBRECHT	88I	PL B210 267	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
HAAS	88	PRL 60 1614	P. Haas <i>et al.</i>	(CLEO Collab.)
ONG	88	PRL 60 2587	R.A. Ong <i>et al.</i>	(Mark II Collab.)
RAAB	88	PR D37 2391	J.R. Raab <i>et al.</i>	(FNAL E691 Collab.)
ADAMOVICH	87	EPL 4 887	M.I. Adamovich <i>et al.</i>	(Photon Emulsion Collab.)
ADLER	87	PL B196 107	J. Adler <i>et al.</i>	(Mark III Collab.)
BARTEL	87	ZPHY C33 339	W. Bartel <i>et al.</i>	(JADE Collab.)
BALTRUSAIT...	86E	PRL 56 2140	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	85B	PRL 54 1976	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	85E	PRL 55 150	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BARTEL	85J	PL 163B 277	W. Bartel <i>et al.</i>	(JADE Collab.)
ADAMOVICH	84	PL 140B 119	M.I. Adamovich <i>et al.</i>	(CERN WA58 Collab.)
ALTHOFF	84G	ZPHY C22 219	M. Althoff <i>et al.</i>	(TASSO Collab.)
DERRICK	84	PRL 53 1971	M. Derrick <i>et al.</i>	(HRS Collab.)
SCHINDLER	81	PR D24 78	R.H. Schindler <i>et al.</i>	(Mark II Collab.)
TRILLING	81	PRPL 75 57	G.H. Trilling	(LBL, UCB) J
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34	1471.	
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)
PERUZZI	77	PRL 39 1301	I. Peruzzi <i>et al.</i>	(LGW Collab.)
PICCOLO	77	PL 70B 260	M. Piccolo <i>et al.</i>	(Mark I Collab.)
PERUZZI	76	PRL 37 569	I. Peruzzi <i>et al.</i>	(Mark I Collab.)

### OTHER RELATED PAPERS

RICHMAN	95	RMP 67 893	J.D. Richman, P.R. Burchat	(UCSB, STAN)
ROSNER	95	CNPP 21 369	J. Rosner	(CHIC)