$N(1650)\ 1/2^-$

$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$ Status: ****

Most of the results published before 1975 were last included in our 1982 edition, Physics Letters **111B** 1 (1982). Some further obsolete results published before 1984 were last included in our 2006 edition, Journal of Physics (generic for all A,B,E,G) **G33** 1 (2006).

N(1650) BREIT-WIGNER MASS

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
1645 to 1670 (≈ 1655) OUR ESTIMATE				
1665 ± 2	SHKLYAR	13	DPWA	Multichannel
1651 ± 6	ANISOVICH	12A	DPWA	Multichannel
1634.7± 1.1	ARNDT	06	DPWA	$\pi N \to \pi N, \eta N$
1650 ±30	CUTKOSKY	80	IPWA	$\pi N \to \pi N$
1670 ± 8	HOEHLER	79	IPWA	$\pi N \to \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1664 ± 2	SHRESTHA	12A	DPWA	Multichannel
1680 ±40	ANISOVICH	10	DPWA	Multichannel
1652 ± 9	BATINIC	10	DPWA	$\pi N \to N\pi, N\eta$
1655 ±15	THOMA	08	DPWA	Multichannel
1651.2± 4.7	ARNDT	04	DPWA	$\pi N \to \pi N, \eta N$
1665 ± 2	PENNER	02C	DPWA	Multichannel
1647 ±20	BAI	01B	BES	$J/\psi \to p\overline{p}\eta$
1689 ±12	VRANA	00	DPWA	Multichannel
1677 ± 8	ARNDT	96	IPWA	$\gamma N \to \pi N$
1667	ARNDT	95	DPWA	$\pi N \to N\pi$
1712	[1] ARNDT	95	DPWA	$\pi N \to N\pi$
1674	LI	93	IPWA	$\gamma N \to \pi N$
1659 ± 9	MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$
1672	MUSETTE	80	IPWA	$\pi^- p \to \Lambda K^0$
1680	SAXON	80	DPWA	$\pi^- p \to \Lambda K^0$
1700	[2] LONGACRE	77	IPWA	$\pi N \to N\pi\pi$
1660	[3] LONGACRE	75	IPWA	$\pi N \to N\pi\pi$

N(1650) BREIT-WIGNER WIDTH

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
110 to 170 (≈ 140) OUR ESTIMATE				
147 ±14	SHKLYAR	13	DPWA	Multichannel
104 ±10	ANISOVICH	12A	DPWA	Multichannel
115.4± 2.8	ARNDT	06	DPWA	$\pi N \to \pi N, \eta N$
167.9± 9.4	GREEN	97	DPWA	$\pi N \to \pi N, \eta N$
150 ±40	CUTKOSKY	80	IPWA	$\pi N \to \pi N$
180 ±20	HOEHLER	79	IPWA	$\pi N \to \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

126 ± 3		SHRESTHA	12A	DPWA	Multichannel
170 ±45		ANISOVICH	10	DPWA	Multichannel
202 ±16		BATINIC	10	DPWA	$\pi N \to N\pi$, $N\eta$
180 ±20		THOMA	08	DPWA	Multichannel
130.6± 7.0		ARNDT	04	DPWA	$\pi N \to \pi N$, ηN
138 ± 7		PENNER	02C	DPWA	Multichannel
145 $^{+80}_{-45}$		BAI	01B	BES	$J/\psi \to p\overline{p}\eta$
202 ±40		VRANA	00	DPWA	Multichannel
160 ±12		ARNDT	96	IPWA	$\gamma N \to \pi N$
90		ARNDT	95	DPWA	$\pi N \to N\pi$
184	[1]	ARNDT	95	DPWA	$\pi N \to N\pi$
225		LI	93	IPWA	$\gamma N \to \pi N$
173 ±12		MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$
179		MUSETTE	80	IPWA	$\pi^- p \to \Lambda K^0$
120		SAXON	80	DPWA	$\pi^- p \to \Lambda K^0$
170	[2]	LONGACRE	77	IPWA	$\pi N \to N\pi\pi$
130	[3]	LONGACRE	75	IPWA	$\pi N \to N\pi\pi$

N(1650) POLE POSITION

REAL PART

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
1640 to 1670 (≈ 1655) OUR ESTIMATE				
1647± 6		ANISOVICH	12A	DPWA Multichannel
1648		ARNDT	06	DPWA $\pi N \to \pi N$, ηN
1670	[4]	HOEHLER	93	ARGD $\pi N \to \pi N$
1640±20		CUTKOSKY	80	IPWA $\pi N \to \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1650		SHKLYAR	13	DPWA	Multichannel
1655		SHRESTHA	12A	DPWA	Multichannel
1670±35		ANISOVICH	10	DPWA	Multichannel
1646± 8		BATINIC	10	DPWA	$\pi N \to N\pi$, $N\eta$
1645±15		THOMA	08	DPWA	Multichannel
1653		ARNDT	04	DPWA	$\pi N \to \pi N$, ηN
1663		VRANA	00	DPWA	Multichannel
1660±10	[5]	ARNDT	98	DPWA	$\pi N \to \pi N$, ηN
1673		ARNDT	95	DPWA	$\pi N \to N\pi$
1689	[1]	ARNDT	95	DPWA	$\pi N \to N\pi$
1657		ARNDT	91	DPWA	$\pi N \to \pi N$ Soln SM90
1648 or 1651	[6]	LONGACRE	78	IPWA	$\pi N \to N\pi\pi$
1699 or 1698	[2]	LONGACRE	77	IPWA	$\pi N \to N\pi\pi$

−2×IMAGINARY PART

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
100 to 170 (≈ 135) OUR ESTIMATE				
103± 8		ANISOVICH	12A	DPWA Multichannel
80		ARNDT	06	DPWA $\pi N \to \pi N$, ηN
163	[4]	HOEHLER	93	ARGD $\pi N \to \pi N$
150±30		CUTKOSKY	80	IPWA $\pi N \to \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

89		SHKLYAR	13	DPWA	Multichannel
123		SHRESTHA	12A	DPWA	Multichannel
170 ± 40		ANISOVICH	10	DPWA	Multichannel
204 ± 17		BATINIC	10	DPWA	$\pi N \to N\pi$, $N\eta$
187 ± 20		THOMA	08	DPWA	Multichannel
182		ARNDT	04	DPWA	$\pi N \to \pi N$, ηN
240		VRANA	00	DPWA	Multichannel
140 ± 20	[5]	ARNDT	98	DPWA	$\pi N \to \pi N$, ηN
82		ARNDT	95	DPWA	$\pi N \to N\pi$
192	[1]	ARNDT	95	DPWA	$\pi N \to N\pi$
160		ARNDT	91	DPWA	$\pi N \to \pi N$ Soln SM90
117 or 119	[6]	LONGACRE	78	IPWA	$\pi N \to N\pi\pi$
174 or 173	[2]	LONGACRE	77	IPWA	$\pi N \to N\pi\pi$

N(1650) ELASTIC POLE RESIDUE

MODULUS $|r|$

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
20 to 50 (≈ 35) OUR ESTIMATE				
24 ± 3	ANISOVICH	12A	DPWA	Multichannel
14	ARNDT	06	DPWA	$\pi N \to \pi N$, ηN
39	HOEHLER	93	ARGD	$\pi N \to \pi N$
60 ± 10	CUTKOSKY	80	IPWA	$\pi N \to \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

19		SHKLYAR	13	DPWA	Multichannel
100		BATINIC	10	DPWA	$\pi N \to N\pi$, $N\eta$
69		ARNDT	04	DPWA	$\pi N \to \pi N$, ηN
22		ARNDT	95	DPWA	$\pi N \to N\pi$
72	[1]	ARNDT	95	DPWA	$\pi N \to N\pi$
54		ARNDT	91	DPWA	$\pi N \to \pi N$ Soln SM90

PHASE θ

VALUE (°)	*DOCUMENT ID*		*TECN*	*COMMENT*
50 to 80 (≈ 70) OUR ESTIMATE				
−75 ± 12	ANISOVICH	12A	DPWA	Multichannel
−69	ARNDT	06	DPWA	$\pi N \to \pi N$, ηN
−37	HOEHLER	93	ARGD	$\pi N \to \pi N$
−75 ± 25	CUTKOSKY	80	IPWA	$\pi N \to \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

−46		SHKLYAR	13	DPWA	Multichannel
−65		BATINIC	10	DPWA	$\pi N \to N\pi$, $N\eta$
−55		ARNDT	04	DPWA	$\pi N \to \pi N$, ηN
29		ARNDT	95	DPWA	$\pi N \to N\pi$
−85	[1]	ARNDT	95	DPWA	$\pi N \to N\pi$
−38		ARNDT	91	DPWA	$\pi N \to \pi N$ Soln SM90

N(1650) INELASTIC POLE RESIDUE

The "normalized residue" is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\pi \rightarrow N(1650) \rightarrow N\eta$

MODULUS (%)	*PHASE (°)*	*DOCUMENT ID*		*TECN*	*COMMENT*
29±3	**134 ± 10**	ANISOVICH	12A	DPWA	Multichannel

Normalized residue in $N\pi \rightarrow N(1650) \rightarrow \Lambda K$

MODULUS (%)	*PHASE (°)*	*DOCUMENT ID*		*TECN*	*COMMENT*
23±9	**85 ± 9**	ANISOVICH	12A	DPWA	Multichannel

Normalized residue in $N\pi \rightarrow N(1650) \rightarrow \Delta\pi$, *D*-wave

MODULUS (%)	*PHASE (°)*	*DOCUMENT ID*		*TECN*	*COMMENT*
23±4	**−30 ± 20**	ANISOVICH	12A	DPWA	Multichannel

N(1650) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

	Mode	Fraction (Γ_i/Γ)
Γ_1	$N\pi$	50–90 %
Γ_2	$N\eta$	5–15 %
Γ_3	ΛK	3–11 %
Γ_4	ΣK	
Γ_5	$N\pi\pi$	10–20 %
Γ_6	$\Delta\pi$	0–25 %
Γ_7	$\Delta(1232)\pi$, *D*-wave	0–25 %
Γ_8	$N\rho$	4–12 %
Γ_9	$N\rho$, S=1/2, *S*-wave	(1.0±1.0) %
Γ_{10}	$N\rho$, S=3/2, *D*-wave	(13.0±3.0) %
Γ_{11}	$N(\pi\pi)^{I=0}_{S\text{-wave}}$	<4 %
Γ_{12}	$N(1440)\pi$	<5 %
Γ_{13}	$p\gamma$	0.04–0.20 %
Γ_{14}	$p\gamma$, helicity=1/2	0.04–0.20 %
Γ_{15}	$n\gamma$	0.003–0.17 %
Γ_{16}	$n\gamma$, helicity=1/2	0.003–0.17 %

N(1650) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$ Γ_1/Γ

VALUE (%)	*DOCUMENT ID*		*TECN*	*COMMENT*
50 to 70 (≈ 60) OUR ESTIMATE				
74 ± 3	SHKLYAR	13	DPWA	Multichannel
51 ± 4	ANISOVICH	12A	DPWA	Multichannel
100	ARNDT	06	DPWA	$\pi N \rightarrow \pi N$, ηN

73.5± 1.1	GREEN	97	DPWA	$\pi N \to \pi N, \eta N$
65 ±10	CUTKOSKY	80	IPWA	$\pi N \to \pi N$
61 ± 4	HOEHLER	79	IPWA	$\pi N \to \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

57 ± 2	SHRESTHA	12A	DPWA	Multichannel
50 ±25	ANISOVICH	10	DPWA	Multichannel
79 ± 6	BATINIC	10	DPWA	$\pi N \to N\pi, N\eta$
70 ±15	THOMA	08	DPWA	Multichannel
100.0	ARNDT	04	DPWA	$\pi N \to \pi N, \eta N$
65 ± 4	PENNER	02C	DPWA	Multichannel
74 ± 2	VRANA	00	DPWA	Multichannel
99	ARNDT	95	DPWA	$\pi N \to N\pi$
27	[1] ARNDT	95	DPWA	$\pi N \to N\pi$
89 ± 7	MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$

$\Gamma(N\eta)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
5 to 15 OUR ESTIMATE				
1 ±2	SHKLYAR	13	DPWA	Multichannel
18 ±4	ANISOVICH	12A	DPWA	Multichannel
1.0±0.6	PENNER	02C	DPWA	Multichannel
6 ±1	VRANA	00	DPWA	Multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

21 ±2	SHRESTHA	12A	DPWA	Multichannel
13 ±5	BATINIC	10	DPWA	$\pi N \to N\pi, N\eta$
15 ±6	THOMA	08	DPWA	Multichannel

$\Gamma(\Lambda K)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
2.9±0.4 OUR AVERAGE Error includes scale factor of 1.2.				
10 ±5	ANISOVICH	12A	DPWA	Multichannel
4 ±1	SHKLYAR	05	DPWA	Multichannel
2.7±0.4	PENNER	02C	DPWA	Multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

8 ±1	SHRESTHA	12A	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \to N(1650) \to \Lambda K$ $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.27 to −0.17 OUR ESTIMATE				
−0.22	BELL	83	DPWA	$\pi^- p \to \Lambda K^0$
−0.22	SAXON	80	DPWA	$\pi^- p \to \Lambda K^0$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \to N(1650) \to \Sigma K$ $(\Gamma_1\Gamma_4)^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

−0.254	LIVANOS	80	DPWA	$\pi p \to \Sigma K$

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620)$ S_{31} coupling to $\Delta(1232)\pi$.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow N(1650) \rightarrow \Delta(1232)\pi$, D-wave $(\Gamma_1\Gamma_7)^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
+0.15 to 0.23 OUR ESTIMATE				
+0.29	[2,7] LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.15	[3] LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.26±0.14	THOMA	08	DPWA	Multichannel
+0.12±0.04	MANLEY	92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(\Delta(1232)\pi, D\text{-wave})/\Gamma_{total}$ Γ_7/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
0 to 25 OUR ESTIMATE				
19±9	ANISOVICH	12A	DPWA	Multichannel
2±1	VRANA	00	DPWA	Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7±2	SHRESTHA	12A	DPWA	Multichannel
10±5	THOMA	08	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow N(1650) \rightarrow N\rho$, S=1/2, S-wave $(\Gamma_1\Gamma_9)^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
±0.03 to ±0.19 OUR ESTIMATE				
+0.17	[2,7] LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
−0.16	[3] LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−0.01±0.09	MANLEY	92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(N\rho, S$=1/2, S-wave$)/\Gamma_{total}$ Γ_9/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
1±1	VRANA	00	DPWA	Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
6±1	SHRESTHA	12A	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow N(1650) \rightarrow N\rho$, S=3/2, D-wave $(\Gamma_1\Gamma_{10})^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
+0.17 to +0.29 OUR ESTIMATE				
+0.29	[2,7] LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.16±0.06	MANLEY	92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(N\rho, S$=3/2, D-wave$)/\Gamma_{total}$ Γ_{10}/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
13±3	VRANA	00	DPWA	Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1	SHRESTHA	12A	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \to N(1650) \to N(\pi\pi)^{I=0}_{S\text{-wave}}$ $(\Gamma_1\Gamma_{11})^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
+0.04 to +0.18 OUR ESTIMATE				
0.00	[2,7] LONGACRE	77	IPWA	$\pi N \to N\pi\pi$
+0.25	[3] LONGACRE	75	IPWA	$\pi N \to N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.12±0.08	MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$

$\Gamma(N(\pi\pi)^{I=0}_{S\text{-wave}})/\Gamma_{total}$ Γ_{11}/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
1±1	VRANA	00	DPWA	Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1	SHRESTHA	12A	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \to N(1650) \to N(1440)\pi$ $(\Gamma_1\Gamma_{12})^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.11±0.06	MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$

$\Gamma(N(1440)\pi)/\Gamma_{total}$ Γ_{12}/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
3±1	VRANA	00	DPWA	Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1	SHRESTHA	12A	DPWA	Multichannel

N(1650) PHOTON DECAY AMPLITUDES

Papers on γN amplitudes predating 1981 may be found in our 2006 edition, Journal of Physics (generic for all A,B,E,G) **G33** 1 (2006).

N(1650) → pγ, helicity-1/2 amplitude $A_{1/2}$

VALUE ($GeV^{-1/2}$)	DOCUMENT ID		TECN	COMMENT
+0.045±0.010 OUR ESTIMATE				
0.033±0.007	ANISOVICH	12A	DPWA	Multichannel
0.055±0.030	WORKMAN	12A	DPWA	$\gamma N \to N\pi$
0.022±0.007	DUGGER	07	DPWA	$\gamma N \to \pi N$
0.033±0.015	CRAWFORD	83	IPWA	$\gamma N \to \pi N$
0.050±0.010	AWAJI	81	DPWA	$\gamma N \to \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.063±0.006	SHKLYAR	13	DPWA	Multichannel
0.030±0.003	SHRESTHA	12A	DPWA	Multichannel
0.060±0.020	ANISOVICH	10	DPWA	Multichannel
0.100±0.035	[8] ANISOVICH	09A	DPWA	$\gamma d \to \eta N(N)$
0.033	DRECHSEL	07	DPWA	$\gamma N \to \pi N$
0.049	PENNER	02D	DPWA	Multichannel
0.069±0.005	ARNDT	96	IPWA	$\gamma N \to \pi N$
0.068±0.003	LI	93	IPWA	$\gamma N \to \pi N$
0.091	WADA	84	DPWA	Compton scattering

$N(1650) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($GeV^{-1/2}$)	DOCUMENT ID		TECN	COMMENT
-0.050 ± 0.020 OUR ESTIMATE				
-0.040 ± 0.010	CHEN	12A	DPWA	$\gamma N \rightarrow \pi N$
-0.055 ± 0.020	[9] ANISOVICH	09A	DPWA	$\gamma d \rightarrow \eta N(N)$
-0.008 ± 0.004	AWAJI	81	DPWA	$\gamma N \rightarrow \pi N$
0.004 ± 0.004	FUJII	81	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.011 ± 0.002	SHRESTHA	12A	DPWA	Multichannel
0.009	DRECHSEL	07	DPWA	$\gamma N \rightarrow \pi N$
-0.011	PENNER	02D	DPWA	Multichannel
-0.015 ± 0.005	ARNDT	96	IPWA	$\gamma N \rightarrow \pi N$
-0.002 ± 0.002	LI	93	IPWA	$\gamma N \rightarrow \pi N$

$N(1650)$ $\gamma p \rightarrow \Lambda K^+$ AMPLITUDES

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $p\gamma \rightarrow N(1650) \rightarrow \Lambda K^+$ (E_{0+} amplitude)

VALUE (units 10^{-3})	DOCUMENT ID		TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •			
7.8 ± 0.3	WORKMAN	90	DPWA
8.13	TANABE	89	DPWA

$p\gamma \rightarrow N(1650) \rightarrow \Lambda K^+$ phase angle θ (E_{0+} amplitude)

VALUE (degrees)	DOCUMENT ID		TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-107 ± 3	WORKMAN	90	DPWA
-107.8	TANABE	89	DPWA

$N(1650)$ FOOTNOTES

[1] ARNDT 95 finds two distinct states.

[2] LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

[3] From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

[4] See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

[5] ARNDT 98 also lists pole residues, which display more model dependence than do the associated pole positions.

[6] LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

[7] LONGACRE 77 considers this coupling to be well determined.

[8] This ANISOVICH 09A amplitude is evaluated at the pole position; the phase is $(25 \pm 20)^\circ$.

[9] This ANISOVICH 09A amplitude is evaluated at the pole position; the phase is $(30 \pm 25)^\circ$.

N(1650) REFERENCES

For early references, see Physics Letters **111B** 1 (1982).

SHKLYAR	13	PR C87 015201	V. Shklyar, H. Lenske, U. Mosel	(GIES)
ANISOVICH	12A	EPJ A48 15	A.V. Anisovich *et al.*	(BONN, PNPI)
CHEN	12A	PR C86 015206	W. Chen *et al.*	(DUKE, GWU, MSST, ITEP+)
SHRESTHA	12A	PR C86 055203	M. Shrestha, D.M. Manley	(KSU)
WORKMAN	12A	PR C86 015202	R. Workman *et al.*	(GWU)
ANISOVICH	10	EPJ A44 203	A.V. Anisovich *et al.*	(BONN, PNPI)
BATINIC	10	PR C82 038203	M. Batinic *et al.*	(ZAGR)
ANISOVICH	09A	EPJ A41 13	A.V. Anisovich *et al.*	(BONN, PNPI, BASL)
THOMA	08	PL B659 87	U. Thoma *et al.*	(CB-ELSA Collab.)
DRECHSEL	07	EPJ A34 69	D. Drechsel, S.S. Kamalov, L. Tiator	(MAINZ, JINR)
DUGGER	07	PR C76 025211	M. Dugger *et al.*	(Jefferson Lab CLAS Collab.)
ARNDT	06	PR C74 045205	R.A. Arndt *et al.*	(GWU)
PDG	06	JP G33 1	W.-M. Yao *et al.*	(PDG Collab.)
SHKLYAR	05	PR C72 015210	V. Shklyar, H. Lenske, U. Mosel	(GIES)
ARNDT	04	PR C69 035213	R.A. Arndt *et al.*	(GWU, TRIU)
PENNER	02C	PR C66 055211	G. Penner, U. Mosel	(GIES)
PENNER	02D	PR C66 055212	G. Penner, U. Mosel	(GIES)
BAI	01B	PL B510 75	J.Z. Bai *et al.*	(BES Collab.)
VRANA	00	PRPL 328 181	T.P. Vrana, S.A. Dytman,, T.-S.H. Lee	(PITT+)
ARNDT	98	PR C58 3636	R.A. Arndt *et al.*	
GREEN	97	PR C55 R2167	A.M. Green, S. Wycech	(HELS, WINR)
ARNDT	96	PR C53 430	R.A. Arndt, I.I. Strakovsky, R.L. Workman	(VPI)
ARNDT	95	PR C52 2120	R.A. Arndt *et al.*	(VPI, BRCO)
HOEHLER	93	πN Newsletter 9 1	G. Hohler	(KARL)
LI	93	PR C47 2759	Z.J. Li *et al.*	(VPI)
MANLEY	92	PR D45 4002	D.M. Manley, E.M. Saleski	(KSA) IJP
Also		PR D30 904	D.M. Manley *et al.*	(VPI)
ARNDT	91	PR D43 2131	R.A. Arndt *et al.*	(VPI, TELE) IJP
WORKMAN	90	PR C42 781	R.L. Workman	(VPI)
TANABE	89	PR C39 741	H. Tanabe, M. Kohno, C. Bennhold	(MANZ)
Also		NC 102A 193	M. Kohno, H. Tanabe, C. Bennhold	(MANZ)
WADA	84	NP B247 313	Y. Wada *et al.*	(INUS)
BELL	83	NP B222 389	K.W. Bell *et al.*	(RL) IJP
CRAWFORD	83	NP B211 1	R.L. Crawford, W.T. Morton	(GLAS)
PDG	82	PL 111B 1	M. Roos *et al.*	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	N. Awaji, R. Kajikawa	(NAGO)
Also		NP B197 365	K. Fujii *et al.*	(NAGO)
FUJII	81	NP B187 53	K. Fujii *et al.*	(NAGO, OSAK)
CUTKOSKY	80	Toronto Conf. 19	R.E. Cutkosky *et al.*	(CMU, LBL) IJP
Also		PR D20 2839	R.E. Cutkosky *et al.*	(CMU, LBL) IJP
LIVANOS	80	Toronto Conf. 35	P. Livanos *et al.*	(SACL) IJP
MUSETTE	80	NC 57A 37	M. Musette	(BRUX) IJP
SAXON	80	NP B162 522	D.H. Saxon *et al.*	(RHEL, BRIS) IJP
HOEHLER	79	PDAT 12-1	G. Hohler *et al.*	(KARLT) IJP
Also		Toronto Conf. 3	R. Koch	(KARLT) IJP
LONGACRE	78	PR D17 1795	R.S. Longacre *et al.*	(LBL, SLAC)
LONGACRE	77	NP B122 493	R.S. Longacre, J. Dolbeau	(SACL) IJP
Also		NP B108 365	J. Dolbeau *et al.*	(SACL) IJP
LONGACRE	75	PL 55B 415	R.S. Longacre *et al.*	(LBL, SLAC) IJP