

**$N(2120) 3/2^-$** 

$$I(J^P) = \frac{1}{2}(3/2^-) \text{ Status: } **$$

## OMITTED FROM SUMMARY TABLE

Before the 2012 *Review*, all the evidence for a  $J^P = 3/2^-$  state with a mass above 1800 MeV was filed under a two-star  $N(2080)$ .

There is now evidence from ANISOVICH 12A for two  $3/2^-$  states in this region, so we have split the older data (according to mass) between a three-star  $N(1875)$  and a two-star  $N(2120)$ .

 **$N(2120)$  BREIT-WIGNER MASS**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2120 OUR ESTIMATE</b>			
2150 ± 60	ANISOVICH	12A	DPWA Multichannel
2060 ± 80	<sup>1</sup> CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
2081 ± 20	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$

 **$N(2120)$  BREIT-WIGNER WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
330 ± 45	ANISOVICH	12A	DPWA Multichannel
300 ± 100	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$ (higher $m$ )
265 ± 40	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$

 **$N(2120)$  POLE POSITION****REAL PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2110 ± 50	ANISOVICH	12A	DPWA Multichannel
2050 ± 70	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$ (higher $m$ )

**−2×IMAGINARY PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
340 ± 45	ANISOVICH	12A	DPWA Multichannel
200 ± 80	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$ (higher $m$ )

 **$N(2120)$  ELASTIC POLE RESIDUE****MODULUS  $|r|$** 

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
13 ± 3	ANISOVICH	12A	DPWA Multichannel
30 ± 20	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$ (higher $m$ )

**PHASE  $\theta$** 

<u>VALUE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
−20 ± 10	ANISOVICH	12A	DPWA Multichannel
0 ± 100	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$ (higher $m$ )

**$N(2120)$  INELASTIC POLE RESIDUE**The “normalized residue” is the residue divided by  $\Gamma_{pole}/2$ .**Normalized residue in  $N\pi \rightarrow N(2120) \rightarrow \Lambda K$** 

<u>MODULUS (%)</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3±1	100 ± 30	ANISOVICH	12A DPWA	Multichannel

**Normalized residue in  $N\pi \rightarrow N(2120) \rightarrow \Sigma K$** 

<u>MODULUS (%)</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2±1.5	-50 ± 40	ANISOVICH	12A DPWA	Multichannel

 **$N(2120)$  DECAY MODES**

Mode
$\Gamma_1$ $N\pi$

 **$N(2120)$  BRANCHING RATIOS**

<u><math>\Gamma(N\pi)/\Gamma_{total}</math></u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u><math>\Gamma_1/\Gamma</math></u>
6±2	ANISOVICH	12A DPWA	Multichannel	
14±7	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$ (higher $m$ )	
6±2	HOEHLER	79 IPWA	$\pi N \rightarrow \pi N$	

 **$N(2120)$  PHOTON DECAY AMPLITUDES** **$N(2120) \rightarrow p\gamma$ , helicity-1/2 amplitude  $A_{1/2}$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.125±0.045	<sup>2</sup> ANISOVICH	12A DPWA	Phase = (-55 ± 20)°

 **$N(2120) \rightarrow p\gamma$ , helicity-3/2 amplitude  $A_{3/2}$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.150±0.060	<sup>2</sup> ANISOVICH	12A DPWA	Phase = (-35 ± 15)°

 **$N(2120)$  FOOTNOTES**<sup>1</sup> CUTKOSKY 80 finds a lower mass  $D_{13}$  resonance, as well as one in this region. Both are listed here.<sup>2</sup> This ANISOVICH 12A value is the complex helicity amplitude at the pole position. **$N(2120)$  REFERENCES**

ANISOVICH	12A	EPJ A48 15	A.V. Anisovich <i>et al.</i>	(BONN, PNPI)
CUTKOSKY	80	Toronto Conf. 19	R.E. Cutkosky <i>et al.</i>	(CMU, LBL)
HOEHLER	79	PDAT 12-1	G. Hohler <i>et al.</i>	(KARLT)