

$\eta'(958)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

$\eta'(958)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
957.78 ± 0.06 OUR AVERAGE				
957.793 ± 0.054 ± 0.036	3.9k	LIBBY	08	CLEO $J/\psi \rightarrow \gamma\eta'$
957.9 ± 0.2 ± 0.6	4800	WURZINGER	96	SPEC 1.68 $pd \rightarrow {}^3\text{He}\eta'$
957.46 ± 0.33		DUANE	74	MMS $\pi^- p \rightarrow n\text{MM}$
958.2 ± 0.5	1414	DANBURG	73	HBC 2.2 $K^- p \rightarrow \Lambda\eta'$
958 ± 1	400	JACOBS	73	HBC 2.9 $K^- p \rightarrow \Lambda\eta'$
956.1 ± 1.1	3415	¹ BASILE	71	CNTR 1.6 $\pi^- p \rightarrow n\eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
957.5 ± 0.2		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
959 ± 1	630	² BELADIDZE	92C	VES 36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
958 ± 1	340	² ARMSTRONG	91B	OMEG 300 $pp \rightarrow pp\eta\pi^+\pi^-$
958.2 ± 0.4	622	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
957.8 ± 0.2	2420	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
956.3 ± 1.0	143	² GIDAL	87	MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
957.4 ± 1.4	535	³ BASILE	71	CNTR 1.6 $\pi^- p \rightarrow n\eta'$
957 ± 1		RITTENBERG	69	HBC 1.7-2.7 $K^- p$

¹ Using all η' decays.² Systematic uncertainty not estimated.³ Using η' decays into neutrals. Not independent of the other listed BASILE 71 η' mass measurement.

$\eta'(958)$ WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
0.198 ± 0.009 OUR FIT					
0.230 ± 0.021 OUR AVERAGE					
0.226 ± 0.017 ± 0.014	2300	CZERWINSKI	10	MMS	$pp \rightarrow pp\eta'$
0.40 ± 0.22	4800	WURZINGER	96	SPEC	1.68 $pd \rightarrow {}^3\text{He}\eta'$
0.28 ± 0.10	1000	BINNIE	79	MMS	0 $\pi^- p \rightarrow n\text{MM}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.20 ± 0.04		BAI	04J	BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

$\eta'(958)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\pi^+\pi^-\eta$	(42.9 \pm 0.7) %	
Γ_2 $\rho^0\gamma$ (including non-resonant $\pi^+\pi^-\gamma$)	(29.1 \pm 0.5) %	
Γ_3 $\pi^0\pi^0\eta$	(22.2 \pm 0.8) %	
Γ_4 $\omega\gamma$	(2.75 \pm 0.23) %	
Γ_5 $\gamma\gamma$	(2.20 \pm 0.08) %	
Γ_6 $3\pi^0$	(2.14 \pm 0.20) $\times 10^{-3}$	
Γ_7 $\mu^+\mu^-\gamma$	(1.08 \pm 0.27) $\times 10^{-4}$	
Γ_8 $\pi^+\pi^-\mu^+\mu^-$	< 2.9 $\times 10^{-5}$	90%
Γ_9 $\pi^+\pi^-\pi^0$	(3.8 \pm 0.4) $\times 10^{-3}$	
Γ_{10} $\pi^0\rho^0$	< 4 %	90%
Γ_{11} $2(\pi^+\pi^-)$	< 2.4 $\times 10^{-4}$	90%
Γ_{12} $\pi^+\pi^-2\pi^0$	< 2.5 $\times 10^{-3}$	90%
Γ_{13} $2(\pi^+\pi^-)$ neutrals	< 1 %	95%
Γ_{14} $2(\pi^+\pi^-)\pi^0$	< 1.9 $\times 10^{-3}$	90%
Γ_{15} $2(\pi^+\pi^-)2\pi^0$	< 1 %	95%
Γ_{16} $3(\pi^+\pi^-)$	< 3.1 $\times 10^{-5}$	90%
Γ_{17} $\pi^+\pi^-e^+e^-$	(2.4 $^{+1.3}_{-1.0}$) $\times 10^{-3}$	
Γ_{18} $\pi^+e^-\nu_e + \text{c.c.}$	< 2.1 $\times 10^{-4}$	90%
Γ_{19} γe^+e^-	< 9 $\times 10^{-4}$	90%
Γ_{20} $\pi^0\gamma\gamma$	< 8 $\times 10^{-4}$	90%
Γ_{21} $4\pi^0$	< 5 $\times 10^{-4}$	90%
Γ_{22} e^+e^-	< 2.1 $\times 10^{-7}$	90%
Γ_{23} invisible	< 5 $\times 10^{-4}$	90%

Charge conjugation (C), Parity (P), Lepton family number (LF) violating modes

Γ_{24} $\pi^+\pi^-$	P, CP	< 6 $\times 10^{-5}$	90%
Γ_{25} $\pi^0\pi^0$	P, CP	< 4 $\times 10^{-4}$	90%
Γ_{26} $\pi^0e^+e^-$	C [a]	< 1.4 $\times 10^{-3}$	90%
Γ_{27} ηe^+e^-	C [a]	< 2.4 $\times 10^{-3}$	90%
Γ_{28} 3γ	C	< 1.0 $\times 10^{-4}$	90%
Γ_{29} $\mu^+\mu^-\pi^0$	C [a]	< 6.0 $\times 10^{-5}$	90%
Γ_{30} $\mu^+\mu^-\eta$	C [a]	< 1.5 $\times 10^{-5}$	90%
Γ_{31} $e\mu$	LF	< 4.7 $\times 10^{-4}$	90%

[a] C parity forbids this to occur as a single-photon process.

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, 2 combinations of partial widths obtained from integrated cross section, and 15 branching ratios uses 43 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 48.0$ for 35 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	0							
x_3	-76	-58						
x_4	-19	-23	4					
x_5	-29	-25	32	-1				
x_6	-24	-18	29	1	9			
x_9	0	-2	-3	-1	-1	-1		
x_{17}	-4	-6	-5	-2	-3	-2	0	
Γ	25	5	-19	5	-71	-5	1	3
	x_1	x_2	x_3	x_4	x_5	x_6	x_9	x_{17}

Mode	Rate (MeV)
Γ_1 $\pi^+ \pi^- \eta$	0.085 \pm 0.004
Γ_2 $\rho^0 \gamma$ (including non-resonant $\pi^+ \pi^- \gamma$)	0.0575 \pm 0.0028
Γ_3 $\pi^0 \pi^0 \eta$	0.0439 \pm 0.0023
Γ_4 $\omega \gamma$	0.0054 \pm 0.0005
Γ_5 $\gamma \gamma$	0.00435 \pm 0.00013
Γ_6 $3\pi^0$	(4.2 \pm 0.4) $\times 10^{-4}$
Γ_9 $\pi^+ \pi^- \pi^0$	(7.5 \pm 0.8) $\times 10^{-4}$
Γ_{17} $\pi^+ \pi^- e^+ e^-$	(4.7 $^{+2.6}_{-1.9}$) $\times 10^{-4}$

$\eta'(958)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$								Γ_5
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT				
4.35 \pm 0.14 OUR FIT								
4.28 \pm 0.19 OUR AVERAGE								
4.17 \pm 0.10 \pm 0.27	2000	⁴ ACCIARRI	98Q L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \gamma$				
4.53 \pm 0.29 \pm 0.51	266	KARCH	92 CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$				
3.61 \pm 0.13 \pm 0.48		⁵ BEHREND	91 CELL	$e^+ e^- \rightarrow e^+ e^- \eta'(958)$				
4.6 \pm 1.1 \pm 0.6	23	BARU	90 MD1	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \gamma$				

4.57±0.25±0.44		BUTLER	90	MRK2	$e^+e^- \rightarrow e^+e^-\eta'(958)$
5.08±0.24±0.71	547	⁶ ROE	90	ASP	$e^+e^- \rightarrow e^+e^-2\gamma$
3.8 ±0.7 ±0.6	34	AIHARA	88C	TPC	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.9 ±0.5 ±0.5	136	⁷ WILLIAMS	88	CBAL	$e^+e^- \rightarrow e^+e^-2\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
4.7 ±0.6 ±0.9	143	⁸ GIDAL	87	MRK2	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.0 ±0.9		⁹ BARTEL	85E	JADE	$e^+e^- \rightarrow e^+e^-2\gamma$

⁴ No non-resonant $\pi^+\pi^-$ contribution found.

⁵ Reevaluated by us using $B(\eta' \rightarrow \rho(770)\gamma) = (30.2 \pm 1.3)\%$.

⁶ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.

⁷ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.

⁸ Superseded by BUTLER 90.

⁹ Systematic error not evaluated.

$\eta'(958) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $\gamma\gamma$ and with the total width is obtained from the integrated cross section into channel(i) in the $\gamma\gamma$ annihilation.

$\Gamma(\gamma\gamma) \times \Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}} \quad \Gamma_5\Gamma_2/\Gamma$

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.27±0.04 OUR FIT				
1.26±0.07 OUR AVERAGE Error includes scale factor of 1.2.				
1.09±0.04±0.13		BEHREND	91	CELL $e^+e^- \rightarrow e^+e^-\rho(770)^0\gamma$
1.35±0.09±0.21		AIHARA	87	TPC $e^+e^- \rightarrow e^+e^-\rho\gamma$
1.13±0.04±0.13	867	ALBRECHT	87B	ARG $e^+e^- \rightarrow e^+e^-\rho\gamma$
1.53±0.09±0.21		ALTHOFF	84E	TASS $e^+e^- \rightarrow e^+e^-\rho\gamma$
1.14±0.08±0.11	243	BERGER	84B	PLUT $e^+e^- \rightarrow e^+e^-\rho\gamma$
1.73±0.34±0.35	95	JENNI	83	MRK2 $e^+e^- \rightarrow e^+e^-\rho\gamma$
1.49±0.13±0.027	213	BARTEL	82B	JADE $e^+e^- \rightarrow e^+e^-\rho\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.85±0.31±0.24	43	BEHREND	83B	CELL $e^+e^- \rightarrow e^+e^-\rho\gamma$

$\Gamma(\gamma\gamma) \times \Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}} \quad \Gamma_5\Gamma_3/\Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.97±0.05 OUR FIT			
0.92±0.06±0.11 ¹⁰ KARCH 92 CBAL $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.95±0.05±0.08	¹¹ KARCH	90	CBAL $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
1.00±0.08±0.10	^{11,12} ANTREASYAN	87	CBAL $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
¹⁰ Reevaluated by us using $B(\eta \rightarrow \gamma\gamma) = (39.21 \pm 0.34)\%$. Supersedes ANTREASYAN 87 and KARCH 90.			
¹¹ Superseded by KARCH 92.			
¹² Using $BR(\eta \rightarrow 2\gamma) = (38.9 \pm 0.5)\%$.			

$\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

$$|\text{MATRIX ELEMENT}|^2 = |1 + \alpha Y|^2 + CX + DX^2$$

X and Y are Dalitz variables; α is complex and C , and D are real-valued. Parameters C and D are not necessarily equal to c and d , respectively, in the generalized parameterization following this one. May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays. Because of different initial assumptions and strong correlations of the parameters we do not average the parameters in the section below.

Re(α) decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.033 \pm 0.005 \pm 0.003$	44k	¹³ ABLIKIM	11 BES3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$-0.072 \pm 0.012 \pm 0.006$	7k	¹⁴ AMELIN	05A VES	$28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
$-0.021 \pm 0.018 \pm 0.017$	6.7k	¹⁵ BRIERE	00 CLEO	$10.6 e^+e^- \rightarrow \eta\pi^+\pi^- X$
$-0.058 \pm 0.013 \pm 0.003$	5.4k	¹⁶ ALDE	86 GAM2	$38 \pi^- p \rightarrow n\eta\pi^0\pi^0$
-0.08 ± 0.03		^{16,17} KALBFLEISCH	74 RVUE	$\eta' \rightarrow \eta\pi^+\pi^-$

¹³ See ABLIKIM 11 for the full correlation matrix.

¹⁴ Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

¹⁵ Assuming $\text{Im}(\alpha) = 0$, $C = 0$, and $D = 0$.

¹⁶ Assuming $C = 0$.

¹⁷ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

Im(α) decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.000 \pm 0.049 \pm 0.001$	44k	¹⁸ ABLIKIM	11 BES3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$0.0 \pm 0.1 \pm 0.0$	7k	¹⁹ AMELIN	05A VES	$28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
$-0.00 \pm 0.13 \pm 0.00$	5.4k	²⁰ ALDE	86 GAM2	$38 \pi^- p \rightarrow n\eta\pi^0\pi^0$
0.0 ± 0.3		^{20,21} KALBFLEISCH	74 RVUE	$\eta' \rightarrow \eta\pi^+\pi^-$

¹⁸ See ABLIKIM 11 for the full correlation matrix.

¹⁹ Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

²⁰ Assuming $C = 0$.

²¹ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

C decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$+0.018 \pm 0.009 \pm 0.003$	44k	²² ABLIKIM	11 BES3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$0.020 \pm 0.018 \pm 0.004$	7k	²³ AMELIN	05A VES	$28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$

²² See ABLIKIM 11 for the full correlation matrix.

²³ Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

D decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.059 \pm 0.012 \pm 0.004$	44k	²⁴ ABLIKIM	11 BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.066 \pm 0.030 \pm 0.015$	7k	²⁵ AMELIN	05A VES	$28 \pi^- A \rightarrow \eta \pi^+ \pi^- \pi^- A^*$
$0.00 \pm 0.03 \pm 0.00$	5.4k	²⁶ ALDE	86 GAM2	$38 \pi^- p \rightarrow n \eta \pi^0 \pi^0$
0		^{26,27} KALBFLEISCH	74 RVUE	$\eta' \rightarrow \eta \pi^+ \pi^-$

²⁴ See ABLIKIM 11 for the full correlation matrix.

²⁵ Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

²⁶ Assuming $C = 0$.

²⁷ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

$\eta'(958) \rightarrow \eta \pi \pi$ DECAY PARAMETERS

$$|\text{MATRIX ELEMENT}|^2 \propto 1 + a Y + b Y^2 + c X + d X^2$$

X and Y are Dalitz variables and a , b , c , and d are real-valued parameters. May be different for $\eta'(958) \rightarrow \eta \pi^+ \pi^-$ and $\eta'(958) \rightarrow \eta \pi^0 \pi^0$ decays. We do not average measurements in the section below because parameter values from each experiment are strongly correlated.

a decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.047 \pm 0.011 \pm 0.003$	44k	²⁸ ABLIKIM	11 BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.066 \pm 0.016 \pm 0.003$	15k	²⁹ BLIK	09 GAM4	$32.5 \pi^- p \rightarrow \eta' n$
$-0.127 \pm 0.016 \pm 0.008$	20k	³⁰ DOROFEEV	07 VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

²⁸ See ABLIKIM 11 for the full correlation matrix.

²⁹ From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay.

³⁰ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

b decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.069 \pm 0.019 \pm 0.009$	44k	³¹ ABLIKIM	11 BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.063 \pm 0.028 \pm 0.004$	15k	³² BLIK	09 GAM4	$32.5 \pi^- p \rightarrow \eta' n$
$-0.106 \pm 0.028 \pm 0.014$	20k	³³ DOROFEEV	07 VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

³¹ See ABLIKIM 11 for the full correlation matrix.

³² From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay.

³³ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

c decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$+0.019 \pm 0.011 \pm 0.003$	44k	34 ABLIKIM	11 BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.107 \pm 0.096 \pm 0.003$	15k	35 BLIK	09 GAM4	$32.5 \pi^- p \rightarrow \eta' n$
$0.015 \pm 0.011 \pm 0.014$	20k	36 DOROFEEV	07 VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

³⁴ See ABLIKIM 11 for the full correlation matrix.

³⁵ From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay.

³⁶ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

d decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.073 \pm 0.012 \pm 0.003$	44k	37 ABLIKIM	11 BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.018 \pm 0.078 \pm 0.006$	15k	38 BLIK	09 GAM4	$32.5 \pi^- p \rightarrow \eta' n$
$-0.082 \pm 0.017 \pm 0.008$	20k	39 DOROFEEV	07 VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

³⁷ See ABLIKIM 11 for the full correlation matrix.

³⁸ From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay. If $c \equiv 0$ from Bose-Einstein symmetry, $d = -0.067 \pm 0.020 \pm 0.003$.

³⁹ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

$\eta'(958)$ β PARAMETER

$|\text{MATRIX ELEMENT}|^2 = (1 + 2\beta Z)$

See the "Note on η Decay Parameters" in our 1994 edition *Physical Review D50* 1173 (1994), p. 1454.

β decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.46 ± 0.22 OUR AVERAGE		Error includes scale factor of 1.4.		
-0.59 ± 0.18	235	BLIK	08 GAMS	$32 \pi^- p \rightarrow \eta' n$
-0.1 ± 0.3		ALDE	87B GAM2	$38 \pi^- p \rightarrow n 3\pi^0$

$\eta'(958)$ BRANCHING RATIOS

$\Gamma(\pi^+ \pi^- \eta) / \Gamma_{\text{total}}$

Γ_1 / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.429 ± 0.007 OUR FIT				

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

$0.424 \pm 0.011 \pm 0.004$ 1.2k ⁴⁰ PEDLAR 09 CLEO $J/\psi \rightarrow \gamma \eta'$

⁴⁰ Not independent of other η' branching fractions and ratios in PEDLAR 09.

$\Gamma(\pi^+ \pi^- \eta(\text{charged decay}))/\Gamma_{\text{total}}$ **0.286 Γ_1/Γ**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.1228\pm0.0020 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.123 \pm 0.014	107	RITTENBERG 69	HBC	1.7–2.7 $K^- p$
0.10 \pm 0.04	10	LONDON 66	HBC	2.24 $K^- p \rightarrow \Lambda 2\pi^+ 2\pi^- \pi^0$
0.07 \pm 0.04	7	BADIER 65B	HBC	3 $K^- p$

$\Gamma(\pi^+ \pi^- \eta(\text{neutral decay}))/\Gamma_{\text{total}}$ **0.714 Γ_1/Γ**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.307\pm0.005 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.314 \pm 0.026	281	RITTENBERG 69	HBC	1.7–2.7 $K^- p$
-------------------	-----	---------------	-----	-----------------

$\Gamma(\rho^0 \gamma(\text{including non-resonant } \pi^+ \pi^- \gamma))/\Gamma_{\text{total}}$ **Γ_2/Γ**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.291\pm0.006 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.287 \pm 0.007 \pm 0.004	0.2k	⁴¹ PEDLAR 09	CLEO	$J/\psi \rightarrow \gamma \eta'$
0.329 \pm 0.033	298	RITTENBERG 69	HBC	1.7–2.7 $K^- p$
0.2 \pm 0.1	20	LONDON 66	HBC	2.24 $K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$
0.34 \pm 0.09	35	BADIER 65B	HBC	3 $K^- p$

⁴¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.

$\Gamma(\rho^0 \gamma(\text{including non-resonant } \pi^+ \pi^- \gamma))/\Gamma(\pi^+ \pi^- \eta)$ **Γ_2/Γ_1**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.677\pm0.017 OUR FIT			

0.683 \pm 0.020 OUR AVERAGE

0.677 \pm 0.024 \pm 0.011	PEDLAR 09	CLE3	$J/\psi \rightarrow \eta' \gamma$
0.69 \pm 0.03	ABLIKIM 06E	BES2	$J/\psi \rightarrow \eta' \gamma$

$\Gamma(\rho^0 \gamma(\text{including non-resonant } \pi^+ \pi^- \gamma))/\Gamma(\pi^+ \pi^- \eta(\text{neutral decay}))$ **$\Gamma_2/0.714\Gamma_1$**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.949\pm0.024 OUR FIT				

0.97 \pm 0.09 OUR AVERAGE

0.70 \pm 0.22	AMSLER 04B	CBAR	$0 \bar{p} p \rightarrow \pi^+ \pi^- \eta$
1.07 \pm 0.17	BELADIDZE 92C	VES	$36 \pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
0.92 \pm 0.14	DANBURG 73	HBC	$2.2 K^- p \rightarrow \Lambda X^0$
1.11 \pm 0.18	JACOBS 73	HBC	$2.9 K^- p \rightarrow \Lambda X^0$

$\Gamma(\pi^0 \pi^0 \eta)/\Gamma_{\text{total}}$ **Γ_3/Γ**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.222\pm0.008 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.235 \pm 0.013 \pm 0.004	3.2k	⁴² PEDLAR 09	CLEO	$J/\psi \rightarrow \gamma \eta'$
-------------------------------	------	-------------------------	------	-----------------------------------

⁴² Not independent of other η' branching fractions and ratios in PEDLAR 09.

$\Gamma(\pi^0 \pi^0 \eta(3\pi^0 \text{ decay}))/\Gamma_{\text{total}}$ **0.321\Gamma_3/\Gamma**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0712±0.0026 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.11 ±0.06	4	BENSINGER	70	DBC	2.2 $\pi^+ d$
------------	---	-----------	----	-----	---------------

$\Gamma(\pi^0 \pi^0 \eta)/\Gamma(\pi^+ \pi^- \eta)$ **\Gamma_3/\Gamma_1**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.517±0.026 OUR FIT			

0.555±0.043±0.013	PEDLAR	09	CLE3	$J/\psi \rightarrow \eta' \gamma$
--------------------------	--------	----	------	-----------------------------------

$\Gamma(\rho^0 \gamma(\text{including non-resonant } \pi^+ \pi^- \gamma))/\Gamma(\pi \pi \eta)$ **\Gamma_2/(\Gamma_1+\Gamma_3)**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.447±0.012 OUR FIT			

0.43 ±0.02 ±0.02	BARBERIS	98C	OMEG	450 $p p \rightarrow p_f \eta' p_s$
-------------------------	----------	-----	------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ±0.15	DAVIS	68	HBC	5.5 $K^- p$
------------	-------	----	-----	-------------

$\Gamma(\omega \gamma)/\Gamma_{\text{total}}$ **\Gamma_4/\Gamma**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0275±0.0023 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0234±0.0030±0.0004	70	⁴³ PEDLAR	09	CLEO	$J/\psi \rightarrow \gamma \eta'$
----------------------	----	----------------------	----	------	-----------------------------------

⁴³ Not independent of other η' branching fractions and ratios in PEDLAR 09.

$\Gamma(\omega \gamma)/\Gamma(\pi^+ \pi^- \eta)$ **\Gamma_4/\Gamma_1**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.064±0.006 OUR FIT				

0.055±0.007±0.001	PEDLAR	09	CLE3	$J/\psi \rightarrow \eta' \gamma$
--------------------------	--------	----	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.068±0.013	68	ZANFINO	77	ASPK	8.4 $\pi^- p$
-------------	----	---------	----	------	---------------

$\Gamma(\omega \gamma)/\Gamma(\pi^0 \pi^0 \eta)$ **\Gamma_4/\Gamma_3**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.124±0.011 OUR FIT			

0.147±0.016	ALDE	87B	GAM2	38 $\pi^- p \rightarrow n 4\gamma$
--------------------	------	-----	------	------------------------------------

$\Gamma(\rho^0 \gamma(\text{including non-resonant } \pi^+ \pi^- \gamma))/[\Gamma(\pi^+ \pi^- \eta) + \Gamma(\pi^0 \pi^0 \eta) + \Gamma(\omega \gamma)]$ **\Gamma_2/(\Gamma_1+\Gamma_3+\Gamma_4)**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.428±0.011 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.25 ±0.14	DAUBER	64	HBC	1.95 $K^- p$
------------	--------	----	-----	--------------

$[\Gamma(\pi^0 \pi^0 \eta(\text{charged decay})) + \Gamma(\omega(\text{charged decay})\gamma)]/\Gamma_{\text{total}}$ **(0.286\Gamma_3+0.89\Gamma_4)/\Gamma**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0880±0.0031 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.045 ±0.029	42	RITTENBERG	69	HBC	1.7–2.7 $K^- p$
--------------	----	------------	----	-----	-----------------

$\Gamma(\pi^+\pi^-\text{ neutrals})/\Gamma_{\text{total}}$ $(0.714\Gamma_1+0.286\Gamma_3+0.89\Gamma_4)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.395±0.004 OUR FIT

••• We do not use the following data for averages, fits, limits, etc. •••

0.4 ±0.1	39	LONDON	66	HBC	2.24 $K^- p \rightarrow \Lambda \pi^+ \pi^-$ neutrals
0.35 ±0.06	33	BADIER	65B	HBC	3 $K^- p$

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.20±0.08 OUR FIT

2.00±0.15 OUR AVERAGE

1.98 ^{+0.31} _{-0.27} ±0.07	114	44 WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
2.00±0.18		45 STANTON	80	SPEC	8.45 $\pi^- p \rightarrow n \pi^+ \pi^- 2\gamma$

••• We do not use the following data for averages, fits, limits, etc. •••

2.25±0.16±0.03	0.3k	46 PEDLAR	09	CLEO	$J/\psi \rightarrow \gamma \eta'$
1.8 ±0.2	6000	47 APEL	79	NICE	15-40 $\pi^- p \rightarrow n 2\gamma$
2.5 ±0.7		DUANE	74	MMS	$\pi^- p \rightarrow n \text{MM}$
1.71±0.33	68	DALPIAZ	72	CNTR	1.6 $\pi^- p \rightarrow n X^0$
2.0 ^{+0.8} _{-0.6}	31	HARVEY	71	OSPK	3.65 $\pi^- p \rightarrow n X^0$

⁴⁴WICHT 08 reports $[\Gamma(\eta'(958) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta' K^+)] = (1.40^{+0.16+0.15}_{-0.15-0.12}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta' K^+) = (7.06 \pm 0.25) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁵Includes APEL 79 result.

⁴⁶Not independent of other η' branching fractions and ratios in PEDLAR 09.

⁴⁷Data is included in STANTON 80 evaluation.

$\Gamma(\gamma\gamma)/\Gamma(\pi^+\pi^-\eta)$ Γ_5/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0513±0.0022 OUR FIT

0.053 ±0.004 ±0.001 PEDLAR 09 CLE3 $J/\psi \rightarrow \eta' \gamma$

$\Gamma(\gamma\gamma)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_5/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0757±0.0033 OUR FIT

0.080 ±0.008 ABLIKIM 06E BES2 $J/\psi \rightarrow \eta' \gamma$

$\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_5/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.099±0.004 OUR FIT

0.105±0.010 OUR AVERAGE Error includes scale factor of 1.9.

0.091±0.009	AMSLER	93	CBAR	0.0 $\bar{p} p$
0.112±0.002±0.006	ALDE	87B	GAM2	38 $\pi^- p \rightarrow n 2\gamma$

$\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta(\text{neutral decay}))$ $\Gamma_5/0.714\Gamma_3$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.139±0.006 OUR FIT

••• We do not use the following data for averages, fits, limits, etc. •••

0.188±0.058	16	APEL	72	OSPK	3.8 $\pi^- p \rightarrow n X^0$
-------------	----	------	----	------	---------------------------------

$\Gamma(\text{neutrals})/\Gamma_{\text{total}}$ $(0.714\Gamma_3+0.09\Gamma_4+\Gamma_5)/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.183±0.006 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.185±0.022	535	BASILE	71	CNTR	1.6 $\pi^- p \rightarrow n X^0$
0.189±0.026	123	RITTENBERG	69	HBC	1.7–2.7 $K^- p$

$\Gamma(3\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	-------------	--------------------	-------------	----------------

2.14±0.20 OUR FIT

3.56±0.22±0.34	309	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$
-----------------------	-----	---------	-----	------	-------------------------------------

$\Gamma(3\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_6/Γ_3

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	-------------	--------------------	-------------	----------------

96± 9 OUR FIT

78±10 OUR AVERAGE

86±19	235	BLIK	08	GAMS	32 $\pi^- p \rightarrow \eta' n$
74±15		ALDE	87B	GAM2	38 $\pi^- p \rightarrow n6\gamma$
75±18		BINON	84	GAM2	30–40 $\pi^- p \rightarrow n6\gamma$

$\Gamma(\mu^+\mu^-\gamma)/\Gamma(\gamma\gamma)$ Γ_7/Γ_5

<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	-------------	--------------------	-------------	----------------

4.9±1.2	33	VIKTOROV	80	CNTR	25,33 $\pi^- p \rightarrow 2\mu\gamma$
----------------	----	----------	----	------	--

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.29	90	⁴⁸ ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
<2.4	90	⁴⁹ NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$

⁴⁸ Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.

⁴⁹ Not independent of measured value of Γ_8/Γ_1 from NAIK 09.

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_8/Γ_1

<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	------------	--------------------	-------------	----------------

<0.5	90	⁵⁰ NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
----------------	----	--------------------	----	------	----------------------------------

⁵⁰ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.3 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_8/Γ_2

<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	------------	--------------------	-------------	----------------

<1.0	90	ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
----------------	----	---------	-----	------	----------------------------------

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

3.8 ± 0.4 OUR FIT

3.8 ± 0.4 OUR AVERAGE

3.83 ± 0.15 ± 0.39	1014	ABLIKIM	12E BES3	J/ψ → γ(π ⁺ π ⁻ π ⁰)
3.7 ^{+1.1} _{-0.9} ± 0.4		⁵¹ NAIK	09 CLEO	J/ψ → γη'

⁵¹ Not independent of measured value of Γ_9/Γ_1 from NAIK 09.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_9/Γ_1

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

8.8 ± 0.9 OUR FIT

8.28 ^{+2.49} _{-2.12} ± 0.04	20	⁵² NAIK	09 CLEO	J/ψ → γη'
---	----	--------------------	---------	-----------

⁵² NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (21 ⁺⁶₋₅ ± 2) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0\rho^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.04	90	RITTENBERG 65	HBC	2.7 K ⁻ p
-----------------	----	---------------	-----	----------------------

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.4	90	⁵³ NAIK	09 CLEO	J/ψ → γη'
<100	90	RITTENBERG 69	HBC	1.7-2.7 K ⁻ p

⁵³ Not independent of measured value of Γ_{11}/Γ_1 from NAIK 09.

$\Gamma(2(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{11}/Γ_1

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<0.6	90	⁵⁴ NAIK	09 CLEO	J/ψ → γη'
----------------	----	--------------------	---------	-----------

⁵⁴ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.4 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<27	90	⁵⁵ NAIK	09 CLEO	J/ψ → γη'
---------------	----	--------------------	---------	-----------

⁵⁵ Not independent of measured value of Γ_{12}/Γ_1 from NAIK 09.

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{12}/Γ_1

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<6	90	⁵⁶ NAIK	09 CLEO	J/ψ → γη'
--------------	----	--------------------	---------	-----------

⁵⁶ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 15 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

$\Gamma(2(\pi^+\pi^-)\text{ neutrals})/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.01	95	DANBURG 73	HBC	2.2 $K^- p \rightarrow \Lambda \chi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.01	90	RITTENBERG 69	HBC	1.7–2.7 $K^- p$

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ **Γ_{14}/Γ**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.002	90	⁵⁷ NAIK 09	CLEO	$J/\psi \rightarrow \gamma \eta'$
<0.01	90	RITTENBERG 69	HBC	1.7–2.7 $K^- p$

⁵⁷ Not independent of measured value of Γ_{14}/Γ_1 from NAIK 09.

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma(\pi^+\pi^-\eta)$ **Γ_{14}/Γ_1**

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	⁵⁸ NAIK 09	CLEO	$J/\psi \rightarrow \gamma \eta'$

⁵⁸ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 11 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

$\Gamma(2(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$ **Γ_{15}/Γ**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.01	95	KALBFLEISCH 64B	HBC	$K^- p \rightarrow \Lambda 2(\pi^+\pi^-)+MM$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.01	90	LONDON 66	HBC	Compilation

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ **Γ_{16}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3.1	90	⁵⁹ ABLIKIM 13U	BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 53	90	⁶⁰ NAIK 09	CLEO	$J/\psi \rightarrow \gamma \eta'$
<500	95	KALBFLEISCH 64B	HBC	$K^- p \rightarrow \Lambda 2(\pi^+\pi^-)$

⁵⁹ Using $B(J/\psi \rightarrow \gamma \eta'(958)) = (5.16 \pm 0.15) \times 10^{-3}$.

⁶⁰ Not independent of measured value of Γ_{16}/Γ_1 from NAIK 09.

$\Gamma(3(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ **Γ_{16}/Γ_1**

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2	90	⁶¹ NAIK 09	CLEO	$J/\psi \rightarrow \gamma \eta'$

⁶¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 3(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 3.0 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

2.4 $^{+1.3}_{-1.0}$ OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.11 ± 0.12 ± 0.14 429 ⁶² ABLIKIM 130 BES3 $J/\psi \rightarrow \gamma\eta'$

2.5 $^{+1.2}_{-0.9}$ ± 0.5 ⁶³ NAIK 09 CLEO $J/\psi \rightarrow \gamma\eta'$

<6 90 RITTENBERG 65 HBC 2.7 K^-p

⁶² Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.

⁶³ Not independent of measured value of Γ_{17}/Γ_1 from NAIK 09.

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_{17}/Γ_1

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.6 $^{+3.0}_{-2.2}$ OUR FIT

5.52 $^{+3.00}_{-2.30} \pm 0.03$ 8 ⁶⁴ NAIK 09 CLEO $J/\psi \rightarrow \gamma\eta'$

⁶⁴ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-e^+e^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (14^{+7}_{-5} \pm 3) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_{17}/Γ_2

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

7.2 ± 0.4 ± 0.5 429 ABLIKIM 130 BES3 $J/\psi \rightarrow \gamma\eta'$

$\Gamma(\pi^+e^-\nu_e + \text{c.c.})/\Gamma(\pi^+\pi^-\eta)$ Γ_{18}/Γ_1

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<5.0 90 ABLIKIM 13G BES3 $J/\psi \rightarrow \phi\eta'$

$\Gamma(\gamma e^+e^-)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.9 90 BRIERE 00 CLEO 10.6 e^+e^-

$\Gamma(\pi^0\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_{20}/Γ_3

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<37 90 ALDE 87B GAM2 38 $\pi^-p \rightarrow n4\gamma$

$\Gamma(4\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_{21}/Γ_3

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<23 90 ALDE 87B GAM2 38 $\pi^-p \rightarrow n8\gamma$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<2.1 90 VOROBYEV 88 ND $e^+e^- \rightarrow \pi^+\pi^-\eta$

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9.5	90	⁶⁵ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
------	----	--------------------	----	---------------------------------------

⁶⁵ Not independent of measured value of Γ_{23}/Γ_1 from NAIK 09.

$\Gamma(\text{invisible})/\Gamma(\gamma\gamma)$ Γ_{23}/Γ_5

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<2.4	90	ABLIKIM	13	BES3 $J/\psi \rightarrow \phi\eta'$
----------------	----	---------	----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.69	90	ABLIKIM	06Q	BES $J/\psi \rightarrow \phi\eta'$
-------	----	---------	-----	------------------------------------

$\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\eta)$ Γ_{23}/Γ_1

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1	90	⁶⁶ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
------	----	--------------------	----	---------------------------------------

⁶⁶ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \text{invisible})/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 5.4 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

< 0.6	90	⁶⁷ ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-$
-----------------	----	-----------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 29	90	⁶⁸ MORI	07A	BELL $\gamma\gamma \rightarrow \pi^+\pi^-$
------	----	--------------------	-----	--

< 3.3	90	⁶⁹ MORI	07A	BELL $\gamma\gamma \rightarrow \pi^+\pi^-$
-------	----	--------------------	-----	--

<800	95	DANBURG	73	HBC $2.2 K^- p \rightarrow \Lambda X^0$
------	----	---------	----	---

<200	90	RITTENBERG	69	HBC $1.7\text{--}2.7 K^- p$
------	----	------------	----	-----------------------------

⁶⁷ ABLIKIM 11G reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.15 \times 10^{-3}$.

⁶⁸ Taking into account interference with the $\gamma\gamma \rightarrow \pi^+\pi^-$ continuum.

⁶⁹ Without interference with the $\gamma\gamma \rightarrow \pi^+\pi^-$ continuum.

$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

< 4×10^{-4}	90	⁷⁰ ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma\pi^0\pi^0$
---	----	-----------------------	-----	--

⁷⁰ ABLIKIM 11G reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.15 \times 10^{-3}$.

$\Gamma(\pi^0\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_{25}/Γ_3

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<45	90	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n4\gamma$
---------------	----	------	-----	--

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	90	BRIERE	00	CLEO 10.6 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<13	90	RITTENBERG	65	HBC 2.7 $K^- p$

$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.4	90	BRIERE	00	CLEO 10.6 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<11	90	RITTENBERG	65	HBC 2.7 $K^- p$

$\Gamma(3\gamma)/\Gamma(\pi^0 \pi^0 \eta)$ Γ_{28}/Γ_3

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.6	90	ALDE	87B	GAM2 38 $\pi^- p \rightarrow n 3\gamma$

$\Gamma(\mu^+ \mu^- \pi^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<6.0	90	DZHELYADIN	81	CNTR 30 $\pi^- p \rightarrow \eta' n$

$\Gamma(\mu^+ \mu^- \eta)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	DZHELYADIN	81	CNTR 30 $\pi^- p \rightarrow \eta' n$

$\Gamma(e\mu)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.7	90	BRIERE	00	CLEO 10.6 $e^+ e^-$

$\eta'(958)$ C-NONCONSERVING DECAY PARAMETER

See the note on η decay parameters in the Stable Particle Particle Listings for definition of this parameter.

DECAY ASYMMETRY PARAMETER FOR $\pi^+ \pi^- \gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.03 ± 0.04 OUR AVERAGE				
-0.019 ± 0.056		AIHARA	87	TPC $2\gamma \rightarrow \pi^+ \pi^- \gamma$
-0.069 ± 0.078	295	GRIGORIAN	75	STRC 2.1 $\pi^- p$
0.00 ± 0.10	103	KALBFLEISCH	75	HBC 2.18 $K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.07 ± 0.08	152	RITTENBERG	65	HBC 2.1-2.7 $K^- p$

$\eta'(958)$ REFERENCES

- ABLIKIM 13 PR D87 012009 M. Ablikim *et al.* (BES III Collab.)
 ABLIKIM 13G PR D87 032006 M. Ablikim *et al.* (BES III Collab.)
 ABLIKIM 13O PR D87 092011 M. Ablikim *et al.* (BES III Collab.)
 ABLIKIM 13U PR D88 091502 M. Ablikim *et al.* (BES III Collab.)
 ABLIKIM 12E PRL 108 182001 M. Ablikim *et al.* (BES III Collab.)
 PDG 12 PR D86 010001 J. Beringer *et al.* (PDG Collab.)
 ABLIKIM 11 PR D83 012003 M. Ablikim *et al.* (BES III Collab.)
 ABLIKIM 11G PR D84 032006 M. Ablikim *et al.* (BES III Collab.)
 CZERWINSKI 10 PRL 105 122001 E. Czerwinski *et al.* (COSY-11 Collab.)
 BLIK 09 PAN 72 231 A.M. Blik *et al.* (IHEP (Protvino))
 Translated from YAF 72 258.
 NAIK 09 PRL 102 061801 P. Naik *et al.* (CLEO Collab.)
 PEDLAR 09 PR D79 111101 T.K. Pedlar *et al.* (CLEO Collab.)
 BLIK 08 PAN 71 2124 A. Blik *et al.* (GAMS-4 π Collab.)
 Translated from YAF 71 2161.
 LIBBY 08 PRL 101 182002 J. Libby *et al.* (CLEO Collab.)
 WICHT 08 PL B662 323 J. Wicht *et al.* (BELLE Collab.)
 DOROFEEV 07 PL B651 22 V. Dorofeev *et al.* (VES Collab.)
 MORI 07A JPSJ 76 074102 T. Mori *et al.* (BELLE Collab.)
 ABLIKIM 06E PR D73 052008 M. Ablikim *et al.* (BES Collab.)
 ABLIKIM 06Q PRL 97 202002 M. Ablikim *et al.* (BES Collab.)
 AMELIN 05A PAN 68 372 D.V. Amelin *et al.* (VES Collab.)
 Translated from YAF 68 401.
 AMSLER 04B EPJ C33 23 C. Amsler *et al.* (Crystal Barrel Collab.)
 BAI 04J PL B594 47 J.Z. Bai *et al.* (BES Collab.)
 BRIERE 00 PRL 84 26 R. Briere *et al.* (CLEO Collab.)
 ACCIARRI 98Q PL B418 399 M. Acciarri *et al.* (L3 Collab.)
 BARBERIS 98C PL B440 225 D. Barberis *et al.* (WA 102 Collab.)
 WURZINGER 96 PL B374 283 R. Wurzinger *et al.* (BONN, ORSAY, SACL+)
 PDG 94 PR D50 1173 L. Montanet *et al.* (CERN, LBL, BOST+)
 AMSLER 93 ZPHY C58 175 C. Amsler *et al.* (Crystal Barrel Collab.)
 BELADIDZE 92C SJNP 55 1535 G.M. Beladidze, S.I. Bitjukov, G.V. Borisov (SERP+)
 Translated from YAF 55 2748.
 KARCH 92 ZPHY C54 33 K. Karch *et al.* (Crystal Ball Collab.)
 ARMSTRONG 91B ZPHY C52 389 T.A. Armstrong *et al.* (ATHU, BARI, BIRM+)
 BEHREND 91 ZPHY C49 401 H.J. Behrend *et al.* (CELLO Collab.)
 AUGUSTIN 90 PR D42 10 J.E. Augustin *et al.* (DM2 Collab.)
 BARU 90 ZPHY C48 581 S.E. Baru *et al.* (MD-1 Collab.)
 BUTLER 90 PR D42 1368 F. Butler *et al.* (Mark II Collab.)
 KARCH 90 PL B249 353 K. Karch *et al.* (Crystal Ball Collab.)
 ROE 90 PR D41 17 N.A. Roe *et al.* (ASP Collab.)
 AIHARA 88C PR D38 1 H. Aihara *et al.* (TPC-2 γ Collab.)
 VOROBYEV 88 SJNP 48 273 P.V. Vorobiev *et al.* (NOVO)
 Translated from YAF 48 436.
 WILLIAMS 88 PR D38 1365 D.A. Williams *et al.* (Crystal Ball Collab.)
 AIHARA 87 PR D35 2650 H. Aihara *et al.* (TPC-2 γ Collab.) JP
 ALBRECHT 87B PL B199 457 H. Albrecht *et al.* (ARGUS Collab.)
 ALDE 87B ZPHY C36 603 D.M. Alde *et al.* (LANL, BELG, SERP, LAPP)
 ANTREASYAN 87 PR D36 2633 D. Antreasyan *et al.* (Crystal Ball Collab.)
 GIDAL 87 PRL 59 2012 G. Gidal *et al.* (LBL, SLAC, HARV)
 ALDE 86 PL B177 115 D.M. Alde *et al.* (SERP, BELG, LANL, LAPP)
 BARTEL 85E PL 160B 421 W. Bartel *et al.* (JADE Collab.)
 ALTHOFF 84E PL 147B 487 M. Althoff *et al.* (TASSO Collab.)
 BERGER 84B PL 142B 125 C. Berger (PLUTO Collab.)
 BINON 84 PL 140B 264 F.G. Binon *et al.* (SERP, BELG, LAPP+)
 BEHREND 83B PL 125B 518 (erratum) H.J. Behrend *et al.* (CELLO Collab.)
 Also PL 114B 378 H.J. Behrend *et al.* (CELLO Collab.)
 JENNI 83 PR D27 1031 P. Jenni *et al.* (SLAC, LBL)
 BARTEL 82B PL 113B 190 W. Bartel *et al.* (JADE Collab.)
 DZHELADIN 81 PL 105B 239 R.I. Dzhelyadin *et al.* (SERP)
 STANTON 80 PL B92 353 N.R. Stanton *et al.* (OSU, CARL, MCGI+)
 VIKTOROV 80 SJNP 32 520 V.A. Viktorov *et al.* (SERP)
 Translated from YAF 32 1005.
 APEL 79 PL 83B 131 W.D. Apel, K.H. Augenstein, E. Bertolucci (KARLK+)
 BINNIE 79 PL 83B 141 D.M. Binnie *et al.* (LOIC)
 ZANFINO 77 PRL 38 930 C. Zanfino *et al.* (CARL, MCGI, OHIO+)
 GRIGORIAN 75 NP B91 232 A. Grigorian *et al.* (+)
 KALBFLEISCH 75 PR D11 987 G.R. Kalbfleisch, R.C. Strand, J.W. Chapman (BNL+)
 DUANE 74 PRL 32 425 A. Duane *et al.* (LOIC, SHMP)

KALBFLEISCH	74	PR D10 916	G.R. Kalbfleisch	(BNL)
DANBURG	73	PR D8 3744	J.S. Danburg <i>et al.</i>	(BNL, MICH) JP
JACOBS	73	PR D8 18	S.M. Jacobs <i>et al.</i>	(BRAN, UMD, SYRA+) JP
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)
APEL	72	PL 40B 680	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)
DALPIAZ	72	PL 42B 377	P.F. Dalpiaz <i>et al.</i>	(CERN)
BASILE	71	NC 3A 371	M. Basile <i>et al.</i>	(CERN, BGNA, STRB)
HARVEY	71	PRL 27 885	E.H. Harvey <i>et al.</i>	(MINN, MICH)
BENSINGER	70	PL 33B 505	J.R. Bensinger <i>et al.</i>	(WISC)
RITTENBERG	69	Thesis UCRL 18863	A. Rittenberg	(LRL) I
DAVIS	68	PL 27B 532	R. Davis <i>et al.</i>	(NWES, ANL)
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IJP
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)
RITTENBERG	65	PRL 15 556	A. Rittenberg, G.R. Kalbfleisch	(LRL, BNL)
DAUBER	64	PRL 13 449	P.M. Dauber <i>et al.</i>	(UCLA) JP
KALBFLEISCH	64B	PRL 13 349	G.R. Kalbfleisch, O.I. Dahl, A. Rittenberg	(LRL) JP
