

$f_2(1810)$


$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

$f_2(1810)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1815±12 OUR AVERAGE				
Error includes scale factor of 1.4. See the ideogram below.				
$1737 \pm 9^{+198}_{-65}$		¹ UEHARA	10A BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
1800 ± 30	40	ALDE	88D GAM4	300 $\pi^-p \rightarrow \pi^-p4\pi^0$
1806 ± 10	1600	ALDE	87 GAM4	100 $\pi^-p \rightarrow 4\pi^0n$
1870 ± 40		² ALDE	86D GAM4	100 $\pi^-p \rightarrow \eta\eta n$
1857^{+35}_{-24}		³ COSTA...	80 OMEG	10 $\pi^-p \rightarrow K^+K^-n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1858^{+18}_{-71}		⁴ LONGACRE	86 RVUE	Compilation
1799 ± 15		⁵ CASON	82 STRC	8 $\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$



¹ Breit-Wigner mass.

² Seen in only one solution.

³ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.


⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

⁵ From an amplitude analysis of the reaction $\pi^+\pi^-\rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.

$f_2(1810)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
197\pm 22 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
228 $^{+21}_{-20}$ 234-153		⁶ UEHARA	10A BELL	10.6 $e^+e^-\rightarrow e^+e^-\eta\eta$
160 \pm 30	40	ALDE	88D GAM4	300 $\pi^-p\rightarrow \pi^-p4\pi^0$
190 \pm 20	1600	ALDE	87 GAM4	100 $\pi^-p\rightarrow 4\pi^0n$
250 \pm 30		⁷ ALDE	86D GAM4	100 $\pi^-p\rightarrow \eta\eta n$
185 $^{+102}_{-139}$		⁸ COSTA...	80 OMEG	10 $\pi^-p\rightarrow K^+K^-n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
388 $^{+15}_{-21}$		⁹ LONGACRE	86 RVUE	Compilation
280 $^{+42}_{-35}$		¹⁰ CASON	82 STRC	8 $\pi^+p\rightarrow \Delta^{++}\pi^0\pi^0$

WEIGHTED AVERAGE
197 \pm 22 (Error scaled by 1.5)



⁶ Breit-Wigner width.

⁷ Seen in only one solution.

⁸ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

⁹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

¹⁰ From an amplitude analysis of the reaction $\pi^+\pi^-\rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.

$f_2(1810)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	
Γ_2 $\eta\eta$	
Γ_3 $4\pi^0$	seen
Γ_4 K^+K^-	
Γ_5 $\gamma\gamma$	seen

$f_2(1810)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_2\Gamma_5/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
$5.2^{+0.9+37.3}_{-0.8-4.5}$	¹¹ UEHARA	10A BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$	

¹¹ Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

$f_2(1810)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AMSLER	02 CBAR	$0.9 \bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$	
not seen	PROKOSHKIN 97	GAM2	$38 \pi^-p \rightarrow \pi^0\pi^0n$	
$0.21^{+0.02}_{-0.03}$	¹² LONGACRE	86 RVUE	Compilation	
0.44 ± 0.03	¹³ CASON	82 STRC	$8 \pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$	

¹² From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

¹³ Included in LONGACRE 86 global analysis.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$				Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
$0.008^{+0.028}_{-0.003}$	¹⁴ LONGACRE	86 RVUE	Compilation	

¹⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

$\Gamma(\pi\pi)/\Gamma(4\pi^0)$					Γ_1/Γ_3
VALUE	DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.75	ALDE	87	GAM4	100	$\pi^- p \rightarrow 4\pi^0 n$

$\Gamma(4\pi^0)/\Gamma(\eta\eta)$					Γ_3/Γ_2
VALUE	DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.8 ± 0.3	ALDE	87	GAM4	100	$\pi^- p \rightarrow 4\pi^0 n$

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$0.003^{+0.019}_{-0.002}$	¹⁵ LONGACRE	86	RVUE	Compilation	
seen	COSTA...	80	OMEG	$10 \pi^- p \rightarrow K^+ K^- n$	
¹⁵ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.					

$f_2(1810)$ REFERENCES

UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
PROKOSHKIN	97	SPD 42 117	Y.D. Prokoshkin <i>et al.</i>	(SERP)
		Translated from DANS 353 323.		
ALDE	88D	SJNP 47 810	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
		Translated from YAF 47 1273.		
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)
COSTA...	80	NP B175 402	G. Costa de Beaugard <i>et al.</i>	(BARI, BONN+)