

$\Lambda_c(2625)^+$

$$I(J^P) = 0(\frac{3}{2}^-) \quad \text{Status: } ***$$

The spin-parity has not been measured but is expected to be $3/2^-$:
this is presumably the charm counterpart of the strange $\Lambda(1520)$.

$\Lambda_c(2625)^+$ MASS


The mass is obtained from the $\Lambda_c(2625)^+ - \Lambda_c^+$ mass-difference measurements below.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|--------|-------------|---------|-------------------------------------|
| 2628.11 ± 0.19 OUR FIT | | | | Error includes scale factor of 1.1. |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 2626.6 ± 0.5 ± 1.5 | 42 ± 9 | ALBRECHT | 93F ARG | See ALBRECHT 97 |

$\Lambda_c(2625)^+ - \Lambda_c^+$ MASS DIFFERENCE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|----------|-------------|---------|---|
| 341.65 ± 0.13 OUR FIT | | | | Error includes scale factor of 1.1. |
| 341.65 ± 0.15 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |
| 341.65 ± 0.04 ± 0.12 | 6.2k | AALTONEN | 11H CDF | $p\bar{p}$ at 1.96 TeV |
| 342.1 ± 0.5 ± 0.5 | 51 | ALBRECHT | 97 ARG | $e^+e^- \approx 10$ GeV |
| 342.2 ± 0.2 ± 0.5 | 245 ± 19 | EDWARDS | 95 CLE2 | $e^+e^- \approx 10.5$ GeV |
| 340.4 ± 0.6 ± 0.3 | 40 ± 9 | FRABETTI | 94 E687 | $\gamma\text{Be}, \bar{E}_\gamma = 220$ GeV |

WEIGHTED AVERAGE
341.65 ± 0.15 (Error scaled by 1.3)



$\Lambda_c(2625)^+$ WIDTH

| VALUE (MeV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------|-------------|---------|-------------------------------|
| <0.97 | 90 | 6.2k | AALTONEN | 11H CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <1.9 | 90 | 245 ± 19 | EDWARDS | 95 CLE2 | $e^+e^- \approx 10.5$ GeV |
| <3.2 | 90 | | ALBRECHT | 93F ARG | $e^+e^- \approx \Upsilon(4S)$ |

$\Lambda_c(2625)^+$ DECAY MODES

$\Lambda_c^+ \pi \pi$ and its submode $\Sigma(2455)\pi$ are the only strong decays allowed to an excited Λ_c^+ having this mass.

| Mode | Fraction (Γ_i/Γ) | Confidence level |
|---|--------------------------------|------------------|
| Γ_1 $\Lambda_c^+ \pi^+ \pi^-$ | [a] $\approx 67\%$ | |
| Γ_2 $\Sigma_c(2455)^{++} \pi^-$ | <5 | 90% |
| Γ_3 $\Sigma_c(2455)^0 \pi^+$ | <5 | 90% |
| Γ_4 $\Lambda_c^+ \pi^+ \pi^-$ 3-body | large | |
| Γ_5 $\Lambda_c^+ \pi^0$ | [b] not seen | |
| Γ_6 $\Lambda_c^+ \gamma$ | not seen | |

[a] Assuming isospin conservation, so that the other third is $\Lambda_c^+ \pi^0 \pi^0$.

[b] A test that the isospin is indeed 0, so that the particle is indeed a Λ_c^+ .

$\Lambda_c(2625)^+$ BRANCHING RATIOS

| $\Gamma(\Sigma_c(2455)^{++} \pi^-)/\Gamma(\Lambda_c^+ \pi^+ \pi^-)$ | | | | | Γ_2/Γ_1 |
|---|-----|-------------|---------|---------------------------|---------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| <0.08 | 90 | EDWARDS | 95 CLE2 | $e^+e^- \approx 10.5$ GeV | |

| $\Gamma(\Sigma_c(2455)^0 \pi^+)/\Gamma(\Lambda_c^+ \pi^+ \pi^-)$ | | | | | Γ_3/Γ_1 |
|--|-----|-------------|---------|---------------------------|---------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| <0.07 | 90 | EDWARDS | 95 CLE2 | $e^+e^- \approx 10.5$ GeV | |

| $[\Gamma(\Sigma_c(2455)^{++} \pi^-) + \Gamma(\Sigma_c(2455)^0 \pi^+)]/\Gamma(\Lambda_c^+ \pi^+ \pi^-)$ | | | | | $(\Gamma_2 + \Gamma_3)/\Gamma_1$ |
|--|-----|------|-------------|------|----------------------------------|
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------|----|----|----------|---------|---|
| <0.36 | 90 | | FRABETTI | 94 E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |
| 0.46 ± 0.14 | | 21 | ALBRECHT | 93F ARG | $e^+e^- \approx \Upsilon(4S)$ |

| $\Gamma(\Lambda_c^+ \pi^+ \pi^- \text{ 3-body})/\Gamma(\Lambda_c^+ \pi^+ \pi^-)$ | | | | | Γ_4/Γ_1 |
|--|-----|------|-------------|------|---------------------|
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------|--|----|----------|---------|-------------------------------|
| 0.54 ± 0.14 | | 16 | ALBRECHT | 93F ARG | $e^+e^- \approx \Upsilon(4S)$ |
|-------------|--|----|----------|---------|-------------------------------|

$\Gamma(\Lambda_c^+ \pi^0)/\Gamma(\Lambda_c^+ \pi^+ \pi^-)$ Γ_5/Γ_1

$\Lambda_c^+ \pi^0$ decay is forbidden by isospin conservation if this state is in fact a Λ_c .

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|------------|--------------------|-------------|----------------------------|
| <0.91 | 90 | EDWARDS | 95 CLE2 | $e^+ e^- \approx 10.5$ GeV |

$\Gamma(\Lambda_c^+ \gamma)/\Gamma(\Lambda_c^+ \pi^+ \pi^-)$ Γ_6/Γ_1

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|------------|--------------------|-------------|----------------------------|
| <0.52 | 90 | EDWARDS | 95 CLE2 | $e^+ e^- \approx 10.5$ GeV |

$\Lambda_c(2625)^+$ REFERENCES

| | | | | |
|----------|-----|---------------|-----------------------------|---------------------|
| AALTONEN | 11H | PR D84 012003 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ALBRECHT | 97 | PL B402 207 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| EDWARDS | 95 | PRL 74 3331 | K.W. Edwards <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 94 | PRL 72 961 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| ALBRECHT | 93F | PL B317 227 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |