

# ϕ(2170)

$$I^G(J^{PC}) = 0^-(1^{--})$$

Observed by AUBERT, BE 06D in the initial-state radiation process  
 $e^+e^- \rightarrow \phi f_0(980)\gamma$ .


## ϕ(2170) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2175 ± 15 OUR AVERAGE</b> Error includes scale factor of 1.6. See the ideogram below.				
2186 ± 10 ± 6	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$
2125 ± 22 ± 10	483	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi \eta \gamma$
2175 ± 10 ± 15	201	<sup>1</sup> AUBERT, BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^- \pi \pi \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2079 ± 13 <sup>+79</sup> <sub>-28</sub>	4.8k	<sup>2</sup> SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^- \pi^+ \pi^- \gamma$
2192 ± 14	116 ± 95	<sup>3</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^- \pi^+ \pi^- \gamma$
2169 ± 20	149 ± 36	<sup>3</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^- \pi^0 \pi^0 \gamma$

<sup>1</sup> From the  $\phi f_0(980)$  component.

<sup>2</sup> From a fit with two incoherent Breit-Wigners.

<sup>3</sup> From the  $K^+K^- f_0(980)$  component.



## $\phi(2170)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>61 ± 18 OUR AVERAGE</b>				
65 ± 23 ± 17	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$
61 ± 50 ± 13	483	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \eta \gamma$
58 ± 16 ± 20	201	<sup>4</sup> AUBERT, BE	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi \pi \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
192 ± 23 <sup>+25</sup> <sub>-61</sub>	4.8k	<sup>5</sup> SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
71 ± 21	116 ± 95	<sup>6</sup> AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
102 ± 27	149 ± 36	<sup>6</sup> AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$
<sup>4</sup> From the $\phi f_0(980)$ component.				
<sup>5</sup> From a fit with two incoherent Breit-Wigners.				
<sup>6</sup> From the $K^+ K^- f_0(980)$ component.				

## $\phi(2170)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $e^+ e^-$	seen
$\Gamma_2$ $\phi \eta$	
$\Gamma_3$ $\phi \pi \pi$	
$\Gamma_4$ $\phi f_0(980)$	seen
$\Gamma_5$ $K^+ K^- \pi^+ \pi^-$	
$\Gamma_6$ $K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-$	seen
$\Gamma_7$ $K^+ K^- \pi^0 \pi^0$	
$\Gamma_8$ $K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$	seen
$\Gamma_9$ $K^{*0} K^\pm \pi^\mp$	not seen
$\Gamma_{10}$ $K^*(892)^0 \bar{K}^*(892)^0$	not seen

## $\phi(2170)$ $\Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\phi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_2 \Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.7 ± 0.7 ± 1.3	483	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \eta \gamma$	
$\Gamma(\phi f_0(980)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_4 \Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>2.5 ± 0.8 ± 0.4</b>	201	<sup>7</sup> AUBERT, BE	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi \pi \gamma$	
<sup>7</sup> From the $\phi f_0(980)$ component.					

### $\phi(2170) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

$$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_3/\Gamma \times \Gamma_1/\Gamma$$

VALUE (units  $10^{-7}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.65 ± 0.15 ± 0.18    4.8k    <sup>8</sup> SHEN    09 BELL    10.6  $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$   
<sup>8</sup> Multiplied by 3/2 to take into account the  $\phi\pi^0\pi^0$  mode. Using  $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$ .

### $\phi(2170)$ BRANCHING RATIOS

$$\Gamma(K^+K^-f_0(980) \rightarrow K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}} \qquad \Gamma_6/\Gamma$$

VALUE    DOCUMENT ID    TECN    COMMENT  
**seen**    AUBERT    07AK BABR    10.6  $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

$$\Gamma(K^+K^-f_0(980) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}} \qquad \Gamma_8/\Gamma$$

VALUE    DOCUMENT ID    TECN    COMMENT  
**seen**    AUBERT    07AK BABR    10.6  $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$

$$\Gamma(K^{*0}K^\pm\pi^\mp)/\Gamma_{\text{total}} \qquad \Gamma_9/\Gamma$$

VALUE    DOCUMENT ID    TECN    COMMENT  
**not seen**    AUBERT    07AK BABR    10.6 GeV  $e^+e^-$

$$\Gamma(K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}} \qquad \Gamma_{10}/\Gamma$$

VALUE    DOCUMENT ID    TECN    COMMENT  
**not seen**    ABLIKIM    10C BES2     $J/\psi \rightarrow \eta K^+\pi^-K^-\pi^+$

### $\phi(2170)$ REFERENCES

ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT, BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)