

$\Sigma(1580) 3/2^-$  $I(J^P) = 1(\frac{3}{2}^-)$  Status: \*

## OMITTED FROM SUMMARY TABLE

Seen in the isospin-1  $\bar{K}N$  cross section at BNL (LI 73, CARROLL 76) and in a partial-wave analysis of  $K^- p \rightarrow \Lambda\pi^0$  for c.m. energies 1560–1600 MeV by LITCHFIELD 74. LITCHFIELD 74 finds  $J^P = 3/2^-$ . Not seen by ENGLER 78 or by CAMERON 78C (with larger statistics in  $K_L^0 p \rightarrow \Lambda\pi^+$  and  $\Sigma^0\pi^+$ ).

Neither OLMSTED 04 (in  $K^- p \rightarrow \Lambda\pi^0$ ) nor PRAKHOV 04 (in  $K^- p \rightarrow \Lambda\pi^0\pi^0$ ) see any evidence for this state.

 $\Sigma(1580)$  MASS

| VALUE (MeV)                                   | DOCUMENT ID                | TECN | COMMENT                          |
|---|----------------------------|------|----------------------------------|
| <b><math>\approx 1580</math> OUR ESTIMATE</b> |                            |      |                                  |
| 1583 $\pm$ 4                                  | <sup>1</sup> CARROLL 76    | DPWA | Isospin-1 total $\sigma$         |
| 1582 $\pm$ 4                                  | <sup>2</sup> LITCHFIELD 74 | DPWA | $K^- p \rightarrow \Lambda\pi^0$ |

 $\Sigma(1580)$  WIDTH

| VALUE (MeV) | DOCUMENT ID                | TECN | COMMENT                          |
|-------------|----------------------------|------|----------------------------------|
| 15          | <sup>1</sup> CARROLL 76    | DPWA | Isospin-1 total $\sigma$         |
| 11 $\pm$ 4  | <sup>2</sup> LITCHFIELD 74 | DPWA | $K^- p \rightarrow \Lambda\pi^0$ |

 $\Sigma(1580)$  DECAY MODES

| Mode                    |
|-------------------------|
| $\Gamma_1$ $N\bar{K}$   |
| $\Gamma_2$ $\Lambda\pi$ |
| $\Gamma_3$ $\Sigma\pi$  |

 $\Sigma(1580)$  BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on  $\Lambda$  and  $\Sigma$  Resonances.

| $\Gamma(N\bar{K})/\Gamma_{\text{total}}$ | DOCUMENT ID                | TECN | COMMENT                 | $\Gamma_1/\Gamma$ |
|--|----------------------------|------|-------------------------|-------------------|
| VALUE                                    |                            |      |                         |                   |
| +0.03 $\pm$ 0.01                         | <sup>2</sup> LITCHFIELD 74 | DPWA | $\bar{K}N$ multichannel |                   |

| $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Lambda\pi$ | DOCUMENT ID                | TECN | COMMENT                            | $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$ |
|--|----------------------------|------|------------------------------------|-----------------------------------|
| VALUE  |                            |      |                                    |                                   |
| not seen   | CAMERON 78C                | HBC  | $K_L^0 p \rightarrow \Lambda\pi^+$ |                                   |
| not seen   | ENGLER 78                  | HBC  | $K_L^0 p \rightarrow \Lambda\pi^+$ |                                   |
| +0.10 $\pm$ 0.02   | <sup>2</sup> LITCHFIELD 74 | DPWA | $K^- p \rightarrow \Lambda\pi^0$   |                                   |

| $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Sigma\pi$ |                         |      |         | $(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$ |
|--|-------------------------|------|---------|--------------------------------------|
| VALUE  | DOCUMENT ID             | TECN | COMMENT |                                      |
| not seen   | CAMERON                 | 78C  | HBC     | $K_L^0 p \rightarrow \Sigma^0 \pi^+$ |
| not seen   | ENGLER                  | 78   | HBC     | $K_L^0 p \rightarrow \Sigma^0 \pi^+$ |
| $+0.03 \pm 0.04$   | <sup>2</sup> LITCHFIELD | 74   | DPWA    | $\bar{K}N$ multichannel              |

### $\Sigma(1580)$ FOOTNOTES

<sup>1</sup> CARROLL 76 sees a total-cross-section bump with  $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}} = 0.06$ .

<sup>2</sup> The main effect observed by LITCHFIELD 74 is in the  $\Lambda\pi$  final state; the  $\bar{K}N$  and  $\Sigma\pi$  couplings are estimated from a multichannel fit including total-cross-section data of LI 73.

### $\Sigma(1580)$ REFERENCES

|            |     |                  |                            |                            |
|------------|-----|------------------|----------------------------|----------------------------|
| OLMSTED    | 04  | PL B588 29       | J. Olmsted <i>et al.</i>   | (BNL Crystal Ball Collab.) |
| PRAKHOV    | 04  | PR C69 042202    | S. Prakhov <i>et al.</i>   | (BNL Crystal Ball Collab.) |
| CAMERON    | 78C | NP B132 189      | W. Cameron <i>et al.</i>   | (BGNA, EDIN, GLAS+) I      |
| ENGLER     | 78  | PR D18 3061      | A. Engler <i>et al.</i>    | (CMU, ANL)                 |
| CARROLL    | 76  | PRL 37 806       | A.S. Carroll <i>et al.</i> | (BNL) I                    |
| LITCHFIELD | 74  | PL 51B 509       | P.J. Litchfield            | (CERN) IJP                 |
| LI         | 73  | Purdue Conf. 283 | K.K. Li                    | (BNL) I                    |