

A Note on Simultaneous Diophantine Approximation in Positive Characteristic

Michael Fuchs

Abstract

In a recent paper, Inoue and Nakada proved a 0-1 law and a strong law of large numbers with error term for the number of coprime solutions of the one-dimensional Diophantine approximation problem in the field of formal Laurent series over a finite base field. In this note, we generalize their results to higher dimensions.

1 Introduction

Let \mathbb{F}_q be a finite field with q elements and denote by $\mathbb{F}_q((T^{-1}))$ the field of formal Laurent series. For $f \in \mathbb{F}_q((T^{-1}))$ let $|f| = q^{\deg f}$ be the valuation induced by the generalized degree function. Set

$$\mathbb{L} = \{f \in \mathbb{F}_q((T^{-1})) : |f| < 1\}.$$

Then, with the restriction of $|\cdot|$ to \mathbb{L} , \mathbb{L} is a compact topological group. Hence, there exists a (unique) translation-invariant probability measure which will be denoted by m .

We are interested in the Diophantine approximation problem

$$\left| f - \frac{P}{Q} \right| < \frac{1}{q^{n+l_n}}, \quad \deg Q = n, \quad Q \text{ monic}, \quad (P, Q) = 1, \quad (1)$$

where $f \in \mathbb{L}$, $P, Q \in \mathbb{F}_q[T]$ with $Q \neq 0$, and l_n is a sequence of non-negative integers (subsequently, we will use (\cdot, \cdot) to denote the gcd, whereas $\langle \cdot, \cdot \rangle$ will be used for pairs).

Concerning the number of solutions of (1), Inoue and Nakada [5] proved the following 0-1 law: the number of solutions is either finite or infinite for almost all $f \in \mathbb{L}$, the latter holding if and only if

$$\sum_{n=0}^{\infty} q^{n-l_n} = \infty.$$

Moreover, the method of proof in [5] also gives a quantitative result under one additional assumption on l_n : if $l_n \geq n$, then the number of solutions of (1) with $\deg Q \leq N$ is given by

$$(1 - q^{-1}) \Psi(N) + \mathcal{O} \left(\Psi(N)^{1/2} (\log \Psi(N))^{3/2+\epsilon} \right),$$

where $\epsilon > 0$ is an arbitrary small constant and $\Psi(N) := \sum_{n \leq N} q^{n-l_n}$.

Key words: formal Laurent series, simultaneous Diophantine approximation, 0-1 law, strong law of large numbers.
2010 Mathematics Subject Classification: 11J61, 11J83, 11K60.

The purpose of this note is to prove generalizations of the above two results to multidimensional Diophantine approximation. Therefore, consider

$$\left| f_j - \frac{P_j}{Q} \right| < \frac{1}{q^{n+l_n^{(j)}}}, \deg Q = n, Q \text{ monic}, (P_j, Q) = 1, j = 1, \dots, d, \quad (2)$$

where $(f_1, \dots, f_d) \in \mathbb{L} \times \dots \times \mathbb{L}$, $P_j, j = 1, \dots, d, Q \in \mathbb{F}_q[T]$ with $Q \neq 0$, and $l_n^{(j)}, j = 1, \dots, d$ are sequences of non-negative integers. Moreover, set $l_n := \sum_{j=1}^d l_n^{(j)}$.

Then, the first result above has the following extension to the multidimensional setting.

Theorem 1. *The number of solutions of (2) is either finite or infinite for almost all $(f_1, \dots, f_d) \in \mathbb{L} \times \dots \times \mathbb{L}$, the latter holding if and only if*

$$\sum_{n=0}^{\infty} q^{n-l_n} = \infty. \quad (3)$$

Moreover, also the second result admits an extension to higher dimensions.

Theorem 2. *Assume that $l_n \geq n$. Then, for almost all (f_1, \dots, f_d) , the number of solutions of (2) with $\deg Q \leq N$ is given by*

$$c_0 \Psi(N) + \mathcal{O}\left(\Psi(N)^{1/2+\epsilon}\right),$$

where $\epsilon > 0$ is an arbitrary small constant, $\Psi(N) := \sum_{n \leq N} q^{n-l_n}$, and

$$c_0 := \sum_{Q_1 \text{ monic}} \dots \sum_{Q_d \text{ monic}} \frac{\mu(Q_1)}{|Q_1|} \dots \frac{\mu(Q_d)}{|Q_d|} \frac{1}{|\text{lcm}(Q_1, \dots, Q_d)|} > 0.$$

Here, $\mu(\cdot)$ denotes the Moebius μ function on $\mathbb{F}_q[T]$.

Remark 1. The constant c_0 will already appear in the proof of Theorem 1. In particular, we will show the claim about the positivity already in the next section (see the proof of Lemma 1 below).

Remark 2. Observe that the error term in the above result for $d = 1$ is weaker than the corresponding one in the result of Inoue and Nakada. The reason for this is that our method is completely different from the approach used by the latter two authors (it is not obvious how to generalize their approach to higher dimensions).

Notation. We will use $[D_1, \dots, D_d]$ to denote the lcm of the polynomials D_1, \dots, D_d . All sums will be over monic polynomials. Logarithms in this paper just take on values ≥ 1 , i.e. $\log_a x$ should be interpreted as $\max\{\log_a x, 1\}$. We will use both Landau's notation $f(x) = \mathcal{O}(g(x))$ as well as Vinogradov's notation $f(x) \ll g(x)$. Finally, ϵ will denote an arbitrary small positive number whose value might change from one appearance to the next.

2 Proof of Theorem 1

First, note that the necessity of (3) for the number of solutions of (2) being infinite follows from a standard application of the Borel-Cantelli lemma. Hence, we only have to focus on the sufficiency part. For this purpose, we use a slight extension of the d -dimensional Duffin-Schaeffer theorem for formal Laurent series due to Inoue [4].

Theorem 3 (Inoue). *Consider*

$$\left| f_j - \frac{P_j}{Q} \right| < \frac{1}{q^{n+l_Q^{(j)}}}, \deg Q = n, Q \text{ monic}, (P_j, Q) = 1, j = 1, \dots, d, \quad (4)$$

where $(f_1, \dots, f_d) \in \mathbb{L} \times \dots \times \mathbb{L}$, $P_j, j = 1, \dots, d, Q$ with $Q \neq 0$, and $l_Q^{(j)}, j = 1, \dots, d$ are sequences of non-negative integers. Assume that

$$\sum_Q q^{-l_Q^{(1)} - \dots - l_Q^{(j)}} = \infty$$

and that for infinitely many N

$$\sum_{\deg Q \leq N} q^{-l_Q^{(1)} - \dots - l_Q^{(j)}} < C \sum_{\deg Q \leq N} q^{-l_Q^{(1)} - \dots - l_Q^{(j)}} \varphi(Q)^d / |Q|^d,$$

where C is some positive constant. Then, (4) has infinitely many solutions for almost all $(f_1, \dots, f_d) \in \mathbb{L} \times \dots \times \mathbb{L}$.

Remark 3. Note that the result in [4] is just stated for the special case $l_Q^{(1)} = \dots = l_Q^{(d)}$. An inspection of the proof, however, shows that the result continues to hold for different approximation functions in every coordinate.

Before we can apply this result, we need a technical lemma.

Lemma 1. *We have,*

$$\sum_{\deg Q=n} \varphi(Q)^d = c_0 q^{n(d+1)} + \mathcal{O}\left(q^{n(d+\epsilon)}\right),$$

where c_0 is as in Theorem 2 and $\varphi(\cdot)$ is Euler's totient function.

Proof. Note that

$$\begin{aligned} \sum_{\deg Q=n} \varphi(Q)^d &= q^{nd} \sum_{\deg Q=n} \left(\sum_{D|Q} \frac{\mu(D)}{|D|} \right)^d \\ &= q^{nd} \sum_{\deg Q=n} \sum_{D_1|Q} \dots \sum_{D_d|Q} \frac{\mu(D_1)}{|D_1|} \dots \frac{\mu(D_d)}{|D_d|} \\ &= q^{nd} \sum_{\deg D_1 \leq n} \dots \sum_{\deg D_d \leq n} \frac{\mu(D_1)}{|D_1|} \dots \frac{\mu(D_d)}{|D_d|} \sum_{[D_1, \dots, D_d] | Q, \deg Q=n} 1. \end{aligned}$$

The latter sum becomes

$$\sum_{[D_1, \dots, D_d] | Q, \deg Q=n} 1 = \begin{cases} q^n / |[D_1, \dots, D_d]|, & \text{if } \deg[D_1, \dots, D_d] \leq n; \\ 0, & \text{otherwise.} \end{cases}$$

Consequently,

$$\begin{aligned} \sum_{\deg Q=n} \varphi(Q)^d &= q^{n(d+1)} \sum_{\deg D_1 \leq n} \dots \sum_{\deg D_d \leq n} \frac{\mu(D_1)}{|D_1|} \dots \frac{\mu(D_d)}{|D_d|} \frac{1}{|[D_1, \dots, D_d]|} \\ &\quad + \mathcal{O}\left(q^{nd} \left(\sum_{\deg D \leq n} \frac{1}{|D|} \right)^d\right) \\ &= q^{n(d+1)} \sum_{\deg D_1 \leq n} \dots \sum_{\deg D_d \leq n} \frac{\mu(D_1)}{|D_1|} \dots \frac{\mu(D_d)}{|D_d|} \frac{1}{|[D_1, \dots, D_d]|} + \mathcal{O}\left(n^d q^{nd}\right). \quad (5) \end{aligned}$$

Next, observe that

$$\begin{aligned} & \left| \sum_{\deg D_1 \leq n} \cdots \sum_{\deg D_d \leq n} \frac{\mu(D_1)}{|D_1|} \cdots \frac{\mu(D_d)}{|D_d|} \frac{1}{|[D_1, \dots, D_d]|} - c_0 \right| \\ & \leq \sum_{\deg D > n} \sum_{[D_1, \dots, D_d] = D} \frac{1}{|D_1 \cdots D_d| \cdot |D|} \leq \sum_{\deg D > n} \frac{\omega(D)^d}{|D|^2}, \end{aligned}$$

where $\omega(D)$ denotes the number of monic divisors of D . Since $\omega(D) = \mathcal{O}(|D|^\epsilon)$ for arbitrary small $\epsilon > 0$ (this is proved as for integers; see page 296 in [1]), we obtain

$$\sum_{\deg D > n} \frac{\omega(D)^d}{|D|^2} \ll \sum_{l=n+1}^{\infty} q^{l(\epsilon d - 1)} \ll q^{n(\epsilon d - 1)}.$$

So, we have

$$\sum_{\deg D_1 \leq n} \cdots \sum_{\deg D_d \leq n} \frac{\mu(D_1)}{|D_1|} \cdots \frac{\mu(D_d)}{|D_d|} \frac{1}{|[D_1, \dots, D_d]|} = c_0 + \mathcal{O}\left(q^{n(-1+\epsilon)}\right).$$

Plugging this into (5) yields the claimed expansion.

What is left to show is that $c_0 > 0$. Therefore, observe that

$$\sum_{\deg Q = n} \varphi(Q)^d \geq \sum_{\deg I = n} \varphi(I)^d = (q^n - 1)^d \sum_{\deg I = n} 1 \gg (q^n - 1)^d q^n / n,$$

where the second and third sum runs over all irreducible polynomials and the last bound is well-known. Hence, $c_0 > 0$ as claimed. ■

Remark 4. For $d = 1$, note that

$$c_0 = \sum_Q \frac{\mu(Q)}{|Q|^2} = \prod_I \left(1 - \frac{1}{|I|^2}\right) = \left(\sum_Q \frac{1}{|Q|^2}\right)^{-1} = 1 - \frac{1}{q}.$$

In this situation even more is known, namely,

$$\sum_{\deg Q = n} \varphi(Q) = \left(1 - \frac{1}{q}\right) q^{2n}.$$

For a proof of the latter claim e.g. see [5].

Now, we can prove our first main result.

Proof of Theorem 1. As already mentioned before, we only have to show that (3) is sufficient for the number of solutions of (2) being infinity. For this purpose, we just have to check the two conditions in Inoue's result. First, note that since $l_Q^{(1)} + \cdots + l_Q^{(d)} = l_{\deg Q}$, we have

$$\sum_{\deg Q \leq N} q^{-l_{\deg Q}} = \sum_{n \leq N} q^{n-l_n}$$

and

$$\sum_{\deg Q \leq N} q^{-l_{\deg Q}} \varphi(Q)^d / |Q|^d = \sum_{n \leq N} q^{-nd-l_n} \sum_{\deg Q = n} \varphi(Q)^d = c_0 \sum_{n \leq N} q^{n-l_n} + \mathcal{O}\left(\sum_{n \leq N} q^{\epsilon n-l_n}\right).$$

Moreover, by Cauchy's inequality

$$\sum_{n \leq N} q^{\epsilon n-l_n} \ll \left(\sum_{n \leq N} q^{n-l_n}\right)^{1/2}.$$

Hence, both conditions are satisfied and our result follows from Inoue's result. ■

3 Proof of Theorem 2

We start with a technical lemma.

Lemma 2. *We have,*

$$\sum_{\deg(D_1), \dots, \deg(D_d) \leq n} \frac{1}{|[D_1, \dots, D_d]|} \ll q^{n\epsilon}.$$

Proof. First note that

$$\begin{aligned} \sum_{\deg(D_1), \dots, \deg(D_d) \leq n} \frac{1}{|[D_1, \dots, D_d]|} &\leq \sum_{\deg(D_1), \dots, \deg(D_d) \leq n} \frac{1}{|[D_1, \dots, D_d]|^{1-\epsilon}} \\ &\leq \sum_{\deg(D_1), \dots, \deg(D_d) \leq n} \frac{|([D_1, \dots, D_{d-1}], D_d)|^{1-\epsilon}}{|[D_1, \dots, D_{d-1}]|^{1-\epsilon} \cdot |D_d|^{1-\epsilon}}. \end{aligned}$$

Next we change the order of summation and obtain

$$\begin{aligned} &\sum_{\deg(D_1), \dots, \deg(D_d) \leq n} \frac{1}{|[D_1, \dots, D_d]|^{1-\epsilon}} \\ &\leq \sum_{\deg D \leq n} \sum_{D|[D_1, \dots, D_{d-1}], \deg D_i \leq n} \frac{1}{|[D_1, \dots, D_{d-1}]|^{1-\epsilon}} \sum_{D|D_d, \deg D_d \leq n} \left(\frac{|D|}{|D_d|} \right)^{1-\epsilon} \\ &\leq \sum_{\deg D \leq n} \sum_{D|[D_1, \dots, D_{d-1}], \deg D_i \leq n} \frac{1}{|[D_1, \dots, D_{d-1}]|^{1-\epsilon}} \sum_{\deg Q \leq n} \frac{1}{|Q|^{1-\epsilon}} \\ &\ll q^{n\epsilon} \sum_{\deg(D_1), \dots, \deg(D_{d-1}) \leq n} \frac{1}{|[D_1, \dots, D_{d-1}]|^{1-\epsilon}} \sum_{D|[D_1, \dots, D_{d-1}]} 1 \\ &= q^{n\epsilon} \sum_{\deg(D_1), \dots, \deg(D_{d-1}) \leq n} \frac{\omega([D_1, \dots, D_{d-1}])}{|[D_1, \dots, D_{d-1}]|^{1-\epsilon}}. \end{aligned}$$

Now, as before, we use the estimate $\omega(D) = \mathcal{O}(|D|^\epsilon)$ for all sufficiently small ϵ . Hence,

$$\sum_{\deg(D_1), \dots, \deg(D_d) \leq n} \frac{1}{|[D_1, \dots, D_d]|^{1-\epsilon}} \ll q^{n\epsilon} \sum_{\deg(D_1), \dots, \deg(D_{d-1}) \leq n} \frac{1}{|[D_1, \dots, D_d]|^{1-2\epsilon}}.$$

Iterating this result proves the claim. \blacksquare

Now, we turn to the proof of Theorem 2. For this purpose, we extend an approach due to Harman (see proof of Theorem 4.4 starting on page 109 in [3]) to higher dimensions.

We first need some notation. Let $\Gamma_1(N) = \lfloor \log_q \Psi(N)^2 \rfloor$ and $\Gamma_2(N) = \lfloor \log_q \Psi(N)^4 \rfloor$. Moreover, consider the following approximation problem

$$\left| f_j - \frac{P_j}{Q} \right| < \frac{1}{q^{n+l_n^{(j)}}}, \quad \deg Q = n, \quad Q \text{ monic}, \quad D_j | (P_j, Q), \quad \deg(P_j, Q) \leq \Gamma_2(N), \quad j = 1, \dots, d, \quad (6)$$

where D_1, \dots, D_d are fixed monic polynomials. For fixed (f_1, \dots, f_d) and Q denote by $s(Q; D_1, \dots, D_d)$ the number of solutions of (6).

We gather some properties of $s(Q; D_1, \dots, D_d)$ needed below.

Lemma 3. *We have,*

$$\mathbb{E} \left(\sum_{M_1 < n \leq M_2} \sum_{\deg Q = n, [D_1, \dots, D_d] | Q} s(Q; D_1, \dots, D_d) \right) \ll \frac{1}{|D_1 \cdots D_d| \cdot |[D_1, \dots, D_d]|} \sum_{M_1 < n \leq M_2} q^{n-l_n}$$

and

$$\mathbb{E} \left(\sum_{M_1 < n \leq M_2} \sum_{\deg Q=n, [D_1, \dots, D_d] | Q} \left(s(Q; D_1, \dots, D_d) - \frac{1}{|D_1 \cdots D_d|} \cdot \frac{1}{q^{l_n}} \right) \right)^2 \ll \frac{\Gamma_2(N)}{|D_1 \cdots D_d| \cdot |[D_1, \dots, D_d]|} \sum_{M_1 < n \leq M_2} q^{n-l_n}$$

for all $M_1 \leq M_2$.

Proof. Both properties are easy extensions of the corresponding properties from the case $d = 1$ (see Proposition 3 and Proposition 4 in [2]). For the reader's convenience, we recall the proof of the first property.

Therefore, observe that $s(Q; D_1, \dots, D_d) \leq s^*(Q; D_1, \dots, D_d)$ where the latter denotes the number of solutions of (6) with the upper bound on the gcd removed. Of course, $s^*(Q; D_1, \dots, D_d) = 0$ if $[D_1, \dots, D_d] \nmid Q$.

Now, for $[D_1, \dots, D_d] | Q$, note that $s^*(Q; D_1, \dots, D_d) = \mathbf{1}_A$ ($\mathbf{1}_A$ denotes an indicator random variable) with

$$A = \bigcup_{P_j | D_j, \deg P_j < n, 1 \leq j \leq d} B(P_1/Q; q^{-n-l_n^{(1)}}) \times \cdots \times B(P_d/Q; q^{-n-l_n^{(d)}}),$$

where $B(f; q^{-n})$ denotes the open ball with center f and radius q^{-n} and the above union is disjoint. Since

$$(m \times \cdots \times m) \left(B(P_1/Q; q^{-n-l_n^{(1)}}) \times \cdots \times B(P_d/Q; q^{-n-l_n^{(d)}}) \right) = q^{-dn-l_n}$$

and consequently

$$m(A) = \frac{1}{|D_1 \cdots D_d|} q^{-l_n},$$

the result follows from elementary properties of the mean. \blacksquare

Next, we prove the following proposition for the number of solutions of (6).

Proposition 1. *For almost all (f_1, \dots, f_d) , the number of solutions of (6) with $\deg Q \leq N$ is given by*

$$\frac{1}{|D_1 \cdots D_d| \cdot |[D_1, \dots, D_d]|} \Psi(N) + E(N; D_1, \dots, D_d),$$

where the second term satisfies

$$\sum_{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_1(N)} E(N; D_1, \dots, D_d) = \mathcal{O} \left(\Psi(N)^{1/2+\epsilon} \right)$$

with $\epsilon > 0$ an arbitrary small constant.

Proof. First note that it suffices to prove our claim for the case where $\Psi(N) \rightarrow \infty$ as $N \rightarrow \infty$ (otherwise, the result is an easy consequence of the Borel-Cantelli lemma). Next, denote by N_k the largest integer with $\Psi(N_k) < k$. It is easy to see that we only have to prove the result for the subsequence N_k .

We are going to need some notation. First, put

$$k = \sum_{j=0}^l a_j 2^j, \quad a_l \neq 0, \quad a_j \in \{0, 1\} \forall j.$$

Define the following set

$$S(k) = \left\{ (i, m) : a_i = 1, m = \sum_{j=i+1}^l a_j 2^{j-i} \right\}.$$

Moreover, denote by

$$u_t = u_t(i, m) = \max \{ n \in \mathbb{N} : \Psi(n) < (m + t)2^i \},$$

where $t \in \{0, 1\}$. Finally, with the notation of Lemma 3, we put

$$E(i, m; D_1, \dots, D_d) = \sum_{u_0 < n \leq u_1} \sum_{\deg Q = n, [D_1, \dots, D_d] | Q} \left(s(Q; D_1, \dots, D_d) - \frac{1}{|D_1 \cdots D_d|} \cdot \frac{1}{q^{l_n}} \right).$$

Then, we obviously have

$$E(N_k; D_1, \dots, D_d) = \sum_{(i, m) \in S(k)} E(i, m; D_1, \dots, D_d).$$

Now, set

$$E(l) := \sum_{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_1(N_{2^{l+1}})} |D_1 \cdots D_d| \sum_{0 \leq i \leq l, m < 2^{l-i+1}} E(i, m; D_1, \dots, D_d)^2.$$

Then, with the estimate from Lemma 3

$$\mathbb{E}E(i, m; D_1, \dots, D_d)^2 \ll \frac{\Gamma_2(N_{2^{l+1}})}{|D_1 \cdots D_d| \cdot |[D_1, \dots, D_d]|} \sum_{u_0 < n \leq u_1} q^{n-l_n},$$

we obtain

$$E(l) \ll 2^l l^2 \sum_{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_1(N_{2^{l+1}})} \frac{1}{|[D_1, \dots, D_d]|} \ll 2^{l(1+\bar{\epsilon})},$$

where the last step follows from Lemma 2 and $\bar{\epsilon}$ will be chosen below. This in turn implies that

$$P \left(E(l) \geq 2^{l(1+\epsilon)} \right) \ll \frac{1}{2^{l(\epsilon-\bar{\epsilon})}},$$

where we choose $\bar{\epsilon} < \epsilon$. Hence, the Borel-Cantelli lemma yields that

$$E(l) < 2^{l(1+\epsilon)}, \quad \text{a.s.}$$

for l large enough.

Finally consider

$$\begin{aligned} & \sum_{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_1(N_k)} E(N_k; D_1, \dots, D_d) \\ & \leq \left(\sum_{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_1(N_k)} \frac{1}{|D_1 \cdots D_d|} \sum_{(i, m) \in S(k)} 1 \right)^{1/2} \cdot (E(r))^{1/2} \\ & \ll 2^{l(1/2+\epsilon)} l^{d+1} \ll 2^{l(1/2+\epsilon)}. \end{aligned}$$

From this the assertion follows. \blacksquare

Now, we can prove our second main result.

Proof of Theorem 2. As in the proof of the proposition, we can assume w.l.o.g. that $\Psi(N) \rightarrow \infty$ as $N \rightarrow \infty$. Then, we again choose N_k as the largest integer with $\Psi(N_k) < k$. As before, it is easy to see that it suffices to prove our claim for the sequence N_k .

Next, we introduce the notation $S(N_k; D_1, \dots, D_d)$ for the number of solutions of (6) with $\deg Q \leq N_k$ (here, (f_1, \dots, f_d) is fixed). Then, by an inclusion-exclusion argument, the number of solutions of (2) with $\deg Q \leq N_k$ is given by

$$\sum_{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_2(N_k)} \mu(D_1) \cdots \mu(D_d) S(N_k; D_1, \dots, D_d),$$

where $\mu(\cdot)$ denotes the Moebius function. We split the sum into two parts A and B according to whether there is an D_i with $\deg D_i > \Gamma_1(N_k)$ or not, respectively.

First, we will consider A . Note that

$$\begin{aligned} \mathbb{E}|A| &\leq \sum_{\substack{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_2(N_k) \\ \deg D_i > \Gamma_1(N_k) \text{ for some } i}} \mathbb{E}S(N_k; D_1, \dots, D_d) \\ &\ll \Psi(N_k) \sum_{\substack{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_2(N_k) \\ \deg D_i > \Gamma_1(N_k) \text{ for some } i}} \frac{1}{|D_1 \cdots D_d|} \cdot \frac{1}{|[D_1, \dots, D_d]|} \\ &\ll \Psi(N_k) \left(\sum_{\deg D_1 > \Gamma_1(N_k)} \frac{1}{|D_1|^2} \right) \cdot \left(\sum_{\deg D_2 \leq \Gamma_2(N_k)} \frac{1}{|D_2|} \right) \cdots \left(\sum_{\deg D_d \leq \Gamma_2(N_k)} \frac{1}{|D_d|} \right) \\ &\ll \frac{(\log \Psi(N_k))^{d-1}}{\Psi(N_k)}, \end{aligned}$$

where we have used Lemma 3. Consequently,

$$P(|A| > (\log \Psi(N_k))^{d+1}) \ll \frac{1}{\Psi(N_k) (\log \Psi(N_k))^2} \ll \frac{1}{k (\log k)^2}.$$

Hence, the Borel-Cantelli lemma implies that for almost all (f_1, \dots, f_d) ,

$$A = \mathcal{O}((\log \Psi(N_k))^{d+1}).$$

So, in view of our claimed result, the main contribution will come from B . Here, we can use the above proposition and obtain

$$B = \Psi(N_k) \sum_{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_1(N_k)} \frac{\mu(D_1) \cdots \mu(D_d)}{|D_1 \cdots D_d| \cdot |[D_1, \dots, D_d]|} + \mathcal{O}\left(\Psi(N_k)^{1/2+\epsilon}\right).$$

Now, as in the proof of Lemma 1

$$\sum_{\deg(D_1), \dots, \deg(D_d) \leq \Gamma_1(N_k)} \frac{\mu(D_1) \cdots \mu(D_d)}{|D_1 \cdots D_d| \cdot |[D_1, \dots, D_d]|} = c_0 + \Psi(N_k)^{\epsilon-2}.$$

Combining all the estimates proves the claimed result. \blacksquare

Acknowledgments

The author acknowledges partial support by National Science Council under the grant NSC-98-2115-M-009-009. Moreover, the authors thanks the anonymous referee for a careful reading.

References

- [1] T. M. Apostol (1976). Introduction to Analytic Number Theory, Springer-Verlag, New York.
- [2] M. Fuchs (2010). Metrical theorems for inhomogeneous Diophantine approximation in positive characteristic, *Acta Arith.*, **141**, 191-208.
- [3] G. Harman (1998). Metric Number Theory, London Mathematical Society Monographs (New Series) **18**, The Clarendon Press Oxford University Press, New York.
- [4] K. Inoue (2003). The metric simultaneous Diophantine approximations over formal power series, *J. Theor. Nombres Bordeaux*, **15**, 151V161.
- [5] K. Inoue and H. Nakada (2003). On metric Diophantine approximation in positive characteristic, *Acta Arith.*, **110**, 205-218.

Michael Fuchs
Department of Applied Mathematics
National Chiao Tung University
1001 Ta Hsue Road
Hsinchu, 300, Taiwan
E-mail: mfuchs@math.nctu.edu.tw