

## Propositional Logic Intro

Note 1

**Proposition:** A statement with a truth value; it is either true or false.

Propositions can be combined to form more complicated expressions, using the following operations:

| Operators              | Quantifiers            | Implication operations                     |
|------------------------|------------------------|--|
| $\wedge$ and           | $\forall$ for all      | Implication $P \Rightarrow Q$              |
| $\vee$ or              | $\exists$ there exists | Inverse $\neg P \Rightarrow \neg Q$        |
| $\neg$ not             |                        | Converse $Q \Rightarrow P$                 |
| $\Rightarrow$ implies  |                        | Contrapositive $\neg Q \Rightarrow \neg P$ |
| $\equiv$ equivalent to |                        |  |

Further, for an implication  $P \Rightarrow Q$  where  $P$  is the *hypothesis* and  $Q$  is the *conclusion*, it is useful to know that  $P \Rightarrow Q \equiv \neg P \vee Q$ . Additionally, observe that any implication is logically equivalent to its contrapositive.

**DeMorgan's Laws:** The following identities can be helpful when simplifying expressions and distributing negations.

- $\neg(P \wedge Q) \equiv \neg P \vee \neg Q$
- $\neg(P \vee Q) \equiv \neg P \wedge \neg Q$
- $\neg(\forall x P(x)) \equiv \exists x(\neg P(x))$
- $\neg(\exists x P(x)) \equiv \forall x(\neg P(x))$

## 1 Propositional Practice

Note 1 Convert the following English sentences into propositional logic and the following propositions into English. State whether or not each statement is true with brief justification.

Recall that  $\mathbb{R}$  is the set of reals,  $\mathbb{Q}$  is the set of rationals,  $\mathbb{Z}$  is the set of integers, and  $\mathbb{N}$  is the set of natural numbers. The notation " $a \mid b$ ", read as " $a$  divides  $b$ ", means that  $a$  is a divisor of  $b$ .

- There is a real number which is not rational.
- All integers are natural numbers or are negative, but not both.
- If a natural number is divisible by 6, it is divisible by 2 or it is divisible by 3.

(d)  $\neg(\forall x \in \mathbb{Q})(x \in \mathbb{Z})$   
 (e)  $(\forall x \in \mathbb{Z})(((2 \mid x) \vee (3 \mid x)) \implies (6 \mid x))$   
 (f)  $(\forall x \in \mathbb{N})((x > 7) \implies ((\exists a, b \in \mathbb{N}) (a + b = x)))$

**Solution:**

(a)  $(\exists x \in \mathbb{R}) (x \notin \mathbb{Q})$ , or equivalently  $(\exists x \in \mathbb{R}) \neg(x \in \mathbb{Q})$ . This is true, and we can use  $\pi$  as an example to prove it.

(b)  $(\forall x \in \mathbb{Z}) (((x \in \mathbb{N}) \vee (x < 0)) \wedge \neg((x \in \mathbb{N}) \wedge (x < 0)))$ . This is true, since we define the naturals to contain all integers which are not negative.

(c)  $(\forall x \in \mathbb{N}) ((6 \mid x) \implies ((2 \mid x) \vee (3 \mid x)))$ . This is true, since any number divisible by 6 can be written as  $6k = (2 \cdot 3)k = 2(3k)$ , meaning it must also be divisible by 2.

(d) It is not the case that all rational numbers are integers. This is true, since we have rational numbers like  $\frac{1}{2}$  that are not integers.

(e) Any integer that is divisible by 2 or 3 is also divisible by 6. This is false: 2 provides the easiest counterexample. Note that this statement is false even though its converse (part c) is true.

(f) If a natural number is larger than 7, it can be written as the sum of two other natural numbers. This is trivially true, since if  $x = a + b$  we can take  $a = x$  and  $b = 0$ .

(Aside: this is a reference to the very weak Goldbach Conjecture (<https://xkcd.com/1310/>)).

## 2 Truth Tables

**Note 1** Determine whether the following equivalences hold, by writing out truth tables. Clearly state whether or not each pair is equivalent.

(a)  $P \wedge (Q \vee P) \equiv P \wedge Q$   
 (b)  $(P \vee Q) \wedge R \equiv (P \wedge R) \vee (Q \wedge R)$   
 (c)  $(P \wedge Q) \vee R \equiv (P \vee R) \wedge (Q \vee R)$

**Solution:**

(a) Not equivalent.

| $P$ | $Q$ | $P \wedge (Q \vee P)$ | $P \wedge Q$ |
|-----|-----|-----------------------|--------------|
| T   | T   | T                     | T            |
| T   | F   | T                     | F            |
| F   | T   | F                     | F            |
| F   | F   | F                     | F            |

(b) Equivalent.

| $P$ | $Q$ | $R$ | $(P \vee Q) \wedge R$ | $(P \wedge R) \vee (Q \wedge R)$ |
|-----|-----|-----|-----------------------|----------------------------------|
| T   | T   | T   | T                     | T                                |
| T   | T   | F   | F                     | F                                |
| T   | F   | T   | T                     | T                                |
| T   | F   | F   | F                     | F                                |
| F   | T   | T   | T                     | T                                |
| F   | T   | F   | F                     | F                                |
| F   | F   | T   | F                     | F                                |
| F   | F   | F   | F                     | F                                |

(c) Equivalent.

| $P$ | $Q$ | $R$ | $(P \wedge Q) \vee R$ | $(P \vee R) \wedge (Q \vee R)$ |
|-----|-----|-----|-----------------------|--------------------------------|
| T   | T   | T   | T                     | T                              |
| T   | T   | F   | T                     | T                              |
| T   | F   | T   | T                     | T                              |
| T   | F   | F   | F                     | F                              |
| F   | T   | T   | T                     | T                              |
| F   | T   | F   | F                     | F                              |
| F   | F   | T   | T                     | T                              |
| F   | F   | F   | F                     | F                              |

### 3 Implication

Note 0  
Note 1

Which of the following implications are always true, regardless of  $P$ ? Give a counterexample for each false assertion (i.e. come up with a statement  $P(x,y)$  that would make the implication false).

- (a)  $\forall x \forall y P(x,y) \implies \forall y \forall x P(x,y)$ .
- (b)  $\forall x \exists y P(x,y) \implies \exists y \forall x P(x,y)$ .
- (c)  $\exists x \forall y P(x,y) \implies \forall y \exists x P(x,y)$ .

**Solution:**

- (a) True. For all can be switched if they are adjacent; since  $\forall x, \forall y$  and  $\forall y, \forall x$  means for all  $x$  and  $y$  in our universe.
- (b) False. Let  $P(x,y)$  be  $x < y$ , and the universe for  $x$  and  $y$  be the integers. Or let  $P(x,y)$  be  $x = y$  and the universe be any set with at least two elements. In both cases, the antecedent is true and the consequence is false, thus the entire implication statement is false.
- (c) True. The antecedent says that there is an  $x$ , say  $x'$ , where for every  $y$ ,  $P(x,y)$  is true. If the antecedent is true, then for every  $y$  one can choose  $x = x'$ , which will make the consequent true.