


2020



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 774210



D 3.8: Policy brief based on current research literature and existing databases targeting adolescents

NIPH
30.04.2020



Deliverable administration and summary			
Due date	30.04.2020		
Submission date	30.04.2020		
Deliverable type	Report		
Contributors:	Name	Organisation	Role / Title
Deliverable Leader	Arnfinn Helleve	NIPH	WP-3 leader
Contributing Author(s)	Anne-Siri Fismen	NIPH	Post doc
	Knut-Inge Klepp		Coordinator
Reviewer(s)	Kate Oldridge-Turner	WCRF	WP-2 leader
	Julianne Williams	WHO	WHO
	Margot Neveux	WOF	WP-9 coordinator
Final review and approval	Arnfinn Helleve	NIPH	WP-3 leader

Document change history				
Version	Release date	Reason for Change	Status (Draft/In-review/Submitted)	Distribution

Dissemination level		
PU	Public	X



Executive Summary

The policy brief is based on current research literature on primary prevention of overweight and obesity in adolescents, and data on overweight/obesity and energy-balance related behaviours among children and adolescents. The brief focus particularly on the socio-economic differences in overweight and obesity among adolescents, and the lack of evidence of effective interventions. The brief concludes with four concrete action points.



Table of content

Executive Summary	3
Introduction.....	5
Deliverable description	5
Objective of deliverable	5
Background.....	5
Description of activities.....	5
Results	6
References.....	9



Introduction

Deliverable description

The deliverable is a policy brief based on a summary of research on effective interventions to prevent overweight and obesity among adolescents and data on energy balance related behaviours.

Objective of deliverable

The objective of the deliverable is to present the current status of research on the effectiveness of interventions targeting adolescents affected by overweight and obesity in the light on data on BMI and energy balance related behaviours from the WHO COSI and HBSC studies in a policy relevant format.

Background

In deliverable D3.1, we made an overview of systematic reviews to summarise evidence from up-to-date reviews of the effectiveness of interventions aimed at preventing overweight and obesity in adolescents aged 10 to 19 years. The deliverable has been further developed with updated literature search and submitted in manuscript format for publication (1).

The WHO European Childhood Obesity Surveillance Initiative (COSI) provides standardized height and weight measurements from at least 250.000 children aged 6-9 years from national representative samples from European countries. The data collection, which started in 2007, has been repeated every three years, and the fifth round of data collection was recently completed (2).

The WHO Health Behaviour in School-aged Children (HBSC) survey has been conducted in nationally representative samples of 11-, 13-, and 15-years-old every fourth year since 1985 in countries and regions across Europe and America. The first study was conducted in four countries, while the most recent survey was conducted in 45 countries (3). The overall dataset from the most recent data collection is not available by the deadline of this deliverable. We therefore used data from the WHO HBSC report (4).

Description of activities

Data from the COSI study and HBSC study show that the inequalities in children and adolescents affected by overweight/obesity observed have persisted in most of the countries over time. A similar pattern is found for dietary behaviour and physical activity.

The results across the 13 systematic reviews included in the overview (1), showed little or no effect on body mass index, or physical activity levels of adolescents, while results from a couple of reviews suggest possibly beneficial effects of public health interventions in adolescents on dietary behaviours

(i.e. consumption of sugar-sweetened beverages). The certainty of evidence was low to very low for all outcomes. Overall, the evidence-base for the effect of primary interventions to prevent overweight and obesity in adolescents is weak. In particular, there is a lack of reviews assessing the impact of environmental interventions targeting adolescents, and reviews addressing social inequality are virtually absent from this body of literature (1).

The policy brief points towards four actions:

- Developing effective and coordinated policy responses which target inequalities, both in overweight and obesity, as well as in the social determinants of health.
- Cultivating a new understanding of the associations between national overweight and obesity prevalence data and national policies.
- Conducting high quality evaluations on the effectiveness of structural and environmental interventions targeting overweight and obesity among adolescents
- Studying the differential effects of interventions targeting overweight and obesity among adolescents on different socio-economic groups

Results

The final policy brief is presented below.

POLICY BRIEF

April 2020



Socio-economic differences in overweight and obesity among European adolescents – lack of evidence of effective interventions

Overweight and obesity rates among adolescents vary considerably between countries, by gender, and by socio-economic status. There is a similar pattern in dietary behavior and physical activity, and the trends shows little improvement over the last decade. An overview of summarized research reveals that there is lack of evidence of effective interventions targeting overweight and obesity among adolescents.

Background

Among young people in Europe, one in every seven adolescents aged 15 years is overweight or obese (1). By 2025 overweight is expected to affect more than 16 million children in the European community (2). Within Europe, overweight and obesity rates among adolescents vary considerably between countries, by gender, and by socio-economic status.

The Health Behaviour in School-aged Children (HBSC) survey is a WHO collaborative cross-national study that monitors the health behaviours, health outcomes and social environments of boys and girls aged 11, 13 and 15 years every four years (3). The most recent (2013/2014) survey was conducted in 42 countries across Europe. Self-reported data on height and weight from the study revealed that the prevalence of obesity is higher among, younger adolescents and those with lower affluence. The inequalities in obesity observed have persisted in most of the countries over time (3).

The primary causes of overweight and obesity are energy-related behaviours – physical activity, sedentary behaviour, eating behaviour and sleep – which contribute to an energy imbalance between calorie intake and energy expenditure. In addition, the obesogenic environments have been identified as the key driver for low levels of physical activity and intake of energy-dense foods by encouraging sedentary lifestyle and offer ready availability of energy-dense, nutrient-poor food (4). The HBSC study also shows differences in dietary behaviours and physical activity between European countries and socio-economic groups.

The majority of European adolescents do not meet the dietary recommendations. Daily consumption of fruit and vegetables are low, and the inequalities between socio-economic groups remained unchanged. The trend for physical activities among European adolescents is similar. The proportion of young people achieving the recommended 60 minutes of moderate to vigorous physical activity (MVPA) daily in 2014 was low across all countries and regions in boys (25%) and girls (15%). In addition to the gender differences, adolescents from higher-affluence families reported to be more physically active than adolescents from lower-affluence families (3).

The WHO-Europe Childhood Obesity Surveillance Initiative (COSI) collect objectively measured height and weight data among 6-9-year-old boys and girls using data collected within most countries of the European Region (5). COSI data shows that children's BMI vary across the different parts of the WHO European region. Higher prevalence of overweight, obesity and severe obesity is reported in boys compared to girls.

COSI data indicate that there is heterogeneity in the association between parental socio-economic status and children's BMI (6). Positive as well as negative associations between socio-economic indicators and overweight and obesity has been observed within the European region. In most countries, the distribution of overweight and obesity by parental socio-economic status has remained stable during the time period 2007-2017 (unpublished data from COSI).

This project has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 774210. This document reflects only the authors' views and the European Commission is not responsible for any use that may be made of the information it contains.



Review of current research

The obesity epidemic is driven by determinants in the physical, social, economic, commercial and cultural environments. Individual level interventions requiring high levels of personal agency may contribute to widening of social inequalities, including inequalities on overweight and obesity.

A recent review of the evidence of effective interventions targeting overweight and obesity among adolescent showed little or no effect on body mass index, or physical activity levels of adolescents, while results from a couple of reviews suggest possibly beneficial effects of public health interventions on dietary behaviours (i.e. consumption of sugar-sweetened beverages). There were a dominance of school-based interventions targeting individual behavioural change (7). Few if any interventions were directed towards the wider community, for example creating more green spaces, improving cycle networks, using nutrient labelling/profiles, or reducing the affordability of unhealthy food and drinks. This is surprising, since it is generally agreed that policy approaches to obesity prevention are required, due to its potential to reach the whole population, reduce inequities, and enable systemic changes, with potential benefits in terms of duration of effects (1).

The review revealed that there is limited evidence of differential effects of interventions on adolescents from different socio-economic groups. This despite the fact that groups with lower SES have the highest and fastest increasing obesity prevalence in Europe.

References

1. WHO Regional Office for Europe (2016). Health Behaviour in School-aged Children (HBSC) study: international report from the 2013/2014 survey. WHO, Copenhagen.
2. World Obesity Federation (2017) Obesity Atlas for the European Union 2017.
3. Inchley J et al. eds. Adolescent obesity and related behaviours: trends and inequalities in the WHO European Region, 2002–2014. Observations from the Health Behaviour in School-aged Children (HBSC) WHO collaborative cross-national study. Copenhagen, WHO Regional Office for Europe, 2017.
4. World Health Organization; 2016. Report of the Commission on Ending Childhood Obesity. Geneva
5. <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/activities/who-european-childhood-obesity-surveillance-initiative-cosi>
6. Lissner et al (2016): Socioeconomic inequalities in childhood overweight: heterogeneity across five countries in the WHO European Childhood Obesity Surveillance Initiative (COSI-2008). *Int J Obes (Lond)*. 2016;40(5):796–802.
7. Flodgren GM et al (2020). Primary prevention of overweight and obesity in adolescents: an overview of systematic reviews. (sub.)

The review concludes that the evidence-base for the effectiveness of interventions to prevent overweight and obesity in adolescents is weak. In addition, the body of evidence is incomplete, since the vast majority of included interventions targeted the individual and were set in schools, while structural and environmental interventions applied in the wider society were lacking. Community and population level interventions might stand a better chance of having a significant impact on the dietary and physical activity behaviour and health of the adolescent population (7).


Action needs


There is a need for:

- Effective and coordinated policy responses targeting inequalities both in overweight and obesity as well as in the social determinants of health.
- A new understanding of the associations between national overweight and obesity prevalence among adolescents and national policies
- High quality evaluations of the effectiveness of structural and environmental interventions targeting overweight and obesity among adolescents
- Studies on differential effects of interventions targeting overweight and obesity among adolescents on different socio-economic groups

ABOUT: CO-CREATE is led by the Norwegian Institute of Public Health and brings together 14 international research and advocacy organisations to work with young people to create, inform and promote policies for obesity prevention.

www.co-create.eu

 [@cocreate_eu](https://www.instagram.com/cocreate_eu)

 [@EU_COCREATE](https://twitter.com/EU_COCREATE)



References

- (1) Flodgren GM et al (2020). Primary prevention of overweight and obesity in adolescents: an overview of systematic reviews. (submitted)
- (2) <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/activities/who-european-childhood-obesity-surveillance-initiative-cosi/about-cosi>
- (3) <http://www.hbsc.org>
- (4) Inchley J et al. eds. Adolescent obesity and related behaviours: trends and inequalities in the WHO European Region, 2002–2014. Observations from the Health Behaviour in School-aged Children (HBSC) WHO collaborative cross-national study. Copenhagen, WHO Regional Office for Europe, 2017.



→ The **CO-CREATE project** has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 774210. The products of the research are the responsibility of the authors: the European Commission is not responsible for any use that may be made of them.

