



Trabajo Original

Valoración nutricional

Assessment of the fat mass index in women recently diagnosed with gynecological tumors

Evaluación del índice de masa grasa en mujeres diagnosticadas recientemente de tumores ginecológicos

Lígia Zampieri de Brito¹, Mariana dos Santos Murra¹, Ricardo Reis¹, Sheilla de Oliveira Faria², Maria Claudia Bernardes Spexoto³

¹Hospital de Amor. Barretos, São Paulo, Brazil. ²Departamento de Medicina Preventiva. Faculdade de Medicina. Universidade de São Paulo (USP). São Paulo, Brazil.

³Faculdade de Ciências da Saúde. Universidade Federal da Grande Dourados. Dourados, Mato Grosso do Sul, Brazil

Abstract

Objectives: to evaluate the nutritional status and body composition of women with gynecological tumors and evaluate the fat mass index (FMI) as a complementary indicator for addressing the nutrition status.

Methods: a cross-sectional study with women recently diagnosed with gynecological tumors. Nutritional status was assessed using conventional anthropometry and the Patient-Generated Subjective Global Assessment. For body composition, bioelectrical impedance was used.

Results: a total of 158 women participated, most of them with excess weight and high body fat. The FMI showed a positive and significant correlation with body mass index, arm circumference, tricipital skinfold, and arm muscle circumference.

Conclusion: women recently diagnosed with gynecological tumors had excess weight and high body fat. The FMI may be a potentially useful indicator to complement the assessment of nutritional status and help the multidisciplinary team to perform early clinical and nutritional interventions.

Keywords:

Nutritional status. Body composition. Gynecologic neoplasms.

Resumen

Objetivos: evaluar el estado nutricional y la composición corporal de mujeres con tumores ginecológicos, y evaluar el índice de masa grasa (IMG) como indicador nutricional complementario.

Métodos: estudio transversal con mujeres diagnosticadas recientemente de tumores ginecológicos. El estado nutricional se evaluó mediante la antropometría convencional y la Evaluación Global Subjetiva Generada por el Paciente. Para la composición corporal se utilizó la impedancia bioeléctrica.

Resultados: participaron 158 mujeres, la mayoría con exceso de peso y grasa corporal alta. El IMG mostró una correlación positiva y significativa con el índice de masa corporal, la circunferencia del brazo, el pliegue cutáneo tricipital y la circunferencia de los músculos del brazo.

Conclusión: las mujeres diagnosticadas recientemente con tumores ginecológicos presentaron exceso de peso y grasa corporal alta. El IMG puede ser un indicador potencialmente útil para complementar la evaluación del estado nutricional y ayudar al equipo multidisciplinario a realizar intervenciones clínicas y nutricionales tempranas.

Palabras clave:

Estado nutricional. Composición corporal. Neoplasias ginecológicas.

Received: 10/04/2021 • Accepted: 22/11/2021

Conflicts of interest: the authors declare no conflicts of interest.

Brito LZ, Murra MS, Reis R, Faria SO, Spexoto MCB. Assessment of the fat mass index in women recently diagnosed with gynecological tumors. *Nutr Hosp* 2022;39(2):348-354

DOI: <http://dx.doi.org/10.20960/nh.03641>

Correspondence:

Maria Claudia Bernardes Spexoto. Faculdade de Ciências da Saúde. Universidade Federal da Grande Dourados. Rodovia Dourados / Itahum, Km 12 - Unidade II. Cidade Universitária. Caixa Postal: 364. 79.804-970. Dourados, Mato Grosso do Sul, Brazil
e-mail: mariaspexoto@ufgd.edu.br

INTRODUCTION

The global cancer burden has grown significantly. Gynecological cancer is most prevalent among obese women living in developing countries and having low socioeconomic status. In Brazil, the expected number of new cases of this neoplasm for each year of the 2020-2022 triennium will be 16,710, with an estimated risk of 16.35 cases per 100,000 women (1).

Among risk factors for cancer inadequate lifestyles stand out, including smoking, high alcohol consumption, inadequate diet, obesity, and body fat, among other environmental factors (2-4). Obesity, a major global epidemic, is considered the most significant preventable risk factor for several types of malignant tumors among adult women; therefore, maintaining a healthy weight is a primary recommendation among cancer prevention entities (5). A study on the proportion of cancer cases attributed to lifestyle in Brazil found that 36.5 % of cervical cancers and 5.7 % of ovarian cancers in Brazil are attributed to an elevated body mass index (BMI) (6).

There are few studies in the literature assessing the nutritional status and body composition of women with gynecological tumors, especially before antineoplastic therapy. Gold-standard methods to estimate body composition in this population, such as magnetic resonance imaging, computed tomography, and DEXA, could be high-cost (12). Therefore, in the context of limited resources, the adoption of complementary indicators in determining nutritional status can be very useful.

Thus, the objectives of this study were: 1) to evaluate the nutritional status and body composition of women with gynecological tumors before starting cancer treatment; 2) to evaluate the fat mass index (FMI) as a complementary indicator for addressing the nutrition status.

MATERIALS AND METHODS

STUDY AND SAMPLE CHARACTERIZATION

This is a cross-sectional study with women recently diagnosed with gynecological tumors, between January and September 2017, at a public hospital in Brazil. The study included women with positive pathology for gynecological cancer (ICD 10 - C52, ICD 10 - C53, ICD 10 - C56, and other correspondents) (7,8), without any previous antineoplastic treatments, aged 20 years or more. Women with mental or cognitive deficits were excluded.

For the sociodemographic characterization we used the economic class according to the Brazilian Economic Classification criteria of the Brazilian Association of Research Companies (ABEP) (9). Clinical stage was classified according to the AJCC 8th edition (10). We collected self-reported diabetes *mellitus* (type 2) and arterial hypertension (11-13).

ASSESSMENT OF NUTRITIONAL STATUS AND BODY COMPOSITION

The following measures were considered: current weight (kg), height (m), body mass index (BMI, kg/m²) (14), arm circumference (AC, cm), tricipital skinfold (TS, mm), and arm muscle circumference (AMC, cm).

AC and TS measurements were determined according to the criteria established by Lohman et al. (15). We adopted the cut-off points proposed by Blackburn and Thornton (16). Nutritional status was also assessed using the Patient-Generated Subjective Global Assessment (PG-SGA), culturally adapted to Portuguese (Brazilian) by Campos and Prado (17), and classified according to Ottery (A: well-nourished; B: mildly/moderately malnourished; C: severely malnourished) (18).

We used the multi-frequency segmented bioimpedance analysis (InBody[®] model 230 equipment) to evaluate body composition. The fat mass index (FMI) was determined using the equation: FMI (kg/m²) = fat mass (kg) / height² (m), and classified according to the cutoff points for women proposed by Kyle et al. (19). Of the 171 women who agreed to participate in the survey, 158 (92.4 %) underwent a bioelectrical impedance test.

We conducted a descriptive analysis of the data. Pearson's correlation coefficient (*r*) was used to estimate the correlation between FMI and conventional anthropometric variables. We considered a strong correlation for values higher than 0.70 (20). We adopted the significance level of *p* < 0.05. The analyses were conducted with the aid of the SPSS software, version 22.

This study followed the rules and guidelines of Good Clinical Practice according to Resolution 466/2012, and was approved by the Research Ethics Committee of the Hospital under the protocol number: 2.042.767. We have no conflicts of interest to declare.

RESULTS

A total of 158 women recently diagnosed predominantly with cervical neoplasia (56.9 %) in a non-advanced stage (I and II, 59.2 %) participated in the study. The average age of the participants was 52.2 ± 15.3 years. Most were married (53.2 %), economic class C (52.6 %) (low economic level), non-smokers (72.8 %), non-alcoholic (68.4 %), and unemployed (62.0 %) (Table I).

Regarding nutritional status, 45.9 % (*n* = 72) were adequate according to AMC, and well nourished (A) (86.1 %, *n* = 136) when assessed by the PG-SGA. As for BMI, most were overweight (71.5 %, *n* = 113) (Table I). The sample was composed, on average, by women with excess weight and high body fat (Table II).

The FMI showed a positive and significant correlation with BMI (*r* = 0.934), AC (*r* = 0.812), TS (*r* = 0.562) and AMC (*r* = 0.747), *p* < 0.001 (Fig. 1).

We identified that women with a very high FMI belonged to the groups with illiterate/incomplete elementary schooling (52.4 %), no work activity (73.0 %), a low economic class (C, 57.4 %), obesity (BMI) (85.7 %) and well nourished (95.2 %) (Table III).

Table I. Sociodemographic, clinical and nutritional status characterization of the participants

Characteristics	n	%
Sociodemographic		
<i>Education</i>		
Illiterate/Incomplete elementary school	73	46.2
Complete elementary school and incomplete middle school	31	19.6
Complete high school and incomplete higher education	40	25.3
Higher education or more	14	8.9
<i>Total</i>	158	100.0
<i>Marital status</i>		
Single	31	19.6
Married	84	53.2
Widowed	26	16.5
Divorced	17	10.8
<i>Total</i>	158	100.0
<i>Work activity</i>		
No	98	62.0
Yes	60	38.0
<i>Total</i>	158	100.0
<i>Economic class*</i>		
A	1	0.6
B	30	19.2
C	82	52.6
D and E	43	27.6
<i>Total</i>	156	100.0
<i>Smoker</i>		
No	115	72.8
Yes	18	11.4
Former smoker	25	15.8
<i>Total</i>	158	100.0
<i>Drinker</i>		
No	108	68.4
Yes	30	19.0
Former drinker	20	12.7
<i>Total</i>	158	100.0
Clinical		
<i>Tumor site^f</i>		
Cervix	87	56.9
Endometrium	39	25.5
Ovary	23	15.0
Vagina	3	2.0
Vulva	1	0.7
<i>Total</i>	153	100.0
<i>Staging^g</i>		
I	45	33.3
II	35	25.9
III	40	29.6
IV	15	11.1
<i>Total</i>	135	100.0
<i>Diabetes (type 2)</i>		
No	135	85.4
Yes	23	14.6
<i>Total</i>	158	100.0

(Continues on next column)

Table I (Cont.). Sociodemographic, clinical and nutritional status characterization of the participants

Characteristics	n	%
<i>Hypertension</i>		
No	96	60.8
Yes	62	39.2
<i>Total</i>	158	100.0
<i>BMI</i>		
Underweight	6	3.8
Adequate	39	24.7
Overweight	57	36.1
Obesity	56	35.4
<i>Total</i>	158	100.0
<i>AMC^s</i>		
Severe malnutrition	4	2.5
Moderate malnutrition	11	7.0
Mild malnutrition	26	16.6
Adequate	72	45.9
Overweight	25	15.9
Obesity	19	12.1
<i>Total</i>	157	100.0
<i>PG-SGA</i>		
Well nourished (A)	136	86.1
Moderate malnutrition (B)	21	13.3
Severely malnourished (C)	1	0.6
<i>Total</i>	158	100.0

BMI: body mass index; AMC: arm muscle circumference; PG-SGA: Patient-Generated Subjective Global Assessment. *2 missing values; [†]5 missing values; [‡]23 missing values; [§]1 missing value.

Table II. Summary measures of the participants' anthropometric variables and body composition

Variables	Mean ± standard deviation
Conventional anthropometry	
Weight (kg)	70.5 ± 17.2
Height (m)	1.6 ± 0.1
Body mass index (kg/m ²)	28.7 ± 6.5
Arm circumference (cm)	30.4 ± 5.2
Tricipital skinfold (mm)	24.8 ± 6.7
Arm muscle circumference (cm)	22.5 ± 4.1
Bioelectrical impedance	
Muscle mass (%)	33.5 ± 5.3
Muscle mass (kg)	22.9 ± 4.3
Fat mass (%)	38.1 ± 9.9
Fat mass (kg)	27.7 ± 12.0




Figure 1.

Correlation between fat mass index (FMI) and conventional anthropometric variables (FMI: fat mass index; BMI: body mass index; AC: arm circumference; TS: triceps skinfold; AMC: arm muscle circumference).

Table III. Classification of the Fat Mass Index (FMI) of the participants considering the sociodemographic, clinical, and nutritional status characteristics

Characteristics	Fat Mass Index (FMI), n (%) (95 % CI)			
	Low	Normal	High	Very high
Sociodemographic				
<i>Education</i>				
Illiterate/Incomplete elementary school	9 (90.0) (70.0-100.0)	10 (33.3) (16.7-50.0)	21 (38.2) (25.5-50.9)	33 (52.4) (41.3-63.5)
Complete elementary school and incomplete middle school	1 (10.0) (0-30.0)	5 (16.7) (6.7-30.0)	5 (9.1) (1.9-18.2)	20 (31.7) (20.6-42.9)
Complete high school and incomplete higher education	-	9 (30.0) (13.3-46.7)	21 (38.2) (25.5-50.9)	10 (15.9) (7.9-25.4)
Higher education or more	-	6 (20.0) (6.7-36.6)	8 (14.5) (5.5-23.6)	-

(Continues on next page)

Table III (Cont.). Classification of the Fat Mass Index (FMI) of the participants considering the sociodemographic, clinical, and nutritional status characteristics

Characteristics	Fat Mass Index (FMI), n (%) (95 % CI)			
	Low	Normal	High	Very high
Sociodemographic				
<i>Marital status</i>				
Single	2 (20.0) (0-50.0)	4 (13.3) (3.3-26.7)	16 (29.1) (18.2-41.8)	9 (14.3) (6.3-23.8)
Married	4 (40.0) (10.0-70.0)	20 (66.7) (50.0-83.3)	30 (54.5) (41.8-67.3)	30 (47.6) (34.9-60.3)
Widow	2 (20.0) (0-50.0)	3 (10.0) (0-23.3)	8 (14.5) (5.5-23.6)	13 (20.6) (11.1-31.7)
Divorced	2 (20.0) (0-50.0)	3 (10.0) (0-20.0)	1 (1.8) (0-5.5)	11 (17.5) (7.9-27.0)
<i>Work activity</i>				
No	5 (50.0) (20.0-80.0)	19 (63.3) (46.7-80.0)	28 (50.9) (38.2-65.5)	46 (73.0) (61.9-84.1)
Yes	5 (50.0) (20.0-80.0)	11 (36.7) (20.0-53.3)	27 (49.1) (34.5-61.8)	17 (27.0) (15.9-38.1)
<i>Economic class*</i>				
A	-	-	-	1 (1.6) (0-4.9)
B	-	7 (23.3) (10.0-40.0)	15 (27.3) (14.5-38.2)	8 (13.1) (4.9-23.0)
C	3 (30.0) (0-60.0)	15 (50.0) (30.0-66.7)	29 (52.7) (40.0-67.3)	35 (57.4) (44.3-70.5)
D and E	7 (70.0) (40.0-100.0)	8 (26.7) (10.0-43.3)	11 (20.0) (9.1-30.9)	17 (27.9) (16.4-39.3)
<i>Smoker</i>				
No	8 (80.0) (50.0-100.0)	24 (80.0) (63.3-93.3)	40 (72.7) (60.0-83.6)	43 (68.3) (55.6-79.4)
Yes	1 (10.0) (0-30.0)	2 (6.7) (0-16.7)	7 (12.7) (5.5-21.8)	8 (12.7) (4.8-22.2)
Former smoker	1 (10.0) (0-30.0)	4 (13.3) (3.3-26.7)	8 (14.5) (5.5-25.4)	12 (19.0) (9.5-28.6)
<i>Drinker</i>				
No	9 (90.0) (70.0-100.0)	21 (70.0) (53.3-86.7)	36 (65.5) (52.7-76.4)	42 (66.7) (55.6-77.8)
Yes	-	8 (26.7) (10.0-43.3)	11 (20.0) (10.9-30.9)	11 (17.5) (9.5-28.6)
Former drinker	1 (10.0) (0-30.0)	1 (3.3) (0-10.0)	8 (14.5) (5.5-23.6)	10 (15.9) (7.9-25.4)
Clinical				
<i>Tumor site[‡]</i>				
Cervix	5 (55.6) (22.2-88.9)	20 (69.0) (51.7-86.2)	34 (63.0) (50.0-75.9)	28 (45.9) (32.8-59.0)
Endometrium	2 (22.2) (0-55.6)	3 (10.3) (0-20.7)	7 (13.0) (5.6-22.2)	27 (44.3) (31.1-57.4)
Ovary	1 (11.1) (0-33.3)	5 (17.2) (3.4-31.0)	12 (22.2) (11.1-33.3)	5 (8.2) (1.6-16.4)
Vagina	1 (11.1) (0-33.3)	-	1 (1.9) (0-5.6)	1 (1.6) (0-4.9)
Vulva	-	1 (3.4) (0-10.3)	-	-
<i>Staging[‡]</i>				
I	2 (20.0) (0-50.0)	5 (20.0) (4.1-36.0)	16 (32.7) (20.4-46.9)	22 (43.1) (29.4-56.9)
II	2 (20.0) (0-50.0)	8 (32.0) (16.0-52.0)	14 (28.6) (16.3-40.8)	11 (21.6) (9.8-33.3)
III	2 (20.0) (0-50.0)	8 (32.0) (16.0-48.0)	16 (32.7) (20.4-46.9)	14 (27.5) (15.7-41.1)
IV	4 (40.0) (10.0-70.0)	4 (16.0) (4.0-32.0)	3 (6.1) (0-14.3)	4 (7.8) (2.0-15.7)
<i>Diabetes (type 2)</i>				
No	10 (100.0)	28 (93.3) (83.3-100.0)	51 (92.7) (83.6-98.2)	46 (73.0) (61.9-82.5)
Yes	-	2 (6.7) (0-16.7)	4 (7.3) (1.8-16.4)	17 (27.0) (17.5-38.1)
<i>Hypertension</i>				
No	7 (70.0) (40.0-90.0)	20 (66.7) (50.0-83.3)	42 (76.4) (63.6-87.3)	27 (42.9) (31.7-55.6)
Yes	3 (30.0) (10.0-60.0)	10 (33.3) (16.7-50.0)	13 (23.6) (12.7-36.4)	36 (57.1) (44.4-68.3)
<i>BMI</i>				
Underweight	6 (60.0) (30.0-90.0)	-	-	-
Adequate	2 (20.0) (0-50.0)	27 (90.0) (80.0-100.0)	9 (16.4) (7.3-27.3)	1 (1.6) (0-4.8)
Overweight	2 (20.0) (0-50.0)	3 (10.0) (0-20.0)	44 (80.0) (67.3-90.9)	8 (12.7) (4.8-20.6)
Obesity	-	-	2 (3.6) (0-9.1)	54 (85.7) (76.2-93.7)

(Continues on next page)

Table III (Cont.). Classification of the Fat Mass Index (FMI) of the participants considering the sociodemographic, clinical, and nutritional status characteristics

Characteristics	Fat Mass Index (FMI), n (%) (95 % CI)			
	Low	Normal	High	Very high
Clinical				
<i>AMC</i> [§]				
Severe malnutrition	-	2 (6.7) (0-16.7)	1 (1.8) (0-5.5)	1 (1.6) (0-4.8)
Moderate malnutrition	5 (50.0) (20.0-80.0)	5 (16.7) (3.3-30.0)	1 (1.8) (0-7.3)	-
Mild malnutrition	3 (30.0) (0-60.0)	12 (40.0) (23.3-56.7)	10 (18.2) (9.1-29.1)	1 (1.6) (0-4.8)
Adequate	2 (20.0) (0-50.0)	9 (30.0) (13.3-46.7)	39 (70.9) (58.2-81.8)	22 (35.5) (24.2-48.3)
Overweight	-	1 (3.3) (0-10.0)	3 (5.5) (0-12.7)	21 (33.9) (22.6-45.2)
Obesity	-	1 (3.3) (0-10.0)	1 (1.8) (0-5.5)	17 (27.4) (17.7-38.7)
<i>PG-SGA</i>				
Well nourished (A)	3 (30.0) (10.0-60.0)	25 (83.3) (70.0-96.7)	48 (87.3) (78.2-94.5)	60 (95.2) (88.9-100.0)
Moderate malnutrition (B)	6 (60.0) (30.0-90.0)	5 (16.7) (3.3-30.0)	7 (12.7) (5.5-21.8)	3 (4.8) (0-11.1)
Severely malnourished (C)	1 (10.0) (0-30.0)	-	-	-

BMI: body mass index; AMC: arm muscle circumference; PG-SGA: Patient-Generated Subjective Global Assessment. *2 missing values; †5 missing values; ‡23 missing values; §1 missing value.

DISCUSSION

This study investigated the nutritional status and body composition of women diagnosed with gynecological tumors without previous antineoplastic treatment, and evaluated their FMI as a complementary indicator for addressing the nutrition status.

In this study, cervical cancer was predominant, corroborating both national and international statistics pointing at it as the most frequent gynecological tumor in the female population (1).

The women in this study were overweight and had high body fat at diagnosis. Although excess body fat and lifelong weight gain can influence the development of some gynecological cancers through inflammatory, metabolic, and hormonal mechanisms (21), its influence on cervical cancer risk has been poorly understood.

Excess body fat can negatively influence treatment and patient quality of life. Patients with morbid obesity and endometrial cancer or ovarian cancer submitted to surgery had a higher number of surgical complications than patients with a BMI < 40.0 kg/m² (22,23). Similarly, a study on the impact of obesity on complications and survival in 2,500 patients with endometrial cancer found that obese women had a higher risk of all-cause mortality (24).

In addition to the impact on performing an adequate and safe surgery, excess body fat can compromise a safe and effective administration of cytotoxic agents. There are limited data on the evaluation of the relationship between obesity and the pharmacokinetics of chemotherapy, specifically regarding volume, in obese patients. The issue of reassessing chemotherapy doses based on body composition in this population should be discussed (25).

The FMI showed a strong and positive correlation with conventional anthropometric measurements such as BMI, AC, and AMC, and a moderate and positive correlation with TS; thus, it can be considered a good indicator for assessing body composition at diagnosis, besides using the lean body mass index. The isolated use of BMI does not reflect body composition in a cancer patient and, therefore, should always be used with other body composition measures (26).

In the present study, most women with a very high FMI were obese (BMI > 30 kg/m²) and had a low economic and educational level. Similarly, in a study that estimated the frequency and sociodemographic distribution of risk and protective factors for chronic diseases in Brazil (27), the frequency of overweight and obesity among women decreased notably with increased education.

This study has some limitations. The study was developed in only one oncology hospital, and its cross-sectional design limits the evaluation of causal relationships. We did not evaluate weight loss or weight gain before the assessment, which could influence body composition. In addition, no data were collected that could validate the FMI measurement as a predictor of complications and other clinical outcomes since that was not the objective of the present study. It is important to point out that it would be interesting to evaluate whether the patients were menopausal or premenopausal, since fat mass is associated with an increased risk of uterine corpus cancer in postmenopausal women (28). We believe that our results may arouse interest in conducting new studies with prospective follow-up, given the importance of nutritional intervention at diagnosis and the implications of overweight in cancer treatment.

CONCLUSION

Women recently diagnosed with malignant gynecological tumors without previous antineoplastic treatments were admitted with overweight and increased FMI.

Besides, FMI showed a good correlation with conventional measures of BMI, AC, and AMC. Thus, FMI may be a good indicator of body composition and a potentially useful tool to complement the assessment of nutritional status. In clinical practice, FMI can help multidisciplinary teams to perform early clinical and nutritional interventions.

REFERENCES

- Instituto Nacional de Cancer José Alencar Gomes da Silva. INCA. Estimativa 2020: incidência de câncer no Brasil. Ministério da Saúde Instituto Nacional de Cancer José Alencar Gomes da Silva; 2019.
- Carter BD, Abnet CC, Feskanich D, Freedman ND, Hartge P, Lewis CE, et al. Smoking and Mortality-Beyond Established Causes. *N Engl J Med* 2015;372(7):631-40. DOI: 10.1056/NEJMsa1407211
- Bagnardi V, Rota M, Botteri E, Tramacere I, Islami F, Fedirko V, et al. Alcohol consumption and site-specific cancer risk: A comprehensive dose-response meta-analysis. *Br J Cancer* 2015;112(3):580-93. DOI: 10.1038/bjc.2014.579
- Lauby-Secretan B, Scoccianti C, Loomis D. Body Fatness and Cancer --Viewpoint of the IARC Working Group. *N Engl J Med* 2016;375(8):794-8. DOI: 10.1056/NEJMs1606602
- Clinton SK, Giovannucci EL, Hursting SD. The World Cancer Research Fund/American Institute for Cancer Research Third Expert Report on Diet, Nutrition, Physical Activity, and Cancer: Impact and Future Directions. *Journal of Nutrition* 2020;150(4):663-71. DOI: 10.1093/jn/nxz268
- Rezende LFM de, Lee DH, Louzada ML da C, Song M, Giovannucci E, Eluf-Neto J. Proportion of cancer cases and deaths attributable to lifestyle risk factors in Brazil. *Cancer Epidemiol* 2019;59:148-57. DOI: 10.1016/j.canep.2019.01.021
- Organização Mundial da Saúde. CID-10 Classificação Estatística Internacional de Doenças e Problemas Relacionados à Saúde. 10a rev. São Paulo: Universidade de São Paulo; 1998.
- OMS-Organização Mundial da Saúde. CID-10 VOL. 1: Classificação Estatística Internacional de Doenças. Organização Mundial da Saúde (OMS). Organização Pan-Americana de Saúde (OPAS); 2008.
- ABEP. Critérios de Classificação Econômica Brasil. In: Alterações na aplicação do Critério Brasil. ABEP; 2018.
- Edge SB, Compton CC. The american joint committee on cancer: The 7th edition of the AJCC cancer staging manual and the future of TNM. *Annals of Surgical Oncology* 2010;17(6):1471-4. DOI: 10.1245/s10434-010-0985-4
- Vrachnis N, Iavazzo C, Iliodromiti Z, Sifakis S, Alexandrou A, Siristatidis C, et al. Diabetes mellitus and gynecologic cancer: molecular mechanisms, epidemiological, clinical and prognostic perspectives. *Arch Gynecol Obstet* 2016;293(2):239-46. DOI: 10.1007/s00404-015-3858-z
- Lees B, Leath CA 3rd. The Impact of Diabetes on Gynecologic Cancer: Current Status and Future Directions. *Curr Obstet Gynecol Rep* 2015;4(4):234-9. DOI: 10.1007/s13669-015-0127-x
- Sun L-M, Kuo H-T, Jeng L-B, Lin C-L, Liang J-A, Kao C-H. Hypertension and subsequent genitourinary and gynecologic cancers risk: a population-based cohort study. *Medicine (Baltimore)* 2015;94(16):e753. DOI: 10.1097/MD.0000000000000753
- WHO. Obesity: preventing and managing the global epidemic [Internet]. Vol. 894, World Health Organization technical report series; 2000. p. 1-253. Available from: <http://books.google.com/books?hl=en&lr=&id=Avn-q0sqv9doC&oi=fnd&pg=PA1&dq=Obesity:+Preventing+and+managing+the+global+epidemic&ots=6UF0cjXY6M&sig=iA1h8eokNhrTKDfX-qIVZaivyyk4%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/11234459>
- Lohman TG, Roche AF, Martorell R. Anthropometric standardization reference manual. Hum Kinet Books; 1988.
- Blackburn GL, Thornton PA. Nutritional assessment of the hospitalized patient. *Med Clin North Am* 1979;63(5):11103-15.
- Duarte Bonini Campos JA, Dias do Prado C. Cross-cultural adaptation of the Portuguese version of the Patient-Generated Subjective Global Assessment. *Nutr Hosp* 2012;27(2):583-9. DOI: 10.1590/S0212-16112012000200035
- Ottery FD. Definition of standardized nutritional assessment and interventional pathways in oncology. *Nutrition* 1996;12(Suppl1):S15-9. DOI: 10.1016/0899-9007(96)90011-8
- Kyle UG, Pirlich M, Lochs H, Schuetz T, Pichard C. Increased length of hospital stay in underweight and overweight patients at hospital admission: A controlled population study. *Clin Nutr* 2005;24(1):133-42. DOI: 10.1016/j.clnu.2004.08.012
- Mukaka MM. Statistics corner: A guide to appropriate use of correlation coefficient in medical research. *Malawi Med J* 2012;24(3):69-71.
- Staley SA, Tucker KR, Clark LH. The Role of Obesity in the Development and Management of Gynecologic Cancer. *Obstet Gynecol Surv* 2020;75(5):308-16. DOI: 10.1097/OGX.0000000000000779
- Smits A, Lopes A, Das N, Kumar A, Cliby W, Smits E, et al. Surgical morbidity and clinical outcomes in ovarian cancer - The role of obesity. *BJOG: An International Journal of Obstetrics and Gynaecology* 2016;123(2):300-8. DOI: 10.1111/1471-0528.13585
- Bouwman F, Smits A, Lopes A, Das N, Pollard A, Massuger L, et al. The impact of BMI on surgical complications and outcomes in endometrial cancer surgery - An institutional study and systematic review of the literature. *Gynecologic Oncology* 2015;139(2):369-76. DOI: 10.1016/j.ygyno.2015.09.020
- Gunderson CC, Java J, Moore KN, Walker JL. The impact of obesity on surgical staging, complications, and survival with uterine cancer: A Gynecologic Oncology Group LAP2 ancillary data study. *Gynecol Oncol* 2014;133(1):23-7. DOI: 10.1016/j.ygyno.2014.01.041
- Desmedt C, Fornili M, Clatot F, Demicheli R, De Bortoli D, Di Leo A, et al. Differential Benefit of Adjuvant Docetaxel-Based Chemotherapy in Patients With Early Breast Cancer According to Baseline Body Mass Index. *J Clin Oncol* 2020;38(25):2883-91. DOI: 10.1200/JCO.19.01771
- Ravasco P. Nutrition in Cancer Patients. *J Clin Med* 2019;8(8):1211. DOI: 10.3390/jcm8081211
- Brasil. Ministério da Saúde. Secretaria de Atenção à Saúde. Secretaria de Vigilância em Saúde. Departamento de Análise em Saúde e Vigilância de Doenças Não Transmissíveis. *Vigilância em Saúde* 2019: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico: estimativas sobre frequência e distribuição sociodemográfica de fatores de risco e proteção para doenças crônicas nas capitais dos 26 estados [Internet]. Brasília; 2020. Available from: <https://portalarquivos.saude.gov.br/images/pdf/2020/Abril/27/vigil-em-brasil-2019-vigilancia-fatores-risco.pdf>
- Yun P, Xia B, Tian X-H, Gong T, Liu A-R, Yuan J-Q, et al. Body composition and risk of major gynecologic malignancies: Results from the UK Biobank prospective cohort. *Cancer Med* 2021;10(13):4522-31. DOI: 10.1002/cam4.3925