

27. Marrie TJ. *Coxiella burnetii* pneumonia. Eur Respir J. 2003;21:713–9. DOI: 10.1183/09031936.03.00095403
28. Ayres JG, Flint N, Smith EG, et al. Post-infection fatigue syndrome following Q fever. QJM. 1998;91:105–23. DOI: 10.1093/qjmed/91.2.105
29. Marmion BP, Shannon M, Maddocks I, Storm P, Penttila I. Protracted debility and fatigue after acute Q fever. Lancet. 1996;347:977–8. DOI: 10.1016/S0140-6736(96)91469-5
30. Smith DL, Ayres JG, Blair I, Burge PS, Carpenter MJ, Caul EO, et al. A large Q fever outbreak in the West Midlands: clinical aspects. Respir Med. 1993;87:509–16. DOI: 10.1016/0954-6111(93)90006-L
31. Raoult D, Levy PY, Dupont HT, Chicheportiche C, Tamalet C, Gastaut JA, et al. Q fever and HIV infection. AIDS. 1993;7:81–6. DOI: 10.1097/00002030-199301000-00012
32. Siegman-Igra Y, Kaufman O, Keysary A, Rzotkiewicz S, Shalit I. Q fever endocarditis in Israel and a worldwide review. Scand J Infect Dis. 1997;29:41–9.
33. Fowler RA, Sanders GD, Bravata DM, Nouri B, Gastwirth JM, Peterson D, et al. Cost-effectiveness of defending against bioterrorism: a comparison of vaccination and antibiotic prophylaxis against anthrax. Ann Intern Med. 2005;142:601–10.
34. Sabbaj J, Hoagland VL, Shih WJ. Multiclinic comparative study of norfloxacin and trimethoprim-sulfamethoxazole for treatment of urinary tract infections. Antimicrob Agents Chemother. 1985;27:297–301.
35. Talan DA, Stamm WE, Hooton TM, Moran DJ, Burke T, Travani A, et al. Comparison of ciprofloxacin (7 days) and trimethoprim-sulfamethoxazole (14 days) for acute uncomplicated pyelonephritis in women: a randomized trial. JAMA. 2000;283:1583–90. DOI: 10.1001/jama.283.12.1583
36. Kocak Z, Hatipoglu CA, Ertem G, Kinikli S, Tufan A, Irmak H, et al. Trimethoprim-sulfamethoxazole induced rash and fatal hematologic disorders. J Infect. 2006;52:e49–52. DOI: 10.1016/j.jinf.2005.05.008
37. See S, Mumford JM. Trimethoprim/sulfamethoxazole-induced toxic epidermal necrolysis. Ann Pharmacother. 2001;35:694–7. DOI: 10.1345/aph.10310
38. Fennelly KP, Davidow AL, Miller SL, Connell N, Ellner JJ. Airborne infection with *Bacillus anthracis*—from mills to mail. Emerg Infect Dis. 2004;10:996–1001.

Address for correspondence: Claire E. Moodie, 754 Centre St #11, Jamaica Plain, MA 02130, USA; email: claire_moodie@hotmail.com

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institutions with which the authors are affiliated.

etymologia

Coxiella [kok" se-el' ə] *burnetii*

Etiologic agent of Q fever, named after American bacteriologist Herald Rea Cox and Australian physician Frank MacFarlane Burnet, who both independently isolated the bacterium in the 1930s. *C. burnetii* belongs in the family *Coxiellaceae*, which consists of gram-negative rods without flagella or a capsule. The bacteria occur in ticks and various vertebrates, including humans.

Source: Dorland's illustrated medical dictionary, 31st edition. Philadelphia: Saunders; 2007; American Veterinary Medical Association. Q fever backgrounder. Available from http://www.avma.org/public_health/biosecurity/qfever_bgnd.asp

**Now that you've read the article,
you can listen to the commentary.**

Prophylaxis after Exposure
to *Coxiella burnetii*

