

# Epidemiologic Changes of Scrub Typhus in China, 1952–2016

Zhongjie Li,<sup>1</sup> Hualei Xin,<sup>1</sup> Junling Sun,<sup>1</sup> Shengjie Lai, Lingjia Zeng, Canjun Zheng, Sarah E. Ray, Nicole Davis Weaver, Liping Wang, Jianxing Yu, Zijian Feng,<sup>2</sup> Simon I. Hay, George F. Gao<sup>2</sup>

Scrub typhus, a miteborne rickettsiosis, has emerged in many areas globally. We analyzed the incidence and spatial–temporal distribution of scrub typhus in China during 1952–1989 and 2006–2016 using national disease surveillance data. A total of 133,623 cases and 174 deaths were recorded. The average annual incidence was 0.13 cases/100,000 population during 1952–1989; incidence increased sharply from 0.09/100,000 population in 2006 to 1.60/100,000 population in 2016. The disease, historically endemic to southern China, has expanded to all the provinces across both rural and urban areas. We identified 3 distinct seasonal patterns nationwide; infections peaked in summer in the southwest, summer-autumn in the southeast, and autumn in the middle-east. Persons  $\geq 40$  years of age and in nonfarming occupations had a higher risk for death. The changing epidemiology of scrub typhus in China warrants an enhanced disease control and prevention program.

Scrub typhus is a life-threatening disease caused by *Orientia tsutsugamushi*, an obligate intracellular bacterium transmitted by the larvae of trombiculid mites (1). Only biting larvae of Asian scrub typhus chiggers (*Leptotrombidium* spp.) can transmit the disease. After the bite of an infective mite, a characteristic necrotic inoculation lesion (an eschar) can develop. The microorganism then spreads through the lymphatic fluid and blood, causing manifestations including fever, headache, rash, lymphadenopathy, and mental changes (1). Without appropriate treatment with specific antimicrobial drugs (e.g., tetracycline, chloramphenicol, doxycycline, or azithromycin),  $>6\%$  of infected patients will die

(2,3). There is no licensed human vaccine to prevent scrub typhus infection.

Globally, scrub typhus is traditionally regarded as a disease endemic to a region called the Asia-Pacific tsutsugamushi triangle, which extends from Pakistan in the west to far eastern Russia in the east to northern Australia in the south. In some countries of Southeast Asia, scrub typhus is a leading cause of treatable nonmalarial febrile illness (4–6). It is estimated that scrub typhus threatens  $>1$  billion persons, causes at least 1 million clinical cases per year, and is associated with substantial mortality rates globally (1,7). The 2010s saw a widespread reemergence of scrub typhus in endemic regions such as India, Korea, Laos, and the Maldives (3,6,8). The recent emergence of scrub typhus in the Arabian Peninsula, Chile, and possibly Kenya suggests wider global distribution of this disease in tropical and subtropical regions, far from the tsutsugamushi triangle (9–12). Although scrub typhus poses the greatest threat to residents of eastern and southern Asia as well as tourists traveling to these regions, it remains a neglected disease globally. The lack of both research and a nationwide surveillance system within many endemic regions have resulted in poorly understood epidemiologic characteristics and disease burden of scrub typhus at global, national, and subnational levels (7). The Global Burden of Disease Study publishes estimates for 333 diseases and injuries, but currently bundles scrub typhus with other neglected tropical diseases, rather than providing specific burden estimates (13).

In China, scrub typhus cases were recorded in the early 1950s, and a disease surveillance system for scrub typhus was established in 1952 (14,15). *Leptotrombidium deliense* and *L. scutellare* mites are the 2 principal vectors transmitting the disease in the country (16). *L. deliense* mites inhabit southern China and emerge in April, peaking in June–August, and decreasing

Author affiliations: Chinese Center for Disease Control and Prevention, Beijing, China (Z. Li, H. Xin, S. Lai, L. Zeng, C. Zheng, L. Wang, J. Yu, Z. Feng, G.F. Gao); Qingdao City Center for Disease Control and Prevention, Shandong, China (H. Xin); University of Southampton, Southampton, UK (S. Lai); University of Washington, Seattle, Washington, USA (S.E. Ray, N.D. Weaver, S.I. Hay); Chinese Academy of Sciences, Beijing (G.F. Gao)

DOI: <https://doi.org/10.3201/eid2606.191168>

<sup>1</sup>These first authors contributed equally to this article.

<sup>2</sup>These senior authors contributed equally to this article.

September–December, whereas *L. scutellare* is widespread in China, emerging annually in October–December, and is the main mite species in northern China (17). Several studies have revealed that the incidence of scrub typhus is rising nationwide (18,19); however, the changing epidemiologic characteristics of scrub typhus at the subnational level and among the subgroups are not fully illustrated. After the SARS outbreak in 2003, a nationwide population-based infectious disease surveillance system was developed in China to collect epidemiologic information at the patient level, enabling study of the epidemiologic features of a disease at the smaller scale of a geographic region and specific population (20,21).

In this study, we systematically collated longitudinal nationwide surveillance data of scrub typhus in China from 1952–1990 and 2006–2016. We explored changes in the disease's epidemiologic characteristics, both spatiotemporally and seasonally, and changes in high-risk population groups at both national and subnational levels.

## Materials and Methods

### National Surveillance of Scrub Typhus

In China, the national surveillance program on scrub typhus (Appendix Tables 1, 2, <https://wwwnc.cdc.gov/EID/article/26/6/19-1168-App1.pdf>) varied through 3 time periods. Initially, mail-based reporting of monthly aggregated data occurred in 1952–1989. Beginning in 1952, data on scrub typhus were reported on a voluntary basis in certain areas of China. In 1955, scrub typhus became a statutorily notifiable disease, and reporting all suspected, probable, and laboratory-confirmed scrub typhus cases was required. Reporting was suspended during 1990–2005; in 1990, scrub typhus was removed from the list of notifiable diseases, due to its relatively low threat. Therefore, nationwide data on the disease are not available from 1990–2005. Internet-based reporting of individual cases occurred in 2006–2016; in 2006, scrub typhus was added as a reporting disease in the National Notifiable Infectious Disease Reporting Information System at the Chinese Centers for Disease Control and Prevention (China CDC). The information on individual cases includes patient demographic information (name, sex, occupation, residence), date of illness onset, case classification (confirmed, probable, suspected), date of diagnosis, date of report, reporter institution, and date of death (if applicable).

The National Health Commission of the People's Republic of China determined that the collection of data on human cases of scrub typhus was part of a

routine public health investigation and was exempt from institutional review board assessment. All data in this study are anonymous so that individual patients cannot be identified.

### Case Definition

Scrub typhus cases have been classified as probable (clinically diagnosed) or confirmed (laboratory confirmed) in accordance with the diagnosis criteria and case classification guidelines issued by the Chinese national health authorities (Appendix Table 2). Probable cases are diagnosed by local experienced physicians according to epidemiologic exposure (travel to a disease-endemic area and contact with chiggers or rodents  $\leq 3$  weeks before the onset of illness) and clinical manifestations (such as high fever, lymphadenopathy, skin rash, and eschars or ulcers). Confirmed cases are probable cases with 1 positive result among the following tests: Weil-Felix test, indirect immunofluorescence antibody assay, PCR, or isolation of *Orientia tsutsugamushi* (22).

### Data Analysis

Our analysis comprised all probable and confirmed human scrub typhus cases in 1952–1989 and 2006–2016 (Appendix Tables 3, 4). We calculated the annual incidence rate by dividing the number of human scrub typhus cases by the corresponding population at the end of a given year. We obtained population data used to calculate incidence rates from the National Bureau of Statistics of China (23). We calculated the case-fatality ratio by dividing the number of human scrub typhus deaths by the number of scrub typhus cases. We used Joinpoint regression models to examine the incidence trends from 1952–2016 and calculated annual percentage changes (24).

We created a heat map of yearly incidence rates (standardized range 0–1) to visualize the long-term change over the 49-year period by province (Appendix Table 5, Figure 3). To assess the changing spatial and temporal patterns of the affected areas from 2006–2016, we plotted the newly affected and previously affected counties and townships by year in both the rural and urban areas where each case resided. In our study, the rural areas refer to the village or rural town, and urban areas refer to the city community or urban town. We identified epidemiologically related regions of scrub typhus based on hierarchical cluster analysis with Ward's minimum variance method, using the Pearson correlation coefficient matrix of average weekly cases between paired provinces as distance.

To quantify the characteristics of scrub typhus season, we fitted multiple linear regression models

to time series of the weekly number of scrub typhus cases during 2006–2016, including harmonic terms representing annual and semiannual periodic cycles as described previously (25–27). We defined scrub typhus season as the period of consecutive weeks in which >95% of the total annual cases occurred and as the smallest number of consecutive weeks. We characterized scrub typhus season by plotting the season onset, peak (the week with the highest case number in the fitted seasonal curve), and offset against the calendar week of the year. We used a multivariable logistic regression model to explore the predictors of death for scrub typhus. We used R statistical software version 3.4.1 (<https://www.r-project.org>) to produce graphs and perform statistical analyses (28) and ArcGIS version 10.2.2 (ESRI, <https://www.arcgis.com>) to plot geographic patterns. We used Joinpoint version 4.5.0.0 (US National Cancer Institute, <https://surveillance.cancer.gov/joinpoint>) to examine incidence trends.

**Results**


**Overall Incidence**

During the 49 years covered in this study (1952–1989 and 2006–2016), a total of 133,623 scrub typhus cases were recorded throughout the country. During 1952–

1989, the occurrence of the disease remained at a stable, low level, with a mean annual incidence rate of 0.13 (95% CI 0.11–0.15) per 100,000 population (Figure 1, panel A). Since 2006, the annual incidence rate has increased sharply, changing from 0.09/100,000 population in 2006 to 1.60/100,000 population in 2016 (>16-fold increase) (Table 1), with an annual percent change of 32% (95% CI 30%–34%) (Appendix Figure 1).

**Demographic Features**

From 2006–2016, the mean age of case-patients was 48 years (range 1 month–98 years). The median age increased from 46 years (interquartile range 27–58 years) in 2006 to 53 years (interquartile range 41–64 years) in 2016. Overall, case-patients ≥45 years of age were the most common (66%); the highest incidence was among those 60–64 years (1.67 cases/100,000 population) and 65–69 years (1.71 cases/100,000 population). However, a subgroup of children <5 years of age had the highest incidence rate in the southwest disease-endemic regions (4.30 cases/100,000 population for Yunnan and Sichuan provinces combined) (Table 1). The sex ratio of male to female changed from 1.09:1 in 2006–2009 to 0.85:1 in 2010–2016. In general, farmers (73%) were the most affected group, with 86% of cases occurring in rural areas.



**Figure 1.** Reported cases and deaths of scrub typhus in China, 1952–2016. A) Aggregated number of cases by year (red bars), annual incidence rate (blue line), and average annual incidence rate (black dashed line) per 100,000 residents. B) Aggregated number of deaths by year (red bars), case-fatality ratio (blue line), and average annual case-fatality ratio (black dashed line). The data from 1990–2005 are missing because surveillance for scrub typhus was suspended during the period.

**Table 1.** Characteristics of reported scrub typhus cases, China, 2006–2016\*

| Characteristic          | 2006          | 2007          | 2008          | 2009          | 2010          | 2011          | 2012          | 2013          | 2014           | 2015           | 2016           |
|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|----------------|----------------|----------------|
| No. cases               | 1,254         | 1,340         | 2,613         | 3,237         | 4,088         | 6,020         | 8,928         | 11,111        | 16,035         | 17,293         | 21,562         |
| No. probable cases      | 1,049         | 1,187         | 2,360         | 2,939         | 3,811         | 5,630         | 8,555         | 10,536        | 15,339         | 16,696         | 20,997         |
| No. confirmed cases     | 205           | 153           | 253           | 298           | 277           | 390           | 373           | 575           | 696            | 597            | 565            |
| Incidence†              | 0.09          | 0.10          | 0.19          | 0.24          | 0.30          | 0.45          | 0.66          | 0.82          | 1.19           | 1.28           | 1.60           |
| Sex                     |               |               |               |               |               |               |               |               |                |                |                |
| M                       | 642<br>(51)   | 711<br>(53)   | 1,385<br>(53) | 1,660<br>(51) | 2,040<br>(50) | 2,884<br>(48) | 4,228<br>(47) | 4,910<br>(44) | 7,222<br>(45)  | 8,021<br>(46)  | 9,871<br>(46)  |
| F                       | 612<br>(49)   | 629<br>(47)   | 1,228<br>(47) | 1,577<br>(49) | 2,048<br>(50) | 3,136<br>(52) | 4,700<br>(53) | 6,201<br>(56) | 8,813<br>(55)  | 9,272<br>(54)  | 11,691<br>(54) |
| Age, y, median (IQR)‡   | 46<br>(27–58) | 46<br>(27–59) | 48<br>(30–60) | 44<br>(25–57) | 46<br>(28–59) | 50<br>(33–61) | 50<br>(36–62) | 51<br>(38–62) | 52<br>(39–63)  | 53<br>(42–64)  | 53<br>(41–64)  |
| <5                      | 86 (7)        | 92 (7)        | 198 (8)       | 279 (9)       | 358 (9)       | 423 (7)       | 606 (7)       | 638 (6)       | 775 (5)        | 681 (4)        | 928 (4)        |
| 5–14                    | 143 (11)      | 139 (10)      | 173 (7)       | 341 (11)      | 344 (8)       | 401 (7)       | 539 (6)       | 629 (6)       | 811 (5)        | 716 (4)        | 899 (4)        |
| 15–44                   | 373<br>(30)   | 402<br>(30)   | 783<br>(30)   | 1,012<br>(31) | 1,200<br>(29) | 1,543<br>(26) | 2,147<br>(24) | 2,589<br>(23) | 3,683<br>(23)  | 3,738<br>(22)  | 4,578<br>(21)  |
| 45–60                   | 397<br>(32)   | 415<br>(31)   | 878<br>(34)   | 1,001<br>(31) | 1,320<br>(32) | 2,035<br>(34) | 3,169<br>(35) | 4,038<br>(36) | 5,791<br>(36)  | 6,339<br>(37)  | 7,797<br>(36)  |
| ≥61                     | 255<br>(20)   | 292<br>(22)   | 581<br>(22)   | 604<br>(19)   | 866<br>(21)   | 1,618<br>(27) | 2,467<br>(28) | 3,217<br>(29) | 4,975<br>(31)  | 5,819<br>(34)  | 7,360<br>(34)  |
| Residence§              |               |               |               |               |               |               |               |               |                |                |                |
| Rural                   | 892<br>(76)   | 1,023<br>(79) | 2,108<br>(83) | 2,758<br>(87) | 3,512<br>(88) | 5,231<br>(88) | 7,725<br>(87) | 9,622<br>(87) | 13,803<br>(87) | 15,058<br>(88) | 18,597<br>(89) |
| Urban                   | 279<br>(24)   | 264<br>(21)   | 424<br>(17)   | 408<br>(13)   | 477<br>(12)   | 718<br>(12)   | 1,170<br>(13) | 1,443<br>(13) | 2,105<br>(13)  | 2,051<br>(12)  | 2,414<br>(11)  |
| Occupation, farmer      | 733<br>(58)   | 859<br>(64)   | 1,700<br>(65) | 2,079<br>(64) | 2,721<br>(67) | 4,250<br>(71) | 6,179<br>(69) | 7,853<br>(71) | 11,842<br>(74) | 13,230<br>(77) | 16,670<br>(77) |
| No. in endemic province | 16            | 19            | 18            | 22            | 20            | 20            | 23            | 21            | 27             | 26             | 25             |
| No. in endemic county   | 222           | 225           | 327           | 372           | 431           | 509           | 582           | 692           | 769            | 809            | 883            |
| No. in endemic township | 616           | 683           | 1,183         | 1,283         | 1,523         | 2,088         | 2,644         | 3,264         | 4,125          | 4,353          | 5,025          |
| No. deaths              | 0             | 1             | 3             | 0             | 11            | 6             | 9             | 4             | 8              | 3              | 12             |

\*Values are no. (%) except as indicated. IQR, interquartile range.

†No. cases/100,000 population.

‡Some columns do not add up to 100% because of rounding.

§Some columns do not add up to equal the total because of missing data.

**Spatial-Temporal Distribution**

During the 28 years from 1952–1979, nearly all cases (99.9%) occurred in the southern provinces of China; sporadic cases were recorded in several northern provinces. Beginning in 1980, the disease began to expand slowly northward and westward; the number of affected provinces gradually increased from 17 during 1952–1989 to all 31 provinces during 2011–2016 (Figure 2). We saw dramatic expansion in both rural and urban areas. In rural areas, the affected townships increased nearly 10-fold, from 422 in 2006 to 4,083 in 2016, and the affected urban townships expanded from 194 in 2006 to 942 in 2016 (Figure 3). As of 2016, a total of 28% (883/3,101) of all counties and 11% (5,025/44,539) of all townships recorded cases of scrub typhus throughout the country (Appendix Figure 2).

During 2006–2016, nearly all cases (93,187/93,481; 99.7%) occurred in the 15 provinces that had a cumulative case count ≥100. Among these provinces, we identified 3 epidemiologic regions using hierarchical

cluster analysis (Figure 4, panels A, B): middle-east (latitude range 31°–41°N, longitude range 105°–125°E); southwest (latitude range 21°–31°N, longitude range 95°–105°E); and southeast China (latitude range 21°–31°N, longitude range 105°–125°E).

**Seasonal Pattern**


Nationally, the seasonal pattern of scrub typhus incidence begins to increase rapidly in May, reaches a peak in October, and falls in November; 94% of cases occurred from May through November in both periods, 1980–1989 and 2006–2016. Heat maps showed consistent seasonal patterns of scrub typhus within each province and between the 2 periods of 1980–1989 and 2006–2016 (Appendix Figure 3), but we observed geographic diversity of seasonal patterns across China. Multiple linear regression models of time series data confirmed and quantified seasonal characteristics of scrub typhus activity in the 3 epidemiologic regions (p<0.001 for all) (Appendix Tables 6, 7). Specifically, the scrub typhus season

in the southeast extended from calendar week 17 (early May) to calendar week 50 (mid-December), lasting for 34 weeks and capturing 95% of the total reported cases from this region. In the southwest, the season extended from calendar week 21 (late May) to calendar week 48 (early December), for 28 weeks of duration, capturing 95% of total reported cases in the region. In the middle-east, scrub typhus season lasted 12 weeks, from calendar week 38 (late September) to calendar week 49 (mid-


December), capturing 97% of total reported cases in the region (Figure 5, panels A, B).

**Risk for Death**

A total of 174 deaths were recorded throughout the country during the 49 years of this study (1952–1989 and 2006–2016). The overall case-fatality ratio was 0.13% (95% CI 0.11%–0.15%); the case fatality ratio from 2006–2016 was significantly lower than that



**Figure 2.** Incidence rate of scrub typhus for each province of China during 1952–1989 and 2006–2016, by time period. Annual average incidence of scrub typhus per 100,000 population in the 31 provinces investigated is shown. The rings contain data for 11 periods studied; the innermost ring shows data for early periods of 1952–1954, and the outermost ring data for 2011–2016. The latitude and longitude of the capital city of each province are shown.



**Figure 3.** Number of townships with scrub typhus cases reported in China during 2006–2016. A) Number of affected townships for each year, divided by those that were affected in previous years (red bars) and those newly affected townships for each year (blue bars). B) Total number of affected townships for each year in rural (red bars) and urban (light pink bars).

from 1952–1989 (0.06% vs. 0.29%;  $p < 0.001$ ). The following subgroups had a higher risk for death: age  $\geq 40$  years (adjusted odds ratio [aOR] 3.43, 95% CI 1.69–7.72), nonfarmer occupation (aOR 3.56, 95% CI 1.97–6.43), and interval of illness onset to diagnosis  $\geq 8$  days (aOR 2.36, 95% CI 1.07–5.19) (Table 2).

**Discussion**


In this study of a longitudinal surveillance dataset spanning 49 years, we found the occurrence of scrub typhus in China has been maintained at a relatively low level from the 1950s–1980s. However, in the 2000s, incidence rates increased in an unprecedented manner, in both the historically endemic areas and in new areas not previously identified as having cases. In addition, profound epidemiologic changes, including geographic expansion, varying age and sex distribution, and diverse seasonal patterns were further discovered during the period 2006–2016.

In China, some novel natural and socioeconomic factors may have contributed to the increased risk for scrub typhus infection. During the past decade, the reduction of industrial land use, recovering forests and wetlands, and the banning of traditional burning of straw within rural areas have provided a favorable setting for the breeding of rodents and mites, increasing the possibility of human exposure, and the forming of new natural foci for scrub typhus in previously unaffected areas (29). In some other scrub typhus–endemic countries, evidence suggests that a

combination of climate change and the expansion of humans into previously uninhabited areas may play roles in both the reemergence and the apparent rising incidence of scrub typhus (30,31).

Globally, the application of improved surveillance mechanisms and enhanced awareness of clinicians have contributed to greater recognition of scrub typhus in some countries, such as India (1), South Korea (1,6), Laos (6), and Japan (3,32). In China, the overall incidence of all notifiable infectious diseases was relatively stable during 2006–2013 (20). We believe changes in surveillance may have played a minor role in the increase of scrub typhus in China; although the reincorporating of the disease in the National Notifiable Infectious Disease Reporting Information System and the issuing of 2009 national guidelines may have influenced disease awareness among reporters while the increased use of diagnostic tools such as PCR enabled identification of more cases, the change in reporting alone could not fully explain the drastic changes in the geographic and demographic distribution of scrub typhus. More investigations into the pathogen, the vectors, the environment, and the hosts driving the changing epidemiology of scrub typhus in the country should be conducted.

In this study, we determined that most cases of scrub typhus occur in rural areas and in older persons. During 1990–2020, in rural areas of China, a large number of young and middle-aged male persons have traveled to metropolitan areas seeking




**Figure 4.** Epidemiologic regions of scrub typhus in China, 2006–2016. A) Epidemiologic regions based on hierarchical clustering, using the Pearson correlation coefficient matrix between average weekly scrub typhus time series of paired provinces that had a cumulative number of cases >100 in 2006–2016 combined. B) Map of identified epidemiologic regions identified by hierarchical clustering, e.g., middle-east (latitude range 31°–41°N and longitude range 105°–125°E), southwest (latitude range 21°–31°N and longitude range 95°–105°E), southeast of China (latitude range 21°–31°N and longitude range 105°–125°E). Other provinces had a combined total of <100 cases in 2006–2016.

better jobs and higher incomes, often leaving their older, younger, or female family members in rural areas. Older family members >60 years of age commonly performed routine farming activities and so are at higher risk for infection. However, the high incidence of scrub typhus cases in patients <5 years of age in 2 specific provinces, Yunnan and Sichuan, should be addressed, and possible risk factors must be further explored.

We discovered that both the number of scrub typhus cases and the number of affected urban areas are rapidly increasing. Because of improved economic and living conditions in China, more persons live in urban areas and travel to rural areas for recreation and entertainment (e.g., fishing, hiking, picnicking, camping, and picking fruit) on the weekends and on holidays. For urban groups susceptible to scrub typhus, frequent outdoor activities in endemic areas increase the risk for infection (30,33). In 2012, an outbreak of scrub typhus was detected in a city park,

implying that the rodent host and mites may inhabit urban settings. In addition, frequent transportation of farmed goods may lead to the migration of rodents carrying infected mites, which may cause the expansion of scrub typhus to more urban and non-typhus-endemic areas (33,34).

Our study revealed varying seasonal patterns of the scrub typhus epidemic in China, which have also previously been reported in certain nearby countries. In Japan, 2 types of seasonal patterns peaked in weeks 18–25 (April–May) and weeks 43–52 (November–December) and were located north and south of 37°N latitude (35); in South Korea, a single seasonal pattern peaking in October–November was reported (8). The appearance of larvae determines the seasonality of scrub typhus. The heterogeneous geographic distribution of *L. deliense* and *L. scutellare* mites may well explain the diverse seasonal patterns of scrub typhus in the southeast, southwest, and middle-east parts of China (17). Because of the complex interactions among



**Figure 5.** Seasonal characteristics of scrub typhus in China, 2006–2016. A) The weekly number of scrub typhus in year 2016, with a fitted seasonal curve superimposed, by epidemiologic regions identified. B) Duration and peak of scrub typhus seasons in epidemiologic regions. Scrub typhus season was defined as starting with the week in which a fast increase of average weekly case numbers began and ending with the week that the cumulative case numbers captured in the interval between the start and end accounted for >95% of the reported cases in 2006–2016. The colored dot in each region indicates the week with the highest predicted number of scrub typhus cases in the fitted seasonal curve (the peak). The colored bar indicates the number of weeks between season start and end (duration).

vectors, hosts, and humans, as well as climatic factors, the seasonal distribution of scrub typhus outbreaks may also vary locally; a spring outbreak of scrub typhus has occurred on an island in Fujian Province (in the southeastern coastal area) during 2000–2005 (36). Knowing the exact timing of scrub typhus activity at both national and subnational levels is important to healthcare providers and health officials who use this data to guide diagnostic testing, conduct disease surveillance, plan vector control measures, respond to outbreaks, and provide antimicrobial treatment to febrile patients. Travel precautions for those entering scrub typhus–endemic regions, and strategies for controlling mites, should be given before the beginning of scrub typhus season.

Because scrub typhus is a disease that is treatable with specific antimicrobial medicine, early and accurate diagnosis is essential to reduce the risk for severe complications and death. The case-fatality rate of scrub typhus in China was reduced from 13%–16% to 0.2%–0.4% after the introduction of antirickettsial drugs (14). However, the absence of definitive signs

and symptoms make difficult the differentiation of scrub typhus from other common febrile diseases, such as murine typhus, typhoid fever, and leptospirosis. Our study reveals that, compared with diagnosis within 2 days after illness onset, diagnosis later than 8 days after the onset of illness poses a 2.36 times higher risk of death for the patient.

The immunofluorescence assay is the reference diagnostic test for scrub typhus recommended by the World Health Organization (37), but the need for standardization has hampered the method’s widespread use. PCR assay targeting on the gene encoding the 56-kDa type-specific antigen is highly sensitive and useful (38,39) but also has limited uses because of the lack of standardized reagents or commercial diagnostic kits. Currently, the Weil-Felix test, despite its lack of sensitivity and specificity (40), is widely accessible throughout China, specifically in conjunction with a previous exposure history or specific eschars of scrub typhus (41). The finding that delayed diagnosis is associated with death highlights the need for research and development of inexpensive, accurate point-of-



**Table 2.** Risk factors associated with death from scrub typhus, China, 2006–2016

| Factor                                  | No. cases | No. deaths | Case-fatality ratio | Crude odds ratio (95% CI) | Adjusted odds ratio (95% CI)* |
|---|-----------|------------|---------------------|---------------------------|-------------------------------|
| Sex                                     |           |            |                     |                           |                               |
| F                                       | 49,907    | 22         | 0.04                | 1.00                      | 1.00                          |
| M                                       | 43,574    | 35         | 0.08                | <b>1.82 (1.07–3.11)</b>   | <b>1.75 (1.02–3.01)</b>       |
| Age group, y                            |           |            |                     |                           |                               |
| <40                                     | 25,261    | 9          | 0.04                | 1.00                      | 1.00                          |
| 40–59                                   | 37,660    | 19         | 0.05                | 1.50 (0.68–3.30)          | <b>2.86 (1.22–6.69)</b>       |
| ≥60                                     | 30,560    | 29         | 0.09                | <b>2.70 (1.27–5.72)</b>   | <b>5.88 (2.55–13.55)</b>      |
| Residence                               |           |            |                     |                           |                               |
| Rural                                   | 80,329    | 37         | 0.05                | 1.00                      | 1.00                          |
| Urban                                   | 11,753    | 20         | 0.17                | <b>3.31 (1.92–5.70)</b>   | <b>1.89 (1.05–3.42)</b>       |
| Region                                  |           |            |                     |                           |                               |
| Middle-east                             | 25,186    | 2          | 0.01                | 1.00                      | 1.00                          |
| Southeast                               | 45,043    | 44         | 0.10                | <b>12.31 (2.98–50.79)</b> | <b>10.05 (2.41–41.86)</b>     |
| Southwest                               | 23,202    | 11         | 0.05                | <b>5.97 (1.32–26.95)</b>  | <b>9.48 (2.05–43.94)</b>      |
| North and west                          | 50        | 0          | 0                   | NA                        | NA                            |
| Occupation                              |           |            |                     |                           |                               |
| Farmer                                  | 68,116    | 24         | 0.04                | 1.00                      | 1.00                          |
| Nonfarmer                               | 25,365    | 33         | 0.13                | <b>3.7 (2.18–6.25)</b>    | <b>3.56 (1.97–6.43)</b>       |
| Time from illness onset to diagnosis, d |           |            |                     |                           |                               |
| <2                                      | 20,775    | 8          | 0.04                | 1.00                      | 1.00                          |
| 2–7                                     | 42,875    | 22         | 0.05                | 1.33 (0.59–2.99)          | 1.67 (0.74–3.77)              |
| >8                                      | 29,831    | 27         | 0.09                | <b>2.35 (1.07–5.18)</b>   | <b>2.36 (1.07–5.19)</b>       |

\*Result from multivariable logistic regression. Boldface type indicates significance ( $p < 0.05$ ). NA, not available.

care diagnostic tests for the acute phase of infection in clinical settings.

Our study has 3 limitations. First, we collected all study data used from a passive surveillance system and, therefore, data quality may be influenced by the availability of diagnostic technology, underreporting, and the completeness and accuracy of the data over the years. Diagnosis and notification of cases may vary across areas, affecting the geographic comparisons of incidence. Second, individual case data with demographic information were not reported from 1952–1989; therefore, we could only analyze population characteristics, case distributions, and death risk factors from the period 2006–2016. Furthermore, because scrub typhus was removed from the notifiable disease list in 1990, the lack of surveillance data during 1990–2005 limits the consecutiveness of epidemic trend analysis. Third, data on *O. tsutsugamushi* strains and on the accurate distribution of *Leptotrombidium* mite species were unavailable in this study, making it impossible for us to present the whole picture of scrub typhus in China involving human cases, pathogens, hosts, and vectors. However, data used in this study were the most comprehensive and reliable data on scrub typhus available at national and subnational levels in China; these nationwide report data demonstrate striking changes in epidemiologic features of scrub typhus in China, highlighting the need to conduct further high-quality investigations to better interpret the findings from passive surveillance data.

Among the dozens of countries with endemic scrub typhus, China is one of the few that has

established a nationwide surveillance system. By using these long-term incidence data, we thoroughly described the epidemiologic change of scrub typhus over time; our findings are a starting point for further studies to explore additional information related to the global disease burden of scrub typhus. This study may also benefit other scrub typhus–endemic regions outside of China; our findings suggest the possibility of similar epidemiologic changes in areas with similar social and ecologic environments.

As a whole, after ≈40 years of low-level transmission, scrub typhus has become a markedly greater threat in China than previously understood, warranting a higher degree of scrutiny and study to inform health policy. Furthermore, the epidemiologic changes resulting from geographic expansion, demographic transition, and multiple seasonal patterns highlight the need to adjust and enhance current disease prevention and control strategies at national and subnational levels.

#### Acknowledgments

We acknowledge staff members of the county-, district-, and province-level Centers for Disease Control and Prevention of China for their assistance in the field investigation and data collection. We also thank Bob Thompson for his help in manuscript improvement.

This study was supported by grants from the Ministry of Science and Technology of China (2018ZX10101002, 2018ZX10713001, 2018ZX10713001, 2018ZX10713001-005, 2018ZX10101002-003-002) and Emergency Response

Mechanism Operation Program from the Chinese Center for Disease Control and Prevention (131031001000015001). S.E.R., N.D.W., and S.I.H. are primarily supported by the Bill & Melinda Gates Foundation (grant no. OPP1132415).

## About the Author

Dr. Zhongjie Li is an epidemiologist at the Chinese Center for Disease Control and Prevention. His research interests include the epidemiology of infectious disease, disease surveillance, and early-warning systems on disease outbreak detection.

## References


- Walker DH. Scrub typhus – scientific neglect, ever-widening impact. *N Engl J Med*. 2016;375:913–5. <https://doi.org/10.1056/NEJMp1608499>
- Taylor AJ, Paris DH, Newton PN. A systematic review of mortality from untreated scrub typhus (*Orientia tsutsugamushi*). *PLoS Negl Trop Dis*. 2015;9:e0003971. <https://doi.org/10.1371/journal.pntd.0003971>
- Bonell A, Lubell Y, Newton PN, Crump JA, Paris DH. Estimating the burden of scrub typhus: a systematic review. *PLoS Negl Trop Dis*. 2017;11:e0005838. <https://doi.org/10.1371/journal.pntd.0005838>
- Kasper MR, Blair PJ, Touch S, Sokhal B, Yasuda CY, Williams M, et al. Infectious etiologies of acute febrile illness among patients seeking health care in south-central Cambodia. *Am J Trop Med Hyg*. 2012;86:246–53. <https://doi.org/10.4269/ajtmh.2012.11-0409>
- Blacksell SD, Sharma NP, Phumratanaprapin W, Jenjaroen K, Peacock SJ, White NJ, et al. Serological and blood culture investigations of Nepalese fever patients. *Trans R Soc Trop Med Hyg*. 2007;101:686–90. <https://doi.org/10.1016/j.trstmh.2007.02.015>
- Mayxay M, Castonguay-Vanier J, Chansamouth V, Dubot-Pères A, Paris DH, Phetsouvanh R, et al. Causes of non-malarial fever in Laos: a prospective study. *Lancet Glob Health*. 2013;1:e46–54. [https://doi.org/10.1016/S2214-109X\(13\)70008-1](https://doi.org/10.1016/S2214-109X(13)70008-1)
- Paris DH, Shelite TR, Day NP, Walker DH. Unresolved problems related to scrub typhus: a seriously neglected life-threatening disease. *Am J Trop Med Hyg*. 2013;89:301–7. <https://doi.org/10.4269/ajtmh.13-0064>
- Lee HW, Cho PY, Moon SU, Na BK, Kang YJ, Sohn Y, et al. Current situation of scrub typhus in South Korea from 2001–2013. *Parasit Vectors*. 2015;8:238. <https://doi.org/10.1186/s13071-015-0858-6>
- Horton KC, Jiang J, Maina A, Dueger E, Zayed A, Ahmed AA, et al. Evidence of *Rickettsia* and *Orientia* infections among abattoir workers in Djibouti. *Am J Trop Med Hyg*. 2016;95:462–5. <https://doi.org/10.4269/ajtmh.15-0775>
- Maina AN, Farris CM, Odhiambo A, Jiang J, Laktabai J, Armstrong J, et al. Q fever, scrub typhus, and rickettsial diseases in children, Kenya, 2011–2012. *Emerg Infect Dis*. 2016;22:883–6. <https://doi.org/10.3201/eid2205.150953>
- Izzard L, Fuller A, Blacksell SD, Paris DH, Richards AL, Aukkanit N, et al. Isolation of a novel *Orientia* species (*O. chuto* sp. nov.) from a patient infected in Dubai. *J Clin Microbiol*. 2010;48:4404–9. <https://doi.org/10.1128/JCM.01526-10>
- Weitzel T, Dittrich S, López J, Phuklia W, Martinez-Valdebenito C, Velásquez K, et al. Endemic scrub typhus in South America. *N Engl J Med*. 2016;375:954–61. <https://doi.org/10.1056/NEJMoa1603657>
- Hay SI, Abajobir AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, et al.; GBD 2016 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017;390:1260–344. [https://doi.org/10.1016/S0140-6736\(17\)32130-X](https://doi.org/10.1016/S0140-6736(17)32130-X)
- Fan MY, Walker DH, Yu SR, Liu QH. Epidemiology and ecology of rickettsial diseases in the People's Republic of China. *Rev Infect Dis*. 1987;9:823–40. <https://doi.org/10.1093/clinids/9.4.823>
- Zheng C, Jiang D, Ding F, Fu J, Hao M. Spatiotemporal patterns and risk factors for scrub typhus from 2007 to 2017 in southern China. *Clin Infect Dis*. 2019;69:1205–11. <https://doi.org/10.1093/cid/ciy1050>
- Luo Y, Yin J. Progress in research of *Orientia tsutsugamushi* and its host and vector [in Chinese]. *Disease Surveillance*. 2019;(10):920–3.
- Wang G-h, Wang C-j, Li B-j, Jiang Z-k, Ding L-y, Wang L. General situation on studies of animal hosts of tsutsugamushi disease in China [in Chinese]. *Chinese Journal of Hygienic Insecticides & Equipments*. 2013;19:370–3.
- Wu YC, Qian Q, Soares Magalhaes RJ, Han ZH, Hu WB, Haque U, et al. Spatiotemporal dynamics of scrub typhus transmission in mainland China, 2006–2014. *PLoS Negl Trop Dis*. 2016;10:e0004875. <https://doi.org/10.1371/journal.pntd.0004875>
- Sun Y, Wei YH, Yang Y, Ma Y, de Vlas SJ, Yao HW, et al. Rapid increase of scrub typhus incidence in Guangzhou, southern China, 2006–2014. *BMC Infect Dis*. 2017;17:13. <https://doi.org/10.1186/s12879-016-2153-3>
- Li Z, Gao GF. Infectious disease trends in China since the SARS outbreak. *Lancet Infect Dis*. 2017;17:1113–5. [https://doi.org/10.1016/S1473-3099\(17\)30579-0](https://doi.org/10.1016/S1473-3099(17)30579-0)
- Wang L, Wang Y, Jin S, Wu Z, Chin DP, Koplan JP, et al. Emergence and control of infectious diseases in China. *Lancet*. 2008;372:1598–605. [https://doi.org/10.1016/S0140-6736\(08\)61365-3](https://doi.org/10.1016/S0140-6736(08)61365-3)
- Chinese Center for Disease Control and Prevention. National guideline of scrub typhus control and prevention [in Chinese]. 2009 [cited 2017 Mar 16]. [http://www.chinacdc.cn/tzgg/200901/t20090105\\_40316.html](http://www.chinacdc.cn/tzgg/200901/t20090105_40316.html)
- National Bureau of Statistics of China. National data [in Chinese]. 2018 [cited 2020 Mar 27]. <http://data.stats.gov.cn/english/index.htm>
- Kim HJ, Fay MP, Feuer EJ, Midthune DN. Permutation tests for Joinpoint regression with applications to cancer rates. *Stat Med*. 2000;19:335–51. [https://doi.org/10.1002/\(SICI\)1097-0258\(20000215\)19:3<335::AID-SIM336>3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1097-0258(20000215)19:3<335::AID-SIM336>3.0.CO;2-Z)
- Yu H, Alonso WJ, Feng L, Tan Y, Shu Y, Yang W, et al. Characterization of regional influenza seasonality patterns in China and implications for vaccination strategies: spatio-temporal modeling of surveillance data. *PLoS Med*. 2013;10:e1001552. <https://doi.org/10.1371/journal.pmed.1001552>
- Naumova EN, Jagai JS, Matyas B, DeMaria A Jr, MacNeill IB, Griffiths JK. Seasonality in six enterically transmitted diseases and ambient temperature. *Epidemiol Infect*. 2007;135:281–92. <https://doi.org/10.1017/S0950268806006698>
- Xing W, Liao Q, Viboud C, Zhang J, Sun J, Wu JT, et al. Hand, foot, and mouth disease in China, 2008–12: an

- epidemiological study. *Lancet Infect Dis*. 2014;14:308–18. [https://doi.org/10.1016/S1473-3099\(13\)70342-6](https://doi.org/10.1016/S1473-3099(13)70342-6)
28. R Core Team. R: A language and environment for statistical computing. Vienna (Austria): R Foundation for Statistical Computing; 2013.
  29. Liu J, Li S, Ouyang Z, Tam C, Chen X. Ecological and socioeconomic effects of China's policies for ecosystem services. *Proc Natl Acad Sci U S A*. 2008;105:9477–82. <https://doi.org/10.1073/pnas.0706436105>
  30. Tsai PJ, Yeh HC. Scrub typhus islands in the Taiwan area and the association between scrub typhus disease and forest land use and farmer population density: geographically weighted regression. *BMC Infect Dis*. 2013;13:191. <https://doi.org/10.1186/1471-2334-13-191>
  31. Li T, Yang Z, Dong Z, Wang M. Meteorological factors and risk of scrub typhus in Guangzhou, southern China, 2006–2012. *BMC Infect Dis*. 2014;14:139. <https://doi.org/10.1186/1471-2334-14-139>
  32. Matsui T, Kramer MH, Mendlein JM, Osaka K, Ohyama T, Takahashi H, et al. Evaluation of national tsutsugamushi disease surveillance – Japan, 2000. *Jpn J Infect Dis*. 2002; 55:197–203.
  33. Park SW, Ha NY, Ryu B, Bang JH, Song H, Kim Y, et al. Urbanization of scrub typhus disease in South Korea. *PLoS Negl Trop Dis*. 2015;9:e0003814. <https://doi.org/10.1371/journal.pntd.0003814>
  34. Yang LP, Liu J, Wang XJ, Ma W, Jia CX, Jiang BF. Effects of meteorological factors on scrub typhus in a temperate region of China. *Epidemiol Infect*. 2014;142:2217–26. <https://doi.org/10.1017/S0950268813003208>
  35. Yoshikura H. Geographical distribution of Japanese spotted fever and tsutsugamushi disease in Japan – possible effect of environmental temperature. *Jpn J Infect Dis*. 2017;70:349–51. <https://doi.org/10.7883/yoken.JJID.2016.274>
  36. Cao M, Guo H, Tang T, Wang C, Li X, Pan X, et al. Spring scrub typhus, People's Republic of China. *Emerg Infect Dis*. 2006;12:1463–5. <https://doi.org/10.3201/eid1209.060257>
  37. Janardhanan J, Trowbridge P, Varghese GM. Diagnosis of scrub typhus. *Expert Rev Anti Infect Ther*. 2014;12:1533–40. <https://doi.org/10.1586/14787210.2014.974559>
  38. Fournier PE, Siritantikorn S, Rolain JM, Suputtamongkol Y, Hoontrakul S, Charoenwat S, et al. Detection of new genotypes of *Orientia tsutsugamushi* infecting humans in Thailand. *Clin Microbiol Infect*. 2008;14:168–73. <https://doi.org/10.1111/j.1469-0691.2007.01889.x>
  39. Bechah Y, Socolovschi C, Raoult D. Identification of rickettsial infections by using cutaneous swab specimens and PCR. *Emerg Infect Dis*. 2011;17:83–6. <https://doi.org/10.3201/eid1701.100854>
  40. Kováčová E, Kazár J. Rickettsial diseases and their serological diagnosis. *Clin Lab*. 2000;46:239–45.
  41. Zheng MH, Shi KQ, Chen YP. Skin lesion in a critically ill man. *BMJ*. 2015;351:h4570. <https://doi.org/10.1136/bmj.h4570>

Address for correspondence: Jianxing Yu, State Key Laboratory of Infectious Disease Prevention and Control, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Changbai Rd 155#, Changping District, Beijing, 102206, China; email: yujianxing@icdc.cn

# EID Podcast: The Red Boy, the Black Cat

Byron Breedlove, managing editor of *Emerging Infectious Diseases*, discusses the June 2019 EID cover artwork, a painting of Don Manuel Osorio Manrique de Zuniga, by Francisco de Goya y Lucientes.



Visit our website to listen:  
<https://go.usa.gov/xysv5>

**EMERGING  
INFECTIOUS DISEASES®**

# Epidemiologic Changes of Scrub Typhus in China, 1952–2016

## Appendix

### Characteristics of Scrub Typhus Seasonality

To characterize the seasonality of scrub typhus, we used the Poisson regression model to estimate the peak timing and amplitude of the annual and semiannual periodicities of scrub typhus. The generic model formula takes the following form, as described previously (1–3).

$$\log(Y_i) = a + b * \cos\left(2\pi * \frac{W_i}{52.17}\right) + c * \sin\left(2\pi * \frac{W_i}{52.17}\right) + d * \cos\left(4\pi * \frac{W_i}{52.17}\right) + e * \sin\left(4\pi * \frac{W_i}{52.17}\right)$$

where  $Y_i$  are the weekly number of scrub typhus cases during 2006–2016 aggregated by epidemiological regions, which were log transformed.  $w_i$  is a running index for week, and  $a$ ,  $b$ ,  $c$ ,  $d$ , and  $e$  are the intercept and seasonal terms to be estimated from the data.

Specifically,

the amplitude of the annual periodicity is estimated as  $\text{AnnAmp} = \sqrt{b^2 + c^2}$ ;

the shift of the annual periodicity is estimated as  $\text{AnnPeakTiming} = -\text{atan}(c/b)$ ;

the amplitude of the semi-annual periodicity is estimated as  $\text{SemiAnnAmp} = \sqrt{d^2 + e^2}$ ; and

the shift of the semi-annual periodicity is estimated as  $\text{SemiAnnPeakTiming} = -\text{atan}(d/e)$ .

**Appendix Table 1.** Summary of laws or regulations related to scrub typhus surveillance and control, mainland China, 1952–2016

| ID | Title   | Approved by  | Issued by                                  | Period enforced   | Notes  |
|----|---|--|--|---|--|
| 1  | The Administrative Measures of Infectious Disease         | The State Council, PRC, on 1955 Jun 1  | The Ministry of Health, PRC on 1955 Jul 5  | 1955 Jul 5–1978 Sep 19  | Scrub typhus was one of the 18 notifiable infectious diseases  |
| 2  | The Administrative Regulation of Acute Infectious Disease | The State Council, PRC   | The Ministry of Health, PRC on 1978 Sep 20 | 1978 Sep 20   | Scrub typhus was one of the 25 notifiable infectious diseases  |
| 3  | The Law on Infectious Disease Control and Prevention      | The Standing Committee of the National People's Congress, PRC, on 1989 Feb 21 and revised on 2004 Aug 28 | The National                               | 1989 Sep 1–2004 Nov 30; the revised version during 2004 Dec 1–2009 Jan. 3 | Scrub typhus was removed from the list of notifiable diseases  |
| 4  | National Scrub Typhus Control and Prevention Guideline    | Issued by Chinese Center for Disease Control and Prevention on 2009 Jan. 4                               |  | 2009 Jan. 4–current   | Scrub typhus was required to be reported in the National Notifiable Infectious Disease Reporting Information System (NNIDRIS), at the Chinese Centers for Disease Control and Prevention (China CDC) |

**Appendix Table 2.** Summary of diagnosis criteria and classification for scrub typhus, mainland China

| Variable                     | 1952–2008  | 2009–present  |
|------------------------------|--|---|
| Criteria or guidelines       | Educational Book on Infectious Diseases (First–eighth edition)   | National Scrub Typhus Control and Prevention Guideline. 2009  |
| Issued by                    | Beijing: People's Medical Publishing House   | Chinese Center for Disease Control and Prevention   |
| Epidemiology linkage         | 1.1 An individual who experienced possible outdoor exposure to mite bites three weeks before the onset of illness, i.e., farming, fishing, camping, and straw collection, during the epidemic season of the disease  |   |
| Clinical description         | 2.1 Sudden high fever accompanied by characteristic eschar or ulcer.<br>2.2 Enlarged lymph nodes, skin rash, splenomegaly, or hepatomegaly.  | 2.1 Fever<br>2.2 Lymphadenopathy<br>2.3 Skin rash<br>2.4 Specific eschars/ulcers  |
| Laboratory tests             | 3.1 An agglutination titer $\geq 1:160$ in the Weil-Felix test using the OXK strain of <i>Proteus mirabilis</i> .<br>3.2 Seroconversion or a four-fold or greater rise in serum IgG antibody titers between acute and convalescent sera detected by using mixed Gilliam, Karp, Kato, and Kawasaki strains of <i>O. tsutsugamushi</i> as diagnostic antigen in indirect immunofluorescence antibody assay (IFA).<br>3.3 The detection of <i>O. tsutsugamushi</i> 56-kDa gene by polymerase chain reaction in clinical specimens.<br>3.4 The isolation of <i>O. tsutsugamushi</i> from clinical specimens. |   |
| Diagnosis and classification | Probable or confirmed cases: a patient with any 3 items among 1.1, 2.1, 2.2, 3.1, 3.2.<br><br>The case was not further classified as probable or confirmed cases.  | Suspected case:<br>(1) a patient with item 1.1, 2.1, plus either 2.2 or 2.3, and was excluded from other diseases. Or<br>(2) a patient with item 2.1, 2.2 and 2.3 during the local epidemic season of scrub typhus (May–November south of Yangtze River and October–November in north of Yangtze River).<br><br>Probable case:<br>(1) a suspected case with item 2.4. or<br>(2) a patient with item of 1.1, 2.1, and 2.4.<br><br>Confirmed case:<br>(1) a probable case with any one of the item 3.1–3.4. or<br>(2) a suspected case with any one of the item 3.2–3.4 |

**Appendix Table 3.** Variables in the aggregated dataset of scrub typhus cases in China, 1952–1989\*

| Variables*               | Aggregated                   | Period    |
|--------------------------|------------------------------|-----------|
| No. cases and no. deaths | By month at national level   | 1952–1989 |
|                          | By year at provincial level  | 1952–1979 |
|                          | By month at provincial level | 1980–1989 |
| Annual incidence rate†   | At national level            | 1952–1989 |
|                          | At provincial level          | 1952–1989 |
| Annual mortality rate†   | At national level            | 1952–1989 |
|                          | At provincial level          | 1952–1989 |

\*The data were aggregated and reported monthly by each province in China, 1952–1989.

†Per 100,000 population

**Appendix Table 4.** Variables in the individual dataset of scrub typhus cases in China, 2006-2016\*

| Variables                      | Definition/Classification                                | Completeness  |
|--------------------------------|--|---------------|
| Identification                 | A unique 8-digital number for each case                  | 100% reported |
| Sex                            | Male and female  | 100% reported |
| Age                            | Interval from the date of birth to onset date of illness | 100% reported |
| Zone code of residence address | Unique 8-digital number at county and township level     | 100% reported |
| Occupation                     | Occupation/status of case patients                       | 100% reported |
| Case classification            | Suspected case   | 100% reported |
|                                | Probable case  |               |
|                                | Confirmed case   |               |
| Date of illness onset          | Date of illness onset                                    | 100% reported |
| Date of diagnosis              | Date of diagnosis as a probable or confirmed case        | 100% reported |
| Date of report                 | First date of reporting to surveillance system           | 100% reported |
| Date of death                  | Date of case-patient death, if applicable                | 100% reported |

\*Data were reported by doctors within 24 hours after diagnosis to the online National Notifiable Infectious Disease Reporting Information System (NNIDRIS) since 2004. NNIDRIS enables all the healthcare institutes across the country to report individual cases of scrub typhus rapidly through the internet to the data center located in the Chinese Center for Disease Control and Prevention.

**Appendix Table 5.** Geographic information for provinces in China

| Province       | Inland or coastal | Region         | Capital city | Latitude, ° | Longitude, ° |
|----------------|-------------------|----------------|--------------|-------------|--------------|
| Heilongjiang   | Inland            | North and west | Harbin       | 45.45       | 126.41       |
| Xinjiang       | Inland            | North and west | Urumqi       | 43.79       | 87.57        |
| Jilin          | Inland            | North and west | Changchun    | 43.52       | 125.19       |
| Liaoning       | Coastal           | Middle-east    | Shenyang     | 41.50       | 123.24       |
| Inner Mongolia | Inland            | North and west | Hohhot       | 40.49       | 111.48       |
| Beijing        | Inland            | Middle-east    | Beijing      | 39.54       | 116.28       |
| Tianjin        | Coastal           | Middle-east    | Tianjin      | 39.09       | 117.11       |
| Ningxia        | Inland            | North and west | Yinchuan     | 38.20       | 106.16       |
| Hebei          | Coastal           | Middle-east    | Shijiazhuang | 38.02       | 114.28       |
| Shanxi         | Inland            | Middle-east    | Tanyuan      | 37.52       | 112.34       |
| Qinghai        | Inland            | North and west | Xining       | 36.38       | 101.45       |
| Shandong       | Coastal           | Middle-east    | Jinan        | 36.38       | 117.00       |
| Gansu          | Inland            | North and west | Lanzhou      | 36.03       | 103.49       |
| Henan          | Inland            | Middle-east    | Zhengzhou    | 34.48       | 113.42       |
| Shaanxi        | Inland            | Middle-east    | Xi'an        | 34.16       | 108.54       |
| Jiangsu        | Coastal           | Middle-east    | Nanjing      | 32.02       | 118.50       |
| Anhui          | Inland            | Middle-east    | Hefei        | 31.51       | 117.18       |
| Shanghai       | Coastal           | Southeast      | Shanghai     | 31.14       | 121.29       |
| Sichuan        | Inland            | Southwest      | Chengdu      | 30.39       | 104.05       |
| Hubei          | Inland            | Southeast      | Wuhan        | 30.37       | 114.21       |
| Zhejiang       | Coastal           | Southeast      | Hangzhou     | 30.14       | 120.09       |
| Tibet          | Inland            | Southwest      | Lhasa        | 29.40       | 91.10        |
| Chongqing      | Inland            | Southeast      | Chongqing    | 29.32       | 106.32       |
| Jiangxi        | Inland            | Southeast      | Nanchang     | 28.41       | 115.52       |
| Hunan          | Inland            | Southeast      | Changsha     | 28.11       | 113.00       |
| Guizhou        | Inland            | Southeast      | Guiyang      | 26.35       | 106.42       |
| Fujian         | Coastal           | Southeast      | Fuzhou       | 26.05       | 119.18       |
| Yunnan         | Inland            | Southwest      | Kunming      | 25.00       | 102.41       |
| Guangdong      | Coastal           | Southeast      | Guangzhou    | 23.08       | 113.15       |
| Guangxi        | Coastal           | Southeast      | Nanning      | 22.48       | 108.20       |
| Hainan         | Coastal           | Southeast      | Haikou       | 20.02       | 110.20       |



**Appendix Table 6.** Model regression results of scrub typhus seasonality in 3 epidemiological regions of China, 2006–2016\*


| Item | Southwest            |       |         | Southeast            |       |         | Middle-east          |       |         |
|------|----------------------|-------|---------|----------------------|-------|---------|----------------------|-------|---------|
|      | Model<br>coefficient | SE    | p value | Model<br>coefficient | SE    | p value | Model<br>coefficient | SE    | p value |
| a    | 3.684                | 0.027 | <0.001  | 4.866                | 0.018 | <0.001  | 1.1539               | 0.047 | <0.001  |
| b    | -1.109               | 0.019 | <0.001  | -1.068               | 0.011 | <0.001  | -0.179               | 0.068 | 0.009   |
| c    | -1.97                | 0.026 | <0.001  | -1.259               | 0.012 | <0.001  | -2.852               | 0.065 | <0.001  |
| d    | 0.243                | 0.015 | <0.001  | 0.208                | 0.008 | <0.001  | -1.286               | 0.044 | <0.001  |
| e    | -0.042               | 0.015 | 0.005   | -0.587               | 0.009 | <0.001  | -3.411               | 0.041 | <0.001  |

\*Items a, b, c, d, and e are the intercept and seasonal terms to be estimated from the data. SE, standard error.


**Appendix Table 7.** Seasonal characteristics of scrub typhus in the 3 epidemiological regions of China, 2006–2016\*

| Epidemiologic<br>region | Intercept | AnnAmpi | AnnPeakTiming | SemiAnnAmpi | SemiAnnPeakTiming |
|-------------------------|-----------|---------|---------------|-------------|-------------------|
| Southwest China         | 3.684     | 2.26    | 0.332         | 0.247       | 0.027             |
| Southeast China         | 4.866     | 1.651   | 0.362         | 0.623       | 0.196             |
| Middle-east China       | 1.1539    | 2.858   | 0.26          | 3.646       | 0.307             |


\*Values provided are in logarithmic scale. AnnAmpi, amplitude of annual periodicity; AnnPeakTiming, shift of the annual periodicity; SemiAnnAmpi, amplitude of the semi-annual periodicity; SemiAnnPeakTiming, shift of the semiannual periodicity.



**Appendix Figure 1.** Temporal trends of incidence rate of scrub typhus in China, 1952–2016. A) Overall trend of overall incidence from 1952–2016. B) Annual percent change (APC) of incidence rate during 1952–1980. C) APC of incidence rate during 1981–1989. (D) APC of incidence rate during 2006–2016. Black dots indicate observed incidence rates; red lines indicates incidence trends; orange shading indicates 95% CI.



**Appendix Figure 2.** Geographic expansion of scrub typhus in China, 2006–2016.



**Appendix Figure 3.** Heat map of provinces with scrub typhus, by epidemiological region and the latitude of the capital city of each province. A) Time series of monthly cases during 1980–1989, standardized by the annual number of cases reported by each province and standardized to a range of 0–1. B) Seasonal distribution of cases of scrub typhus by province, plotted as the mean value of the proportion of cases in each month of the year from 1980–1989. (C) Time series of weekly cases during 2006–2016, standardized by the annual number of cases reported by each province and standardized to a range of 0–1. (D) Seasonal distribution of cases of scrub typhus by province, plotted as the mean value of the proportion of cases in each week of the year from 2006–2016. Asterisk (\*) indicates provinces in which the total number of cases in 2006–2016 combined was <100.

## References

- <jrn>1. Yu H, Alonso WJ, Feng L, Tan Y, Shu Y, Yang W, et al. Characterization of regional influenza seasonality patterns in China and implications for vaccination strategies: spatio-temporal modeling of surveillance data. *PLoS Med*. 2013;10:e1001552. [PubMed](#)  
<https://doi.org/10.1371/journal.pmed.1001552></jrn>
- <jrn>26. Naumova EN, Jagai JS, Matyas B, DeMaria A Jr, MacNeill IB, Griffiths JK. Seasonality in six enterically transmitted diseases and ambient temperature. *Epidemiol Infect*. 2007;135:281–92. [PubMed](#) <https://doi.org/10.1017/S0950268806006698></jrn>
- <jrn>27. Xing W, Liao Q, Viboud C, Zhang J, Sun J, Wu JT, et al. Hand, foot, and mouth disease in China, 2008–12: an epidemiological study. *Lancet Infect Dis*. 2014;14:308–18. [PubMed](#)  
[https://doi.org/10.1016/S1473-3099\(13\)70342-6](https://doi.org/10.1016/S1473-3099(13)70342-6)</jrn>