

Transition Operators Entering Neutrinos Double Electron Capture to Excited Nuclear States

J. D. Vergados^{1,2} *

¹ *Max Planck für Kernphysik, Saupfercheckweg 1, 69120 Heidelberg, Germany and*
² *CERN theory Division, Geneva, Switzerland*

(Dated: December 2, 2021)

We construct the effective transition operators relevant for neutrinoless double electron capture leading to final nuclear states different than 0^+ . From the structure of these operators we see that, if such a process is observed experimentally, it will be very helpful in singling out the very important light neutrino mass contribution from the other lepton violating mechanisms.

PACS numbers: 14.60.Pq; 13.15.+g; 23.40.Bw; 29.40.-n; 29.40.Cs

INTRODUCTION

The nuclear double beta decay is process considered very long time ago[1]. It can occur whenever the ordinary (single) beta decay is forbidden due to energy conservation or greatly suppressed due to angular momentum mismatch. It could proceed with the emission of two neutrinos or in a neutrinoless mode. The $2\nu\beta\beta$ -decay is just an example of a second order weak interaction. The exotic neutrinoless double beta decay ($0\nu\beta\beta$ -decay) is the most interesting since it violates lepton number by two units. It was first considered by Furry [2] more than half a century ago as soon it was realized that the neutrino might be a Majorana particle. To-day, seventy years later, $0\nu\beta\beta$ -decay:

$$(A, Z) \rightarrow (A, Z + 2) + e^- + e^-, \quad (0\nu \beta\beta \text{ decay}), \quad (1)$$

$\Delta = Q$, continues to be one of the most interesting processes, since:

- It will establish whether there exist lepton violating processes in the Universe.
- It is perhaps the only process, which can decide whether the neutrinos are Majorana particles, i.e. the particle coincides with its charge conjugate, or Dirac particles, if it does not.
- It will aid in settling the outstanding question of the absolute scale of the neutrino masses, especially if this scale turns out to be very small. Neutrino oscillation experiments can only determine the mass squared differences.

For recent reviews see [3–6].

If this process is allowed other related processes in which the charge of the nucleus is decreased by two units may also occur, if they happen to be allowed by energy and angular momentum conservation laws, e.g.

$$(A, Z) \rightarrow (A, Z - 2) + e^+ + e^+ \quad (0\nu \text{ positron emission}), \quad (2)$$

$$\Delta = Q - 4m_e c^2,$$

$$(A, Z) + e^- \rightarrow (A, Z - 2) + e^+ \quad (0\nu \text{ electron positron conversion}), \quad (3)$$

$$\Delta = Q - 2m_e c^2,$$

$$(A, Z) + e^- + e^- \rightarrow (A, Z - 2) + \text{xrays} \quad (0\nu \text{ double electron capture}), \quad (4)$$

$\Delta = Q$, where Δ is the available energy.

* Permanent address: Theoretical Physics Division, University of Ioannina, Ioannina, Gr 451 10, Greece

The last process is always possible, whenever (3) is. It has been first considered long time ago [7, 8] as a two step process: In the first step the two neutral atoms, (A, Z) and the excited $(A, Z-2)$, get admixed via the lepton number violating interaction. In the second step the $(A, Z-2)$ atom de-excites emitting two hard x-rays and the nucleus, if it is found in an excited state, de-excites emitting γ -rays. The life time expected was very long, since the above mixing amplitude was tiny compared to the energy difference of the two atoms involved. It has recently, however, been gaining in importance [9–12] after ion Penning traps [13] made it possible to accurately determine the Q values, which gave rise to the presence of resonances, which in turn could lead to an increase of the width by many orders of magnitude [9]. Decays to excited states are in some cases possible and provide additional experimental information, e.g. γ rays following their de-excitation.

Furthermore neutrinoless double electron capture is of experimental interest, since one expects the reaction to be essentially free of the two neutrino background, which plagues ordinary neutrinoless double beta decay. It has, also recently been realized [10] that the resonance condition can be fulfilled for transitions to excited states. Thus, e.g. in the case of $^{156}\text{Dy} \rightarrow ^{156}\text{Gd}$, such excited states can be $0^-, 1^-, 0^+, 1^+$ and 2^+ .

Before proceeding further in examining neutrinoless double electron capture, we note the following:

- Transitions to $0^-, 1^-$ states have not been considered previously in standard double beta decay, since they are energetically forbidden. Such transitions can naturally occur, however, in the presence of right handed currents, arising by interference between the left handed and right handed currents, i.e. of the type $j_L \times j_R$, with an amplitude is essentially independent of the neutrino mass. We will show that this process can occur in double electron capture, even though the electrons are bound.
- Transitions to 0^+ states, both the ground states and excited states, can occur both in the mass mechanism and the $j_L \times j_R$. The latter contribution in ordinary double beta decay occurs since the produced electrons have sufficiently high momenta. In double electron capture instead of the electron momentum one can exploit the nuclear recoil term.
- transitions to 1^+ and 2^+ can occur only, if the electron momentum is sufficiently high, as in neutrinoless double beta decay, or by retaining the nuclear recoil term in the hadronic current. In double electron capture this can naturally happen, if one of the electrons is captured from a p-orbit.

In the present paper we will derive the effective transition operators relevant for neutrinoless double electron capture, emphasizing the mode which is independent of the neutrino mass. The neutrino mass contribution has already previously examined, see e.g. the recent work and references therein [9].

NUCLEAR TRANSITION OPERATORS LEADING TO NEGATIVE PARITY STATES

To leading order such transitions cannot occur in neutrino mass mechanism. In the $j_L \times j_R$ case, however, the chiralities involved are such that in the intermediate neutrino propagator one picks the momentum, not the mass. The neutrino momentum leads to the following:

- The time component of the momentum leads to an average energy difference $E_{n,1} - E_{n,2}$, where $E_{n,i} = \langle E_n \rangle - M_i + E_e(i)$, with $E_e(i)$ the electron energy, $i = 1, 2$. This can lead to positive parity transitions in double beta decay, but it is not the dominant contribution.
- the space part of the momentum leads to an effective operator which proportional to $\hat{r} \frac{d}{dr} \frac{1}{r}$ with $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$, $r = |\mathbf{r}_1 - \mathbf{r}_2|$. This naturally leads to negative parity transitions.

¹ After completion of this work, two papers have been published [14, 15], which show that the resonance condition can be met for such excited states

Positive parity transitions can, of course, occur in neutrinoless double beta decay via this mechanism by retaining in the dipole term in the expansion of the essentially distorted plane wave function of the two electrons, which leads to the term $i(\mathbf{p}_1 \cdot \mathbf{r}_1 + \mathbf{p}_2 \cdot \mathbf{r}_2) \hat{r}_{12} \frac{d}{dr} \frac{1}{r}$. This possibility does not exist in neutrinoless double electron capture. We will, however, examine alternate possibilities in the next section.

The effective lepton violating transition operator, within the closure approximation over the intermediate states, which holds for neutrinoless double beta decay and is expected to be even better in neutrinoless double electron capture, can be cast in the form:

$$\mathcal{M} = \frac{1}{\sqrt{2}} \left(\frac{G_F}{\sqrt{2}} \right)^2 2 \frac{2i}{8\pi R_0^2} u_{\mu\nu k}^R \sum_{n,m} \phi_{n_1,0}(r_n) \phi_{n_2,0}(r_m) J_L^\mu(n) J_L^\nu(m) \hat{r}_{n,m}^k f(x_{n,m}) + L \leftrightarrow R \quad (5)$$

with $\phi_{n_1,0}(r_n) \phi_{n_2,0}(r_m)$ ($r_n = r_1$, $r_m = r_2$) the bound s -electron wave functions and

$$u_{\mu\nu k}^R = \bar{u}(p_1) \gamma_\mu \gamma_k \gamma_\nu (1 - \gamma_5) u^c(p_2), \quad u_{\mu\nu k}^L = \bar{u}(p_1) \gamma_\mu \gamma_k \gamma_\nu (1 + \gamma_5) u^c(p_2) \quad (6)$$

$$J_L^\mu(n) = \tau_-(n)(g_V, -g_A \sigma_n), \quad J_L^\nu(m) = \tau_-(m)(g_V, +g_A \sigma_m) \quad (7)$$

$$f(x_{n,m}) = \frac{d}{dx} \frac{1}{x} J(\delta_0), \quad x = x(n, m) = |\mathbf{r}_n - \mathbf{r}_m|/R_0 \quad (8)$$

$$J(\delta_0) = \frac{2}{\pi} \int_0^\infty \frac{\sin y}{y + \delta_0} dy, \quad \delta_0 = (\prec E_n \succ - M_i) |\mathbf{r}_n - \mathbf{r}_m| \quad (9)$$

in double electron capture:

$$\delta_0 \approx 0, \quad J(\delta_0) \approx 1 \text{ and } f(x) \approx -\frac{1}{x^2}$$

For non relativistic electrons one finds that the effective transition operator can be cast in the form:

1. An operator associated with the time structure of the leptonic current:

$$\ell_0 = \bar{u}(p_1) \gamma_0 \gamma_5 u^c(p_2), \quad (10)$$

which is

$$\tilde{\Omega}_0 = \frac{G_F^2}{2\sqrt{2}} \frac{i}{\pi R_0^2} 4g_A^2 \sum_{n,m} \phi_{n_1,0}(r_n) \phi_{n_2,0}(r_m) \tau_-(n) \tau_-(m) \hat{r}_{nm} \cdot \omega_0(n, m) f(x_{n,m}), \quad \omega_0(n, m) = i(\sigma_n \times \sigma_m) \quad (11)$$

which can cause $0^+ \rightarrow 0^-$ transitions.

2. An operator associated with the space structure of the leptonic current:

$$\vec{\ell} = \bar{u}(p_1) \vec{\gamma} u^c(p_2) \quad (12)$$

that is

$$\tilde{\Omega}_1 = \frac{G_F^2}{2\sqrt{2}} \frac{i}{\pi R_0^2} g_V g_A \sum_{n,m} \phi_{n_1,0}(r_n) \phi_{n_2,0}(r_m) 2\tau_-(n) \tau_-(m) [\hat{r}_{nm} \times \omega_1(n, m)] f(x_{n,m}), \quad \omega_1(n, m) = (\sigma_n - \sigma_m) \quad (13)$$

Since the $0s$ electron wave functions vary slowly inside the nucleus one can replace them by there average value and write:

$$\tilde{\Omega}_i = \Lambda_{\text{eff}} \langle (0s)^2 a_B^3 \rangle \Omega_i, \quad \Lambda_{\text{eff}} = \frac{G_F^2}{2\sqrt{2}} \frac{i}{\pi R_0^2 a_B^3} \quad (14)$$

where the Λ_{eff} has dimensions of energy and the average is over the radial part alone (the integral over the angles is unity). The dimensionless effective transition operators now become:

$$\Omega_0 = 4g_A^2 \sum_{n,m} \tau_-(n)\tau_-(m)\hat{r}_{nm}\omega_1(n,m)f(x_{n,m}), \quad \omega_0(n,m) = i(\sigma_n \times \sigma_m) \quad (15)$$

$$\Omega_1 = 2g_V g_A \sum_{n,m} \tau_-(n)\tau_-(m)\hat{r}_{nm} \times \omega_1(n,m)f(x_{n,m}), \quad \omega_1(n,m) = \sigma_n - \sigma_m \quad (16)$$

One can show that averaging over the initial polarizations and summing over the final ones one gets

$$|ME|^2 = 2 \sum_{i=0}^1 | \langle f | |\Omega_i^k| | 0^+ \rangle |^2, \quad k = i = J_f = \text{rank of the operator} \quad (17)$$

where the factor of 2 comes from the leptonic current, $k = J_f$ is the rank of the operator and the double bar indicates its standard reduced ME, which can cause $0^+ \rightarrow 1^-$ transitions. The mixing V^2 [14] between the two atoms takes the form:

$$V^2 = \Lambda_{\text{eff}}^2 \langle (0s)^2 a_B^3 \rangle^2 |ME|^2 \quad (18)$$

TRANSITIONS TO POSITIVE PARITY STATES IN ELECTRON CAPTURE

We will distinguish two cases:

The captured electrons are in s states

As we have mentioned the $0^+ \rightarrow 0^+$ transitions associated with double electron capture have already been studied during the last few years. The contribution arising from the neutrino mass independent mechanism has not been studied since the electron wave function cannot provide the extra parity change. This, however, can be accomplished via the recoil term in the nucleon current². In other words one of the hadronic currents is of the form discussed above, but the other is of the form [16]:

$$J_L^\mu(n) = \tau_-(n)(g_V \mathbf{D}_n, -g_A C_n), \quad J_R^\nu(m) = \tau_-(m)(g_V \mathbf{D}_m, +g_A C_m) \quad (19)$$

with

$$C_n = \frac{1}{2m_N} \sigma_n \cdot (q_n - 2P_n), \quad \mathbf{D} = \frac{1}{2m_N} [(q_n - 2P_n) = i\sigma_n \times q_n] \quad (20)$$

Where m_N and P_n are the nucleon mass and momentum of nucleon n , while q_n is the momentum transfer to that nucleon.

The procedure is a bit tedious but one can proceed as above to arrive at

² It is interesting to note that in the case of neutrinoless double beta decay for the mass independent contribution one needs a correction either from the dipole term of the electron wave function or the nucleon recoil. Both are first order corrections. The nucleon recoil term has not been extensively studied even though it has been suggested by Kotani *al* long time ago [16]

1. One operator associated with the leptonic current:

$$\vec{\ell} = \bar{u}(p_1)\vec{\gamma}\gamma_5 u^c(p_2) \quad (21)$$

which is associated with an operator appropriate for $0^+ \rightarrow 1^+$ transitions, namely:

$$\tilde{\Omega}_1^{\text{recoil}} = \frac{G_F^2}{2\sqrt{2}} \frac{i}{\pi R^2} 2 \sum_{n,m} \phi_{n_1,0}(r_n) \phi_{n_2,0}(r_m) \tau_-(n) \tau_-(m) (\vec{\omega}_{nm} + \vec{\omega}'_{nm}) I(x(n,m)) \quad (22)$$

with

$$\vec{\omega}_{nm} = g_V g_A \hat{r}_{nm} (C_n - C_m) \quad (23)$$

$$\vec{\omega}'_{mn} = g_V^2 i (\mathbf{D}_n - \mathbf{D}_m) \times \hat{r}_{n,m} \quad (24)$$

2. One operator associated with the leptonic current:

$$\ell_5 = \bar{u}(p_1)\gamma_5 u^c(p_2) \quad (25)$$

which is suitable for $0^+ \rightarrow 0^+$ transitions, namely:

$$\tilde{\Omega}_0^{\text{recoil}} = \frac{G_F^2}{2\sqrt{2}} \frac{i}{\pi R_0^2} 2 \sum_{n,m} \phi_{n_1,0}(r_n) \phi_{n_2,0}(r_m) \tau_-(n) \tau_-(m) (\omega_{nm} + \omega'_{nm}) \quad (26)$$

$$\omega_{nm} = g_V g_A \hat{r}_{nm} \cdot [i(-\sigma_n \times D_m + \sigma_m \times D_n)] \quad (27)$$

and

$$\omega'_{nm} = g_A^2 \hat{r}_{nm} \cdot [-\sigma_n C_m + \sigma_m C_n] \quad (28)$$

Under the factorization approximation one can write:

$$\tilde{\Omega}_i^{\text{recoil}} = \Lambda_{\text{eff}} \prec (0s)^2 a_B^3 \succ \Omega_i^{\text{recoil}}, \quad i = 0, 1 \quad (29)$$

$$\Omega_1^{\text{recoil}} = 2 \sum_{n,m} \tau_-(n) \tau_-(m) (\vec{\omega}_{nm} + \vec{\omega}'_{nm}) I(x(n,m)), \quad (30)$$

$$\Omega_0^{\text{recoil}} = 2 \sum_{n,m} \tau_-(n) \tau_-(m) (\omega_{nm} + \omega'_{nm}) I(x(n,m)) \quad (31)$$

and then use Eqs (17) and (18).

There remains the problem of transforming the operators C_n and \mathbf{D}_n into coordinate space. The operator \mathbf{q}_n is local and leads to the operator $-i \frac{1}{R_0} \hat{r} \frac{d}{dx}$, where $\mathbf{x} = \mathbf{r}/R_0$, \mathbf{r} being the relative distance of the two nucleons. The operator P_n is non local. Thus one finds:

$$P_n \rightarrow -i \frac{1}{R_0} \left(\overleftarrow{\nabla}_n - \overrightarrow{\nabla}_n \right) \quad (32)$$

where the operator ∇_n operates with respect to \mathbf{x}_n on the nuclear wave functions on the bra and ket as indicated by the arrows. As a simple example we will rewrite the operator the operator $\hat{r}_{nm}(C_n - C_m)$ in coordinate space. We find:

$$\begin{aligned} \hat{r}_{nm}(C_n - C_m) &= \frac{-i}{2M_n R_0} \hat{r}_{nm} \left\{ (\hat{r}_{nm} \cdot \sigma_n - \hat{r}_{nm} \cdot \sigma_m) \frac{df}{dx} \right. \\ &\quad \left. - \left[\left(\overleftarrow{\nabla}_n f(x) - f(x) \overrightarrow{\nabla}_n \right) \cdot \sigma_n - \left(\overleftarrow{\nabla}_m f(x) - f(x) \overrightarrow{\nabla}_m \right) \cdot \sigma_m \right] \right\} \end{aligned} \quad (33)$$

This can be rewritten by showing the tensor character of the various components as:

$$\begin{aligned}
\hat{r}_{nm}(C_n - C_m) = & \frac{-i}{2M_n R_0} \left\{ \left[-\frac{1}{\sqrt{3}}(\sigma_n - \sigma_m) - \frac{2\sqrt{2}}{\sqrt{3}} \left([\hat{r}_{nm} \times \hat{r}_{nm}]^2 \otimes (\sigma_n - \sigma_m) \right) \right] \frac{df}{dx} \right. \\
& - \frac{1}{\sqrt{3}} \left(\hat{r}_{nm} \cdot \left(\overleftarrow{\nabla}_n f(x) - f(x) \overrightarrow{\nabla}_n \right) \sigma_n - \hat{r}_{nm} \cdot \left(\overleftarrow{\nabla}_m f(x) - f(x) \overrightarrow{\nabla}_m \right) \sigma_m \right) \\
& - \frac{2\sqrt{2}}{\sqrt{3}} \left(\left[\hat{r}_{nm} \otimes \left(\overleftarrow{\nabla}_n f(x) - f(x) \overrightarrow{\nabla}_n \right) \right]^2 \otimes \sigma_n \right. \\
& \left. \left. - \left[\hat{r}_{nm} \otimes \left(\overleftarrow{\nabla}_m f(x) - f(x) \overrightarrow{\nabla}_m \right) \right]^2 \otimes \sigma_m \right) \right\} \quad (34)
\end{aligned}$$

In the case of the local scalar spatial contribution it is understood that:

$$\frac{dI}{dx} \rightarrow \frac{\delta(x)}{4\pi x^2} \quad (35)$$

The overall tensor rank is $J = 1$, the spin rank is one, but it contains both scalar and tensor components in the orbital sector. It can thus lead to $0^+ \rightarrow 1^+$ transitions. The decomposition of the other operators, if needed, can be done in a similar fashion. It is quite tedious but straightforward.

The scale of the nucleon recoil contribution is set by $(1/2m_N R_0) = 0.07A^{-1/3}$. In the case of neutrinoless double beta decay the square of this scale should be compared with the ratio of the phase space of this $j_L \otimes j_R$ going through the electron momenta to that of the mass term.

One of the electrons is in a p orbit.

If the resonance condition is met for such an electron, one may consider this possibility, even though the probability for finding a p -electron inside the nucleus is tiny. The factorization approximation should not be used in this case and the electron wave function must be part of the nuclear transition operator. The average over the nucleus of the square of the electron wave function is, however, still a good measure of the scale of the expected nuclear matrix elements. Thus combining the above operator for the decays to negative parity states we find that one can have transitions to 0^+ , 1^+ and 2^+ . The spin structure of the operator is still:

$$\omega_0 = i\sigma_n \otimes \sigma_m \text{ or } \omega_1 = \sigma_n - \sigma_m \quad (36)$$

i.e. antisymmetric of rank one. There appears to exist a wealth of orbital contributions.

Indeed let us assume that the p electron has total angular momentum j and the initial angular momentum of the atom is J . The structure of effective transition operator can be cast in the form:

$$\omega_0 \cdot \hat{r} \ell_0 \rightarrow U(1, 1/2, j, 0, 1/2, 1/2, 1, 0, J) \delta_{J,1} \left[\omega_0 \cdot \hat{r} \vec{V} \right] \ell_0 \quad (37)$$

for the time component and

$$\begin{aligned}
(\omega_1 \cdot \hat{r}) \cdot \vec{\ell} \rightarrow & U(1, 1/2, j, 0, 1/2, 1/2, 1, 0, J) U(1, 1, 0, 1, 1, J, J, 0, J) \\
& \left[(\omega_1 \times \hat{r}) \otimes \vec{V} \right]^J \vec{\ell} \cdot \vec{s} \quad (38)
\end{aligned}$$

for the space component ($\vec{\ell}$ is the leptonic current not to be confused with the orbital angular momentum operator). $U(\dots)$ are the usual unitary nine-j symbols needed for the re-coupling of the angular momenta involved and \vec{V} essentially is the two electron orbital wave function:

$$\vec{V} = \frac{1}{\sqrt{4\pi}} \frac{\sqrt{3}}{\sqrt{4\pi}} \frac{1}{\sqrt{2}} \left(A'' \mathbf{x}_2 + A' \mathbf{x}_1 \right) \quad (39)$$

with

$$A'' = \frac{R_0}{r_m} (\phi_{n_1,0}(r_n), \phi_{n_2,1}(r_m)), \quad A' = \frac{R_0}{r_n} (\phi_{n_1,1}(r_n), \phi_{n_2,0}(r_m)) \quad (40)$$

with ϕ_{n_i,ℓ_i} , $i = 1, 2$ the radial electron wave function. We have chosen to express the length in units of the nuclear radius R_0 , i.e. $\mathbf{x} = \mathbf{r}/R_0$, with compensating factors in the definition of A' and A'' , a procedure quite standard in the evaluation of the nuclear ME.

We found it convenient to go into the relative $\mathbf{x} = \mathbf{x}_n - \mathbf{x}_m$ and center of mass $2\mathbf{X} = \mathbf{x}_n + \mathbf{x}_m$ coordinates and thus we get:

$$\vec{V} = \frac{1}{4\pi} (A(n, m)\mathbf{x} + B(n, m)\mathbf{X}), \quad A(n, m) = \sqrt{3} \frac{1}{\sqrt{2}} \frac{(A' - A'')}{2}, \quad B(n, m) = \sqrt{3} \frac{1}{\sqrt{2}} (A' + A'') \quad (41)$$

The operator \vec{V} must be symmetric under the interchange of particles n and m to yield a total operator which is overall symmetric under the exchange of the particles n and m . So only the B-terms survive.

The next step is to perform an additional re-coupling bringing together the two spacial parts. One then finds:

1. For the time component:

$$\omega_0 \cdot \hat{r} \ell_0 \rightarrow \frac{B(n, m)}{4\pi} \sum_L C_0(j, L) [\omega_0 \otimes T^L(\hat{r}, \vec{R})]^J \delta_{J,1} \ell_0 \quad (42)$$

with

$$C_0(3/2, L) = -\frac{1}{\sqrt{2}} C_0(1/2, L) = \left(\frac{\sqrt{2}}{3\sqrt{3}}, -\frac{1}{\sqrt{6}}, -\frac{\sqrt{2}}{\sqrt{15}} \right) \text{ for } L = 0, 1, 2$$

and

$$T^0 = \hat{r} \cdot \mathbf{X}, \quad T^1 = \hat{r} \times \mathbf{X}, \quad T^2 = [\hat{r} \otimes \mathbf{X}]^2 \quad (43)$$

2. For the space component:

$$\omega_1 \times \hat{r} \ell_0 \rightarrow \frac{B(n, m)}{4\pi} \sum_L C_2(j, J, L) [\omega_1 \otimes T^L(\hat{r}, \vec{R})]^J \vec{\ell} \cdot \vec{s} \quad (44)$$

where the coefficients $C_2(j, J, L)$ are given in table I

By combining Eqs (11) and (42) as well as Eqs (13) and (44) we obtain the complete operators relevant for the decay of 0^+ to positive parity states with $J = 0, 1$ and 2.

To simplify matters one may seek an approximation familiar from the $0s$ case, i.e. to take the radial part of the electron wave function outside of the integrals and replace it with its average value over the nucleus. Thus:

$$\tilde{\Omega}_i = \Lambda_{\text{eff}} \sqrt{\langle a_B^6 B^2 \rangle} \succ \Omega_i, \quad \Lambda_{\text{eff}} = \frac{G_F^2}{2\sqrt{2}} \frac{1}{\pi R_0^2 a_B^3}, \quad B \leftrightarrow B(n, m) \quad (45)$$

where a_B is Bohr radius and, again, the averaging is over the radial part alone. Thus the effective transition operators now become:

$$\Omega_0 = 4g_A^2 \sum_{n,m} \tau_-(n) \tau_-(m) f(x_{n,m}) \sum_L C_0(j, L) [\omega_0 \otimes T^L(\hat{r}, \hat{R})]^J \delta_{J,1} \quad (46)$$

$$\Omega_1 = 2g_V g_A \sum_{n,m} \tau_-(n) \tau_-(m) f(x_{n,m}) \sum_L C_2(j, J, L) [\omega_1 \otimes T^L(\hat{r}, \hat{R})]^J \quad (47)$$

In fact it will be adequate to consider

$$\langle B^2 \rangle \longrightarrow \langle (R_{n,0})^2 \rangle \langle 3 \left(\frac{R_0}{r} R_{n,1} \right)^2 \rangle$$

where $R_{n,\ell}$ are the radial parts of the bound electron wave functions. Note that the p wave function involved in the averaging has been stripped of an extra power of r , since the \vec{r} has been incorporated into the nuclear matrix element.

The expression for the nuclear ME is the same with that of Eq. (17), where $k = J = J_f$ is the rank of the operator and the coupling of the two atoms is given by:

$$V^2 = \Lambda_{\text{eff}}^2 \langle a_B^6 B^2 \rangle |ME|^2 \quad (48)$$

We will check the factorization approximation using realistic atomic $3s$ and $4p_{3/2}$ wave functions provided to us by Shabaev [17], which were obtained by standard methods (see, e.g., Grant [18] and Bratsev *et al* [19]).

- Using the density profile shown in Fig. 1, corresponding to a phenomenological normalized to unity Woods-Saxon density, we find:

$$\langle a_B^3 (R_{3s})^2 \rangle = 2.4 \times 10^3, \quad \langle 3a_B^3 \left(\frac{R_0}{r} R_{4p_{3/2}} \right)^2 \rangle = 3.2 \times 10^{-2} \longrightarrow$$

$$\langle 3 \left(\frac{R_0}{r} R_{4p_{3/2}} \right)^2 \rangle / \langle (R_{3s})^2 \rangle = 1.3 \times 10^{-5}$$

- Using the density profile corresponding to a simplified S.M. wave function shown in Fig. 2, we find:

$$\langle a_B^3 (R_{3s})^2 \rangle = 1.4 \times 10^4, \quad \langle 3a_B^3 \left(\frac{R_0}{r} R_{4p_{3/2}} \right)^2 \rangle = 1.8 \times 10^{-1} \longrightarrow$$

$$\langle 3 \left(\frac{R_0}{r} R_{4p_{3/2}} \right)^2 \rangle / \langle (R_{3s})^2 \rangle = 1.4 \times 10^{-5},$$

Had we not used the factor $3(R_0/r)^2$ the p -integrals would have been quite a bit smaller. The agreement in the ratio of $4p$ to $3s$ probabilities between the two nuclear densities is quite good. Furthermore, using a simple $0s$ HO wave function, we find that the factorization approximation works to within 10%. More detailed computations using realistic and more accurate shell model wave functions are currently underway. The results, with and without employing the factorization approximation, will be published elsewhere [20]. The nuclear matrix elements leading to the $^{156}\text{Dy} \rightarrow ^{156}\text{Gd}$ leading to sharp resonances [15] $0^-, 1^-, 0^+, 1^+$ and 2^+ will also be calculated [20], using realistic wave functions both for the initial and the final nuclear states.

DISCUSSION

As we have mentioned the observation of neutrinoless double decay will be a great event in particle and nuclear physics. It will establish the existence of flavor violating interactions and it will demonstrate that, independently of the dominant mechanism causing it, the neutrinos are Majorana particles. More importantly, it is expected to complement the great successes on neutrino oscillations by determining the elusive up to now absolute scale of neutrino mass. For this last great contribution to become unambiguous, it will be necessary to determine what fraction of this decay is due to the light neutrino mass compared to all the other competing mechanisms [3]. That is a very difficult




Figure 1: The (normalized) Woods-Saxon density profile employed in obtaining the average electron wave functions.




Figure 2: The nuclear S.M. density profile employed in in obtaining the average electron wave functions and checking the factorization approximation.

task indeed. It may become feasible, if there exist data in as many targets as possible. In fact it has recently been shown that, within the mass terms alone, even if both left and right handed currents are involved, this is, in principle, possible [21, 22]. Provided, of course, that experimental data involving a sufficient number of targets become available. It will be much more difficult to achieve this, if the mass independent contribution involving the $j_L \times j_R$ contribution becomes important. In this direction the observation of double electron capture will be very useful. More specifically we have seen that:

Table I: The coefficients $C_2(j, J, L)$ discussed in the text

j	J	L	$C_2(j, J, L)$
1	0	1	$-\frac{1}{2}$
1	1	0	$-\frac{\sqrt{2}}{9}$
1	1	1	$-\frac{1}{9}$
1	1	1	$-\frac{1}{2\sqrt{6}}$
1	1	2	$\frac{\sqrt{5}}{3}$
1	1	0	$-\frac{1}{9}$
1	1	1	$-\frac{1}{9}$
1	1	1	$-\frac{1}{4\sqrt{3}}$
1	1	2	$\frac{\sqrt{5}}{6}$
2	1	1	$\frac{1}{4}$
2	2	2	$\frac{\sqrt{3}}{2}$

1. The neutrino mass term can proceed in neutrinoless double electron capture as in ordinary double beta decay. For the latter to be practical, of course, the resonance condition must be met.
2. Neutrinoless transitions to $0^+ \rightarrow J^\pm \neq 0^+$ can occur only if there exist right handed interactions through the right left interference in the leptonic sector ($j_L \otimes j_R$). Then:
 - The unusual transitions to the negative parity 0^- and 1^- states can naturally occur in double electron capture, since they are energetically allowed, if the resonance condition is met. So such an observation will unambiguously determine the neutrino mass independent lepton violating parameter, which is hard to get in neutrinoless double beta decay.
 - Transitions to positive parity states $\neq 0^+$ in ordinary double beta decay can occur either through the dipole component of the electron wave functions, the most favored scenario, or if one of the hadronic currents involved contains the parity changing nucleon recoil terms.
 - Transitions to positive parity states with angular momentum less than 3 in neutrinoless double electron capture can proceed if a) one of the hadronic currents involves the recoil term, while both electrons are in s-states or b) more naturally via the negative parity operator involving ordinary nucleon currents, if one s-electron and one p-electron are involved in the capture. The probability of finding a p-electron in the nucleus is, however, quite small, but this disadvantage maybe overcome by the collaborative effect of the expected large nuclear matrix elements and the resonance condition for capture to 2^+ .
 - The transition operators involved have a reasonable structure, but nuclear matrix elements of such operators have not yet been calculated for any system. We have no reason to expect them to be suppressed. The nuclear structure calculations involving such nuclear systems may be a true challenge to nuclear theorists.

It is clear from the above that the neutrinoless double electron capture can distinguish between the mass term contribution and the neutrino mass independent $j_L \times j_R$ interference term. It can do it more efficiently than ordinary beta decay due to the possibility of the negative parity nuclear transitions. It cannot, of course, distinguish between the various mass terms (light neutrino, heavy neutrino, SUSY contribution etc), but neither can ordinary double beta decay. This can only be done by exploiting the results of at least four different experiments (leading to 0^+), or a suitable combination of a sufficient number of both ground state and excited 0^+ state transitions. In this multiple experiment analysis, double electron capture can be come a very useful partner.

Acknowledgments

The author is indebted Professor Shabaev for providing the needed atomic wave functions and to Dr S. Eliseev and Professor Y. Novikov for discussions and useful comments. The hospitality of Professor K. Blaum in the Kernphysik, MPI-Hd and of the Theory Division at CERN are happily acknowledged.

Bibliography

-
- [1] Goepfert-Mayer, Phys. Rev. **48**, 512 (1935).
 - [2] W. Furry, Phys. Rev. **56**, 1184 (1939).
 - [3] J. D. Vergados, Phys. Rep. **361**, 1 (2002).
 - [4] F. Avignone, S. Elliott, and J. Engel, Rev. Mod. Phys. **80**, 481 (2008), arXiv:0708.1033 (nuc-th).

- [5] H. Ejiri, Phys. Rep. **338**, 265 (2000).
- [6] W. RODEJOHANN, Neutrinoless Double Beta Decay and Particle Physics, International Journal of Modern Physics E, arXiv:1106.1334 [hep-ph].
- [7] J. D. Vergados, Nuc. Phys. B. **218**, 109 (1983).
- [8] J. Bernabeu, A. de Rujula, and C. Jarlskog, Phys. Rev. C **223**, 15 (1983).
- [9] F. Simkovic and M. Krivoruchenko, Phys. Part. Nuc. **6**, 298 (2009).
- [10] S. Eliseev *et al*, Multiple-resonance phenomenon in neutrinoless double-electron capture, private communication.
- [11] S. Eliseev *et al.*, Phys. Rev. Lett. **106**, 052504 (2011).
- [12] F. Belli *et al*, arXiv:1104.5671 (nuc-ex); F. Belli *et al*, arXiv:1103.5359 (nuc-ex).
- [13] K. Blaum, Phys. Rep. **425**, 1 (2006).
- [14] M. I. Krivoruchenko, F. Simkovic, D. Frekers, and A. Faessler, Nuc. Phys. A **859**, 140 (2011), arXiv:1012.1204 [hep-ph].
- [15] S. Eliseev *et al.*, Phys. Rev. C **84**, 012501 (2011).
- [16] M. Doi, T. Kotani, H. Nishiura, K. Okuda, and E. Takasugi, Prog. Theor. Phys. **66**, 1739 (1981).
- [17] V. M. Shabaev and Ilya Tupitsin, Private communication.
- [18] I. Grant, Advances in Physics **19**, 747 (1970).
- [19] V. Bratsev *et al.*, Bull. Acad. Sci, USSR, Phys. Ser. **41**, 173 (1977).
- [20] O. Civitarese, J. Suhonen and J.D.Vergados, to be published.
- [21] F. Simkovic, J. Vergados, and A. Faessler, Phys. Rev. D **82**, 113015 (2010), arXiv:1006.0571 [hep-ph].
- [22] A. Faessler, A. Meroni, S. T. Petcov, F. Simkovic, and J. D. Vergados, Phys. Rev. D **83**, 113003 (2011), arXiv:1103.2434 [hep-ph].