

based on LHCb-CONF-2014-004
September 26, 2014

Improved constraints on γ : CKM2014 update

TILL MORITZ KARBACH¹

on behalf of the LHCb collaboration

I report on an updated combination of all currently available tree-level measurements of the CKM angle γ from LHCb. This combination currently gives LHCb's most precise value of $\gamma = (73_{-10}^{+9})^\circ$ obtained from $B \rightarrow DK$ -like decays. For the first time in a single experiment, the uncertainty has dropped below 10° . When using $B \rightarrow D\pi$ decays in addition, the situation becomes non-Gaussian which makes the confidence intervals more difficult to interpret.

PRESENTED AT

8th International Workshop on the CKM Unitarity Triangle
(CKM 2014), Vienna, Austria, September 8-12, 2014

¹CERN, Geneva, Switzerland

1 Introduction

The CKM angle $\gamma = \arg[-V_{ud}V_{ub}^*/(V_{cd}V_{cb}^*)]$ is a central parameter describing the CKM matrix. It can be measured using two different approaches. One is using loop-induced decays, which can give rise to effects of New Physics. The other approach is the topic of these proceedings and is purely based on tree level decays. This last property turns γ into one of the corner stones of CKM physics.

At LHCb, γ is measured using a large variety of decay channels. The best sensitivity is achieved through a combination of measurements that determine γ along with several other hadronic parameters. The input measurements provide sensitivity to γ through the interference of $b \rightarrow u$ and $b \rightarrow c$ amplitudes, as described in more detail in Ref. [1]. Determining the additional hadronic parameters from data results in a small systematic uncertainty on the measurement of γ . The presented update [2] improves over the previous combinations [1, 3] by adding more decay channels and updating selected channels to the full available dataset of 3 fb^{-1} . Tab. 1 gives the input measurements considered in this combination. Among the inputs is the first constraint of γ from the time-dependent analysis of $B_s^0 \rightarrow D_s^\mp K^\pm$ [4]. Being the only time-dependent method considered so far at LHCb, it is affected by very different experimental systematic uncertainties, which bases the γ average on a healthy range of measurements.

Table 1: LHCb Input measurements. The symbol h denotes either a kaon or a pion, and the acronyms denote the initials of authors first proposing the measurements in Refs. [5–14].

$B^+ \rightarrow Dh^+, D \rightarrow hh$	GLW/ADS	1 fb^{-1}	[15]
$B^+ \rightarrow Dh^+, D \rightarrow K\pi\pi\pi$	ADS	1 fb^{-1}	[16]
$B^+ \rightarrow DK^+, D \rightarrow K_s^0 hh$	model-independent GGSZ	3 fb^{-1}	[17]
$B^+ \rightarrow DK^+, D \rightarrow K_s^0 K\pi$	GLS	3 fb^{-1}	[18]
$B^0 \rightarrow DK^{*0}, D \rightarrow hh$	GLW/ADS	3 fb^{-1}	[19]
$B_s^0 \rightarrow D_s^\mp K^\pm$	time-dependent	1 fb^{-1}	[4]

2 Two combinations

Two combinations are prepared, referred to as the “robust” and “full” combinations. The robust combination only contains observables measured in $B \rightarrow DK$ -like systems, which are the traditional channels being used to measure γ . Compared to $B \rightarrow D\pi$ -like systems they offer larger interference, and therefore are very robust against systematic effects. The robust combination provides the main result for γ .

The full combination adds information from the $B \rightarrow D\pi$ system. This is motivated by the fact that LHCb has a large set of γ -sensitive observables in $B \rightarrow D\pi$ decays,

although their sensitivity to γ is suppressed compared to the $B \rightarrow DK$ -like decays. The reason for this is that the amplitude ratio, which governs the interference effects and therefore the sensitivity, is expected to be a factor ≈ 15 smaller than for the $B \rightarrow DK$ -like systems. At the same time, the available data samples are usually a factor 10 larger in $B \rightarrow D\pi$ decays. The full combination is more sensitive to several effects. The most prominent of these is $D^0-\bar{D}^0$ mixing [20–22], which we fully correct for in both combinations, taking into account the D^0 decay time acceptances of the individual measurements. We also correct at first order for possible CP violation in the D^0 system using information from the Heavy Flavor Averaging Group (HFAG) [23]. Furthermore, in order to ensure our combination is not sensitive to mixing and CP violation in K^0 decays [24], we exclude observables where such effects may be non-negligible. Due to the larger data samples, the relative impact of the systematic uncertainties is larger than in the robust combination, although both of the current combinations are still statistically limited. Finally, a small value of the amplitude ratio $r_B^{D\pi}$ that is not significantly different from zero, is known to affect the coverage of the frequentist methods used here.

3 Statistical Procedure

The combination follows a frequentist treatment described in detail in Ref. [1], in which the nuisance parameters are kept at their best fit values (known as the “plug-in” method). The results obtained using $B \rightarrow DK$ -like decays alone are also cross-checked using a Bayesian approach assuming a flat prior.

4 Results

The result of the robust combination is given in Tab. 2, and illustrated in Figs 1 and Fig 2. Both frequentist and Bayesian intervals agree very well. The coverage was tested for the frequentist interval and found to be accurate. The fit probability is 89.4%. The robust frequentist values are regarded as the nominal result: $\gamma = (73_{-10}^{+9})^\circ$.

Table 2: Confidence and credibility regions and central values for γ extracted from the robust combination.

	Observable	Central value	Intervals	
			68%	95%
Frequentist	$\gamma[^\circ]$	72.9	[63.0, 82.1]	[52.0, 90.5]
Bayesian	$\gamma[^\circ]$	71.9	[61.9, 81.8]	[50.9, 91]




Figure 1: 1 – CL curves for the robust combination (left), and for both robust and full combinations (right).




Figure 2: The 1D posterior PDF in the robust combination. Dark and light regions show the 68% and 95% probability intervals, respectively.

The full combination is more difficult to interpret. It exhibits a sharp maximum at $\gamma = 78.9^\circ$, and a secondary maximum at a value similar to the maximum of the robust combination (Tab. 3). This sharp maximum corresponds to an unexpectedly large value of the amplitude ratio $r_B^{D\pi}$, $r_B^{D\pi} = 0.027$. This is regarded as a fluctuation, but nevertheless enhances the impact of the $B^+ \rightarrow D\pi^+$ system. It makes the situation visibly non-Gaussian. As a consequence, the 68% CL interval is misleadingly small, while at 95% CL the intervals agree well between both combinations. This is illustrated in Fig. 1 (right), which shows both robust and full combinations in the same plot. The coverage of the intervals of the full combination was also tested, and as expected from the fact that $r_B^{D\pi}$ is still consistent with zero, the intervals were found to undercover a

bit, i.e. the reported intervals are too small. It is expected that the ambiguity in $r_B^{D\pi}$, which causes the non-Gaussian behavior, could be resolved by a GGSZ-type analysis of $B^+ \rightarrow D\pi^+$ decays.

Table 3: Confidence intervals and central values for the full combination. The two columns correspond to the two minima found by the fit. The most probable value is given in the left column, corresponding to a large value of $r_B^{D\pi}$.

quantity	full	
γ ($^\circ$)	78.9	72.8
68% CL ($^\circ$)	[71.5, 84.7]	
95% CL ($^\circ$)	[54.6, 91.4]	
$r_B^{D\pi}$	0.027	0.006
68% CL	[0.016, 0.034]	[0.005, 0.007]
95% CL	[0.001, 0.040]	

5 Discussion

Following the presentation there was a discussion about the following points:

- Why is the $B \rightarrow DK$ -like “robust” combination regarded as the nominal result, rather than the more complete “full” combination? — At this point, LHCb doesn’t have enough data to measure $r_B^{D\pi}$ significantly different from zero, which affects the statistical coverage of the “full” combination. The result is still correct, only much more difficult to interpret. But the $B^+ \rightarrow D\pi^+$ system is a perfect test bed to show up the obstacles on the way to a high precision measurement of γ in $B^+ \rightarrow DK^+$.
- What drives the large $r_B^{D\pi}$ minimum? — It is driven by two analyses taken together: $B^+ \rightarrow Dh^+$, $D \rightarrow hh$ [15] and $B^+ \rightarrow Dh^+$, $D \rightarrow K\pi\pi\pi$ [16]. If either one is dropped, the fit converges to a lower value of $r_B^{D\pi}$.
- Does LHCb plan to make a GGSZ type measurement in $B^+ \rightarrow D\pi^+$? — Yes.
- Why did the precision on γ decrease by so much when the auxiliary inputs were updated? — This was tracked down to a sizable shift in the parameter $R_D(K\pi)$ in the latest HFAG average compared to previous ones.

6 Conclusion

Observables measured by LHCb that have sensitivity at tree-level to the CKM angle γ , along with supplementary information from other experiments, are combined to

determine an improved constraint on γ . The effect of $D^0-\bar{D}^0$ mixing on the decay rate is taken into account, with consideration of the experimental decay time acceptances of the individual measurements. When all observables are combined, we find

$$\gamma = (73_{-10}^{+9})^\circ, \quad (1)$$

using $B \rightarrow DK$ -like decays only. This result is more precise than the combination of the results of the B factories Babar and Belle [25].

References

- [1] LHCb collaboration, R. Aaij *et al.*, *A measurement of the CKM angle γ from a combination of $B^\pm \rightarrow Dh^\pm$ analyses*, Phys. Lett. **B726** (2013) 151, [arXiv:1305.2050](#).
- [2] LHCb collaboration, *Improved constraints on γ : CKM2014 update*, Sep, 2014. LHCb-CONF-2014-004.
- [3] LHCb collaboration, R. Aaij *et al.*, *Improved constraints on γ from $B^\pm \rightarrow DK^\pm$ decays including first results on 2012 data*, LHCb-CONF-2013-006 (2013), presented at Beauty 2013, Bologna, Italy, 8–12 April 2013.
- [4] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP asymmetry in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays*, [arXiv:1407.6127](#), submitted to JHEP.
- [5] M. Gronau and D. Wyler, *On determining a weak phase from CP asymmetries in charged B decays*, Phys. Lett. **B265** (1991) 172.
- [6] M. Gronau and D. London, *How to determine all the angles of the unitarity triangle from $B^0 \rightarrow DK_S^0$ and $B_s^0 \rightarrow D\phi$* , Phys. Lett. **B253** (1991) 483.
- [7] D. Atwood, I. Dunietz, and A. Soni, *Enhanced CP violation with $B \rightarrow KD^0$ (\bar{D}^0) modes and extraction of the CKM angle γ* , Phys. Rev. Lett. **78** (1997) 3257, [arXiv:hep-ph/9612433](#).
- [8] D. Atwood, I. Dunietz, and A. Soni, *Improved methods for observing CP violation in $B^\pm \rightarrow KD$ and measuring the CKM phase γ* , Phys. Rev. **D63** (2001) 036005, [arXiv:hep-ph/0008090](#).
- [9] A. Bondar, *Proceedings of BINP special analysis meeting on Dalitz analysis*, 24.–26. Sep, 2002. Unpublished.
- [10] A. Giri, Y. Grossman, A. Soffer, and J. Zupan, *Determining γ using $B^\pm \rightarrow DK^\pm$ with multibody D decays*, Phys. Rev. **D68** (2003) 054018, [arXiv:hep-ph/0303187](#).

- [11] Y. Grossman, Z. Ligeti, and A. Soffer, *Measuring gamma in $B^\pm \rightarrow K^\pm(KK^*)(D)$ decays*, Phys. Rev. **D67** (2003) 071301, [arXiv:hep-ph/0210433](#).
- [12] R. Fleischer, *New strategies to obtain insights into CP violation through $B_s^0 \rightarrow D_s^\pm K^\mp, D_s^{*\pm} K^\mp, \dots$ and $B^0 \rightarrow D^\pm \pi^\mp, D_s^\pm \pi^\mp, \dots$ decays*, Nucl. Phys. **B671** (2003) 459, [arXiv:hep-ph/0304027](#).
- [13] R. Aleksan, I. Dunietz, and B. Kayser, *Determining the CP violating phase γ* , Z. Phys. **C54** (1992) 653.
- [14] I. Dunietz and R. G. Sachs, *Asymmetry between inclusive charmed and anticharmed modes in B^0, \bar{B}^0 decay as a measure of CP violation*, Phys. Rev. **D37** (1988) 3186.
- [15] LHCb collaboration, R. Aaij *et al.*, *Observation of CP violation in $B^\pm \rightarrow DK^\pm$ decays*, Phys. Lett. **B712** (2012) 203, Erratum *ibid.* **B713** (2012) 351, [arXiv:1203.3662](#).
- [16] LHCb collaboration, R. Aaij *et al.*, *Observation of the suppressed ADS modes $B^\pm \rightarrow [\pi^\pm K^\mp \pi^+ \pi^-]_D K^\pm$ and $B^\pm \rightarrow [\pi^\pm K^\mp \pi^+ \pi^-]_D \pi^\pm$* , Phys. Lett. **B723** (2013) 44, [arXiv:1303.4646](#).
- [17] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP violation and constraints on the CKM angle γ in $B^\pm \rightarrow DK^\pm$ with $D \rightarrow K_S^0 \pi^+ \pi^-$ decays*, [arXiv:1407.6211](#), submitted to Nucl. Phys. B.
- [18] LHCb collaboration, R. Aaij *et al.*, *A study of CP violation in $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D\pi^\pm$ decays with $D \rightarrow K_S^0 K^\pm \pi^\mp$ final states*, Phys. Lett. **B733** (2014) 36, [arXiv:1402.2982](#).
- [19] LHCb collaboration, R. Aaij *et al.*, *Measurements of CP violation parameters in $B^0 \rightarrow DK^{*0}$ decays*, [arXiv:1407.8136](#), submitted to Phys. Rev. D.
- [20] C. C. Meca and J. P. Silva, *Detecting new physics contributions to the D^0 - anti- D^0 mixing through their effects on B decays*, Phys. Rev. Lett. **81** (1998) 1377, [arXiv:hep-ph/9807320](#).
- [21] J. P. Silva and A. Soffer, *Impact of $D^0 - \bar{D}^0$ mixing on the experimental determination of γ* , Phys. Rev. **D61** (2000) 112001, [arXiv:hep-ph/9912242](#).
- [22] M. Rama, *Effect of $D - \bar{D}$ mixing in the extraction of γ with $B^- \rightarrow D^0 K^-$ and $B^- \rightarrow D^0 \pi^-$ decays*, Phys. Rev. **D89** (2014) 014021, [arXiv:1307.4384](#).
- [23] Heavy Flavor Averaging Group, Y. Amhis *et al.*, *Averages of b-hadron, c-hadron, and τ -lepton properties as of early 2012*, [arXiv:1207.1158](#), updated results and plots available at: <http://www.slac.stanford.edu/xorg/hfag/>.

- [24] Y. Grossman and M. Savastio, *Effects of $K^0 - \bar{K}^0$ mixing on determining γ from $B^\pm \rightarrow DK^\pm$* , JHEP **1403** (2014) 008, arXiv:1311.3575.
- [25] A. J. Bevan *et al.*, *The Physics of the B Factories*, arXiv:1406.6311.