

## Dark Matter searches with the ATLAS Detector

---

**Valerio Ippolito\***, on behalf of the ATLAS Collaboration

*Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA, United States of America*

*E-mail:* [valerio.ippolito@cern.ch](mailto:valerio.ippolito@cern.ch)

The presence of a non-baryonic Dark Matter component in the Universe is inferred from the observation of its gravitational interaction. If Dark Matter interacts weakly with the Standard Model particles it may be produced at the Large Hadron Collider (LHC), escaping detection and leaving large missing transverse momentum as its signature. New results from the Dark Matter search programme of the ATLAS experiment are presented, based on LHC proton-proton collision data collected at a center-of-mass energy of 13 TeV.

*The European Physical Society Conference on High Energy Physics  
5-12 July  
Venice, Italy*

---

\*Now at INFN Sezione di Roma.



## 1. Introduction


Investigating the particle nature of Dark Matter (DM), whose existence is supported by strong observational evidence[1], is one of the main goals of the ATLAS experiment at the Large Hadron Collider (LHC). ATLAS searches are based on various theoretical frameworks beyond the Standard Model (SM), such as simplified models[2, 3, 4, 5] which provide rather generic descriptions of the DM-SM interaction, or supersymmetric models. In this document, the most recent search results based on proton-proton collisions at 13 TeV are presented. The ATLAS detector is described elsewhere[6].

## 2. $E_T^{\text{miss}} + \text{jets}$

The highest sensitivity to scenarios where pairs of weakly-interacting DM particles are produced through the  $s$ -channel exchange of a mediator is provided by the final state with large missing transverse momentum,  $E_T^{\text{miss}}$ , recoiling against a high- $p_T$  jet originating from the hadronisation of a gluon from initial-state radiation (“mono-jet”). The ATLAS search[7] is based on  $36.1 \text{ fb}^{-1}$  of collision data collected in 2015 and 2016. Events passing a  $E_T^{\text{miss}}$  trigger are selected if they have at least one jet reconstructed with the anti- $k_T$  algorithm[8] with radius parameter  $R = 0.4$ , with  $p_T > 250 \text{ GeV}$  and pseudorapidity  $|\eta| < 2.4$ ,  $E_T^{\text{miss}} > 250 \text{ GeV}$  and no electron or muon. Up to three additional jets with  $p_T > 30 \text{ GeV}$  are allowed; the difference in azimuthal angle between  $E_T^{\text{miss}}$  and any jets must be  $\Delta\phi > 0.4$ .

The dominant backgrounds to this search come from the  $Z(\nu\nu) + \text{jets}$  process and from the  $W(\mu\nu) + \text{jets}$  and  $W(\tau\nu) + \text{jets}$  processes, where the muon is not reconstructed or the  $\tau$  lepton decays hadronically or into a neutrino and a charged lepton which escapes detection. Next-to-leading-order (NLO) QCD corrections and NLO electroweak corrections, supplemented by Sudakov logarithms at two loops, are applied to the simulated samples for the  $W/Z + \text{jets}$  processes as a function of the boson  $p_T$ , following the prescription of Ref. [9]. A simultaneous binned likelihood fit to the  $W/Z p_T$  distribution in four control regions enriched in  $W(\mu\nu) + \text{jets}$ ,  $W(e\nu) + \text{jets}$ ,  $t\bar{t}$  and  $Z(\mu\mu) + \text{jets}$  events is used to derive a common normalisation factor for all the  $W/Z + \text{jets}$  backgrounds, and a separate normalisation factor for top-quark processes. Uncertainties due to the different effect of higher-order corrections to  $Z(\nu\nu) + \text{jets}$ ,  $W(\ell\nu) + \text{jets}$  and  $Z(\ell\ell) + \text{jets}$  processes are taken into account across the full  $E_T^{\text{miss}}$  spectrum, and have an impact of about 1% on the total background uncertainty in the signal region. Additional contributions come from the uncertainty on the muon and electron selection efficiencies and from the uncertainty on the jet energy scale and resolution. The total background uncertainty varies between 2% and 7%; no significant excess with respect to the SM prediction is observed.

Results are interpreted in terms of simplified models of DM-SM interaction, with the  $s$ -channel exchange of an axial-vector or vector mediator, with a coupling  $g_q = 1/4$  to SM quarks and a coupling  $g_\chi = 1$  to DM. Exclusion contours at 95% CL are presented in Fig. 1 for the two cases: mediator masses up to 1.55 TeV are excluded for low values of the DM mass. The case of pseudo-scalar interactions, with a mediator exchanged in the  $s$ -channel through a top-quark loop, was also considered, but the analysis has not yet enough sensitivity to a mediator coupling  $g = 1$ .



**Figure 1:** Observed and expected exclusion contours as a function of the DM mass  $m_\chi$  and the mediator mass  $m_{Z_{A(V)}}$ , for the simplified model with an axial-vector (left) or vector (right) mediator. From Ref. [7].


### 3. $Z(\nu\nu) + \text{jets}/Z(\ell\ell) + \text{jets}$ cross-section ratio

Similar results can be obtained by comparing the measured differential  $Z(\nu\nu) + \text{jets}/Z(\ell\ell) + \text{jets}$  cross-section ratios, corrected for detector effects, to predictions from new physics models. The ATLAS measurement[10], based on  $3.2 \text{ fb}^{-1}$  of collision data collected in 2015, studies both a mono-jet and a vector-boson fusion (VBF) topology. The measured ratio is expressed as a function of the boson  $p_T$  and, in the case of the VBF topology, the mass of the di-jet system and the difference in azimuthal angle between the two jets. Figure 2(a) shows such ratio as a function of the boson  $p_T$  in the mono-jet topology. A re-interpretation of these results in the context of the axial-vector model of Sec. 2, under the assumption that the detector effects are similar for both the SM backgrounds and this simplified model, is shown in Fig. 2(b).

### 4. $E_T^{\text{miss}} + Z(ee/\mu\mu)$


Final states with a  $Z$  boson decaying into electrons or muons and missing transverse momentum constitute another clear signature for the production of DM, where the  $Z$  boson may come for example from initial state radiation. The ATLAS search[11] is based on  $36.1 \text{ fb}^{-1}$  of data collected in 2015 and 2016. Events are selected with a pair of electrons or muons with invariant mass compatible with the decay from a  $Z$  boson, recoiling against  $E_T^{\text{miss}} > 90 \text{ GeV}$ .

The dominant background, coming from  $ZZ$  production, is estimated from simulation; data-driven techniques are used to determine the  $WZ$  background, as well as the contribution from non-resonant backgrounds. The main uncertainties come from the modelling of the  $ZZ$  process, lepton momentum scale and resolution and uncertainties on the lepton reconstruction and identification efficiencies and jet energy scales.



**Figure 2:** Left: measured ratio between the fiducial cross-sections for the  $Z(vv) + \text{jets}$  and  $Z(\ell\ell) + \text{jets}$  processes,  $R_{\text{miss}}$ , as a function of the boson  $p_T$ , for the mono-jet topology. Right: observed and expected 95% CL exclusion contours as a function of the DM mass  $m_\chi$  and the mediator mass  $m_A$ , for the simplified model with an axial-vector mediator; expected results are compared to a mono-jet search performed with the same data. From Ref. [10].

No significant excess is observed with respect to the SM prediction. Fig. 3 shows the 95% CL exclusion contours for the simplified model with an axial-vector or vector mediator: mediator masses up to 560 GeV are excluded for low values of the DM mass.




**Figure 3:** Observed and expected exclusion contours as a function of the DM mass  $m_\chi$  and the mediator mass  $m_{\text{med}}$ , for the simplified model with an axial-vector (left) or vector (right) mediator. From Ref. [11].


## 5. Summary

Fig. 4 compares all ATLAS search results sensitive to the considered simplified model, including both searches for DM production in final states with  $E_T^{\text{miss}}$  and direct searches for mediator decays into two jets. Results for axial-vector (vector) interactions with this specific choice of the couplings  $g_q$  and  $g_\chi$  are expressed in terms of 95% CL limits on the spin-dependent (spin-independent) DM-proton (DM-nucleon) interaction cross-section, as a function of the assumed DM

mass. Complementary results are obtained with respect to the ones from direct detection experiments for light DM.



(a)



(b)

**Figure 4:** Observed limit on the DM-proton (top) or DM-nucleon (bottom) interaction cross-section, as a function of the DM mass, for the simplified model with an axial-vector (top) or vector (bottom) mediator. From Ref. [12].

## References

- [1] G. Bertone et al., *Particle Dark Matter: Evidence, Candidates and Constraints*, *Phys. Rept.* **405** (2005) 279-390.
- [2] J. Abdallah J. et al., *Simplified Models for Dark Matter Searches at the LHC*, *Phys. Dark Univ.* **9-10** (2015) 8-23.
- [3] D. Abercrombie et al., *Dark Matter Benchmark Models for Early LHC Run-2 Searches: Report of the ATLAS/CMS Dark Matter Forum*, arXiv:1507.00966 [hep-ex].
- [4] A. Boveia et al., *Recommendations on presenting LHC searches for missing transverse energy signals using simplified s-channel models of dark matter*, arXiv:1603.04156 [hep-ex].
- [5] A. Albert et al., *Recommendations of the LHC Dark Matter Working Group: Comparing LHC searches for heavy mediators of dark matter production in visible and invisible decay channels*, arXiv:1703.05703 [hep-ex].
- [6] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, *JINST* **3** (2008) S08003.
- [7] ATLAS Collaboration, *Search for dark matter and other new phenomena in events with an energetic jet and large missing transverse momentum using the ATLAS detector*, ATLAS-CONF-2017-060 (<http://cdsweb.cern.ch/record/2273876>).
- [8] M. Cacciari et al., *The anti- $k_r$  jet clustering algorithm*, *JHEP* **04** (2008) 063.
- [9] J. M. Lindert et al., *Precise predictions for  $V + jets$  dark matter backgrounds*, arXiv:1705.04664 [hep-ph].
- [10] ATLAS Collaboration, *Measurement of detector-corrected observables sensitive to the anomalous production of events with jets and large missing transverse momentum in  $pp$  collisions at  $\sqrt{s} = 13$  TeV using the ATLAS detector*, arXiv:1707.03263 [hep-ex] (submitted to Eur. Phys. J. C).
- [11] ATLAS Collaboration, *Search for an invisibly decaying Higgs boson or dark matter candidates produced in association with a Z boson in  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector*, arXiv:1708.09624 [hep-ex] (submitted to Phys. Lett. B).
- [12] ATLAS Collaboration,  
<https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/CombinedSummaryPlots/EXOTICS/>