

CROSS SECTION OF HADRON PRODUCTION IN $\gamma\gamma$ COLLISIONS AT LEP

Maria Novella Kienzle-Focacci
Université de Genève
24,quai Ernest Ansermet CH-1211 Genève 4
Switzerland



Abstract

The reaction $e^+e^- \rightarrow e^+e^-\gamma^*\gamma^* \rightarrow e^+e^- \text{ hadrons}$ is analysed using data collected by the L3 detector during the LEP runs at $\sqrt{s} = 130\text{-}140$ GeV and $\sqrt{s} = 161$ GeV. The cross sections $\sigma(e^+e^- \rightarrow e^+e^- \text{ hadrons})$ and $\sigma(\gamma\gamma \rightarrow \text{hadrons})$ are measured in the interval $10 \leq W_{\gamma\gamma} \leq 75$ GeV for the first time. The energy dependence of the real photons total cross section, $\sigma(\gamma\gamma \rightarrow \text{hadrons})$, is consistent with the universal Regge behaviour of total hadronic cross sections.




Figure 1: a) The cross section $d\sigma(e^+e^- \rightarrow e^+e^- \text{hadrons})/dW_{\gamma\gamma}$ as measured at $\sqrt{s}=133$ GeV and at $\sqrt{s}=161$ GeV. b) Total cross section $\gamma\gamma \rightarrow \text{hadrons}$. The continuous line is the Regge fit described in the text. The two components: the rapidly decreasing Reggeon part and the slow rising component due to Pomeron exchange are indicated with a dashed line.

Two-photon interactions are the dominant source of hadron production at LEP2. In the two high energy runs at $\sqrt{s}=133$ ($L=4.98 \text{ pb}^{-1}$) and $\sqrt{s}=161$ GeV ($L=10.37 \text{ pb}^{-1}$), a total of 32000 events of anti-tagged two-photon interaction $e^+e^- \rightarrow e^+e^- \text{hadrons}$ were observed in the L3 detector, with $W_{\text{vis}} \geq 5$ GeV. The effective mass of the hadron system (W_{vis}) is calculated by using well reconstructed tracks (≥ 3) and clusters measured in all the calorimeters.

The detailed features of the events: angular and momentum distributions, energy deposited in the calorimeters and visible mass are rather well reproduced by the recent generators PYTHIA [1] and PHOJET [2]. From the visible mass the two-photon center of mass energy $W_{\gamma\gamma}$ is derived by using an efficiency and acceptance matrix calculated with the two generators.

The cross section $\sigma(e^+e^- \rightarrow e^+e^- \text{hadrons})$ for $\langle Q^2 \rangle \geq 0.025 \text{ GeV}^2$ is measured in the interval $5 \leq W_{\gamma\gamma} \leq 75$ GeV (Fig.1a). The real photon total cross section $\sigma(\gamma\gamma \rightarrow \text{hadrons})$ is also derived from the data (Fig.1b). The $\sigma(\gamma\gamma \rightarrow \text{hadrons})$ cross section is dominated by soft $\gamma\gamma$ interactions, where the photon behaves like a hadron. The increase with energy of this cross section is characteristic of Pomeron exchange. The universal Regge parametrisation of A. Donnachie and P.V. Landshoff [3] $\sigma_{\text{tot}} = A s^\epsilon + B s^{-\eta}$ and the energy dependence fixed by the world average hadronic total cross sections [4] $\epsilon = 0.0790 \pm 0.0011$, $\eta = 0.4678 \pm 0.0059$ reproduce well the data ($\chi^2/\text{d.o.f.} = 3/8$) with parameters $A = 173 \pm 7$, $B = 519 \pm 125$, obtained with a fit which takes into account the bin-to-bin systematic errors.

References

- [1] G.A. Schuler and T. Sjöstrand, Nucl.Phys.B **407** (1993) 539 ; Z. Phys. C **73** (1997) 677.
- [2] R. Engel and J. Ranft, Z. Phys. C **66** (1995) 203 ; Phys.Rev.D **54** (1996) 4246.
- [3] A.Donnachie and P.V. Landshoff, Phys.Lett.B **296** (1992) 227.
- [4] Review of Particle Physics, Phys.Rev.D **54** (1996) 192.