

BR

IFUP-TH-46/94

High energy single particle states in the continuum

See 9440

Angela Bonaccorso\*

*Istituto Nazionale di Fisica Nucleare, Sezione di Pisa, 56100 Pisa, Italy*  
*E-mail: Decnet 39186::BONAC ; Internet BONAC@THPISA.DIFI.UNIPI.IT*

Abstract

This paper discusses the origin of the high energy bumps in the inclusive spectra measured in one neutron transfer to the continuum reactions between heavy ions. Our calculations support the interpretation of such structures as due to the population of single particle states of high spin.

PACS number(s):25.70.Hi,24.30.Gd

\*Address for September 1994-June 1995: Institut de Physique Nucléaire,  
IN2P3-CNRS, 91406 Orsay, Cedex, France.



UNIVERSITÀ DEGLI STUDI DI PISA  
DIPARTIMENTO DI FISICA



Istituto Nazionale di Fisica Nucleare  
Sezione di Pisa



CERN LIBRARIES, GENEVA



## 1 Introduction

In this paper we discuss single particle states in the continuum by analyzing a series of experimental results on one-neutron transfer in heavy-ion reactions. The Bonaccorso-Brink model is used for the calculation of the inclusive spectra and for the extraction of the background due to elastic and inelastic breakup. Also it allows the decomposition of the cross section into the contributions of the various final angular momenta. Such a decomposition is used to study the single particle properties of the continuum states populated by transfer. We calculate the spectral distribution, escape and spreading widths of the states in two different ways. One method uses an optical model description of the interaction of the transferred neutron with the target. The other method is based on the semiclassical calculation of the neutron-target S-matrix and of the widths of the single particle target states.

The experimental inclusive spectra of all targets show some structures up to about  $E_x \sim 10\text{MeV}$  and the comparison between the calculated and the experimental cross sections suggests that they are due to the population of continuum single particle states. The higher energy part of the spectra varies according to the incident energy and to the projectile-target combination. If the incident energies are in the range  $E_{inc} \approx 40 - 50\text{MeV/nucleon}$  and provided the initial bound state energy of the neutron in the projectile is of the order of several MeV, then the matching conditions allow the population by transfer of high lying single particle states like the  $1m_{21/2}$  in  $^{208}\text{Pb}$  and the  $1j_{15/2}$  in  $^{91}\text{Zr}$  and the experimental spectra show indeed bumps at the corresponding energies predicted by the shell model. This rather unexpected feature is consistent with the results of the calculations which give values for the widths of the single particle states smaller than their spacing. Such a situation occurs when the resonance energy is below the top of the barrier formed by the real plus centrifugal potentials and at the same time there is a small surface imaginary potential describing the absorption of the neutron on the target. When these conditions are not satisfied, as in the  $^{48}\text{Ca}$  case, the single particle states in the target are quite damped and the transfer spectra are dominated by the breakup.

This paper is organized in the following way. In Sec.2 we give a brief review of the status of the experimental results, then in Sec.3 we resume the transfer to the continuum model and we introduce the semiclassical methods used to study the

microscopic structure of the bumps. The S-matrix calculations are discussed in Sec.4 and the widths in Section 5. Sec.6 contains the discussion of both the experimental and theoretical results and finally in Sec.7 we give our conclusions.

## 2 Experimental review

In this section we review some of the experimental data on neutron transfer to the continuum in heavy-ion reactions for which we have made also a theoretical analysis. Data taken up to 1988 were discussed in [1]. More recent data are published in [2, 3, 4, 5, 6, 7, 8, 9] and in Sec.6 we will show some of them together with our calculations. Many experiments have targets in the region of lead but also  $^{48}\text{Ca}$ ,  $^{90}\text{Zr}$ ,  $^{120}\text{Sn}$  and  $^{64}\text{Ni}$  have been studied.

One characteristic common to all inclusive spectra is that there are structures up to an excitation energy of about  $E_x \sim 10\text{MeV}$ . At higher excitation energies some spectra show a structureless continuum. This happens for light projectiles like  $\alpha$ -particles and/or when the incident energy is less than about  $E_{inc} = 30\text{MeV/nucleon}$ . However there are other spectra in which the situation is rather different because they show also structures at  $E_x \sim 20 - 40\text{MeV}$  superimposed to a large continuum. This happens in the reactions with heavy projectiles like  $^{20}\text{Ne}$  and  $^{40}\text{Ar}$  and when the incident energy is of the order of  $E_{inc} \sim 40\text{MeV/nucleon}$ .

From the experimental point of view the nature of the low energy structures has been studied recently by detecting their decay via neutron emission. This has been possible in some experiments in which not only the inclusive spectra were measured but also the coincidence between the neutrons emitted by the excited residual nucleus and the ejectile. The reaction studied were  $^{208}\text{Pb}(\alpha, ^3\text{He})^{209}\text{Pb}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7] and  $^{48}\text{Ca}(^{20}\text{Ne}, ^{19}\text{Ne})^{48}\text{Ca}$  at  $E_{inc} = 48\text{MeV/nucleon}$ [8]. The analysis of the angular correlation in those data has shown that while the reaction on  $^{48}\text{Ca}$  is dominated by the breakup the reaction on  $^{208}\text{Pb}$  is dominated by the absorption of the transferred neutron on the high spin continuum states of the target.

In this paper we will discuss four reactions whose inclusive cross sections show just one dominant bump centered around  $E_x \sim 10\text{MeV}$   
 $^{208}\text{Pb}(\alpha, ^3\text{He})^{209}\text{Pb}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7];

$^{90}\text{Zr}(\alpha, ^3\text{He})^{91}\text{Zr}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7];

$^{120}\text{Sn}(\alpha, ^3\text{He})^{121}\text{Sn}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7];

$^{64}\text{Ni}(\alpha, ^3\text{He})^{65}\text{Ni}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7].

Then the following reactions which have inclusive cross sections showing high energy structures superimposed to a large continuum

$^{207}\text{Pb}(^{20}\text{Ne}, ^{19}\text{Ne})^{208}\text{Pb}$  at  $E_{inc} = 48\text{MeV/nucleon}$ [6, 9];

$^{208}\text{Pb}(^{40}\text{Ar}, ^{39}\text{Ar})^{209}\text{Pb}$  at  $E_{inc} = 41\text{MeV/nucleon}$ [6, 9];

$^{90}\text{Zr}(^{20}\text{Ne}, ^{19}\text{Ne})^{91}\text{Zr}$  at  $E_{inc} = 40\text{MeV/nucleon}$ [3].

Finally a reaction on  $^{48}\text{Ca}$  that shows some low energy structures and then a large structureless continuum

$^{48}\text{Ca}(^{20}\text{Ne}, ^{19}\text{Ne})^{48}\text{Ca}$  at  $E_{inc} = 48\text{MeV/nucleon}$ [8].

### 3 Theory

We give now a brief account of the theory behind the transfer calculations.

In the Bonaccorso-Brink model[10, 11, 12, 13] the neutron transfer probability from a definite single particle state of energy  $\varepsilon_i$ , angular momentum  $l_i$  and spin  $j_i$  in the projectile to a final continuum state of energy  $\varepsilon_f$  within an interval  $d\varepsilon_f$  in the target is given by the sum of the transfer probabilities to each possible final  $j_f$ -state in the energy bin  $d\varepsilon_f$ .

$$\begin{aligned} \frac{dP}{d\varepsilon_f} &= \sum_{j_f} \langle |1 - S_{j_f}|^2 \rangle B(j_f, j_i) \\ &= \sum_{j_f} (|1 - \langle S_{j_f} \rangle|^2 + T_{j_f}) B(j_f, j_i) \end{aligned} \quad (3.1)$$

where

$$T_{j_f} = 1 - |\langle S_{j_f} \rangle|^2. \quad (3.2)$$

$\langle S_{j_f} \rangle$  is the optical model S-matrix which describes the re-scattering of the neutron on the target. The sum in Eq.(3.1) is over all possible final angular momenta corresponding to the given final energy.  $B(j_f, j_i)$  is an elementary transfer probability

which depends on the details of the initial and final states, on the energy of relative motion and on the distance of closest approach between the two nuclei.

In Ref.[11] it was shown that the first term in Eq.(3.1), proportional to  $|1 - \langle S_{j_f} \rangle|^2$ , gives the elastic breakup spectrum while the second term proportional to the transmission coefficient  $T$  gives the absorption spectrum. The elastic breakup gives some part of the physical background of the spectra. The way of calculating it has been described in previous publications [12, 9]. The inelastic breakup comes from the absorption term from which it can be extracted according to the method described in Ref.[13]. It contributes also to the background. Finally we remind the reader that the way to obtain the cross section  $d\sigma/d\varepsilon_f$  from the transfer probability given by Eq.(3.1) was shown in Ref.[12].

In this paper we discuss the results of the calculated inclusive cross section and we show also the contribution to the absorption term of Eq.(3.1), as a function of the energy, of a number of relevant  $j_f$ -single-particle final states. Since each of these individual terms due to a final state of definite angular momentum is normalized by the transmission coefficient from the optical model describing the neutron-target interaction, then it contains the same physical information as the structure function of the state as defined in [1] and in [16].

Our model requires the calculation of the optical model S-matrix which describes the re-scattering of the neutron on the target. In the next section we will discuss two approaches to such a calculation. First by calculating the S-matrix with an energy dependent optical potential we shall be able to reproduce the experimental spectra, to explain the structure of the bumps in terms of the single particle states contributing and also to estimate the resonance energy for each  $j_f$ -state. On the other hand we will calculate also the S-matrix near resonance with a semiclassical formula which contains a Lorentzian shape for the resonance with an explicit dependence on the width. This calculation, although less accurate than the calculation with the optical model S-matrix, is useful to disentangle the single particle properties of the final states from the resonant behavior of the cross section which, in some cases, might be due only to an optimum Q-value effect.

## 4 S-matrix calculations

### 4.1 *Optical model*

An important part of the calculation is the evaluation of the optical model S-matrix which is responsible for the convergence of the sum over final angular momenta appearing in Eq.(3.1) [12, 11] and for the continuum energy distribution of each final state of definite angular momentum.

The form Eq.(3.1) of the transfer probability shows that the transfer process happens through a re-scattering on the target of the neutron emitted from the projectile. The S-matrix appearing in Eq.(3.1) describes in fact the neutron- target interaction. Such an interaction can be described by the nucleon-nucleus optical potential which is energy dependent. One has to take properly into account this dependence for the description of transfer to the continuum. In fact the energy spectrum of a neutron transfer reaction, corresponds to a collection of information on neutron scattering on the target in a range of incident energies corresponding to the neutron final continuum energies in the target. The peculiarity of the transfer process is that since the neutron comes from inside a nucleus it carries a momentum and an angular momentum higher than those it could have had if it were free. This allows it to match the high angular momentum resonance states of the target which are in the continuum.

In this paper we have used the following form of the real plus spin-orbit neutron-target optical potential

$$V(r) = V_R(\varepsilon_f)f(X_R) + \left(\frac{\hbar}{m_\pi c}\right)^2 V_{so} \frac{1}{r} \frac{d}{dr} f(X_R) l \cdot \sigma \quad (4.1)$$

while the imaginary part contains a volume and a surface term given by

$$W(r, \varepsilon_f) = W_V(\varepsilon_f)f(X_I) - 4a_I W_S(\varepsilon_f) \frac{d}{dr} f(X_I). \quad (4.2)$$

In both the above equations  $f(X_\alpha)$  has a Woods-Saxon shape

$$f(X_\alpha) = \left[1 + \exp \frac{r - R_\alpha}{a_\alpha}\right]^{-1} \quad (4.3)$$

with  $R_\alpha = r_\alpha A^{1/3}$ ,  $\alpha = R, I$  for the real and imaginary parts of the potential respectively, and  $A$  is the target mass number.

In order to reproduce the experimental spectra the depths of the real and imaginary parts of the optical potential have to be energy dependent. The parameterization of such an energy dependence has been discussed in detail in Refs. [9, 12] as it plays a key role in the analysis of the inclusive cross sections. In the next subsection we describe in detail the different parameterizations used to calculate the reactions studied in this paper.

## 4.2 Optical potential parameterization

The nucleus which has been more widely studied both from the experimental and from the theoretical point of view is  $^{208}\text{Pb}$ . To get the energy dependence of the strength of the imaginary potential we followed two different parameterizations of the neutron- $^{208}\text{Pb}$  optical potential given by Mahaux and Sartor [15, 16]. Refs.[15, 16] contain also a very detailed study of the resonance states of  $^{208}\text{Pb}$ .

We have found that the parameterization of Ref.[15] gives a good description of the high energy part of the spectra and we have used it for the calculations relative to the reaction  $^{207}\text{Pb}(^{20}\text{Ne}, ^{19}\text{Ne})^{208}\text{Pb}$  at  $E_{inc} = 48\text{MeV/nucleon}$  together with the real potential of Ref.[9] while we have used both the real and imaginary potential of [16] with a spin-orbit strength  $V_{so} = 7\text{MeV}$  for the reaction  $^{208}\text{Pb}(\alpha, ^3\text{He})^{209}\text{Pb}$  at  $E_{inc} = 30\text{MeV/nucleon}$  because it gives a better description of the spectra at low excitation energy.

In Ref.[9] we gave the sequence of bound and resonance states obtained in the real plus spin-orbit part of the potential when the real depth is taken  $V_R = -45.8\text{MeV}$ . The sequence of the states shown in Ref.[9] agrees with the single particle levels obtained for  $^{208}\text{Pb}$  by using the quasiparticle-phonon model [1].

For the reactions having  $^{48}\text{Ca}$  and  $^{90}\text{Zr}$  as a target we used the parameterizations suggested in [16] and [17] respectively. For the other two targets  $^{120}\text{Sn}$  and  $^{64}\text{Ni}$  we have constructed our own parameterization since there is no one in the literature. In both cases we used an optical potential having only a surface imaginary part. In Fig.1 the full lines show the strengths of the real and imaginary part used for  $^{120}\text{Sn}$  while the dashed line shows the real potential strength used for  $^{64}\text{Ni}$ . The imaginary part parameterization was the same as that suggested in [16], corrected for the appropriate Fermi energy which we took equal to  $E_F = -8\text{MeV}$ .



### 4.3 Semiclassical $S$ -matrix

The transfer to high angular momentum resonances that we would like to discuss in detail now is a peripheral process dominated by barrier penetration effects. These effects are important for potentials which are weakly absorbing at the barrier, the so called surface transparent potentials. When the two nuclei are at the distance of closest approach there is an effective barrier between them such that the neutron can tunnel to a final continuum state in the target.

The resonance states in the target that we are concerned with here are states like the  $1k_{17/2}$  in lead. They have positive energy which is however below the barrier formed by the real plus centrifugal potential plus spin-orbit potential as it is shown in Fig.2. As we will show they are responsible for the bumps in the inclusive spectra. In such a situation one can try to calculate the  $S$ -matrix which appears in Eq.(3.1) by using a semiclassical approximation based on a WKB evaluation of the barrier penetration probability.

The formulas used in the following are due to Brink and Takigawa [19] and they are also discussed in Ref.[20]. The  $S$ -matrix is given by

$$S = e^{2i\delta_1} \left[ \frac{1 + N \exp(2iS_{32})}{N + \exp(2iS_{32})} \right]. \quad (4.4)$$

The quantity  $\delta_1$  in Eq.(4.4) is the external phase shift which could be complex if there were direct reactions. We take it zero since we are interested in discussing only the absorptive part of the scattering. Also

$$N^2 = 1 + \exp(-2Q). \quad (4.5)$$

In Eq.(4.5)  $Q$  is a WKB barrier penetration integral and

$$P = 1 - 1/N^2 = \frac{1}{1 + \exp(2Q)} \quad (4.6)$$

is a barrier penetration factor. The quantity  $S_{32}$  is defined in Appendix A. It is a WKB integral between turning points in the pocket behind the centrifugal barrier. It has a real part and an imaginary part corresponding to absorption so we can write

$$\exp(2iS_{32}) = A \exp(2i\psi) \quad (4.7)$$

where  $A$  and  $\psi$  are real and  $A < 1$ .

We can get resonances by choosing  $0 < A < 1$  and taking some suitable energy dependence for  $\psi$ . Then

$$1 - |S|^2 = P \times \frac{1 - A^2}{1 + 2A\sqrt{1 - P} \cos(2\psi) + A^2(1 - P)}. \quad (4.8)$$

One gets a resonance when  $\cos(2\psi) = -1$ . Expanding about the resonance position,  $\cos(2\psi) \approx -1 + \alpha(\varepsilon_f - \varepsilon_{res})^2$  gives

$$1 - |S|^2 = P \times \frac{1 - A^2}{(1 - A\sqrt{1 - P})^2} \times \frac{(\Gamma/2)^2}{(\varepsilon_f - \varepsilon_{res})^2 + (\Gamma/2)^2}. \quad (4.9)$$

A simple form for  $P$  would be to take a centrifugal barrier with a sharp cut-off at a radius  $R$ . We put  $f = l(l + 1)/(kR)^2$ . Then

$$Q = kR \left[ \sqrt{f} \ln \left( \sqrt{f} + \sqrt{f - 1} \right) - \sqrt{f - 1} \right] \quad (4.10)$$

if  $f > 1$  (below the barrier) and  $Q = 0$  if  $f < 1$  (above the barrier). With the penetration factor given by  $P = \exp(-2Q)$  which is appropriate for a sharp cut-off barrier.

A more accurate calculation of  $P$  would require the evaluation of the barrier penetration integral [20]

$$Q = \int_b^c |\gamma(r)| dr \quad (4.11)$$

at several energies around the resonance energy. This is not necessary for the calculation of Eq.(4.9) while it will be done for the calculation of the escape widths as we shall see in the next section. In Eq.(4.11)  $b$  and  $c$  are the outer and outermost turning points and

$$\gamma^2(r) = \left| \frac{2m}{\hbar^2} (V(r) - \varepsilon_{res}) + \frac{l(l + 1)}{r^2} \right| \quad (4.12)$$

with  $V(r)$  given by Eq.(4.1).

## 5 Width

### 5.1 *Escape width*

In Eq.(4.9)  $\Gamma$  is the total width of the resonance. In Appendix A we show that it is approximately given by [20, 21]

$$\Gamma \approx \Gamma_0 + \Gamma^\downarrow \quad (5.1)$$

where

$$\Gamma_0 = \hbar P/T \quad (5.2)$$

is the escape width which is due to the decay of the resonance by barrier penetration and

$$\Gamma^\downarrow \sim 2 \langle W \rangle \quad (5.3)$$

is the spreading width defined in terms of the expectation value of the imaginary potential. The spreading width and the definition of  $\langle W \rangle$  will be discussed in the next section.

In Eq.(5.2)  $P$  is the barrier penetration factor discussed in the previous section and  $T$  is the period of the radial motion of the particle in the real potential pocket inside the barrier[14]

$$T = \frac{2m}{\hbar^2} \int_a^b \frac{1}{|\gamma(r)|} dr \quad (5.4)$$

where  $a$  is the internal turning point.

Eq.(5.1) corresponds to the usual definition of the single-particle width as the sum of an escape width and a spreading width. However one has to keep in mind that this distinction is somehow arbitrary because the direct decay of the particle which is represented by the escape width is modified by the coupling to other modes of excitation of the nucleus which are described by the spreading width. Therefore the evaluation of the escape width is very sensitive to the model used.

The definition of the escape width given in Eq.(5.2) takes into account only the interaction of the particle with a static potential. The quasiparticle-phonon model used instead in Ref.[22] treats the single-particle potential as a dynamic quantity calculated within the Random Phase Approximation and we expect that the escape widths calculated in this way would be larger than ours.

The interesting point about Eq.(5.2) is that although it is very easy to calculate it gives results quite accurate, comparable to those obtained in Ref.[22] and in Ref.[23]. However it is important to notice that in all methods the values of the escape width depend very much on the parameters used to define the single particle potential. Our method is mainly sensitive to the value of the radius parameter which determines the position of the top of the barrier. Small changes in the potential give rise to changes in the turning points in Eqs.(4.11) and (5.4) and the overall effect can be a variation of a factor two or more on the value of the escape width. Then the agreement between various methods refers only to the order of magnitude.

Another advantage of Eq.(5.2) is that it can be used to estimate the escape widths of states of very high angular momentum like the  $1l_{19/2}$  and  $1m_{21/2}$  in lead which are interesting for the present discussion and that could not be studied with the methods of [22, 23]. Also it allows to understand the variation of the escape width with the resonance energy. To discuss this point we show in Fig. 2 the real plus centrifugal plus spin orbit potentials relative to  $l_f = 7$ ,  $l_f = 8$ ,  $l_f = 9$  and  $l_f = 10$  respectively. The resonance energies indicated correspond to the peaks of the  $l$ -terms shown in Fig.4 and they are given also in Table 3.

It is easy to understand that increasing the resonance energies the barrier penetration range decreases thus giving a larger barrier penetration factor  $P$ . At the same time the period increases because the particle makes a wider orbit inside the potential. But while the variations of  $P$  are of an order of magnitude in the case of  $l_f = 7$  and  $l_f = 8$ , the variation of  $T$  are only few percent, therefore the escape width increases by raising the resonance energy for a given  $l_f$ -state. This discussion is illustrated by the values of  $\Gamma_0$ ,  $P$  and  $T$  given in Table 1.  $P$  was calculated from Eqs.(4.6) and (4.11) and  $T$  from Eq.(5.4). The possible values of the resonance energies given in Table 1 for each  $l_f$ -state are taken around the values given in Table 3 and they are all below the top of their respective potential barriers as shown in Fig.2.

The interesting point is that while the low-lying resonances like the  $1k_{17/2}$  have very small escape width, other states of higher angular momentum and higher resonance energy have larger escape widths although they still maintain quite strong single particle characteristics as we shall see in the following sections.

The calculations in this section have been done by using MATHEMATICA [24].

Table 1: Escape and spreading widths from Eqs.(5.2) and (5.6)

$l_f$	$\varepsilon_{res}(\text{MeV})$	P	T( $10^{-22}\text{sec}$ )	$\Gamma_0(\text{MeV})$	$\Gamma^\dagger(\text{MeV})$
7	3	$0.10 \cdot 10^{-3}$	1.67	$0.44 \cdot 10^{-3}$	3.36
7	5	$0.38 \cdot 10^{-2}$	1.74	$0.15 \cdot 10^{-1}$	4.15
7	7	$0.35 \cdot 10^{-1}$	1.91	0.12	5.04
7	9	0.15	1.96	0.50	5.49
8	4	$0.30 \cdot 10^{-4}$	1.46	$0.13 \cdot 10^{-3}$	3.88
8	6	$0.78 \cdot 10^{-3}$	1.54	$0.30 \cdot 10^{-2}$	4.75
8	8	$0.71 \cdot 10^{-2}$	1.63	$0.28 \cdot 10^{-1}$	5.42
8	10	$0.36 \cdot 10^{-1}$	1.68	0.14	6.09
9	14	0.05	1.51	0.20	6.99
9	16	0.13	1.69	0.49	7.29
9	18	0.27	1.79	0.97	7.51
9	20	0.45	2.20	1.34	7.36
10	18	0.04	1.34	0.19	8.53
10	20	0.09	1.41	0.45	8.39
10	22	0.20	1.55	0.86	8.75
10	24	0.35	1.72	1.35	8.69

## 5.2 Spreading width

The second term in Eq.(5.1) corresponds to the spreading width. We have defined it in terms of the average imaginary potential inside the barrier to be consistent with the discussion in Sec.4 where we described the neutron target interaction by an optical potential. However it is important to point out that equation (5.3) is quite generic and that there is not an unique prescription for calculating  $\langle W \rangle$ . In Appendix A we derive Eq.(5.1) within the semiclassical approximation to the S-matrix and we show that in order to be consistent with the approximation, the spreading width has to be defined as the time average of the imaginary potential

$$\Gamma^\downarrow = -2 \frac{\int_a^b W(r(t)) dt}{\tau}. \quad (5.5)$$

Another possible definition of the average potential giving the spreading width can be obtained in the framework of the nuclear many body theory [25]. In that approach the spreading width, also called damping width, is related to the imaginary part of the self energy which appears in the definition of the Green function of an excitation [16, 25]. The definition given in Ref.[25] is the following

$$\Gamma^\downarrow = -2 \int \rho(r) W(r) d^3r. \quad (5.6)$$

where  $\rho(r)$  is the single particle density given by  $\rho(r) = |\psi(r)|^2$  and  $\psi(r)$  is the single particle wave function. In Appendix B it is shown that if one uses for  $\psi(r)$  the semiclassical form of the wave function, then Eq.(5.5) and Eq.(5.6) are equivalent.

There are two important characteristics of the spreading widths that we wish to study, namely their dependence on the energy and on the angular momentum. Eq.(5.5) depends on the angular momentum in several ways. One is that the classical orbit along which we are taking the time integral of  $W$  is in fact angular momentum dependent. We should have written  $r_\lambda(t)$  where  $\lambda = l + 1/2$  is the classical angular momentum corresponding to the quantum angular momentum  $l$ . Then the angular momentum enters also in the centrifugal potential and as a consequence the period of the classical orbit inside the real potential pocket and the turning points will depend on  $l$ . The energy dependence of  $\Gamma^\downarrow$  is due mainly to the fact that the strength of  $W$  is energy dependent and also to the dependence of the period and of the turning points on the resonance energy.

On the other hand Eq.(5.6) depends on the angular momentum because of the dependence on the single particle wave function  $\psi_{lm}(r)$  and on the energy because both  $\psi$  and  $W$  are energy dependent. However in Ref.[25] it was argued that the density of single particle orbitals  $\rho(r)$  which appears in Eq.(5.6) could be taken constant in the vicinity of the absorptive region. This assumption is justified by the fact that the imaginary potential varies appreciably only on the surface of the nucleus where the radial parts of the wave functions of the high spin single particle states that we are studying in this paper are also peaked. Then according to Ref.[25] one can write

$$\Gamma^+(\varepsilon_f) = 8\pi \bar{\rho} \int r^2 dr W(r, \varepsilon_f) \quad (5.7)$$

where  $\bar{\rho} \sim 0.08 fm^{-3}/A$  is an average density of the single-particle orbitals in the vicinity of the absorptive region. The value chosen is half the nuclear matter density. This approximation allows us to use the Brown and Rho [18] parameterization for the volume integral of the optical potential which appears in eq.(5.7)

$$[r^2]_W^{BR} = \frac{4\pi}{A} \int r^2 dr W(r, \varepsilon_f) = -b_2 \frac{(\varepsilon_f - E_F)^2}{(\varepsilon_f - E_F)^2 + r_2^2} \quad (5.8)$$

According to Mahaux and Sartor [15] the values of the parameters in Eq.(5.8) for the neutron- $^{208}Pb$  potential are  $b_2 = 70.95 MeV fm^3$ ,  $r_2 = 10.87 MeV$  and  $E_F = -5.65 MeV$ . In Ref.[15] it was shown that the energy dependence of  $W_V$  and  $W_S$  saturates around  $40 MeV$ , then according to Eq.(5.7) the widths  $\Gamma^+(\varepsilon_f)$  are constant for energies in the continuum larger than  $40 MeV$  and the single particle effects disappear. The widths obtained from Eqs.(5.7) and (5.6) are given in Table 2a. Our values are in agreement with those given in Refs.[1, 25].

Furthermore in Table 1 we give, for several angular momenta and energies, the spreading widths obtained from Eq.(5.5). It is interesting to notice that there is some dependence on the energy for each fixed angular momentum but there is not much dependence on the angular momentum for a fixed energy. However there is a continuous smooth increase of the width increasing the energy and the widths obtained with this method agree within 20% with the widths of Table 2a. One could get a better agreement just by choosing a slightly different value of the constant density of orbitals in Eq.(5.7).

The choice of the most appropriate  $\bar{\rho}$  is discussed in Appendix C where we show

Table 2a: Spreading width as a function of the continuum energy in  $^{208}Pb$

$\varepsilon_f(\text{MeV})$	$\Gamma^\dagger(\text{MeV})$	$\varepsilon_f(\text{MeV})$	$\Gamma^\dagger(\text{MeV})$	$\varepsilon_f(\text{MeV})$	$\Gamma^\dagger(\text{MeV})$	$\varepsilon_f(\text{MeV})$	$\Gamma^\dagger(\text{MeV})$
1	3.1	11	7.9	21	9.7	31	10.4
2	3.7	12	8.2	22	9.8	32	10.5
3	4.4	13	8.5	23	9.9	33	10.5
4	5.0	14	8.7	24	10.0	34	10.6
5	5.5	15	8.9	25	10.1	35	10.6
6	6.1	16	9.1	26	10.2	36	10.6
7	6.5	17	9.2	27	10.2	37	10.7
8	6.9	18	9.3	28	10.3	38	10.7
9	7.3	19	9.5	29	10.3	39	10.7
10	7.6	20	9.6	30	10.4	40	10.7

the radial dependence of the imaginary potential and of the radial wave functions of some high spin resonance states which have been calculated numerically. With those wave functions it is possible to calculate Eq.(5.6) exactly and to compare the spreading widths obtained in this way to those obtained using the approximation of a constant density of orbitals.

The reason why we wish to study in detail the validity of Eq.(5.6) is that together with Eqs.(5.7) and (5.8) it gives a very simple way of estimating the spreading widths by relating them to the parameterizations of the imaginary potential. The calculation of Eq.(5.5) requires instead the numerical calculation of the turning points, of the period of the radial motion, of the classical trajectory of the particle within the real potential and finally of the time integral of the imaginary potential.

## 6 Results and Discussion

In this section we discuss the results of the calculations made for the reactions listed in Sec.2 and we compare them to the corresponding experimental spectra .

First we discuss the group of reactions made at  $E_{inc} = 30\text{MeV/nucleon}$  and hav-



**Table 2b:** Spreading width as a function of the continuum energy in  $^{91}\text{Zr}$

$\varepsilon_f(\text{MeV})$	$\Gamma^\dagger(\text{MeV})$
1	9.7
5	10.14
10	10.67
15	11.21
20	11.74
25	12.28
30	12.81
35	13.35
40	13.88

ing the  $\alpha$ -particle as a projectile. These reactions are easier to analyze because there is only one initial state in the projectile to be taken into account. For a discussion of the initial state effect we refer the reader to Refs. [9] and [12].

Fig.3a shows the experimental and calculated spectra of the reaction  $^{208}\text{Pb}(\alpha, ^3\text{He})^{209}\text{Pb}$  at  $E_{inc} = 30\text{MeV/nucleon}$ . The full curve (a) is the calculated inclusive cross section which is arbitrarily normalized to the experimental spectrum from [7] while the full curve (b) is the breakup part of the spectrum multiplied by a factor ten. We give also the energy distribution of the cross section corresponding to the absorption term in Eq.(3.1) for some fixed values of the final angular momentum namely  $l_f = 6 - 10$ . For each  $l_f$  the contributions corresponding to  $j_f = l_f \pm 1/2$  were summed. This is possible because the transfer probability Eq.(3.1) contains an incoherent sum of final  $j_f$ -values. We remind that for this calculation we have used the optical potential of Ref.[16].

In Fig.3b we show again some of the calculations of Fig.3a made with the optical model S-matrix (full curves) while the dotted curves were obtained using the semi-classical S-matrix, Eq.(4.9), discussed in Sec.4. The width appearing in Eq.(4.9) is the total width but from the values of the escape and spreading widths given in Tables 1 and 2a respectively we see that the escape widths are negligible compared

Table 3: Resonance energies in  $^{208}Pb$

$l_f$	6	7	8	9	10
$\epsilon_{res}(\text{MeV})$	3.2	4.8	5.5	12	19

to the spreading widths and therefore we shall use Eq.(4.9) with  $\Gamma \approx \Gamma^\dagger$ . We have used Eq.(4.9) with  $A=0.93$  for  $l_f = 8$ ,  $A=0.9$  for  $l_f = 9$ , and  $A=0.85$  for  $l_f = 10$ . The parameter  $A$  gives the magnitude of the S-matrix. Values of  $A$  close to one are appropriate for a weak absorptive potential. The values of  $\Gamma^\dagger$  are given in Table 2 and were obtained from Eqs.(5.7) and (5.8). The resonance energies in Eq.(4.9) were taken as the peak energies of the  $l_f$ -resonant terms obtained from the optical model calculation.

In Fig.3b we notice that the absolute values of the cross sections calculated with the two different methods agree in all cases. This proves that the semiclassical S-matrix is a good approximation to the optical model S-matrix for the kind of reactions we are concerned with in this paper. According to Eq.(4.9) the single particle resonance states have a Lorentzian energy distribution around the resonance energy. By looking at Fig.3a we notice that for the  $l_f = 8$ ,  $l_f = 9$  and  $l_f = 10$  resonances also the optical model calculation gives an energy distribution very close to a Lorentzian. Therefore in all cases the Lorentzian approximation to the spectral distribution of the single particle states in the continuum seems to be correct. Also the values the widths obtained from Eq.(5.7) agree with the exact values implicitly contained in the optical model S-matrix and we can say that Eq.(5.7) gives a very reasonable estimate of the spreading widths of the single particle states.

In Fig.3c we show the spectrum of the reaction  $^{90}Zr(\alpha, ^3He)^{91}Zr$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7]. The full curve superimposed to the experimental spectrum is our calculation of the inclusive cross section while the full curve in the lower part of the figure is the calculated breakup. We show also the individual  $l_f = 3 - 8$  final angular momentum contributions from the optical model calculation. One notices that the big bump in the experimental spectrum can be attributed to the combined effects of the  $l_f = 5 - 6$  final angular momenta corresponding to the population of the  $1h_{9/2}$  and  $1i_{13/2}$  states in  $^{91}Zr$ . The rising of the  $l_f = 5$  contribution towards negative final

energies is due to the presence of the  $1h_{11/2}$  bound state which has a large probability of being populated since in this reaction the spin matching conditions [26] require  $j_i = l_i + 1/2 \rightarrow j_f = l_f + 1/2$ .

Fig.3d refers to the reaction  ${}^{64}\text{Ni}(\alpha, {}^3\text{He}){}^{65}\text{Ni}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7]. The notation is analogous to the previous figure and in this case the bump can be attributed to the population of the  $l_f = 4 - 5$  final states.

Finally in Fig.3e we show the results of the reaction  ${}^{120}\text{Sn}(\alpha, {}^3\text{He}){}^{121}\text{Sn}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7]. In this case the spectrum looks quite different from the previous ones because it contains a large breakup contribution indicated by the dotted curve. There is a peak at about  $\varepsilon_f = 1\text{MeV}$  due to the  $l_f = 6$   $j_f = 13/2$  single particle state which is the only one strongly populated. The  $l_f = 7$  contribution is spread over all final energies thus showing that the state  $1j_{15/2}$  is completely damped in  ${}^{121}\text{Sn}$ .

We turn now to the discussion of the reactions made at high incident energy.

Fig.4 refers to the reaction  ${}^{207}\text{Pb}({}^{20}\text{Ne}, {}^{19}\text{Ne}){}^{208}\text{Pb}$  at  $E_{inc} = 48\text{MeV/nucleon}$ . The experimental spectrum is compared to the calculated inclusive cross section. The details of such a calculation were explained in Ref.[9] and in the previous section. The discrepancy between experiment and calculation for energies larger than  $50\text{MeV}$  is due to a cut in the experimental spectrum [9]. In the lower part of the figure we show also the contributions to the cross section of the terms  $l_f = 8 - 13$ .

It is interesting to notice that the experimental spectrum has three bumps at  $E_x \sim 14\text{MeV}$ ,  $E_x \sim 25\text{MeV}$ , and  $E_x \sim 34\text{MeV}$ . At the same excitation energies also the calculated spectrum has bumps and from the  $l_f$ -decomposition in the low part of the figure we notice that the bumps correspond to the maximum value of the  $l_f = 8$ ,  $l_f = 10$  and  $l_f = 11$  contributions. However because of the cut in the experimental spectrum it is not so clear that the  $l_f = 11$  state gives rise to an isolated bump. The  $l_f = 8$  term shows two peaks. The first one corresponds to the  $1k_{17/2}$  state, while the second one around  $E_x \sim 20\text{MeV}$  corresponds to the state  $1k_{15/2}$ .

The dotted curve shows the elastic breakup which is calculated from the first term of Eq.(3.1). This is a very small contribution to the total inclusive cross section in the low energy part of the spectrum and it has a maximum around  $E_x = 50\text{MeV}$ . Its effect is just to enhance the  $l_f = 10 - 11$  contributions.

To our knowledge this is the first clear example of the persistence of single particle

effects in the excitation energy region  $E_x = 20 - 40 \text{ MeV}$ . This is possible because in this energy range the widths of the  $l_f$ -resonances are still smaller than the spacing between them. At higher excitation energies this is not true anymore. In fact the  $l_f = 12$  contribution has a total width of about  $13 \text{ MeV}$  while the spacing between the  $l_f = 12$  and  $l_f = 13$  maxima is about  $10 \text{ MeV}$ . We remind that the widths of the  $l_f$ -resonances calculated with the optical model S-matrix depend on both the real as well as on the imaginary parts of the optical potential and then they are the sum of the escape plus spreading widths. The optical potential used for this calculation[9] is different from the potential used for the previous reaction on lead[16]. The main difference is that for final energies larger than  $10 \text{ MeV}$  the real potential of Ref.[16] is shallower than ours for about  $3 \text{ MeV}$ . As a result the peaks of the  $l_f = 9$  and  $l_f = 10$  terms in Fig.3a are shifted by about  $4 \text{ MeV}$  with respect to the results shown in Fig.4 and resumed in Table 3, but the peak of the  $l_f = 8$  contribution in Fig.3a fits very well the bump in the experimental spectrum.

Then from the analysis of Fig.4 we can conclude that the transfer to the continuum inclusive spectrum of  $^{208}\text{Pb}$  at incident energy  $E_{inc} = 48 \text{ MeV}$  is still dominated by absorption effects on single particle high-angular momentum states of the target. This is due to various combined facts. One is that the single particle damping is not very big for excitation energies lower than  $40 \text{ MeV}$  and also the escape probability of the particle from the potential pocket via barrier penetration is small. Furthermore there are good matching conditions between the initial and final energies and angular momenta of the neutron in the projectile and target respectively and this enhances the transfer probability.

Fig.5 shows the experimental and calculated spectra of the reaction  $^{208}\text{Pb}(^{40}\text{Ar}, ^{39}\text{Ar})^{209}\text{Pb}$  at  $E_{inc} = 41 \text{ MeV/nucleon}$ [6, 9]. In Ref.[9] we showed that this reaction is dominated by the transfer from the  $1d_{3/2}$  initial state in  $^{40}\text{Ar}$  (solid curve superimposed to the experimental spectrum). The  $l_f$ -decomposition in the lower part of the figure refers to the transfer cross section from this initial state only. As in the previous reaction the  $l_f = 8$  term (full curve) gives rise to two peaks. The individual  $1k_{17/2}$  contribution is shown separately by the dotted curve. The second peak in the full curve is due to the  $1k_{15/2}$  and it is higher than the first one because in this case the spin matching condition requires  $j_i = l_i - 1/2 \rightarrow j_f = l_f - 1/2$ . The  $l_f = 9$  (dashed line) contribution has quite a large width and its contribution does not

give a definite bump in the experimental spectrum, rather it is responsible together with the  $l_f = 10$  contribution shown by the dotdashed curve, for the second bump in the experimental spectrum. This result confirms the analysis done for the previous reaction namely that the experimental spectra contain evidence for the population by transfer of the  $l_f = 10$  state in  $^{208}Pb$  and  $^{209}Pb$ .

Fig.6 shows the calculated inclusive spectrum of the reaction  $^{90}Zr(^{20}Ne, ^{19}Ne)^{91}Zr$  at  $E_{inc} = 40MeV/nucleon$ . From the angular momentum decomposition in the lower part of the figure one notices that the first bump is due to the population of the  $1i_{13/2}$  and  $1h_{9/2}$  states, while the second bump is due to the  $1j_{15/2}$  state. The experimental spectrum for this reaction has been measured [3] and it does show two bumps at the same energies  $E_x = 16MeV$  and  $E_x = 30MeV$  predicted by our calculations.

Therefore we can conclude that the resonant behavior of the transfer inclusive spectra so far analyzed does correspond in fact to an absorption of the neutron into states of the target which have definite angular momentum and a Lorentzian distribution around the resonance energy. We can predict the widths of these states with good accuracy by using Eqs.(5.1), (5.2), (5.7). Our analysis predicts the persistence of single particle effects in lead up to  $l_f = 10$  at  $E_x \approx 26MeV$ . However the states  $l_f = 11$  and  $l_f = 12$  give also a resonant behavior around  $E_x \simeq 32MeV$  and  $E_x \simeq 42MeV$  respectively as shown in Fig.4, but such energies are very close the top of their respective potential barriers which are at  $30MeV$  and  $35MeV$ . Then the treatment of these states as single particle potential resonances cannot be justified so well and we conclude that the resonant behavior is due only to an optimum Q-value effect. In  $^{91}Zr$  single particle states are populated up to  $E_x = 30MeV$ .

Finally Fig.7 shows the results of the reaction  $^{48}Ca(^{20}Ne, ^{19}Ne)^{48}Ca$  at  $E_{inc} = 48MeV/nucleon$ [8]. The full curve superimposed to the experimental spectrum is the calculated inclusive cross section. The dotted curve is the total breakup which in this case gives the dominant contribution to the spectrum. This result is consistent with the  $l_f$ -decomposition shown in the lower part of the figure. Only the  $l_f = 4$  contribution, centered around  $E_x = 9MeV$  seems to have a single particle behavior. All the other contributions have widths larger than their spacing and we have checked that their resonance energies are higher than the top of the corresponding real plus centrifugal potentials. Then also in this case the resonant behavior of the individual  $l_f$ -terms does not correspond to the population of a single particle state but rather

to an energy matching effect. The experimental spectrum does not show any isolated bump, but just a structureless continuum with a maximum corresponding to the maximum breakup probability.

## 7 Conclusions

In this paper we have studied the microscopic origin of the resonant structures shown by several one-neutron transfer to the continuum experimental spectra. It has been demonstrated that they are due to the population of single particle states of positive energy and high angular momentum.

The single particle nature of the excitations has been clarified by introducing a semiclassical S-matrix which describes the rescattering of the neutron on the target. Near resonance the semiclassical S-matrix exhibits a Lorentzian shape with an explicit dependence on the width. We have shown that within the semiclassical theory the width can be written as a sum of an escape width plus a spreading width. The physical meaning of these two terms has been discussed and we have given some prescription to calculate them.

The spreading width has been analyzed in great detail because it gives the largest contribution to the total width. Three different ways of calculating it have been proposed and the numerical values obtained agree within 20%. The spreading widths increase smoothly with the neutron energy in the continuum following the energy dependence of the imaginary part of the optical potential. It was also shown that the dependence on the angular momentum is negligible.

Our calculations suggest that the low energy bump shown by the spectra of the reactions made at  $E_{inc} = 30 \text{ MeV/nucleon}$  is due to the population of single particle states of high angular momentum but low final energy.

On the other hand the results of our analysis seem to suggest that single particle effects in lead and zirconium persist at excitation energies as high as  $20-30 \text{ MeV}$ . The reason is that states like the  $1m_{21/2}$  in lead or the  $1j_{15/2}$  in zirconium, have resonance energies which are still below the top of the barrier formed by the real plus centrifugal plus spin-orbit potentials and their escape probabilities by barrier penetration are small. Furthermore the strength of the imaginary part of the neutron- $^{208}\text{Pb}$  optical

potential which depends on the mixing between the single particle states and other modes of excitation increases up to about  $E_x = 40\text{MeV}$  where it starts to saturate. As a consequence the spreading widths below  $40\text{MeV}$  remain smaller than the spacing between the resonances which then can still be considered isolated. Because of these properties the single particle resonances are preferred modes of excitation in a transfer reaction to targets like  $^{208}\text{Pb}$  and  $^{90}\text{Zr}$ .

This point has been illustrated by the analysis of the reactions  $^{207}\text{Pb}(^{20}\text{Ne}, ^{19}\text{Ne})^{208}\text{Pb}$  at  $E_{inc} = 48\text{MeV/nucleon}$ ,  $^{208}\text{Pb}(^{40}\text{Ar}, ^{39}\text{Ar})^{209}\text{Pb}$  at  $E_{inc} = 41\text{MeV/nucleon}$ [6, 9] and  $^{90}\text{Zr}(^{20}\text{Ne}, ^{19}\text{Ne})^{91}\text{Zr}$  at  $E_{inc} = 40\text{MeV/nucleon}$ [3]. It has been shown that the  $l_f = 10$  final state corresponding to the  $1m_{21/2}$  single particle resonance centered around  $\varepsilon_f \sim 20\text{MeV}$  in  $^{208}\text{Pb}$  can be populated by one neutron transfer with a probability higher than underlying states and the experimental spectrum shows a bump in correspondence to this state, thus indicating that the single particle damping is not complete yet. The estimated escape width for this state is about  $1\text{MeV}$  which is the same order of magnitude as the spreading width and one could hope to detect a noticeable direct decay of this state by performing an experiment similar to the one of [7]. A similar result holds for the  $1j_{15/2}$  state in  $^{91}\text{Zr}$  at  $\varepsilon_f \sim 18\text{MeV}$ .

The fact that at  $E_{inc} = 40 - 50\text{MeV/nucleon}$  transfer reactions on heavy targets dominate the inclusive cross section suggests that the description of the nucleus by a mean field is still valid. This remark might be useful for the understanding of the fragmentation process which is the dominant reaction mechanism at higher incident energies.

All the above remarks hold true because, in the reactions we have described, the background due to breakup is small. This is due to the fact that the initial bound state energy of the neutron in the projectile is large, of the order of  $10 - 20\text{MeV}$ , and the incident energy per nucleon is of the same order of magnitude such that the energy matching condition gives a most favorite final energy which is in the range of persistence of single particle states. In the case of the reactions on  $^{120}\text{Sn}$  and  $^{48}\text{Ca}$  instead, the most favorite final energies do not correspond to any single particle resonance state of the targets and the spectra are dominated by the breakup.

The results would again be different for a reaction having a projectile with a loosely bound neutron like  $^{11}\text{Be}$  or if the incident energy were much higher[11]. In those cases also the inclusive spectra would be dominated by the direct breakup

component.

### Acknowledgments

It is my pleasure to thank David Brink for numerous enlightening discussions on the semiclassical S-matrix; Sydney Galès, Simone Fortier and their group for a long correspondence and for allowing me to use several unpublished experimental spectra. Also I would like to mention the kindness of Claude Mahaux in providing me with the numerical values of some optical potential strengths and of Massimo Di Toro in reading an early version of this paper.



## Appendix A

In this appendix we derive Eq.(5.1) given in section (5.1)

First we give the definitions of  $S_{32}$  and  $A$  which appear in Eq.(4.7)

$$\exp(2iS_{32}) = A \exp(2i\psi). \quad (\text{A.1})$$

$S_{32}$  is a WKB integral defined as

$$S_{32} = \int_a^b |\gamma(r)| dr \quad (\text{A.2})$$

where

$$\gamma^2(r) = \left| \frac{2m}{\hbar^2} (V(r) + iW(r) - \varepsilon_{res}) + \frac{l(l+1)}{r^2} \right| \quad (\text{A.3})$$

In Eq.(A.2)  $a$  and  $b$  are the inner and outer turning points which appear when the real potential has three turning points. Also

$$A = \exp(2ImS_{32}) = \exp\left(\frac{2}{\hbar} \int_a^b W(r(t)) dt\right) \quad (\text{A.4})$$

where  $W(r(t))$  is the imaginary part of the potential which appears in Eq.(A.3).

If the imaginary part of the potential is small one can expand the exponential in the above definition of  $A$  and then get

$$A \approx 1 + 2\bar{W} \quad (\text{A.5})$$

also

$$A^{-1/2} = 1 - \bar{W} \quad (\text{A.6})$$

where

$$\bar{W} = \frac{1}{\hbar} \int_a^b W(r(t)) dt \quad (\text{A.7})$$

The step between Eq.(4.8) and Eq.(4.9) is made by expanding about the resonance position  $\cos(2\psi) \approx -1 + \alpha(\varepsilon_f - \varepsilon_{res})^2$ . The expansion coefficient  $\alpha$  is given by

$$\alpha = 2 \left( \frac{\partial S_{32}}{\partial \varepsilon_f} \right)^2 \Big|_{\varepsilon_f = \varepsilon_{res}} = 2 \left( \frac{\tau}{\hbar} \right)^2 \quad (\text{A.8})$$

where  $\tau$  is half the period of the classical orbit inside the real potential pocket defined by Eq.(5.4).

Then Eq.(4.9) is obtained from Eq.(4.8) if one defines a width  $\Gamma$  as

$$\Gamma = \sqrt{2} \left( \frac{1 - A\sqrt{1-P}}{\sqrt{A\alpha\sqrt{1-P}}} \right) \quad (\text{A.9})$$

where  $P$  is the barrier penetration factor defined in section (4.3).

Now since  $P < 1$  we expand  $\sqrt{1-P}$  and use also the approximated forms of  $A$  and  $\sqrt{A}$  given above which are valid when the imaginary potential is small. Then we get for  $\Gamma$

$$\begin{aligned} \Gamma &= \sqrt{\frac{2}{\alpha}} \frac{1 - A + \frac{AP}{2}}{\sqrt{A}} \\ &= \sqrt{\frac{2}{\alpha}} \left( \frac{P}{2} - 2\bar{W} + P\bar{W} \right) (1 - \bar{W}) \\ &\approx \sqrt{\frac{2}{\alpha}} \left( -2\bar{W} + \frac{P}{2} \right) \\ &= \frac{\hbar P}{T} - 2\frac{\hbar}{\tau}\bar{W} \\ &= \Gamma_0 + 2 \langle W \rangle \end{aligned} \quad (\text{A.10})$$

In getting the above result we neglected terms of the order  $P\bar{W}$  and we defined also  $\Gamma_0 = \hbar P/T$  (cf. Eq.(5.2)). Finally we defined the time average of the imaginary potential as

$$\langle W \rangle = - \frac{\int_a^b W(r(t)) dt}{\tau} \quad (\text{A.11})$$

the minus sign in the above equation is correct since in this paper we use a negative strength for the imaginary potential.

## Appendix B

In this appendix we show the equivalence between Eq.(5.5) and Eq.(5.6). We start from

$$\Gamma^\downarrow = -2 \int \rho(r) W(r) d^3r. \quad (\text{B.1})$$

where  $\rho(r)$  is the single particle density given by  $\rho(r) = |\psi(r)|^2$  and  $\psi(r)$  is the single particle wave function which depends on the quantum numbers  $l$  and  $m$  as well as on the single particle energy. In the following we omit those indices for brevity. Let

$$|\psi(r)|^2 = \left| \frac{\phi(r)}{r} Y_{lm}(\Omega) \right|^2 \quad (\text{B.2})$$

and let us take for the radial wave function  $\phi(r)$  its semiclassical form

$$\phi(r) = \frac{C}{\sqrt{\gamma}} \sin(\text{Re } S_{32} + \frac{\pi}{4}) \quad (\text{B.3})$$

where  $S_{32}$  is the WKB integral defined in Appendix A and the normalization constant  $C$  is defined in terms of the classical period as

$$|C|^2 = \frac{2m}{\hbar} \frac{1}{\tau} \quad (\text{B.4})$$

then we insert these values in Eq.(B.1) and take out of the integral a factor 1/2 which is the average value of the sinus function. Next we perform the angular integration which gives one. We approximate also the local momentum  $\gamma \sim mv/\hbar$  where  $v$  is the average velocity inside the potential pocket, in this way we can change the variable of integration from  $dr$  to  $dt = dr/v$  and we finally obtain

$$\Gamma^\downarrow = -2 \frac{\int_a^b W(r(t)) dt}{\tau} \quad (\text{B.5})$$

## Appendix C

We discuss now under which conditions it is justified to approximate Eq.(B.1) by taking a constant value for the density of orbitals  $\rho(r) \sim \bar{\rho}$  outside the integral. In

**Table 4:** Spreading widths from Eq.(5.6) with the exact wave functions.

nucleus	$l_f$	$\varepsilon_{res}(\text{MeV})$	$\Gamma^\downarrow(\text{MeV})$
$^{208}\text{Pb}$	5	4.02	1.90
	7	6.82	2.01
	8	6.22	3.12
$^{91}\text{Zr}$	6	8.62	10.17
	7	17.8	11.14

Fig.8 we show the radial dependence of the imaginary part of the optical potential used for  $^{208}\text{Pb}$  which according to Eq.(4.2) is a sum of a volume and a surface term with the energy dependent strengths calculated at  $\varepsilon_f = 6.8\text{MeV}$ . Fig.9 shows instead  $R(r) = |\phi(r)/r|^2$  for three resonance states in  $^{209}\text{Pb}$ , the  $2h_{11/2}$ ,  $1k_{17/2}$  and  $1j_{13/2}$ .  $R(r)$  is the modulus square of the radial part of the wave function Eq.(B.2) normalized such that  $\int |\phi(r)|^2 dr = 1$ . One notices that while the  $2h_{11/2}$  state has a node and it has its main peak inside the nuclear surface, the  $1k_{17/2}$  and  $1j_{13/2}$  state are instead peaked at the surface where also the imaginary potential gives its main contribution. In this case the radius of the nucleus is  $R = 7.2\text{fm}$ . Therefore it seem reasonable to take the density of orbitals constant outside the integral when we are in a region of continuum energies where the shell model predicts only states of high angular momentum whose wave functions have no nodes and are peaked at the nuclear surface. For the three states discussed here we calculated the spreading widths from Eq.(B.1) with the exact wave functions shown in Fig.9 and the resulting values are shown in Table 4. They are smaller by about a factor 1.85 than the values given in Table 2. In Table 4 we give also some values of  $\Gamma^\downarrow$  calculated for  $^{91}\text{Zr}$ . In this case the agreement with the values obtained from Eq.(5.7) is very good.

Unfortunately it is rather difficult to get convergence in the numerical solution of the Schrödinger equation for states of angular momentum higher than those discussed here, otherwise one could always calculate Eq.(B.1) exactly. Then we suggest that if one is interested only in an estimate of the spreading widths having in mind that their most important characteristic is the energy dependence, Eq.(5.7) can be used together with an appropriate density of orbitals and with a good parameterization of the volume integral of the imaginary part of the optical potential.

Finally we notice that the value of the resonance energies are different than those given in Table 3, this is because the values given here come from the solution of the Schrödinger equation while those of Table 3 correspond to the peaks of the l-resonant terms of the transfer probability Eq.(3.1).

## References

- [1] S.Galès, Ch.Stoyanov and A.I.Vdovin, Phys.Rep.**166**,125 (1988) and references therein.
- [2] S.Fortier, S.Gàles, S.M.Austin, W.Benenson, G.M.Crawley, C.Djalali, J.S.Winfield and G.Yoo, Phys.Rev.**C41**,2689 (1990) .
- [3] D.Beaumel, Y.Blumenfeld, Ph.Chomaz, N.Frascaria , J.P.Garron, J.C.Roynette , T.Suomijärvi, J.Barrette, B.Berthier, B.Fernandez, J.Gastebois, W.Mittig, 1987, Annual Report IPN-Orsay, pag. 41 .
- [4] C.P.Massolo, S.Fortier, S.Galès, F.Azaiez, E.Gerlic, J.Guillot, H.Hourani, H.Langevin-Joliot, J.M.Maison, J.P.Shapira, and G.M.Crawley, Phys.Rev.**C43**, 1687 (1991).
- [5] G.H.Yoo, G.M.Crawley, N.A.Orr, J.S.Winfield, J.E.Finck, S.Galès, Ph.Chomaz, I.Lhenry and T.Suomijärvi, Phys. Rev. **C47**,1200 (1993).
- [6] I.Lhenry. Ph. D. Thesis, IPNO-T92-01, Orsay April 1992 and I.Lhenry, T.Suomijärvi, Y.Blumenfeld, Ph.Chomaz, N.Frascaria, J.P.Garron, J.C.Roynette, J.A.Scarpaci, D.Beaumel, S.Fortier, S.Gàles, H.Laurent, A.Gillibert, G.Crawley, J.Finck, G.Yoo and J.Barreto, to be published.
- [7] D.Beaumel, S.Fortier, S.Galès , J.Guillot, H.Langevin-Joliot, H.Laurent , J.M.Maison, J.Vernotte, J.Bordewijck, S.Brandenburg, A.Krasznahorkay, G.M.Crawley, C.P.Massolo and M.Renteria , IPNO DRE 93-29 preprint, and private communication.
- [8] H.Laurent, D.Beaumel, S.Fortier, S.Galès, I.Lhenry, J.M.Maison, Y.Blumenfeld, N.Frascaria, J.Guillot, J.C.Roynette, J.A.Scarpaci, T.Suomijärvi, A.Gillibert, Contributions to the ' N-N Collisions V Conference', Taormina, Italy, 1994. Edited by M.Di Toro, P.Piattelli and P.Sapienza, Pag.31.

- [9] A.Bonaccorso, I.Lhenry and T.Suomijärvi, Phys.Rev. **C49**, 329 (1994).
- [10] A.Bonaccorso and D.M.Brink, Phys. Rev. **C38**,1776 (1988).
- [11] A.Bonaccorso and D.M.Brink, Phys.Rev. **C43**, 299 (1991).
- [12] A.Bonaccorso and D.M.Brink, Phys.Rev. **C44**, 1559 (1991)
- [13] A.Bonaccorso and D.M.Brink, Phys.Rev. **C46**, 700 (1992).
- [14] L.D.Landau and E.M.Lifshitz, Quantum Mechanics, Pergamon Press London (1958), Ch. VII.
- [15] C.Mahaux and R.Sartor, Nucl.Phys. **A493**, 157 (1989).
- [16] C.Mahaux and R.Sartor, Adv. Nucl.Phys. **20**, 1 (1991).
- [17] C.Mahaux and R.Sartor, Nucl.Phys. **A568**, 1 (1994) .
- [18] G.E.Brown and M.Rho, Nucl.Phys.**A372** 397 (1981).
- [19] D.M.Brink and N.Takigawa, Nucl.Phys. **A309**, 159,( 1977).
- [20] D.M.Brink, 'Semiclassical Methods in Nucleus-Nucleus Scattering' , Cambridge University Press, 1985, pag.97, and private communication.
- [21] A.Bohr and B.R.Mottelson, 'Nuclear Structure', Vol.1, W.A.Benjamin, Inc., 1969, New York, Amsterdam, Pag. 446.
- [22] N.Van Giai and Ch.Stoyanov, Phys.Lett. **B272**, 178 (1991)
- [23] J.Bang, F.A.Gareev, I.V.Puzynin and R.M.Jamalejev, Nucl. Phys. **A261**, 59 (1976).
- [24] 'Mathematica' by Stephen Wolfram, Addison-Wesley Publishing Company, Inc., 1988.
- [25] G.Bertsch, P.Bortignon, and R.Brogia , Rev.Mod.Phys. **55**, 289 (1983).
- [26] A.Bonaccorso,D.M.Brink and L.Lo Monaco,J.Phys.G **13**,1407 (1987).

## Figure Captions

**Fig.1.** Energy parameterization of the optical potential strengths for  $^{120}\text{Sn}$  (full curves) and  $^{64}\text{Ni}$  (dashed curve).

**Fig.2.** Resonance particle states for  $^{209}\text{Pb}$  in their respective potentials given by the sum of a Woods-Saxon plus spin-orbit term plus centrifugal potential.

**Fig.3a.** Inclusive spectrum of the reaction  $^{208}\text{Pb}(\alpha, ^3\text{He})^{209}\text{Pb}$  at

$E_{inc} = 30\text{MeV/nucleon}$ [7]. The curve (a) is our total cross section, the curve (b) is the breakup multiplied by a factor 10. The other curves are the individual  $l_f = 6 - 10$  contributions to the absorption as indicated on the picture.

**Fig.3b.** The same as Fig.3a . In this case the dotted curves correspond to the calculations made with the semiclassical S-matrix.

**Fig.3c .** Spectrum of the reaction  $^{90}\text{Zr}(\alpha, ^3\text{He})^{91}\text{Zr}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7]. The full curve superimposed to the experimental spectrum is our calculation of the inclusive cross section while the full curve in the lower part of the figure is the calculated breakup. The individual  $l_f = 3 - 8$  final angular momentum contributions are also shown as indicated in the lower part of the figure.

**Fig.3d.**  $^{64}\text{Ni}(\alpha, ^3\text{He})^{65}\text{Ni}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7]. The notation is analogous to the previous figure and in this case the bump can be attributed to the population of the  $l_f = 4 - 5$  final states.

**Fig.3e.** Results of the reaction  $^{120}\text{Sn}(\alpha, ^3\text{He})^{121}\text{Sn}$  at  $E_{inc} = 30\text{MeV/nucleon}$ [7]. The full curve superimposed to the experimental spectrum is the calculated inclusive cross section. The breakup contribution is indicated by the dotted curve. The  $l_f = 4 - 7$  contributions are also shown.

**Fig.4.** Inclusive spectrum of the reaction  $^{207}\text{Pb}(^{20}\text{Ne}, ^{19}\text{Ne})^{208}\text{Pb}$  at

$E_{inc} = 48\text{MeV/nucleon}$ [9]. The full curve superimposed to the experimental spectrum is the result of our calculation. The dotted curve shows the calculated breakup while the curves labeled  $l_f = 8 - 13$  show individual contributions to the absorption cross section of several final angular momenta.



**Fig.5.** Inclusive spectrum of the reaction  $^{208}\text{Pb}(^{40}\text{Ar}, ^{39}\text{Ar})^{209}\text{Pb}$  at


$E_{inc} = 41\text{MeV/nucleon}$ [9]. The full curve superimposed to the experimental spectrum is the result of our calculation for the cross section due to transfer from the  $1d_{3/2}$  initial state in Ar. In the lower part of the figure the dotted curve shows the contribution of the  $1k_{17/2}$  final state. The solid curve is the total contribution due to  $l_f = 8$ . The second peak is due to the  $1k_{15/2}$  state. The dashed line is the contribution of  $l_f = 9$  and the dotdashed line is for  $l_f = 10$ .

**Fig.6 .** Calculated inclusive spectrum of the reaction  $^{90}\text{Zr}(^{20}\text{Ne}, ^{19}\text{Ne})^{91}\text{Zr}$  at  $E_{inc} = 40\text{MeV/nucleon}$ . The final angular momentum decomposition is shown in the lower part of the figure .


**Fig.7.** Results of the reaction  $^{48}\text{Ca}(^{20}\text{Ne}, ^{19}\text{Ne})^{48}\text{Ca}$  at  $E_{inc} = 48\text{MeV/nucleon}$ [8]. The full curve superimposed to the experimental spectrum is the calculated inclusive cross section. The dotted curve is the total breakup. The final angular momentum decomposition is shown in the lower part of the figure .

**Fig.8.** Radial dependence of the imaginary part of the optical potential from Eq.(4.2) with the strength appropriate to  $\varepsilon_f = 6.8\text{MeV}$ .

**Fig.9.**  $R(r) = |\phi(r)/r|^2$  for three resonance states in  $^{209}\text{Pb}$ , the a)  $1k_{17/2}$  , b)  $1j_{13/2}$  and c)  $2h_{11/2}$ .  $R(r)$  is the modulus square of the radial part of the wave function Eq.(B.2).



$^{209}\text{Pb}$



$^{208}\text{Pb}(\alpha, ^3\text{He})^{209}\text{Pb}$   $E_{\text{inc}} = 30\text{MeV/nucleon}$




Fig. 3a






Fig 9

$^{90}\text{Zr}(\alpha, ^3\text{He})^{91}\text{Zr}$   $E_{\text{inc}} = 120\text{MeV}$




$^{64}\text{Ni}(\alpha, ^3\text{He})^{65}\text{Ni}$


$E_\alpha = 120\text{MeV} ; \theta_\alpha = 0^\circ$




$^{120}\text{Sn}(\alpha, ^3\text{He})^{121}\text{Sn}$   $E_{\text{inc}} = 120\text{MeV}$








$^{208}\text{Pb}(^{40}\text{Ar}, ^{39}\text{Ar})^{209}\text{Pb}$   $E_{\text{inc}} = 41\text{MeV/nucleon}$



$^{90}\text{Zr}(^{20}\text{Ne}, ^{19}\text{Ne})^{91}\text{Zr}$   $E_{\text{inc}} = 40\text{MeV/nucleon}$







Fig 7

Fig. 3







Fig 9a



F1196




Fig 9c