

AC

ICHEP 0100

SUNY BING 5/27/94r

Submitted to Pa 01, 09


P1 10, 02

TESTS FOR LEPTONIC CP VIOLATION IN TAU DECAYS

Charles A. Nelson,* Minseob Kim and Hui-Chun Yang
Department of Physics, SUNY at Binghamton
Binghamton, New York 13902-6000 U.S.A.

Abstract

At the Z^0 or a B factory, there are two tests for non-CKM-type leptonic CP violation in the $\tau \rightarrow \rho\nu(a_1\nu)$ decay channel by inclusion of $\rho(a_1)$ polarimetry. By CP invariance, the moduli ratio of, and the phase difference between, the two helicity amplitudes for $\tau^- \rightarrow \rho^-\nu(a_1^-\nu)$ decay should equal those for $\tau^+ \rightarrow \rho^+\bar{\nu}(a_1^+\bar{\nu})$ decay. Formulas are given for a L-handed ν_τ , and also for an arbitrary mixture of ν_L and ν_R neutrino helicities. Statistical errors are listed for both the case that the τ^- momentum direction is not known, and when it is known via a silicon vertex detector.



* cnelson @ bingvmb.cc.binghamton.edu

Introduction

After almost 30 years, the fundamental origin and significance of the observed CP and T violations in kaon decays is still a mystery. On the other hand, it is possible to use new collider data to rigorously search for CP and T violations in the tau lepton decays.¹ While such an observation would be surprising, nevertheless we won't be sure of its absence unless we search for it. Spin-correlation effects plus decay polarimetry enable^{2,3} such a search in $e^-e^+ \rightarrow Z^0$, or $\gamma^* \rightarrow \tau^-\tau^+$.

As shown in Ref. 1, by use of ρ polarimetry, there are two tests for "non-CKM-type" leptonic CP violation in $\tau \rightarrow \rho\nu$ decay.

This is easily seen because by rotational invariance there are two independent helicity amplitudes for $\tau^- \rightarrow \rho^- \nu_\tau$ decay

$$A(-1, -1/2) = |A(-1, -1/2)| e^{i\phi_{-1}^a}, \quad A(0, -1/2) = |A(0, -1/2)| e^{i\phi_0^a} \quad (1)$$

assuming a L-handed ν_τ . The CP-conjugate decay $\tau^+ \rightarrow \rho^+ \bar{\nu}_\tau$ depends on

$$B(1, 1/2) = |B(1, 1/2)| e^{i\phi_1^b}, \quad B(0, 1/2) = |B(0, 1/2)| e^{i\phi_0^b} \quad (2)$$

assuming a R-handed $\bar{\nu}_\tau$. By CP invariance $B(\lambda_{\bar{\rho}}, \lambda_{\bar{\nu}}) = A(-\lambda_{\bar{\rho}}, -\lambda_{\bar{\nu}})$. The two tests are that the phase difference and moduli ratio for the two amplitudes for $\tau^- \rightarrow \rho^- \nu_\tau$ decay must equal those for the CP-conjugate decay. That is

$$\beta_a = \beta_b \quad (1st \text{ test}) \quad (3)$$

where $\beta_a \equiv \phi_{-1}^a - \phi_0^a$, $\beta_b \equiv \phi_1^b - \phi_0^b$; and

$$r_a = r_b \quad (2nd \text{ test}) \quad (4)$$

in terms of the moduli ratios

$$r_a \equiv \frac{|A(-1, -1/2)|}{|A(0, -1/2)|}, \quad r_b \equiv \frac{|B(1, 1/2)|}{|B(0, 1/2)|} \quad (5)$$

It is important to realize that any leptonic-CKM t-type phases will equally affect the $A(-1, -1/2)$ and $A(0, -1/2)$ amplitudes. Therefore, they will cancel out in β_a and in r_a . Hence, $\beta_a = \beta_b$ and $r_a = r_b$ test for a non-CKM-type leptonic CP violation.

In the standard lepton model (pure V-A and no CP violation), $\beta_a = \beta_b = 0$ and the moduli ratio $r_a = r_b \approx \sqrt{2} m_\rho/m_\tau = 0.613$. These two tests, (3) and (4), should be compared with the classic CP test for partial width asymmetry of CP-conjugate reactions:

$$A_\Gamma \equiv \frac{\Gamma - \bar{\Gamma}}{\Gamma + \bar{\Gamma}} \quad (6)$$

where, e.g. $\Gamma = \Gamma(\tau^- \rightarrow \rho^- \nu_\tau)$ and $\bar{\Gamma} = \bar{\Gamma}(\tau^+ \rightarrow \rho^+ \bar{\nu}_\tau)$. For τ two-body decay modes, the denominator of (6) is known to (1 to 4)%, so (at best) we know $A_\Gamma \sim$ (1 to 4)% whereas we find (see below) that the fractional uncertainty of the moduli can be measured to the $(\delta r_a)/r_a \sim$ (0.1 to 1)% level from data, respectively, at γ^* energies (at the Z^0).

Contents of this Paper

This conference contribution extends the analysis of Ref. 1 in three ways:

(a) Ref. 1 considered the $\tau \rightarrow \rho \nu$ decay mode. Here the two tests for non-CKM-type leptonic CP violation are extended to the $\tau \rightarrow a_1 \nu$ mode.

(b) Ref. 1 assumed a L-handed ν_τ for the $\tau \rightarrow \rho \nu$ mode. Here formulas are given for a mixture of V-A and V+A couplings, and for both left-handed and right-handed neutrinos in $\tau^- \rightarrow \rho^- \nu$ ($a_1^- \nu$) decay.

(c) Ref. 1 assumed that the τ^- momentum direction was only known kinematically up to two possible directions. However, by a silicon vertex detector, the τ^- momentum direction may be known at a B factory. Here we obtain and discuss the improvement in the statistical errors for the two tests for the $\tau \rightarrow \rho \nu$ mode when the τ^- momentum direction is measured.

Formulas for $\tau^- \rightarrow \rho^- \nu$ including both $V \pm A$, and both ν helicities.

Including both ν_L and ν_R helicities and using a "compact boldface formalism," we find the composite decay density matrix for $\tau^- \rightarrow \rho^- \nu \rightarrow (\pi^- \pi^0) \nu$ is

$$\mathbf{R} = \begin{pmatrix} \mathbf{R}_{++} & e^{i\phi_1^\tau} r_{+-} \\ e^{-i\phi_1^\tau} r_{-+} & \mathbf{R}_{--} \end{pmatrix} \quad (7)$$

The diagonal elements are

$$\begin{aligned} \mathbf{R}_{\pm\pm} &= n_a [1 \pm f_a \cos \theta_1^\tau] \\ &\mp (1/\sqrt{2}) \sin \theta_1^\tau \sin 2\tilde{\theta}_a [\cos(\tilde{\phi}_a - \beta_a) |A(0, -1/2)| |A(-1, -1/2)| \\ &- \cos(\tilde{\phi}_a + \beta_a^R) |A(0, 1/2)| |A(1, 1/2)|] \end{aligned} \quad (8)$$

and

$$\begin{aligned}
r_{+-} &= (r_{-+})^* \\
&= \mathbf{n}_a \mathbf{f}_a \sin \theta_1^\tau \\
&+ (1/\sqrt{2}) \sin 2 \tilde{\theta}_a \{ [\cos \theta_1^\tau \cos (\tilde{\phi}_a - \beta_a) + i \sin (\tilde{\phi}_a - \beta_a)] |A(0, -1/2)| |A(-1, -1/2)| \\
&- [\cos \theta_1^\tau \cos (\tilde{\phi}_a + \beta_a^R) + i \sin (\tilde{\phi}_a + \beta_a^R)] |A(0, 1/2)| |A(1, 1/2)| \} \quad (9)
\end{aligned}$$

Note that the two observable phase differences are

$$\beta_a \equiv \phi_{-1}^a - \phi_0^a \quad (10a)$$

$$\beta_a^R \equiv \phi_1^a - \phi_0^{aR} \quad (10b)$$

In Eqs. (8-9),

$$\begin{pmatrix} \mathbf{n}_a \\ \mathbf{n}_a \mathbf{f}_a \end{pmatrix} = \cos^2 \tilde{\theta}_a (|A(0, -1/2)|^2 \pm |A(0, 1/2)|^2) \pm \frac{1}{2} \sin^2 \tilde{\theta}_a (|A(-1, -1/2)|^2 \pm |A(1, 1/2)|^2) \quad (11)$$

Similarly, for the conjugate process $\tau^+ \rightarrow \rho^+ \bar{\nu} \rightarrow (\pi^+ \pi^0) \bar{\nu}$, including both

$\bar{\nu}_R$ and $\bar{\nu}_L$ helicities,

$$\bar{\mathbf{R}} = \begin{pmatrix} \bar{\mathbf{R}}_{++} & e^{i\phi_2^\tau} \bar{\mathbf{r}}_{+-} \\ e^{-i\phi_2^\tau} \bar{\mathbf{r}}_{-+} & \bar{\mathbf{R}}_{--} \end{pmatrix} \quad (12)$$

where

$$\begin{aligned}
\mathbf{R}_{\pm\pm} &= \mathbf{n}_b (1 \mp \mathbf{f}_b \cos \theta_2^\tau) \\
&\pm (1/\sqrt{2}) \sin \theta_2^\tau \sin 2 \tilde{\theta}_b [\cos (\tilde{\phi}_b + \beta_b) |B(0, 1/2)| |B(1, 1/2)| \\
&- \cos (\tilde{\phi}_b - \beta_b^L) |B(0, -1/2)| |B(-1, -1/2)|] \quad (13)
\end{aligned}$$

$$\begin{aligned}
\bar{r}_{+-} &= (\bar{r}_{-+})^* \\
&= -\mathbf{n}_b \cdot \mathbf{f}_b \sin \theta_2^\tau \\
&- (1/\sqrt{2}) \sin 2 \tilde{\theta}_b \{ [\cos \theta_2^\tau \cos (\tilde{\phi}_b + \beta_b) + i \sin (\tilde{\phi}_b + \beta_b)] |B(0, 1/2)| |B(1, 1/2)| \\
&- [\cos \theta_2^\tau \cos (\tilde{\phi}_b - \beta_b^L) + i \sin (\tilde{\phi}_b - \beta_b^L)] |B(0, -1/2)| |B(-1, -1/2)| \} \quad (14)
\end{aligned}$$

In Eqs. (13-14),

$$\beta_b \equiv \phi_1^b - \phi_0^b \quad (15a)$$

$$\beta_b^L \equiv \phi_{-1}^b - \phi_0^{bL} \quad (15b)$$

and

$$\begin{pmatrix} \mathbf{n}_b \\ \mathbf{n}_b \cdot \mathbf{f}_b \end{pmatrix} = \cos^2 \tilde{\theta}_b (|B(0, 1/2)|^2 \pm |B(0, -1/2)|^2) \pm \frac{1}{2} \sin^2 \tilde{\theta}_b (|B(1, 1/2)|^2 \pm |B(-1, -1/2)|^2) \quad (16)$$

The full "Stage 2 Spin-Correlation" function (S2SC) is given by

$$I_7 = I_7(\mathbf{R} \rightarrow \mathbf{R}, \bar{\mathbf{R}} \rightarrow \bar{\mathbf{R}}) \quad (17)$$

where the I_7 function on the right-hand side is given in the next equation from Ref. 1. The simpler I_7 of Ref. 1 assumed a L-handed ν in $\tau^- \rightarrow \rho^- \nu$, and a R-handed $\bar{\nu}$ in $\tau^+ \rightarrow \rho^+ \bar{\nu}$.

$$\begin{aligned}
I_7 &= I(E_1, E_2, \phi; \tilde{\theta}_1, \tilde{\phi}_1; \tilde{\theta}_2, \tilde{\phi}_2) \\
&= \sum_{h_1, h_2} |T(h_1, h_2)|^2 R_{h_1, h_1} \bar{R}_{h_2, h_2} \\
&+ e^{i\phi} T(++) T^*(- -) r_{+-} \bar{r}_{+-} + e^{-i\phi} T(- -) T^*(+ +) r_{+-} \bar{r}_{+-} \quad (18)
\end{aligned}$$

where $T(\lambda_1, \lambda_2)$ are the helicity amplitudes⁴ describing $Z^0, \gamma^* \rightarrow \tau \tau^+$.

Similarly, the simpler 4 (5) variable S2SC functions are

$$I_{4,5} = I_{4,5} + (\lambda_R)^2 I_{4,5} (\rho \rightarrow \rho^R) + (\bar{\lambda}_L)^2 I_{4,5} (\bar{\rho} \rightarrow \bar{\rho}^L) \\ + (\lambda_R \bar{\lambda}_L)^2 I_{4,5} (\rho \rightarrow \rho^R, \bar{\rho} \rightarrow \bar{\rho}^L) \quad (19)$$

where the ratios of the R-handed to L-handed $\tau^- \rightarrow \rho^- \nu$ moduli (and vice versa for $\tau^+ \rightarrow \rho^+ \bar{\nu}$) are

$$\lambda_P \equiv \frac{|A(0, 1/2)|}{|A(0, -1/2)|} \quad (20a)$$

$$\bar{\lambda}_L \equiv \frac{|B(0, -1/2)|}{|B(0, 1/2)|} \quad (20b)$$

In Eq. (19), the 4-variable S2SC of Ref. 1 is

$$I(E_{\rho^-}, E_{\rho^+}; \tilde{\theta}_1, \tilde{\theta}_2) = |\Gamma(+ -)|^2 \rho_{++} \bar{\rho}_{..} \\ + |\Gamma(- +)|^2 \rho_{..} \bar{\rho}_{++} + |\Gamma(++)|^2 \rho_{++} \bar{\rho}_{++} + |\Gamma(--)|^2 \rho_{..} \bar{\rho}_{..} \quad (21)$$

with the integrated composite decay density matrix for $\tau^- \rightarrow \rho^- \nu_\tau \rightarrow (\pi^- \pi^0) \nu_\tau$ with τ^- helicity $\lambda_1 = h/2$

$$\rho_{hh} = (1 + h \cos \theta_1^\tau) [\cos^2 \omega_1 \cos^2 \tilde{\theta}_1 + 1/2 \sin^2 \omega_1 \sin^2 \tilde{\theta}_1] \\ + (r_a^2/2) (1 - h \cos \theta_1^\tau) [\sin^2 \omega_1 \cos^2 \tilde{\theta}_1 + 1/2 (1 + \cos^2 \omega_1) \sin^2 \tilde{\theta}_1] \\ + h (r_a/\sqrt{2}) \cos \beta_a \sin \theta_1^\tau \sin 2 \omega_1 [\cos^2 \tilde{\theta}_1 - 1/2 \sin^2 \tilde{\theta}_1] \quad (22)$$

For the CP conjugate process with τ^+ with helicity $\lambda_2 = h/2$,

$$\bar{\rho}_{h,h} = \rho_{-h,-h} \text{ (subscripts } 1 \rightarrow 2, a \rightarrow b) \quad (23)$$

The additional ρ^R and $\bar{\rho}^L$ needed for Eq. (19) are defined (and given) by

$$\rho_{\pm\pm}^R \equiv \frac{1}{2\pi} \int_0^{2\pi} d\tilde{\phi}_1^{(\pm)} \frac{R_{\pm\pm}^R}{|A(0, 1/2)|^2} \\ = \rho_{-h,-h} (r_a \rightarrow r_a^R, \beta_a \rightarrow \beta_a^R) \quad (24)$$

with β_a^R given in Eq. (10b), and

$$r_a^R \equiv \frac{|A(1, 1/2)|}{|A(0, 1/2)|} . \quad (25)$$

Also

$$\begin{aligned} \bar{\rho}_{\pm\pm}^L &\equiv \frac{1}{2\pi} \int_0^{2\pi} d\tilde{\phi}_2^{(\pm)} \frac{\bar{R}_{\pm\pm}^L}{|B(0, -1/2)|^2} \\ &= \bar{\rho}_{-h-h} (r_b \rightarrow r_b^L, \beta_b \rightarrow \beta_b^L) \end{aligned} \quad (26)$$

with β_b^L of Eq. (15b) and

$$r_b^L \equiv \frac{|B(-1, -1/2)|}{|B(0, -1/2)|} . \quad (27)$$

For the 5-variable S2SC, the additional formulas are

$$\begin{aligned} \rho_{\pm\mp}^R &\equiv \frac{1}{2\pi} \int_0^{2\pi} d\tilde{\phi}_1^{(\pm)} \frac{R_{\pm\mp}^R}{|A(0, 1/2)|^2} \\ &= (\rho_{\mp\pm}^R)^* \end{aligned} \quad (28)$$

$$\rho_{+-}^R = -\rho_{-+} (r_a \rightarrow r_a^R, \beta_a \rightarrow -\beta_a^R) \quad (29)$$

and

$$\begin{aligned} \bar{\rho}_{\pm\mp}^L &\equiv \frac{1}{2\pi} \int_0^{2\pi} d\tilde{\phi}_2^{(\pm)} \frac{\bar{r}_{\pm\mp}^L}{|B(0, -1/2)|^2} \\ &= (\bar{\rho}_{\mp\pm}^L)^* \end{aligned} \quad (30)$$

$$\bar{\rho}_{+-}^L = -\bar{\rho}_{-+} (r_b \rightarrow r_b^L, \beta_b \rightarrow -\beta_b^L) . \quad (31)$$

See Ref. 1 for the definitions of ρ_{+-} and $\bar{\rho}_{+-}$.

Additional $\nu_R / \bar{\nu}_L$ Tests for CP Violation

There are two tests of "non-CKM" type leptonic CP violation if R-handed ν (and L-handed $\bar{\nu}$) exist:

$$\beta_a^R = \beta_b^L \quad (\text{1st } v_R / \bar{v}_L \text{ test})$$

$$r_a^R = r_b^L \quad (\text{2nd } v_R / \bar{v}_L \text{ test})$$

where the phase differences are defined by Eqs. (10b, 15b) and the moduli ratios by Eqs. (25, 27).

In the case of both $(V \mp A)$ couplings and possibly $m_\nu \neq 0$, the $\tau^- \rightarrow \rho^- \nu$ amplitudes for $\lambda_\nu = -1/2$ are

$$A(0, -1/2) = g_L \left(\frac{E_\rho + q_\rho}{m_\rho} \right) \sqrt{m_\tau (E_\nu + q_\rho)} - g_R \left(\frac{E_\rho - q_\rho}{m_\rho} \right) \sqrt{m_\tau (E_\nu - q_\rho)} \quad (32a)$$

$$A(-1, -1/2) = g_L \sqrt{2m_\tau (E_\nu + q_\rho)} - g_R \sqrt{2m_\tau (E_\nu - q_\rho)} \quad (32b)$$

For $\lambda_\nu = 1/2$ they are

$$A(-1, 1/2) = 0$$

$$A(0, 1/2) = -g_L \left(\frac{E_\rho - q_\rho}{m_\rho} \right) \sqrt{m_\tau (E_\nu - q_\rho)} - g_R \left(\frac{E_\rho + q_\rho}{m_\rho} \right) \sqrt{m_\tau (E_\nu + q_\rho)} \quad (33a)$$

$$A(1, 1/2) = -g_L \sqrt{2m_\tau (E_\nu - q_\rho)} - g_R \sqrt{2m_\tau (E_\nu + q_\rho)} \quad (33b)$$

Note that $g_{L,R}$ respectively denote the chirality $(V \mp A)$ of the $\tau^- \rightarrow \rho^- \nu$ coupling whereas $\lambda_\nu = \mp 1/2$ denotes the handedness of the (massive) tau neutrino.

Formulas for $\tau^- \rightarrow a_1^- \nu$ including both $V \pm A$, and both ν helicities.

First, note that in kinematically describing the $\tau^- \rightarrow a_1^- \nu \rightarrow (\pi_1^- \pi_2^- \pi_3^+)$ mode, one can use the normal to the $(\pi_1^- \pi_2^- \pi_3^+)$ decay triangle in place of the π^- momentum direction of $\rho^- \rightarrow \pi_1^- \pi_2^0$ of the $\tau^- \rightarrow \rho^- \nu$ decay mode. Then, the various S2SC functions given above still hold, Eqs. (17) and (19).

Including both v_L and v_R helicities, we find composite decay density matrices for the $\tau^- \rightarrow a_1^- \nu \rightarrow (\pi_1^- \pi_2^- \pi_3^+) \nu$ decay sequence⁵

$$\mathbf{R}^\nu = S_1^+ \mathbf{R}^+ + S_1^- \mathbf{R}^- \quad (34)$$

where \mathbf{R}^\pm have the same form as Eq. (12) except the elements have " \pm " superscripts (see below). S_1^\pm describe $a_1^- \rightarrow \pi_1^- \pi_2^- \pi_3^+$. When the 3-body Dalitz plot is integrated over, only the S_1^+ term remains. In Eq. (34), the \mathbf{R}^+ matrix elements are

$$\mathbf{R}_{\pm\pm}^+ = \{\text{Eq. (8) except } (\frac{1}{\sqrt{2}}) \rightarrow (-\frac{1}{\sqrt{2}})\} \quad (35a)$$

$$\begin{aligned} \mathbf{r}_{\pm\pm}^+ &= (\mathbf{r}_{\mp\mp}^+)^* \\ &= \{\text{Eq. (9) except } (\frac{1}{\sqrt{2}}) \rightarrow (-\frac{1}{\sqrt{2}})\} \end{aligned} \quad (35b)$$

with

$$\begin{pmatrix} \mathbf{n}_a \\ \mathbf{n}_a \mathbf{f}_a \end{pmatrix} = \sin^2 \tilde{\theta}_a (|\mathbf{A}(0, -1/2)\rangle^2 \pm |\mathbf{A}(0, 1/2)\rangle^2) \pm (1 - \frac{1}{2} \sin^2 \tilde{\theta}_a) (|\mathbf{A}(-1, -1/2)\rangle^2 \pm |\mathbf{A}(1, 1/2)\rangle^2) \quad (36)$$

Similarly, the \mathbf{R}^- matrix elements are

$$\begin{aligned} \mathbf{R}_{\pm\pm}^- &= -\mathbf{n}_a^- (1 \mp \frac{\mathbf{f}_a^-}{\lambda} \cos \theta_q^\tau) \\ &\mp \sqrt{2} \sin \theta_1^\tau \sin \tilde{\theta}_a [\cos(\tilde{\phi}_a - \beta_a) |\mathbf{A}(0, -1/2)\rangle |\mathbf{A}(-1, -1/2)\rangle \\ &+ \cos(\tilde{\phi}_a + \beta_a^R) |\mathbf{A}(0, 1/2)\rangle |\mathbf{A}(1, 1/2)\rangle] \end{aligned} \quad (37a)$$

with

$$\begin{pmatrix} \mathbf{n}_a^- \\ \mathbf{n}_a^- \mathbf{f}_a^- \end{pmatrix} = \cos \tilde{\theta}_a (|\mathbf{A}(-1, -1/2)\rangle^2 \mp |\mathbf{A}(1, 1/2)\rangle^2), \quad (37b)$$

and

$$\begin{aligned} \mathbf{r}_{\pm\pm}^- &= (\mathbf{r}_{\mp\mp}^-)^* \\ &= \sin \theta_1^\tau \cos \tilde{\theta}_a (|\mathbf{A}(-1, -1/2)\rangle^2 + |\mathbf{A}(1, 1/2)\rangle^2) \\ &+ \sqrt{2} \sin \tilde{\theta}_a \{[\cos \theta_1^\tau \cos(\tilde{\phi}_a - \beta_a) + i \sin(\tilde{\phi}_a - \beta_a)] |\mathbf{A}(0, -1/2)\rangle |\mathbf{A}(-1, -1/2)\rangle \\ &+ [\cos \theta_1^\tau \cos(\tilde{\phi}_a + \beta_a^R) + i \sin(\tilde{\phi}_a + \beta_a^R)] |\mathbf{A}(0, 1/2)\rangle |\mathbf{A}(1, 1/2)\rangle\} \end{aligned} \quad (38)$$

For the conjugate decay sequence, $\tau^+ \rightarrow a_1^+ \bar{\nu} \rightarrow (\pi_1^+ \pi_2^+ \pi_3^-) \bar{\nu}$,

$$\bar{\mathbf{R}}^{\bar{\nu}} = \bar{\mathbf{S}}_1^+ \bar{\mathbf{R}}^+ + \bar{\mathbf{S}}_2^- \bar{\mathbf{R}}^- \quad (39)$$

The $\bar{\mathbf{R}}^+$ matrix elements (see Eq. (12)) are

$$\bar{\mathbf{R}}_{\pm\pm}^+ = \{ \text{Eq. (13) except } (\frac{1}{\sqrt{2}}) \rightarrow (-\frac{1}{\sqrt{2}}) \} \quad (40a)$$

$$\begin{aligned} \bar{\mathbf{r}}_{+..}^+ &= (\bar{\mathbf{r}}_{-..}^+)^* \\ &= \{ \text{Eq. (14) except } (\frac{1}{\sqrt{2}}) \rightarrow (-\frac{1}{\sqrt{2}}) \} \end{aligned} \quad (40b)$$

with

$$\begin{pmatrix} \mathbf{n}_b \\ \mathbf{n}_b \mathbf{f}_b \end{pmatrix} = \sin^2 \tilde{\theta}_b (|\mathbf{B}(0, 1/2)\rangle^2 \pm |\mathbf{B}(0, -1/2)\rangle^2) \pm (1 - \frac{1}{2} \sin^2 \tilde{\theta}_b) (|\mathbf{B}(1, 1/2)\rangle^2 \pm |\mathbf{B}(-1, -1/2)\rangle^2) \quad (41)$$

The $\bar{\mathbf{R}}^-$ matrix elements are

$$\begin{aligned} \bar{\mathbf{R}}_{\pm\pm}^- &= \mathbf{n}_b^- (1 \pm \frac{\mathbf{f}_b^-}{\hat{\mathbf{L}}} \cos \theta_2^\tau) \\ &\mp \sqrt{2} \sin \theta_2^\tau \sin \tilde{\theta}_b [\cos (\tilde{\phi}_b + \beta_b) |\mathbf{B}(0, 1/2)\rangle |\mathbf{B}(1, 1/2)\rangle \\ &+ \cos (\tilde{\phi}_b - \beta_b^L) |\mathbf{B}(0, -1/2)\rangle |\mathbf{B}(-1, -1/2)\rangle] \end{aligned} \quad (42a)$$

with

$$\begin{pmatrix} \mathbf{n}_b^- \\ \mathbf{n}_b \mathbf{f}_b^- \end{pmatrix} = \cos \tilde{\theta}_b (|\mathbf{B}(1, 1/2)\rangle^2 \mp |\mathbf{B}(-1, -1/2)\rangle^2), \quad (42b)$$

$$\begin{aligned} \bar{\mathbf{r}}_{+..}^- &= (\bar{\mathbf{r}}_{-..}^-)^* \\ &= \sin \theta_2^\tau \cos \tilde{\theta}_b (|\mathbf{B}(1, 1/2)\rangle^2 + |\mathbf{B}(-1, -1/2)\rangle^2) \\ &+ \sqrt{2} \sin \tilde{\theta}_b \{ [\cos \theta_2^\tau \cos (\tilde{\phi}_b + \beta_b) + i \sin (\tilde{\phi}_b + \beta_b)] |\mathbf{B}(0, 1/2)\rangle |\mathbf{B}(1, 1/2)\rangle \\ &+ [\cos \theta_2^\tau \cos (\tilde{\phi}_b - \beta_b^L) + i \sin (\tilde{\phi}_b - \beta_b^L)] |\mathbf{B}(0, -1/2)\rangle |\mathbf{B}(-1, -1/2)\rangle \} \end{aligned} \quad (43)$$

Ideal Statistical Errors

$\tau \rightarrow \rho^- \nu$ mode with L-handed ν :

Tables 1, 2 and 3 list the ideal statistical errors⁶ of Ref. 1 for the CP and $\tilde{\mathbf{T}}_{\text{FS}}$ discrete symmetry tests. Here $\tilde{\mathbf{T}}_{\text{FS}}$ is the approximate time-reversal-operation which holds only if possible final-state-interactions are

neglected. Such effects are indeed negligible in the usual V-A, $m_{\nu\tau} = 0$ lepton model. By \tilde{T}_{FS} the decay amplitude (A or B above) is purely real.

See conference contribution ICHEP-0099 for further discussion^{1,7} these statistical errors.

$\tau \rightarrow a^- \nu$ mode with L-handed ν :

Tables 4, 5, 6 list the analogous ideal statistical errors for the CP and \tilde{T}_{FS} discrete symmetry tests in the case of the $\tau \rightarrow a_1 \nu$ decay mode.

Improvement from measurement of τ^- momentum direction:

As discussed above, by use of a silicon vertex detector it may be possible to uniquely determine the τ^- momentum direction. Table 7 shows the improvement for the 2 tests for "non-CKM" type CP-violation in $\tau \rightarrow \rho \nu$ decay.

Conclusions About Ideal Statistical Errors

At the Z^0 , 10^7 Z^0 's are assumed, and at each γ^* energy we assumed 10^7 $\tau^- \tau^+$ pairs. Notice that in the measurement of the phase differences at γ^* energies, versus at the Z^0 , there is not as much improvement as would be expected due to the increase in the number of events. This is because in using ρ -polarimetry (or a_1 -polarimetry) a Wigner-rotation is involved in going from the center of mass frame's ρ -observables (or a_1 -observables) to the respective τ rest frame's ρ -observables (or a_1 -observables). For instance, see Tables 3 and 6.

τ spin correlations are necessary to measure β_a at γ^* energies; at the Z^0 , without using spin-correlations there would be an extra suppression factor of $\langle P_\tau \rangle = -0.138$.

Since the direction of the initial e^- beam has been integrated out, there is no obvious source for a violation of \hat{T}_{FS} invariance for the S2SC processes considered here. For instance, unlike in $K_{\ell 3}$ decays, since ν_τ is only weakly interacting there is no "old physics" source for electromagnetic rescattering of the ν_τ and the ρ^- (or a_1^-).

The tables for the a_1 decay modes show approximately the same patterns as those for the ρ decay modes obtained earlier in Ref. 1 and shown here in the first three tables. However, the net sensitivities differ---the sensitivity for the $\beta_a = \beta_b$ test is about 10 times worse in the a_1 mode, but the **normalized** sensitivity is about the same for the $r_a = r_b$ test for both the ρ mode and for the a_1 mode. "Normalized sensitivity" refers to the value of the fractional error $\{ \sigma(r_a) / r_a \}$.

For measurement of β_a at γ^* energies, knowledge of the τ momentum direction improves the sensitivity by about a factor of $(1/\sqrt{2} = 0.707)$ which is what would be expected by statistics. However, there is a small (about 10%) improvement in the measurement of r_a by measurement of the τ momentum direction.

In conclusion, at γ^* energies one can perform the 1st test, $\beta_a = \beta_b$, to about the 0.5% level, and the 2nd test, $r_a = r_b$, to about the 0.1% level by the ρ decay mode. For the a_1 , the sensitivity for the 1st test is about 10 times worse, but is about the same for the 2nd test.

Acknowledgment

One of us (C.A.N.) thanks the members of the theory and experimental groups at Argonne, Cornell, and Fermilab for their hospitality and intellectual stimulation. This work was partially supported by the U.S. Department of Energy, Contract No. DE-FG02, 96ER40291.

References

1. C.A. Nelson, et. al., "Stage 2 Spin-Correlation Functions: Tests for Non-CKM Type Leptonic CP Violation in $\tau \rightarrow \rho\nu$ Decay," SUNY BING 8/16/93, Phys. Rev. D. (to appear); SUNY BING 7/19/92; and in "Second Workshop on Tau Lepton Physics," p. 353, ed. K.K. Gan (World Sci., 1993).
2. Work on polarimetry and spin-correlation methods in tau lepton processes includes Y.-S. Tsai, Phys. Rev. D4, 2821 (1971); S.Y. Pi and A.I. Sanda, Ann. Phys. (N.Y.) 106, 171 (1977); J. Babson and E. Ma, Phys. Rev. D26, 2497 (1982); H. Kühn and F. Wagner, Nucl. Phys. B236, 16 (1984); C.A. Nelson, Phys. Rev. Lett. 62, 1347 (1989); Phys. Rev. D40, 123 (1989); 41, 2327 (E) (1990); W. Fetscher, Phys. Rev. D43, 1544 (1990); A. Rouge, Z. Phys. C48, 75 (1990); K. Hagiwara, A.D. Martin, and D. Zeppenfeld, Phys. Lett. B235, 198 (1990); W. Bernreuther and O. Nachtmann, Phys. Rev. Lett. 63, 2787 (1989); C.A. Nelson, Phys. Rev. D41, 2805 (1990); D43, 1465 (1991); S. Goozovat and C.A. Nelson, Phys. Lett. B267, 128 (1991); Phys. Rev. D44, 2818 (1991); J. Bernabeu, N. Rius and A. Pich, Phys. Lett. B257, 219 (1991); W. Bernreuther, G.W. Botz, O. Nachtmann and P. Overmann, Z. Phys. C52, 567 (1991); B.K. Bullock, K. Hagiwara and A.D. Martin, Nucl. Phys. B395, 499 (1993); H. Thurn and H. Kolanoski, DESY 93-056, and B. Ananthanarayan and S.D. Rindani, Lusanne/Ahmedabad report (1993).
3. Spin correlation and polarimetry effects are included in the KORALB Monte Carlo for γ^* energies, see S. Jadach and Z. Was, Comput. Phys. Commun. 64, 267 (1991) and June 1993 update.
4. See, e.g., C.A. Nelson, Phys. Rev. D43, 1465 (1991).
5. Work on the $\tau \rightarrow a_1\nu$ decay mode includes the measurements of V-A versus V+A coupling (the chiral polarization parameter) by the ARGUS collaboration, Phys. Lett. B250, 164 (1990). See also M. Feindt, Z. Phys. C48, 681 (1990); J.H. Kuhn and A. Santamaria, Z. Phys. C48, 445 (1990); N.A. Tornquist, HU-TFT-92-50; and P. Privitera, IEKP-KA/93-02, /93-01.
6. We calculate ideal statistical errors as in C.A. Nelson, Phys. Rev. D40, 123 (1989). For instance, for a 2-variable correlation we would distribute N events ideally over a 2-dimensional i_j -grid according to the theoretical result, $I(x, y) = Z_0(x, y) + a Z_1(x, y)$; The ideal error in bin i_j is $\sigma_{i_j} = \sqrt{I(x_i, y_j)}$. By χ^2 minimization, the "ideal statistical error" in the measurement of "a" is $\sigma_a = \{ \sum_{i_j} [Z_1(x_i, y_j) / \sigma_{i_j}]^2 \}^{1/2}$.
7. C.A. Nelson, SUNY BING 4/30/94 and ICHEP-0099.

Table Captions

Table 1: At $E_{\text{cm}} = M_Z$, ideal statistical errors for two tests for CP violation in $\tau \rightarrow \rho\nu$ by the simpler S2SC function $I(E_1, E_2, \tilde{\theta}_1, \tilde{\theta}_2)$, see Eq. (21), for the sequential decay $Z^0 \rightarrow \tau^-\tau^+$ with $\tau^- \rightarrow \rho^-\nu \rightarrow (\pi^-\pi^0)\nu$ and $\tau^+ \rightarrow \rho^+\bar{\nu}, \pi^+\bar{\nu}, \text{ or } \ell^+ \nu_\ell \bar{\nu}_\tau$. We use $10^7 Z^0$ events.

Table 2: At $E_{\text{cm}} = 10 \text{ GeV}$ and 4 GeV respectively, ideal statistical errors for two tests for CP violation in $\tau \rightarrow \rho\nu$ by the simpler S2SC function, Eq. (21), for the decay of an off-mass-shell photon $\gamma^* \rightarrow \tau^-\tau^+$ with $\tau^- \rightarrow \rho^-\nu \rightarrow (\pi^-\pi^0)\nu$, and $\tau^+ \rightarrow \rho^+\bar{\nu}, \pi^+\bar{\nu}, \text{ or } \ell^+ \nu_\ell \bar{\nu}_\tau$. We use $10^7 \gamma^* \rightarrow \tau^-\tau^+$ events.

Table 3: Ideal statistical errors for CP/T violation tests based on the full S2SC function of Eq. (18) for the $\{\rho^-\rho^+\}$ sequential decay mode. Note that $\tilde{\beta} \equiv \beta_a - \beta_b$ and $\beta' \equiv \beta_a + \beta_b$.

Table 4: Ideal statistical errors for CP tests for $\tau^- \rightarrow a_1^-\nu \rightarrow (\pi_1^-\pi_2^-\pi_0^+)\nu$ at Z^0 from the simpler $I(E_1, E_2, \tilde{\theta}_1, \tilde{\theta}_2)$.

Table 5: Same as Table 4 except at $E_{\text{cm}} = 10 \text{ GeV}$ and 4 GeV .

Table 6: Ideal statistical errors for CP/T violation tests based on full S2SC for $\{a_1^-, a_1^+\}$ sequential decay mode.

Table 7: Percentage improvement for tests for $\tau \rightarrow \rho\nu$ mode (compare Table 3) when τ^- direction is known, e.g. via silicon vertex detector.

TABLE 1

$E_{cm} = M_Z$ Mode	Number of events.	Ideal statistical errors	
		$\sigma(r_a)$	$\sigma(\beta_a^2)$
$\{\rho^- \rho^+\}$	20,302	0.0065	$(12^\circ)^2$
$\{\rho^- \pi^+\}$	9,847	0.0091	$(12^\circ)^2$
$\{\rho^- \ell^+\}$	29,074	0.0056	$(15^\circ)^2$
Sum of above modes	59,223	0.0039 [0.6%]	$(10^\circ)^2$

TABLE 2

Mode	Number of events.	$E_{cm} = 10 \text{ GeV}$		$E_{cm} = 4 \text{ GeV}$	
		$\sigma(r_a)$	$\sigma(\beta_a^2)$	$\sigma(r_a)$	$\sigma(\beta_a^2)$
$\{\rho^- \rho^+\}$	605,127	.0012	$(5.5^\circ)^2$.0011	$(8.8^\circ)^2$
$\{\rho^- \pi^+\}$	293,527	.0017	$(5.9^\circ)^2$.0016	$(9.1^\circ)^2$
$\{\rho^- \ell^+\}$	866,658	.0010	$(7.5^\circ)^2$.0010	$(11.5^\circ)^2$
Sum of above modes	1,765,312	.0007 [0.1%]	$(4.7^\circ)^2$.0007 [0.1%]	$(7.3^\circ)^2$

TABLE 3

E_{cm}	Number of $\{\rho^- \rho^+\}$ events	Ideal Statistical Errors		
		$\sigma(\tilde{\beta})$	$\sigma(\beta')$	$\sigma(\beta_a)$
M_Z	20,302	1.88°	3.15°	1.84°
10 GeV	605,127	0.43°	0.74°	0.42°
4 GeV	605,127	0.86°	1.13°	0.71°

TABLE 4

$E_{cm} = M_Z$ Mode	Number of events.	Ideal statistical errors	
		$\sigma(r_a)$	$\sigma(\beta_a^2)$
$\{a_1^- a_1^+\}$	2,718	0.019	$(41^\circ)^2$
$\{a_1^- \rho^+\}$	7,428	0.011	$(22^\circ)^2$
$\{a_1^- \pi^+\}$	3,603	0.016	$(24^\circ)^2$
$\{a_1^- \ell^+\}$	10,638	0.009	$(29^\circ)^2$
Sum of above modes	24,387	0.0062 [0.6%]	$(18^\circ)^2$

TABLE 5

Mode	Number of events.	$E_{cm} = 10 \text{ GeV}$		$E_{cm} = 4 \text{ GeV}$	
		$\sigma(r_a)$	$\sigma(\beta_a^2)$	$\sigma(r_a)$	$\sigma(\beta_a^2)$
$\{a_1^- a_1^+\}$	81,000	.0035	$(21^\circ)^2$.0035	$(26^\circ)^2$
$\{a_1^- \rho^+\}$	221,400	.0021	$(10^\circ)^2$.0021	$(15^\circ)^2$
$\{a_1^- \pi^+\}$	107,388	.0030	$(11^\circ)^2$.0030	$(15^\circ)^2$
$\{a_1^- \ell^+\}$	317,070	.0018	$(14^\circ)^2$.0018	$(19^\circ)^2$
Sum of above modes	726,858	.0011 [0.1%]	$(5^\circ)^2$.0012 [0.1%]	$(12^\circ)^2$

TABLE 6

E_{cm}	Number of $\{a_1^- a_2^+\}$ events	Ideal Statistical Errors		
		$\sigma(\tilde{\beta})$	$\sigma(\beta')$	$\sigma(\beta_a)$
M_Z	2,718	20°	32°	17°
10 GeV	81,000	4°	6°	5°
4 GeV	81,000	8°	10°	8°

TABLE 7

E_{cm}	Number of { $\rho^- \rho^+$ } events	<u>Percentage Improvement by τ^- Direction</u>	
		$\sigma(r_a)$	$\sigma(\beta_a)$
10 GeV	605,127	7%	27%
4 GeV	605,127	12%	26%