

March 12, 2021

Spectroscopy in beauty decays at the LHCb experiment

Tatiana Ovsianikova[†] on behalf of the LHCb collaboration
*Institute for Theoretical and Experimental Physics, NRC Kurchatov Institute,
B. Cheremushkinskaya 25, Moscow, 117218, Russia.*

Abstract

The beauty hadron decays is unique laboratory to study charmonium and charmonium-like states, such as the $\chi_{c1}(3872)$ meson, other exotic states and the tensor D -wave $\psi_2(3823)$ states. However the nature of many exotic charmonium-like candidates are still unknown. The most recent LHCb results related to b-hadron decays to charmonium states and obtained using large data samples collected during the Run 1 and Run 2 periods are presented. This includes the most precise determination of the mass and width of the $\chi_{c1}(3872)$ state using the $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$ decays, observation of a resonant structure denoted as $X(4740)$ in the $J/\psi \phi$ mass spectrum from $B_s^0 \rightarrow J/\psi \pi^+ \pi^- K^+ K^-$ decays and the precise measurement of the B_s^0 meson mass.

Presented at XXVII Cracow EPIPHANY Conference on future of particle physics

[†]E-mail: Tatiana.Ovsianikova@cern.ch

1 Introduction

In the last two decades a plethora of new results in the charmonium spectra have been obtained in the beauty decays studies. A lot of the conventional and exotic charmonium resonances are observed such as $\chi_{c1}(3872)$, $\chi_{c1}(4700)$ and $P_c(4312)^+$ and conventional $\psi_2(3823)$ state. The LHCb experiment has collected high statistics during Run 1 and Run 2 periods that allows us to perform many precise measurements of the branching fractions of B- and B_s^0 -meson decays and searches for new decays and states. The results described below are based on the data samples collected by the LHCb experiment in proton-proton (pp) collisions at the Large Hadron Collider from 2011 to 2018 with centre-of-mass energies of $\sqrt{s} = 7, 8$ and 13 TeV.

2 Study of the $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+ K^-$ decays

Candidates of the $B_s^0 \rightarrow J/\psi \pi^+ \pi^- K^+ K^-$ decays are reconstructed via $J/\psi \rightarrow \mu^- \mu^+$ and selected using based on kinematics, particle identification and topology [1]. The yields of $B_s^0 \rightarrow J/\psi \pi^+ \pi^- K^+ K^-$ decays via the $B_s^0 \rightarrow \psi(2S)\phi$ and $B_s^0 \rightarrow \chi_{c1}(3872)\phi$ and $B_s^0 \rightarrow J/\psi K^{*0} \bar{K}^{*0}$ chains are determined using three-dimensional unbinned extended maximum-likelihood fits. The observed signal yield for the $B_s^0 \rightarrow \chi_{c1}(3872)\phi$ decays is 154 ± 15 which corresponds to a statistical significance more than 10 standard deviations. The fit to the mass distribution for the signal channel is shown in figure 1.




Figure 1: Distributions of the (left) $J/\psi \pi^+ \pi^- K^+ K^-$ and (right) $J/\psi \pi^+ \pi^-$ mass for selected $B_s^0 \rightarrow \chi_{c1}(3872)\phi$ candidates (points with error bars) [1]. The red filled area corresponds to the $B_s^0 \rightarrow \chi_{c1}(3872)\phi$ signal. The orange line is the total fit.

In addition, the decays $B_s^0 \rightarrow \chi_{c1}(3872)K^+ K^-$ where the $K^+ K^-$ pair does not originate from a ϕ meson, is studied using a two-dimensional unbinned extended maximum-likelihood fit which is performed to corresponding mass distributions. The observed yield of signal decays is 378 ± 33 , that is significantly larger than the yield of the $B_s^0 \rightarrow \chi_{c1}(3872)\phi$ decays, indicating a significant $B_s^0 \rightarrow \chi_{c1}(3872)K^+ K^-$ contribution. A narrow ϕ component can be separated from the non- ϕ components using an unbinned maximum-likelihood fit to the background-subtracted and efficiency-corrected $K^+ K^-$ mass distribution. The fraction of the $B_s^0 \rightarrow \chi_{c1}(3872)K^+ K^-$ signal component is found to be $(38.9 \pm 4.9)\%$. Using the obtained signal yields and fractions for described channels and corresponding efficiency




Figure 2: Background-subtracted K^+K^- mass distribution for selected $B_s^0 \rightarrow \chi_{c1}(3872)K^+K^-$ candidates (points with error bars) [1]. The orange line is the total fit.

ratios the following branching fractions are calculated:

$$\frac{\mathcal{B}_{B_s^0 \rightarrow \chi_{c1}(3872)\phi} \times \mathcal{B}_{\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-}}{\mathcal{B}_{B_s^0 \rightarrow \psi(2S)\phi} \times \mathcal{B}_{\psi(2S) \rightarrow J/\psi \pi^+ \pi^-}} = (2.42 \pm 0.23 \pm 0.07) \times 10^{-2},$$

$$\frac{\mathcal{B}_{B_s^0 \rightarrow J/\psi K^{*0} \bar{K}^{*0}} \times \mathcal{B}_{K^{*0} \rightarrow K^+ \pi^-}}{\mathcal{B}_{B_s^0 \rightarrow \psi(2S)\phi} \times \mathcal{B}_{\psi(2S) \rightarrow J/\psi \pi^+ \pi^-} \times \mathcal{B}_{\phi \rightarrow K^+ K^-}} = 1.22 \pm 0.03 \pm 0.04,$$

$$\frac{\mathcal{B}_{B_s^0 \rightarrow \chi_{c1}(3872)(K^+ K^-)_{\text{non-}\phi}}}{\mathcal{B}_{B_s^0 \rightarrow \chi_{c1}(3872)\phi} \times \mathcal{B}_{\phi \rightarrow K^+ K^-}} = 1.57 \pm 0.32 \pm 0.12,$$

where the first uncertainty is statistical and the second is systematic. The result for $B_s^0 \rightarrow \chi_{c1}(3872)\phi$ decay is found to be in a good agreement with the result by the CMS collaboration [2] but is more precise.

Four tetraquark candidates have been observed by the LHCb collaboration using an amplitude analysis of the $B^+ \rightarrow J/\psi \phi K^+$ decays [3, 4]. A search of the exotic states in the $J/\psi \phi$ spectrum is performed using the $B_s^0 \rightarrow J/\psi \pi^+ \pi^- \phi$ decays. The $B_s^0 \rightarrow J/\psi \pi^+ \pi^- \phi$ candidates are determined with two-dimensional unbinned extended maximum-likelihood fit to the $J/\psi \pi^+ \pi^- K^+ K^-$ and $K^+ K^-$ mass distributions.

The background-subtracted $J/\psi \phi$ mass spectrum of $B_s^0 \rightarrow J/\psi \pi^+ \pi^- \phi$ candidates are shown in figure 3. It shows a prominent structure at a mass around $4.74 \text{ GeV}/c^2$. Since the regions of $\psi(2S)$ and $\chi_{c1}(3872)$ resonance masses are vetoed and no sizeable contributions from decays via other narrow charmonium states are observed in the background-subtracted $J/\psi \pi^+ \pi^-$ mass spectrum, this structure cannot be explained by cross-feed from the $J/\psi \pi^+ \pi^-$ mass spectrum. Moreover no such structure is seen in non- ϕ region of the $K^+ K^-$ mass. However the $\phi \pi^+ \pi^-$ spectrum exhibits significant deviations from the phase-space distribution, indicating possible presence of excited ϕ states, referred to as ϕ^* states hereafter. The decays $B_s^0 \rightarrow J/\psi \phi^*$ via intermediate $\phi(1680)$, $\phi(1850)$ or $\phi(2170)$ states [5] are studied using simulated samples and no peaking structures are observed. Under the assumption that the observed structure, referred to as $X(4740)$




Figure 3: Background-subtracted $J/\psi \phi$ mass distribution for the selected $B_s^0 \rightarrow J/\psi \pi^+ \pi^- \phi$ signal candidates (points with error bars) [1]. The red filled area corresponds to the $B_s^0 \rightarrow X(4740)\pi^+ \pi^-$ signal. The orange line is the total fit.

hereafter, has a resonant nature, its mass and width are determined through an unbinned extended maximum-likelihood fit. The fit result is superimposed in figure 3. The obtained signal yield is 175 ± 39 and corresponds to a statistical significance above 5.3 standard deviations. The mass and width for the $X(4740)$ state are found to be

$$\begin{aligned} m_{X(4740)} &= 4741 \pm 6 \pm 6 \text{ GeV}/c^2, \\ \Gamma_{X(4740)} &= 53 \pm 15 \pm 11 \text{ MeV}. \end{aligned}$$

The observed parameters qualitatively agree with those of the $\chi_{c1}(4700)$ state observed by the LHCb collaboration in references [3, 4]. The obtained mass also agrees with the one expected for the $2^{++} c\bar{s}\bar{s}$ tetraquark state [6].

The B_s^0 decays to the $\psi(2S)K^+K^-$ final states characterize the relatively small energy release allowing precise measurement of the B_s^0 meson mass. The mass of the B_s^0 meson is determined from an unbinned extended maximum-likelihood fit to the $\psi(2S)K^+K^-$ mass distribution. The improvement in the B_s^0 mass resolution and significant decrease of the systematic uncertainties is achieved by imposing a constraint on the reconstructed mass of the $J/\psi \pi^+ \pi^-$ system to the known $\psi(2S)$ meson mass [5]. The measured value of the B_s^0 meson mass is found to be

$$m_{B_s^0} = 5366.98 \pm 0.07 \pm 0.13 \text{ MeV}/c^2,$$

that is the most precise single measurement of this quantity.

3 Study of the $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$ decays

The search of the spin-2 component of the D -wave charmonium triplet, the $\psi_2(3823)$ state, is performed with $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$ decays [7, 8]. To extract the B^+ candidates, a

multivariate classifier algorithm based on a decision tree with gradient boosting is applied. For signal yield determinations of the $B^+ \rightarrow (\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) K^+$, $B^+ \rightarrow (\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-) K^+$ and $B^+ \rightarrow (\psi_2(3823) \rightarrow J/\psi \pi^+ \pi^-) K^+$, a simultaneous unbinned extended maximum-likelihood fit to the $m_{J/\psi \pi^+ \pi^- K^+}$ and $m_{J/\psi \pi^+ \pi^-}$ variables is performed. The signal yield for the $B^+ \rightarrow \psi_2(3823) K^+$ decays is determined to be 137 ± 26 which correspond to statistical significance above 5.1 standard deviations. Large signal yield for the $B^+ \rightarrow \psi(2S) K^+$ signal, 4230 ± 70 , allows for the precise measurement of the mass and width of the $\chi_{c1}(3872)$ state. For the first time the non-zero Breit–Wigner width is observed for the $\chi_{c1}(3872)$ state with significance more than 5 standard deviations and its measured value is:

$$\Gamma_{\chi_{c1}(3872)} = 0.96_{-0.18}^{+0.19} \pm 0.21 \text{ MeV}.$$

The upper limit for the Breit–Wigner width of $\psi_2(3823)$ is improved and its value is set to be $\Gamma_{\psi_2(3823)} < 5.2$ (6.6) MeV, for 90 (95)% C.L. The mass splitting between the states are found to be

$$\begin{aligned} \delta m_{\psi_2(3823)}^{\chi_{c1}(3872)} &= 47.50 \pm 0.53 \pm 0.13 \text{ MeV}/c^2, \\ \delta m_{\psi(2S)}^{\psi_2(3823)} &= 137.98 \pm 0.53 \pm 0.14 \text{ MeV}/c^2, \\ \delta m_{\psi(2S)}^{\psi_2(3823)} &= 185.49 \pm 0.06 \pm 0.03 \text{ MeV}/c^2, \end{aligned}$$

The results Breit–Wigner mass of the $\chi_{c1}(3872)$ state are in good agreement with an independent analysis of inclusive $b \rightarrow \chi_{c1}(3872) X$ decays [9]. The binding energy of the $\chi_{c1}(3872)$ state is derived from the mass splitting and its value is found to be $\delta E = 0.12 \pm 0.13$ MeV. It is consistent with zero within uncertainties, that are currently dominated by the uncertainty for the neutral and charged kaon mass measurements [10,11].

The measured yields of the $B^+ \rightarrow \chi_{c1}(3872) K^+$, $B^+ \rightarrow \psi_2(3823) K^+$ and $B^+ \rightarrow \psi(2S) K^+$ signal decays allow for a precise determination of the ratios of the branching fractions:

$$\begin{aligned} \frac{\mathcal{B}_{B^+ \rightarrow \psi_2(3823) K^+} \times \mathcal{B}_{\psi_2(3823) \rightarrow J/\psi \pi^+ \pi^-}}{\mathcal{B}_{B^+ \rightarrow \chi_{c1}(3872) K^+} \times \mathcal{B}_{\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-}} &= (3.56 \pm 0.67 \pm 0.11) \times 10^{-2}, \\ \frac{\mathcal{B}_{B^+ \rightarrow \psi_2(3823) K^+} \times \mathcal{B}_{\psi_2(3823) \rightarrow J/\psi \pi^+ \pi^-}}{\mathcal{B}_{B^+ \rightarrow \psi(2S) K^+} \times \mathcal{B}_{\psi(2S) \rightarrow J/\psi \pi^+ \pi^-}} &= (1.31 \pm 0.25 \pm 0.04) \times 10^{-3}, \\ \frac{\mathcal{B}_{B^+ \rightarrow \chi_{c1}(3872) K^+} \times \mathcal{B}_{\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-}}{\mathcal{B}_{B^+ \rightarrow \psi(2S) K^+} \times \mathcal{B}_{\psi(2S) \rightarrow J/\psi \pi^+ \pi^-}} &= (3.69 \pm 0.07 \pm 0.06) \times 10^{-2}. \end{aligned}$$

4 Conclusion

A study of b -meson decays $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$ and $B_s^0 \rightarrow J/\psi \pi^+ \pi^- K^+ K^-$ is made using the Run 1 and Run 2 data, collected with the LHCb detector [1, 7]. The reported results include the observation of the non-zero width of the $\chi_{c1}(3872)$ state; the most precise measurement of the masses of the $\chi_{c1}(3872)$ and $\psi_2(3823)$ states; the most precise measurement of several ratios of branching fractions of the B^+ and B_s^0 mesons decays; the most precise single measurement of the B_s^0 meson mass and the observation of a new structure, denoted as the X(4740) state, in the $J/\psi \phi$ mass spectrum.

References

- [1] LHCb collaboration, R. Aaij *et al.*, *Study of $B_s^0 \rightarrow J/\psi \pi^+ \pi^- K^+ K^-$ decays*, JHEP **2** (2021) 024 LHCb-PAPER-2020-035.
- [2] CMS collaboration, A. M. Sirunyan *et al.*, *Observation of the $B_s^0 \rightarrow X(3872)\phi$ decay*, Phys. Rev. Lett. **125** (2020) 152001, arXiv:2005.04764.
- [3] LHCb collaboration, R. Aaij *et al.*, *Amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ decays*, Phys. Rev. **D95** (2017) 012002, arXiv:1606.07898.
- [4] LHCb collaboration, R. Aaij *et al.*, *Observation of exotic $J/\psi \phi$ structures from amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ decays*, Phys. Rev. Lett. **118** (2017) 022003, arXiv:1606.07895.
- [5] Particle Data Group, P. A. Zyla *et al.*, *Review of particle physics*, Prog. Theor. Exp. Phys. **2020** (2020) 083C01.
- [6] D. Ebert, R. N. Faustov, and V. O. Galkin, *Excited heavy tetraquarks with hidden charm*, Eur. Phys. J. **C58** (2008) 399, arXiv:0808.3912.
- [7] LHCb collaboration, R. Aaij *et al.*, *Study of the $\psi_2(3823)$ and $\chi_{c1}(3872)$ states in $B^+ \rightarrow (J/\psi \pi^+ \pi^-) K^+$ decays*, JHEP **08** (2020) 123, arXiv:2005.13422.
- [8] D. Pereima, *Search for new decays of beauty particles in the LHCb experiment*, PhD thesis, NRC Kurchatov Institute – ITEP, Moscow, 2020, CERN-THESIS-2020-204.
- [9] LHCb collaboration, R. Aaij *et al.*, *Study of the line shape of the $\chi_{c1}(3872)$ meson*, arXiv:2005.13419, to appear in Phys. Rev. D.
- [10] A. Tomaradze *et al.*, *High precision measurement of the masses of the D^0 and K_S^0 mesons*, Phys. Rev. **D89** (2014) 031501.
- [11] LHCb collaboration, R. Aaij *et al.*, *Precision measurement of D meson mass differences*, JHEP **06** (2013) 065, arXiv:1304.6865.