

# Overview of the latest ALICE UPC and photonuclear results\*

SIMONE RAGONI, FOR THE ALICE COLLABORATION

Creighton University,  
2500 California Plz, Omaha,  
NE 68178, United States, USA  
`simone.ragoni@cern.ch`

*Received December 5, 2024*

Ultra-peripheral collisions (UPC) are events characterised by large impact parameters between the two projectiles, larger than the sum of their radii. In UPCs, the protons and ions accelerated by the LHC do not interact via the strong interaction and can be regarded as sources of quasireal photons. Using the Run 2 data, the ALICE Collaboration has carried out various measurements of different final-state systems, such as exclusive four pion photoproduction as well as photoproduction of  $K^+K^-$  pairs, measured for the first time in ultra-peripheral collisions. In addition, vector meson production in Pb–Pb provides the unique opportunity to carry out an analogy of the double-slit experiment at femtometre scales, owing to the interference between the production sources of the two lead nuclei. These results and prospects for UPC measurements using Run 3 data will be presented.

## 1. Introduction

Vector meson photoproduction in p–Pb and Pb–Pb ultra-peripheral collisions (UPCs) [1] is actively being studied at the CERN LHC. In these events, a photon from one of the two nuclei interacts with a colourless object from the other nucleus, resulting in the production of a vector meson. The ALICE Collaboration has studied  $J/\psi$  [2, 3],  $\psi'$ ,  $\rho^0$  [4],  $K^+K^-$  [5] and more light vector-meson photoproduction. The interest for these processes is growing since they shed light on nuclear shadowing, gluon saturation, and more recently gluonic hotspots, and their dependence on energy [6, 7].

---

\* Presented at “Diffraction and Low- $x$  2024”, Trabia (Palermo, Italy), September 8-14, 2024.

## 2. ALICE results using Run 2 (2015–2018) data

ALICE has recently presented results for exclusive charged  $4\pi$  photoproduction [8]. The  $4\pi$  invariant mass distribution is fitted using either a single Breit-Wigner or a combination of contributions representing excited  $\rho^0$  states, i.e.  $\rho(1450)$  and  $\rho(1700)$ . The cross sections are compared to theoretical models [9] and are shown in Fig. 1. The computation for the combination of  $\rho(1450)$  and  $\rho(1700)$  (upper panel) is in better agreement with the data. The exclusive  $K^+K^-$  photoproduction results [5] show




Fig. 1. Comparison of the  $\rho(1450)$  and  $\rho(1700)$  cross sections (upper panel), and single-resonance cross section (lower panel), as extracted from the fits to the invariant-mass distribution of exclusive  $4\pi$  production [8] and comparison with the theoretical predictions for a one and two-resonance model [9].

that the sample is a cocktail of resonant and non-resonant contributions, as shown in Fig. 2. The  $K^+K^-$  invariant mass distribution is measured for states above  $1.1 \text{ GeV}/c^2$ , away from the  $\phi(1020)$  peak, clear sign that the energy loss in the tracking material is too significant for the decay kaons from the  $\phi(1020)$  to be able to reach the Time Projection Chamber (TPC) using the Inner Tracking System (ITS) that was installed in ALICE during Run 1 and 2. A new ITS was installed during the Long Shutdown 2 [10], with reduced material budget and higher precision.

The ALICE Collaboration has also recently provided new results concerning the impact-parameter dependent azimuthal anisotropy in UPCs, which was measured in coherent  $\rho^0$  production [11]. The results are shown in Fig. 3. The amplitude  $a_2$  of the modulation increases as the impact parameter becomes smaller, which is achieved in UPCs by separating the data set in neutron emission classes [12]. From the  $0n0n$  to the  $XnXn$  class, the impact parameters lower from a median of about 49 fm to about 18 fm [11, 13].

### 3. ALICE future opportunities using Run 3 and 4 data

In Run 3 (2022-2026) and Run 4 (starting in 2030) ALICE will collect significantly higher amounts of data [14], also in previously inaccessible rapidity regions, owing to the installation of new detectors, i.e. Muon Forward Tracker (MFT) in Run 3 and Forward Calorimeter (FoCal) in Run 4. The effects of the introduction of a continuous readout, are particularly evident using UPC selections. While in Run 2 the sample contained about fifty thousand  $\pi^+\pi^-$  candidates in the invariant-mass region of the  $\rho^0$ , as shown in Fig. 4(a), which were used in [4] and [11], the data set collected in Run




Fig. 2. Cross sections for  $K^+K^-$  photoproduction as a function of the  $K^+K^-$  invariant mass as measured by ALICE [5]. The data are compatible with a cocktail of resonant and non-resonant contributions.




Fig. 3. Impact-parameter dependent azimuthal anisotropy measured in coherent  $\rho^0$  photoproduction [11]. The amplitude  $a_2$  of the modulation increases as the impact parameter becomes smaller.

3 is already order of magnitudes larger in a similar invariant-mass region with UPC selections, as shown in Fig. 4(b). More precise and even more (multi-)differential measurements are expected with the new data sample. In addition, the increased acceptance brought by the addition of MFT and




Fig. 4. Transverse momentum distribution of  $\pi\pi$  pairs selected in Run 2 (left panel) [4] and Run 3 (right panel), for UPC measurements in ALICE.

FoCal will allow access to observables that are expected to significantly contribute to observation of the onset of the gluon saturation regime, as described in [15]. FoCal, which will be installed during Long Shutdown 3, will provide sensitivity to charmonia through their decay to dielectrons, as shown in Fig. 5. This figure is an ALICE simulation produced using events from STARlight [16], where the  $J/\psi$  and  $\psi'$  peaks are clearly visible. Finally, the ALICE detector in Run 3 and 4 should be able to collect enough statistics to measure e.g. light-by-light scattering [17, 18],  $\gamma\gamma \rightarrow \gamma\gamma$ , and the anomalous magnetic moment in the tau sector [19],  $\gamma\gamma \rightarrow \tau\tau$ , allowing for measurements beyond the Standard Model with implications for axion-like particles (ALPs) [17] and SUSY [19]. New opportunities will also arise through the usage of machine learning in UPCs, such as anomaly detection through autoencoders for the detection of exotic hadrons such as tetraquarks [20] and pentaquarks, as described in [21].

#### 4. Conclusions

The UPC program in ALICE brought about interesting new results, which cover a large range of phenomena, such as nuclear shadowing, gluon saturation and gluonic hotspots. The most recent measurements were however limited by their statistical uncertainty. The new incoming Run 3 and 4 data sets will provide abundant amounts of high quality data, allowing




Fig. 5. ALICE simulations using STARlight [16] showing the potential of FoCal to measure photoproduced charmonia decaying into dielectrons.

for more differential measurements. These large data sets will also permit the measurement of the  $\gamma\gamma \rightarrow \gamma\gamma$  and  $\gamma\gamma \rightarrow \tau\tau$  processes in ALICE, giving prospects for beyond the Standard Model investigations. The new detectors, i.e. MFT and FoCal in Run 3 and 4, respectively, will in addition allow access to new and exciting rapidity regions.

## REFERENCES

- [1] A.J. Baltz, *The Physics of Ultraperipheral Collisions at the LHC*, *Phys. Rept.* **458** (2008) 1 [0706.3356].
- [2] ALICE collaboration, *Energy dependence of coherent photonuclear production of  $J/\psi$  mesons in ultra-peripheral Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV*, *JHEP* **10** (2023) 119 [2305.19060].
- [3] ALICE collaboration, *Exclusive and dissociative  $J/\psi$  photoproduction, and exclusive dimuon production, in p-Pb collisions at  $s_{NN}=8.16$  TeV*, *Phys. Rev. D* **108** (2023) 112004 [2304.12403].
- [4] ALICE collaboration, *Coherent photoproduction of  $\rho^0$  vector mesons in ultra-peripheral Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV*, *JHEP* **06** (2020) 035 [2002.10897].
- [5] ALICE collaboration, *Photoproduction of  $K^+K^-$  Pairs in Ultraperipheral Collisions*, *Phys. Rev. Lett.* **132** (2024) 222303 [2311.11792].
- [6] ALICE collaboration, *First measurement of the  $|t|$  dependence of incoherent  $J/\psi$  photonuclear production*, *Phys. Rev. Lett.* **132** (2024) 162302 [2305.06169].

- [7] J. Cepila, J.G. Contreras, M. Matas and A. Ridzikova, *Incoherent  $J/\psi$  production at large  $|t|$  identifies the onset of saturation at the LHC*, *Phys. Lett. B* **852** (2024) 138613 [2312.11320].
- [8] ALICE collaboration, *Exclusive four pion photoproduction in ultraperipheral Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV*, 2404.07542.
- [9] M. Klusek-Gawenda and J.D. Tapia Takaki, *Exclusive Four-pion Photoproduction in Ultra-peripheral Heavy-ion Collisions at RHIC and LHC Energies*, *Acta Phys. Polon. B* **51** (2020) 1393 [2005.13624].
- [10] ALICE collaboration, *Upgrade of the ALICE ITS detector*, *Nucl. Instrum. Meth. A* **1032** (2022) 166632 [2111.08301].
- [11] ALICE collaboration, *Measurement of the impact-parameter dependent azimuthal anisotropy in coherent  $\rho^0$  photoproduction in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV*, *Phys. Lett. B* **858** (2024) 139017 [2405.14525].
- [12] V. Guzey, M. Strikman and M. Zhalov, *Disentangling coherent and incoherent quasielastic  $J/\psi$  photoproduction on nuclei by neutron tagging in ultraperipheral ion collisions at the LHC*, *Eur. Phys. J. C* **74** (2014) 2942 [1312.6486].
- [13] A.J. Baltz, S.R. Klein and J. Nystrand, *Coherent vector meson photoproduction with nuclear breakup in relativistic heavy ion collisions*, *Phys. Rev. Lett.* **89** (2002) 012301 [nucl-th/0205031].
- [14] Z. Citron et al., *Report from Working Group 5: Future physics opportunities for high-density QCD at the LHC with heavy-ion and proton beams*, *CERN Yellow Rep. Monogr.* **7** (2019) 1159 [1812.06772].
- [15] A. Bylinkin, J. Nystrand and D. Tapia Takaki, *Vector meson photoproduction in UPCs with FoCal*, *J. Phys. G* **50** (2023) 5 [2211.16107].
- [16] S. Klein, J. Nystrand, J. Seger, Y. Gorbunov and J. Butterworth, *STARlight: A Monte Carlo simulation program for ultra-peripheral collisions of relativistic ions*, *Computer Physics Communications* **212** (2017) 258.
- [17] N.A. Burmasov, *Prospects of light-by-light scattering measurements and axion-like particle searches at the LHC*, *St. Petersburg Polytech. Univ. J. Phys. Math.* **16** (2023) 308.
- [18] N. Burmasov, E. Kryshen, P. Buehler and R. Lavička, *Upcgen: A Monte Carlo simulation program for dilepton pair production in ultra-peripheral collisions of heavy ions*, *Comput. Phys. Commun.* **277** (2022) 108388 [2111.11383].
- [19] P. Bühler, N. Burmasov, R. Lavička and E. Kryshen, *Feasibility study of tau-lepton anomalous magnetic moment measurements with ultra-peripheral collisions at the LHC*, *EPJ Web Conf.* **262** (2022) 01021.
- [20] V.P. Gonçalves and B.D. Moreira, *Fully - heavy tetraquark production by  $\gamma\gamma$  interactions in hadronic collisions at the LHC*, *Phys. Lett. B* **816** (2021) 136249 [2101.03798].
- [21] S. Ragoni, J. Seger and C. Anson, *Zero-bias new particle searches using autoencoders in UPCs and diffractive events*, 2411.00903.