

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

Proposal to the ISOLDE and Neutron Time-of-Flight Committee

Neutron transfer on the intruder ^{79}Zn isomer to probe the
 $N = 50$ shell gap in ^{80}Zn

January 8, 2025

A. Gottardo¹, L. P. Gaffney², D. K. Sharp³, P. Aguilera⁴, G. Andreetta¹, F. Angelini¹,
M. Balogh¹, J. Benito¹, G. Benzoni⁵, S. Bottoni⁵, A. Bracco⁵, D. Brugnara¹, P. Butler²,
F. Camera⁵, S. Carollo⁴, G. Corbari⁵, F. C. L. Crespi⁵, G. De Angelis¹, B. P. Kay⁷,
M. Labiche⁸, I. Lazarus⁸, S. Leoni⁵, M. Luciani⁵, F. Galtarossa⁴, N. Marchini⁹,
R. Menegazzo⁴, D. Mengoni⁴, B. Million⁵, A. Nannini⁹, R. Page², J. Pellumaj¹,
A. Gadea¹⁰, R. M. Perez-Vidal¹⁰, S. Pigliapoco⁴, E. Pilotto¹, C. Porzio⁶, R. Raabe¹¹,
F. Recchia⁴, K. Rezyunkina⁴, M. Rocchini⁹, E. Sahin¹², D. Stramaccioni¹,
J. J. Valiente-Dobon^{1,10}, O. Wieland⁵, L. Zago⁵, Y. Kucuk¹³, I. Boztosun¹³,
H. Dapoand¹³ the ISS collaboration

¹INFN, Laboratori Nazionali di Legnaro, Legnaro, Italy; ²University of Liverpool, Liverpool, UK;
³University of Manchester, Manchester, UK; ⁴University of Padova and INFN Padova, Padova,
Italy; ⁵University of Milano and INFN Milano, Milano, Italy; ⁶CERN, Geneva, Switzerland;
⁷Argonne National Laboratory, Lemont, USA; ⁸STFC, Daresbury Laboratory, Daresbury, UK;
⁹INFN Firenze, Firenze, Italy; ¹⁰IFIC, Valencia, Spain; ¹¹IKS, KU Leuven, Leuven, Belgium;
¹²University of Oslo, Oslo, Norway; ¹³Turkish Accelerator and Radiation Laboratory (TARLA),
Golbasi/Ankara

Spokespersons:

A. Gottardo (andrea.gottardo@lnl.infn.it), L. P. Gaffney (lpg@liverpool.ac.uk)

Contact person: P. MacGregor (patrick.macgregor@cern.ch)

Abstract:

Shape coexistence around ^{78}Ni is key to understanding shell evolution around one of the most neutron-rich doubly-magic nucleus known. The reduction of the $N = 50$ gap from $Z = 40$ to $Z = 32$ and the appearance of low-lying, deformed, intruder states close to $Z = 28$ makes it necessary to probe the evolution of the gap as well as of the intruder configurations in ^{80}Zn , only two protons above the ^{78}Ni core. We propose to perform a (d,p) neutron transfer on the ground $9/2^+$ and isomeric $1/2^+$ states of a ^{79}Zn beam. The exclusive differential cross sections to the states of interest in ^{80}Zn will be measured in the ISS setup. We foresee to obtain the energy of the $5^+, 6^+$ states, which are built by breaking the shell gap, as well as of the intruder 0_2^+ state. Its overlap with the ^{79}Zn $1/2^+$ intruder isomer will provide crucial information on the wave function of deformed configurations in this region.

Summary of requested shifts: 21 shifts + 3 shifts for ISS setup



1 Physics Motivation

The first spectroscopy study of the doubly-magic ^{78}Ni isotope pointed out one of the main issues found investigating the $N = 50$ isotonic chain from $Z = 40$ to $Z = 28$: the appearance of intruder, deformed configurations at low energy, pointing to shape coexistence [1, 2, 3]. Indeed, the $N = 49$ isotones offer a paradigmatic example of the lowering in energy of intruder configurations next to a shell closure. Along the isotonic chain, $1/2^+$ and $5/2^+$ states appear at low energies, as shown in Fig. 1: they are *a-priori* one particle-two holes (1p-2h) excitations across the $N = 50$ gap from the $\nu g_{9/2}$ to the $\nu s_{1/2}d_{5/2}$ shells above the shell closure [4]. Their rapid lowering in energy from about 1.5 MeV in ^{87}Sr to only ~ 500 keV in ^{81}Ge , and their re-increasing to ~ 950 keV in ^{79}Zn , has been interpreted as the conventional behaviour of intruder states which ought to have a minimum in energy at the mid of the proton shell $Z = 28 - 40$ due to the increase of quadrupole correlations [4]. A recent work proposed that the $1/2^+$ isomer in ^{79}Zn is the bandhead of a deformed $K = 1/2$ intruder band, to which also the other intruder state $5/2^+$ belongs [9].




Figure 1: The intruder states in $N = 49$ isotopes. Picture from Ref. [4].

Intruder 2p-2h 0^+ states are predicted also in the $N = 50$ nuclei [5]. Even if several excited 0^+ states are known in ^{88}Sr , ^{86}Kr and ^{84}Se , tentatively also in ^{82}Ge , their experimental identification as intruder 2p-2h states is often uncertain. In any case, the yrare 0^+ levels do feature a decrease in energy towards the mid proton shell at $Z = 34 - 32$.

The $N = 50$ shell closure was also probed with mass measurements which have shown that the $\nu g_{9/2} - d_{5/2}$ gap has a parabolic behaviour with a minimum at $Z = 32$ and a re-increase in ^{80}Zn [6]. The size of the $N = 50$ gap can also be studied measuring the energy of the excited states which represent a breaking of the shell, namely $5^+, 6^+$ and $13/2^+$ states in even and odd $N = 50$ isotones, respectively [7]. The energy of these states, known until $Z = 31$ in ^{81}Ga , shows that the shell-gap decreases in agreement with the mass measurements, but no re-increase is spotted in ^{81}Ga , towards $Z = 30$ [7], as shown in Fig. 2.

Considering the state of the present knowledge of nuclear structure in the ^{78}Ni region, it is clear that spectroscopy of medium-spin and intruder states in ^{80}Zn is much needed. Only the yrast $2^+, 4^+$ levels are known, with with only a few other observed states having an uncertain spin assignment [12, 13]. The energy of the $5^+, 6^+$ states will allow one to probe the $N = 50$ gap, while the energy of the intruder 0^+ state(s) (at least the yrare 0_2^+) will help to understand how the quadrupole correlations are impacting nuclear structure when approaching $Z = 28$. The nature of the wave function of the intruder 0_2^+ state in ^{80}Zn is intertwined with another major spectroscopic feature of this region: the rapid lowering in energy of the $\nu s_{1/2}$ shell above $N = 50$ from $Z = 40$ to $Z = 32$: at $Z = 30$ it should be almost degenerate with the $\nu d_{5/2}$ shell [8]. This, in turn, brings up a question: is the intruder yrare 0_2^+ state in ^{80}Zn a 2p-2h




Figure 2: Evolution of the $N = 50$ shell gap from mass measurements as well as spectroscopic data. The position of the $N = 50$ core-breaking states in ^{80}Zn is not known.

neutron excitation of the type $(g_{9/2})^{-2} - (d_{5/2})^2$ or $(g_{9/2})^{-2} - (s_{1/2})^2$ or some mixing of the two configurations? The answer to this bears crucial information on the deformation of the intruder states towards ^{78}Ni itself, where the degeneracy of the $\nu d_{5/2}$ and $\nu s_{1/2}$ shells should be more marked, with the possibility even of an inversion [8]. Laser spectroscopy assigned to the ^{79}Zn $1/2^+$ intruder isomer a main $\nu(g_{9/2})^{-2}(s_{1/2})^1$ component through the g-factor measurement [2]. From a naive shell-model point-of-view it is then expected that adding a neutron to this state will create an intruder 0^+ in ^{80}Zn with a dominant $\nu(g_{9/2})^{-2}(s_{1/2})^2$ wave function. In the calculations presented in Ref. [9], the $1/2^+$ and $5/2^+$ states in ^{79}Zn feature a quite mixed gds wave function, although with a predominance of the $\nu s_{1/2}$ component for the $1/2^+$ state, predicted with a deformation of $\beta = 0.22$, compatible with the isomer shift [2]. The 0_2^+ state in ^{80}Zn has a predicted similar mixing of different shells above $N = 50$ in its wave function; its deformation is calculated of the same magnitude of the ^{79}Zn intruder states [9].

The aim here is to study these aspects at the same time with a direct neutron transfer measurement $^{79}\text{Zn}^{gs,1/2^+}(d,p)^{80}\text{Zn}$. We will take advantage of the fact that in ^{79}Zn the intruder 1p-2h $1/2^+$ state is a long-living isomer, and thus the ISOLDE ^{79}Zn beam is composed of both the ground state and the isomeric state [2, 9]. The transfer on the ^{79}Zn $9/2^+$ ground state will allow one to populate the ^{80}Zn $5^+, 6^+$ states with favored $\ell = 0, 2$ transfers, while the (d,p) reaction on the ^{79}Zn $1/2^+$ intruder isomeric state will have a large cross section for the population of the intruder 0_2^+ state in ^{80}Zn with an $\ell = 0$ transfer. Figure 3 presents a schematic view of the shell-model structures involved in the neutron transfers on the two ^{79}Zn states composing the ^{79}Zn ISOLDE beam.

1.1 ^{79}Zn Coulex at Miniball

The Coulomb excitation of a $^{79}\text{Zn}^{gs,1/2^+}$ beam was successfully performed at Miniball during the run IS646 [10]. Figure 4 shows the spectrum and the deduced ^{79}Zn partial level scheme resulting from Coulomb excitation from both the ground state and the $1/2^+$ isomer. A previous β -decay work already built a level scheme on top of the $1/2^+$ level, finding almost no connecting transitions between the ground state and intruder isomer bands [11]. The preliminary results




Figure 3: Schematic shell-model view of the proposed reaction(s). The left panel shows the neutron transfer on the ^{79}Zn ground state to the predicted $5^+, 6^+$ states in ^{80}Zn . The right panel shows the transfer from the ^{79}Zn intruder isomeric $1/2^+$ state to the predicted intruder 2p-2h 0_2^+ in ^{80}Zn . Only the main wave function components are illustrated.

from IS646 show a large $E2$ strength built on the $1/2^+$ state, also connecting it to the other intruder $5/2^+$ state, suggesting a significant mixing between the $s_{1/2}$ and $d_{5/2}$ wave functions. The proposed neutron transfer measurement will help to clarify this aspect, probing the amount of $\ell = 0$ transfer to the intruder 0_2^+ level in ^{80}Zn .




Figure 4: On the left: Coulomb excitations and γ -ray transitions observed during the ^{79}Zn Coulex experiment at ISOLDE. On the right: γ -ray spectrum measured by Miniball during the ^{79}Zn Coulex run.

2 Proposed measurement

The intruder $1/2^+$ state in ^{79}Zn is a long living isomer which is about 7% of the total ISOLDE ^{79}Zn beam [9, 10]. In the recent Coulex experiment IS646, a ^{79}Zn accelerated beam intensity of about $8 \cdot 10^4$ pps was obtained [10]. We plan to use the ISS device to measure the differential cross section of the $^{79}\text{Zn}^{gs,1/2^+}(\text{d,p})$ reaction to the different excited states of ^{80}Zn .

The ^{80}Zn excited states as well as the predicted spectroscopic factors were calculated using an interaction derived from the the GWB Hamiltonian from the Oxbash library. The interaction has a ^{66}Ni core and comprises as active shells the full *gds* neutron space across $N = 50$ as well as the $f_{5/2}p_{3/2}p_{1/2}$ shells between $Z = 28$ and $Z = 40$. This interaction can reproduce the intruder states along the $N = 49$ isotonic chain, the $2^+, 4^+, 5^+, 6^+$ energy and $B(E2)$ along the lower-mass $N = 50$ isotones. The $^{79,80}\text{Zn}$ state energies and the (d,p) spectroscopic factors were obtained from a diagonalization with the Antoine code, allowing up to 4p-4h excitations across $N = 50$. In general, the results are qualitatively similar to the interaction in Ref. [9]¹. The wave functions of the 0_2^+ and $5_{1,2}^+, 6_1^+$ states in ^{80}Zn are constituted by about two and one neutron holes in the $g_{9/2}$ shell, respectively. The missing neutrons are excited mainly to the *sd* shells above $N = 50$. Similarly to the calculations in Ref. [9], the breaking of the $N = 50$ core also implies a change in the protons wave function above $Z = 28$, with some depletion of the $\pi f_{5/2}$ shell and more proton occupancy in the $\pi p_{3/2}, p_{1/2}$ shells, compared to the spherical ground state.

The calculated spectroscopic factors were then used as a basis for DWBA calculations with the FRESKO code. The obtained differential cross sections were in turn used for the experiment simulation using the NPTOOL software. Table 1 shows a summary of the predicted cross sections and counting rates for the excited states of ^{80}Zn we will populate with a significant yield. The final rates includes the ISS response function. The DWBA calculations were performed at different beam energies, from 4 MeV/u to 10 MeV/u, and the best rates are obtained at a beam energy of 6 MeV/u. The CD_2 target thickness is a compromise between excitation energy resolution and statistics, which in our case works out at $300 \mu\text{g}/\text{cm}^2$.

^{80}Zn state	Excitation Energy (MeV)	Transferred ℓ	SF	Beam pps	σ (mb)	ISS counts per shift	ISS counts 21 shifts
5_1^+	3.0	0	0.1	$^{79}\text{Zn}^{gs}$	1.2	36	750
		2	0.7	$7.4 \cdot 10^4$	1.3		
6_1^+	3.2	0	0.3	$^{79}\text{Zn}^{gs}$	2.5	35	740
				$7.4 \cdot 10^4$			
5_2^+	3.6	2	0.8	$^{79}\text{Zn}^{gs}$	0.7	10	220
0_2^+	2.2	0	0.4	$^{79}\text{Zn}^{1/2^+}$	1.6	2	40
				$5.6 \cdot 10^3$			

Table 1: Estimation of collected statistics for a $^{79}\text{Zn}^{gs,1/2^+}$ beam at 6 MeV/u and an intensity of $8 \cdot 10^4$ pps. Other ^{80}Zn states are predicted to have negligible cross sections. Predictions on spectroscopic factors (SF) are based on a shell model allowing up to 4p-4h across $N = 50$.

¹The $N = 50$ gap was tuned to reproduce the 0_2^+ energy calculated in Ref. [9].

In Fig. 5 we present the predicted differential cross sections compared with the ISS angular coverage in a configuration where the target to array distance is (-80—580mm) and field settings is 2T. The angular coverage is sufficient to differentiate the different transferred angular momenta.




Figure 5: Predicted differential cross section for a DWBA calculation performed with the FRESKO code. The shaded area is the ISS angular coverage in the configuration described in the text.

Figure 6 presents the output of the Monte Carlo simulation for particle spectroscopy in ISS. The 5_1^+ , 6_1^+ levels are not well separated, while the 5_2^+ is clearly distinguished. Since the two 5_1^+ , 6_1^+ levels represent both a break of the $N = 50$ core and their predicted energy is anyway within the uncertainties of the shell model, their resolution is not crucial for the measurement. The intruder 0_2^+ state clearly stands out: the low percentage of ^{79}Zn in the isomeric state is counterbalanced by a relatively large cross section for the $\ell = 0$ transfer.

Finally, we foresee the use of the zero-degree ionization chamber to check beam contamination and clean the fusion-evaporation background. In the previous IS646 run with ^{79}Zn beam we observed only a minor contamination of the order of 2% from a higher Z , likely Kr.

In total 21 shifts are requested to gather the required statistics, plus 3 shifts for ISS setup, two with stable beam and one with radioactive beam.

References

- [1] R. Taniuchi *et al.*, Nature 569, 53–58 (2019)
- [2] X.F. Yang *et al.*, Phys.Rev.Lett. 116 18, 182502 (2016)
- [3] F. Nowacki *et al.*, Phys. Rev. Lett. 117, 272501 (2016)




Figure 6: Simulation of the excitation energy spectrum in ISS. The blue line represents transfer on the ^{79}Zn ground state, while the red line transfer on the ^{79}Zn $1/2^+$ isomer. The right panel show the 5_1^+ , 6_1^+ states not well resolved, and the 5_2^+ level resolved. The left panel shows, in red, a zoom in the low energy region where the peak corresponding to the predicted intruder 0_2^+ state is simulated. Predicted at 2.1 MeV, it appears at 1.2 MeV because the excitation energy has been determined assuming transfer from the ^{79}Zn ground state.

- [4] C. Wraith *et al.*, Phys.Lett.B 771, 385-391 (2017)
- [5] A. Gottardo *et al.*, Phys. Rev. Lett. 116, 182501 (2016)
- [6] S. Baruah *et al.*, Phys. Rev. Lett. 101, 262501 (2008)
- [7] J. Dudouet *et al.*, Phys. Rev. C 011301(R) (2019)
- [8] C. Delafosse *et al.*, Phys. Rev. Lett. 121 19, 192502 (2018)
- [9] L. Nies *et al.*, Phys. Rev. Lett. 131, 222503 (2023)
- [10] F. Angelini, A. Gottardo *et al.*, IS646 https://indico.cern.ch/event/1442908/contributions/6184073/attachments/2976160/5239124/ISOLDE_WS_79Zn_gottardo.pdf
- [11] M.C. Delattre, PhD thesis, https://theses.hal.science/tel-01453119v1/file/69822_DELATTRE_2016_diffusion.pdf
- [12] Y. Shiga *et al.*, Phys. Rev. C 93, 024320 (2016)
- [13] J. Van de Walle *et al.*, Phys. Rev. Lett. 99, 142501 (2007)

3 Details for the Technical Advisory Committee

3.1 General information

Describe the setup which will be used for the measurement. If necessary, copy the list for each setup used.

- Permanent ISOLDE setup: *ISS*
 - To be used without any modification
 - To be modified: *Short description of required modifications.*
- Travelling setup (*Contact the ISOLDE physics coordinator with details.*)
 - Existing setup, used previously at ISOLDE: *Specify name and IS-number(s)*
 - Existing setup, not yet used at ISOLDE: *Short description*
 - New setup: *Short description*

3.2 Beam production

For any inquiries related to this matter, reach out to the target team and/or RILIS (please do not wait until the last minute!). For Letters of Intent focusing on element (or isotope) specific beam development, this section can be filled in more loosely.

- Requested beams:

Isotope	Production yield in focal point of the separator (/μC)	Minimum required rate at experiment (pps)	$t_{1/2}$
⁷⁹ Zn	$\geq 3 \cdot 10^5$	$\geq 7 \cdot 10^4$	746 ms
Isotope 2			
Isotope 3			

- Full reference of yield information: *extrapolated from IS646 in Sep. 24*
- Target - ion source combination:
- RILIS? *Yes*
 - Special requirements:
- Additional features?
 - Neutron converter: *Yes*
 - Other: *quartz transfer line*
- Expected contaminants: *Isotopes with similar A/q from EBIS, few % of the ⁷⁹Zn beam*
- Acceptable level of contaminants: *(few% of the ⁷⁹Zn beam, either stable or radioactive)*
- Can the experiment accept molecular beams? *Not relevant*
- Are there any potential synergies: *IS743 (⁷⁸Zn beam)*

3.3 HIE-ISOLDE

For any inquiries related to this matter, reach out to the ISOLDE machine supervisors (please do not wait until the last minute!).

- HIE ISOLDE Energy: (~ 6 MeV/u);
 - Precise energy determination required
 - Requires stable beam from REX-EBIS for calibration/setup? *Isotope?*
- REX-EBIS timing
 - Slow extraction
 - Other timing requests
- Which beam diagnostics are available in the setup? Ionization chamber, Faraday cup, active beam collimator
- What is the vacuum level achievable in your setup?

3.4 Shift breakdown

The beam request only includes the shifts requiring radioactive beam, but, for practical purposes, an overview of all the shifts is requested here. Don't forget to include:

- Isotopes/isomers for which the yield need to be determined
- Shifts requiring stable beam (indicate which isotopes, if important) for setup, calibration, etc. Also include if stable beam from the REX-EBIS is required.

An example can be found below, please adapt to your needs. Copy the table if the beam time request is split over several runs.

Summary of requested shifts: 21 shifts of ^{79}Zn for physics plus one shift of ^{79}Zn for ISS setup plus two shifts with stable beam for ISS setup before the experiment

With protons	Requested shifts
Optimization of experimental setup using ^{79}Zn	1
Data taking, ^{79}Zn	21
Without protons	Requested shifts
Stable beam from REX-EBIS (BEFORE run)	2

3.5 Health, Safety and Environmental aspects

3.5.1 Radiation Protection

- If radioactive sources are required:
 - Purpose? Silicon detector calibration
 - Isotopic composition? α source for detector calibration (148Gd, 239Pu, 142Am, 244Cm)
 - Activity? 1 kBq
 - Sealed/unsealed? Unsealed
- For collections:
 - Number of samples?
 - Activity/atoms implanted per sample?
 - Post-collection activities? (*handling, measurements, shipping, etc.*)

3.5.2 Only for traveling setups

- Design and manufacturing
 - Consists of standard equipment supplied by a manufacturer
 - CERN/collaboration responsible for the design and/or manufacturing
- Describe the hazards generated by the experiment:

Domain	Hazards/Hazardous Activities	Description
Mechanical Safety	Pressure	<input type="checkbox"/> [pressure] [bar], [volume][l]
	Vacuum	<input type="checkbox"/>
	Machine tools	<input type="checkbox"/>
	Mechanical energy (moving parts)	<input type="checkbox"/>
	Hot/Cold surfaces	<input type="checkbox"/>
Cryogenic Safety	Cryogenic fluid	<input type="checkbox"/> [fluid] [m3]
Electrical Safety	Electrical equipment and installations	<input type="checkbox"/> [voltage] [V], [current] [A]
	High Voltage equipment	<input type="checkbox"/> [voltage] [V]
Chemical Safety	CMR (carcinogens, mutagens and toxic to reproduction)	<input type="checkbox"/> [fluid], [quantity]
	Toxic/Irritant	<input type="checkbox"/> [fluid], [quantity]
	Corrosive	<input type="checkbox"/> [fluid], [quantity]
	Oxidizing	<input type="checkbox"/> [fluid], [quantity]
	Flammable/Potentially explosive atmospheres	<input type="checkbox"/> [fluid], [quantity]
	Dangerous for the environment	<input type="checkbox"/> [fluid], [quantity]
Non-ionizing radiation Safety	Laser	<input type="checkbox"/> [laser], [class]
	UV light	<input type="checkbox"/>
	Magnetic field	<input type="checkbox"/> [magnetic field] [T]

Workplace	Excessive noise	<input type="checkbox"/>	
	Working outside normal working hours	<input type="checkbox"/>	
	Working at height (climbing platforms, etc.)	<input type="checkbox"/>	
	Outdoor activities	<input type="checkbox"/>	
Fire Safety	Ignition sources	<input type="checkbox"/>	
	Combustible Materials	<input type="checkbox"/>	
	Hot Work (e.g. welding, grinding)	<input type="checkbox"/>	
Other hazards			