

AB



A MEASUREMENT OF $R = \sigma_L/\sigma_T$ IN DEEP INELASTIC
NEUTRINO-NUCLEON SCATTERING AT THE TEVATRON

by

A Bodek, U K Yang, P Auchincloss, P de Barbaro, H Budd, D A Harris,
W K Sakumoto, W C Lefmann, R Steiner, M J Oreglia, B A Schumm,
R A Johnson, M Nussbaum, L Perera, M Vakili, C G Arroyo, A O Bazarko,
J Conrad, J H Kim, B J King, S Koutsoliotas, C McNulty, S R Mishra
Z Quintas, A Romosan, F J Sciulli, W G Seligman, M H Shaevitz,
Spentzouris, R G Stern, R H Bernstein, G Koizumi, M J Lamm,
Marsh, K S McFarland, T Bolton, W B Lowery, D Naples, R B Drucker,
T Kinnel, W H Smith, and P Nienaber
(A. Bodek for the CCFR/NuTeV Collaboration)



CERN LIBRARIES, GENEVA

swg636

This report was prepared as an account of work sponsored by the United States government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assume any legal responsibility or liability for the accuracy, completeness, or usefulness of any information, apparatus, product of process disclosed or represented that its use would not infringe privately-owned rights.

7/96

UNIVERSITY OF ROCHESTER
DEPARTMENT OF PHYSICS AND ASTRONOMY
ROCHESTER, NEW YORK, 14627

A MEASUREMENT OF $R = \sigma_L/\sigma_T$ IN DEEP INELASTIC NEUTRINO-NUCLEON SCATTERING AT THE TEVATRON

A Bodek, U K Yang, P Auchincloss, P de Barbaro, H Budd, D A Harris,
W K Sakamoto, W C Lefmann, R Steiner, M J Oreglia, B A Schumm,
R A Johnson, M Nussbaum, L Perera, M Vakili, C G Arroyo,
A O Bazarko, J Conrad, J H Kim, B J King, S Koutsoliotas, C McNulty,
S R Mishra, Z Quintas, A Romosan, F J Sciulli, W G Seligman,
M H Shaevitz, Spentsouris, E G Stern, R H Bernstein, G Koisumi,
M J Lamm, Marsh, K S McFarland, T Bolton, W B Lowery, D Naples,
R B Drucker, T Kinnel, W H Smith, and P Nienaber
(A. Bodek for the CCFR/NuTeV Collaboration)

Measurements of neutrino-nucleon and antineutrino-nucleon differential cross sections using the CCFR neutrino detector at Fermilab have been used to extract preliminary values of $R = \sigma_L/\sigma_T$ in the kinematic region $0.01 < x < 0.6$, and $4 < Q^2 < 300 \text{ GeV}^2$. The new data provide the first measurements of R in the $x < 0.1$ region. The x and Q^2 dependence of R is compared with a QCD based fit to previous data. The QCD fit, which provides an estimate of R in small x region where R has not been previously measured, is in good agreement with the new CCFR data.

1 Introduction

The ratio $R = \sigma_L/\sigma_T$ of the longitudinal and transverse absorption cross sections in deep inelastic lepton-nucleon scattering provides information about the transverse momentum and spin of the nucleon constituents. Within the theory of quantum chromodynamics (QCD), the nucleon constituents are spin 1/2 quarks and spin 1 gluons. In leading order QCD, $R = 0$, since the quarks have no transverse momentum. In the next to leading order formalism (NLO), to first order in α_s , R is non-zero because of transverse momentum associated with gluon emission¹. The NLO QCD prediction is given by an integral over the quark and gluon distribution and is proportional to α_s . A measurement of R is a test of perturbative QCD at large x , and a clean probe of the gluon density at small x where the quark contribution is small.

Poor knowledge of R , especially at small x , results in uncertainties in the structure functions extracted from deep inelastic scattering cross sections. The most precise previous measurements of R are from SLAC electron scattering experiments², and are at high x and low Q^2 , where non-perturbative contributions (e.g. target mass and higher twist effect) are important. Therefore, extraction of structure functions at large Q^2 or small x must rely on good

measurements of R , or on reasonable estimates of R in the kinematic region where R has not been measured.

We report here on a new extraction of R from measurements of neutrino and antineutrino differential cross sections in the CCFR neutrino detector. We compare the data (particularly at small x) to the prediction of a QCD based fit to R by Bodek, Rock and Yang³. Extrapolation of such a QCD based fit to smaller values of x can be also used to obtain estimates of R in the HERA region where R has not been measured.

2 R calculation within QCD

In this section we provide a brief description of the QCD based model of Bodek, Rock and Yang (BRY). They have used a calculation⁴ of the QCD contribution to R to order α_s^2 (NNLO). The contribution of massive heavy quark (charm) effects to R has been included within the photon gluon fusion process⁵. These higher order QCD corrections and heavy charm quark effects are important at small x . In addition, the SLAC data on R indicate that there are additional non-perturbative $1/Q^2$ and $1/Q^4$ contributions at low Q^2 ^{2,3}. One of these contributions, target mass effects, dominates at low Q^2 and large x . A comparison of the calculation (including both the heavy charm contribution to order α_s^2 and the Georgi-Politzer target mass corrections⁶) to the SLAC data indicates that additional small higher twist contribution at low Q^2 are required³. A parameterization of the higher twist effects at low Q^2 should be constrained such that R approaches zero at the photoproduction limit ($Q^2 = 0$). BRY find that a simple empirical parameterization of the higher twist contribution to R (of the form $A[(Q^2 - B)/(Q^4 + C)]$) fits the SLAC data very well. The best BRY fit to the all previous data^{2,7} (for $Q^2 > 0.4$ GeV²), using the GRV94⁸ parton distribution for the QCD contribution, yields $A = 0.37$, $B = 0.82$, and $C = 0.54$ (χ^2/dof is 21.7/19). Figure 1 shows some of the previous data and the model predictions calculated using various parton distributions. All curves include QCD to order α_s^2 , the Georgi-Politzer target mass correction and the empirical small higher twist contribution. The curves labeled with (h) also include the effects of the charm quark mass. Also shown for comparison are the extrapolation of a previous empirical fit (Rworld) for R used by the SLAC, CCFR, and NMC collaborations. The QCD model predictions with various parton distributions^{8,9} all yield similar results for R at large x (> 0.1). At small x , the predictions of the model using either the MRSD' and the MRSA distributions are very different because of the different gluon distributions. Note that target mass effects at small x are negligible, and the small empirical higher twist contribution is assumed to be independent of




Figure 1: R at fixed x vs Q^2 : A comparison of the BRY prediction of R calculated using various parton distributions (including target mass and empirical higher twist) to data. A massive heavy (charm) quark contribution is included in the MRS(h) and GRV94(h) prediction. Also shown is R_{world} , which is a previous empirical parameterization of R data in the SLAC region.

x . The model calculated with the GRV94 parton distributions (solid curve, figure 1) is probably the best available estimate for extracting F_2 in regions where R has not been measured. We proceed to test the models shown in figure 1 against the new CCFR data in the section below.

3 A Preliminary CCFR measurement of R

The data sample includes two CCFR data runs (E744 and E770) collected using the Fermilab Tevatron Quad-Triplet neutrino beam. The wide-band beam is composed of ν_μ and $\bar{\nu}_\mu$ with energies up to 600 GeV. The CCFR neutrino detector¹⁰ consists of an unmagnetized steel-scintillator target calorimeter instrumented with drift chambers. The hadron energy resolution is $\Delta E/E = 0.85/\sqrt{E}(\text{GeV})$. The target is followed by a solid iron toroidal magnet muon spectrometer which measures muon momentum with a resolution $\Delta p/p = 0.11$. The relative flux at different energies is obtained from the events with low hadron energy, $\nu < 20$ GeV, and is normalized so that the neutrino total cross




Figure 2: Preliminary differential cross section (statistical error only) for neutrino[a] and antineutrino[b] at $E_\nu = 150$ GeV: Preliminary CCFR data (solid circles) are compared to the QCD LO prediction and previous CDHSW data (open squares).

section equals the world average $\sigma^{\nu N}/E = (0.674 \pm 0.014) \times 10^{-38} \text{ cm}^2/\text{GeV}$, and $\sigma^{\bar{\nu}N}/\sigma^{\nu N} = 0.504 \pm 0.005$.

Preliminary values of R have been extracted from the differential cross sections in neutrino charged current scattering. The differential cross sections are written as

$$\frac{d^2\sigma}{E dx dy} = \frac{G_F^2 M}{\pi} \left[F_2(x, Q^2) \left(1 - y + \frac{y^2}{2(1+R)} \right) \pm x F_3(x, Q^2) \left(y - \frac{y^2}{2} \right) \right] \quad (1)$$

where $y = \nu/E_\nu$ is the inelasticity, and $R = \sigma_L/\sigma_T = F_2(1+Q^2/\nu^2)/2xF_1 - 1$. In the quark-parton model, $2xF_1^{\nu, \bar{\nu}} = q + \bar{q}$ and $xF_3^{\nu, \bar{\nu}} = (q - \bar{q}) \pm 2(s - c)$ on an isoscalar target.

The data sample consists of 1,280,000 ν_μ and 270,000 $\bar{\nu}_\mu$ events after fiducial and kinematic cuts ($P_\mu > 15$ GeV, $\theta_\mu < .150$, $\nu > 10$ GeV, $Q^2 > 1$ GeV², and $E_\nu > 30$ GeV). Dimuon events are removed because of the ambiguous identification of the leading muon for high y events. The data are corrected

by Monte Carlo for cuts, acceptance and resolution smearing. The differential cross sections are determined in bins of x , y and E_ν ($0.01 < x < 0.60$, $0.05 < y < 0.95$, and $30 < E_\nu < 360$ GeV). Figure 2 shows the differential cross sections at $E_\nu = 150$ GeV. The differential cross sections agree with a LO QCD model which predicts a quadratic y dependence at small x for neutrino and anti-neutrino and a flat y distribution at high x for the neutrino cross section. A comparison of CCFR and CDHSW differential cross sections⁷ indicates good agreement at high x , and poor agreement at small x , as shown in Figure 2.

The values of R are extracted from linear fits to F vs ϵ at fixed x and Q^2 bins through the following relation,

$$F(x, Q^2, \epsilon) = \frac{\pi(1-\epsilon)}{y^2 G_F^2 M E} \left[\frac{d^2 \sigma^\nu}{dx dy} + \frac{d^2 \sigma^{\bar{\nu}}}{dx dy} \right] = 2xF_1(x, Q^2) [1 + \epsilon R(x, Q^2)] \quad (2)$$

which assumes $x F_3^\nu = x F_3^{\bar{\nu}}$. Here $\epsilon \simeq 2(1-y)/(1+(1-y)^2)$ is the polarization of virtual W boson. Corrections for electroweak radiative effects (Bardin), the W boson propagator, and isoscalar target (the 6.85% excess of neutron over proton in iron) are applied. The correction for $\Delta x F_3 = x F_3^\nu - x F_3^{\bar{\nu}}$ (which is important at small x) requires a knowledge of the strange sea and charm production mass effects (e.g. slow rescaling). The strange sea distribution, extracted to NLO from CCFR dimuon data¹¹ has been used. The fits for R at each x and Q^2 have reasonable χ^2 . Figure 3 shows one of the linear fits. The values of R with statistical errors at fixed x vs Q^2 are shown in Figure 4. The new data, which extends a decade lower in x , are in agreement with the QCD based model. The values of R are not sensitive to the normalization of the absolute flux. The dominant uncertainty at very small x and low Q^2 is from the non-zero value of $\Delta x F_3$ because of uncertainties in the slow rescaling formalism and the level of the strange sea. Therefore, the values of R in the kinematic region $x < 0.1$ and $Q^2 < 4$ GeV² are not shown at present.

Better modelling of $\Delta x F_3$ at low Q^2 and small x , and more precise data from the 1996-97 run of NuTeV (FNAL E815) experiment with sign selected neutrino beams are expected to yield better determinations of R at small x .

References

1. Altarelli G and Martinelli G 1978 *Phys. Lett.* **76B** 89
2. Tao L H *et al.*, (E140X collab.), *SLAC-PUB-6737 to be published in 1996 Z. Phys. C*;
Dasu S *et al.*, 1994 *Phys. Rev. Lett.* **60**; Whitlow L W *et al.*, 1990 *Phys. Lett.* **250B** 193




Figure 3: $F(x, Q^2, \epsilon)$ vs ϵ fit at $x = 0.045$ and $Q^2 = 5.01 \text{ GeV}^2$: The errors on F are only statistical errors.

3. Bodek A, Rock S and Yang U K *UR-1355, to be published in 1996 Z. Phys. C*
4. Zijlstra E and van Neerven W L 1992 *Nucl. Phys. B* **383** 525
5. Laenen E, Riemersma S, Smith J and van Neerven W L 1993 *Nucl. Phys. B* **392** 162
6. Georgi H and Politzer H D 1976 *Phys. Rev. D* **14** 1829
7. Aubert J J *et al.*, (EMC collab.) 1987 *Nucl. Phys. B* **293** 740; Benvenuti A C *et al.*, (BCDMS collab.) 1990 *Phys. Lett.* **237B** 592; Berge P *et al.*, (CDHSW collab.) 1991 *Z. Phys. C* **49**
8. Gluck M, Reya E, and Vogt A 1995 *Z. Phys. C* **67** 433
9. Martin A D, Stirling W J and Roberts R G 1994 *Phys. Rev* **50** 6734
10. Sakumoto W K *et al.*, (CCFR collab.) 1990 *NIM A* **294** 179; King B J *et al.*, 1991 *NIM A* **302** 254
11. Bazarko A O *et al.*, (CCFR collab.) 1995 *Z. Phys. C* **65** 189




Figure 4: R at fixed x vs Q^2 : Preliminary CCFR data are compared with other data and with the Bodek, Rock and Yang QCD based fit with various parton distributions.

