

**CENTRALITY DEPENDENCE OF  $K^+$  PRODUCED IN PB+PB  
COLLISIONS AT 158 GEV PER NUCLEON**

SONJA KABANA FOR THE NA52 COLLABORATION:

R ARSENECU, H P BECK, K BORER, S KABANA, R KLINGENBERG, G LEHMANN,  
R MOMMSEN, U MOSER, K PRETZL, J SCHACHER, R SPIWOKS, M WEBER

*Laboratory for High Energy Physics, University of Bern, Sidlerstrasse 5, CH-3012 Bern,  
Switzerland (E-mail: sonja.kabana@cern.ch)*

K ELSENER, K D LOHMANN

*CERN, CH-1211 Geneva 23, Switzerland*

C BAGLIN, A BUSSIÈRE, J P GUILLAUD

*CNRS-IN2P3, LAPP Annecy, F-74941 Annecy-le-Vieux, France*

T LINDÉN, J TUOMINIEMI

*Dept. of Physics and Helsinki Institute of Physics, University of Helsinki, PO Box 9,  
FIN-00014 Helsinki, Finland*

PH GORODETZKY

*PCC-College de France, 11 place Marcelin Berthelot, 75005 Paris, France*

The NA52 collaboration searches for a discontinuous behaviour of charged kaons produced in Pb+Pb collisions at 158 A GeV as a function of the impact parameter, which could reveal a hadron to quark-gluon plasma (QGP) phase transition. The  $K^+$  yield is found to grow proportional to the number of participating ('wounded') nucleons  $N$ , above  $N=100$ . Previous NA52 data agree with the above finding and show a discontinuous behaviour in the kaon centrality dependence near  $N=100$ , marking the onset of strangeness enhancement -over e.g. p+A data at the same  $\sqrt{s}$ - in a chemically equilibrated phase.

## 1 Introduction

The hadron to quark-gluon plasma phase transition predicted by QCD <sup>2</sup> may occur and manifest itself in ultrarelativistic heavy ion collisions through discontinuities in the energy density dependence of relevant observables. A major example of such a discontinuity is seen in the  $J/\Psi/DY$  ratio <sup>3</sup>. The NA52 collaboration searches for discontinuities in strangeness production measuring charged kaons as a function of the impact parameter. Results from the 1995 NA52 run are published in <sup>4</sup>. We report here on new preliminary results from the 1998 run of the NA52 experiment, on  $K^+$  at rapidity 4.1 and trans-

verse momentum near 0 produced in Pb+Pb collisions at 158 A GeV <sup>5</sup>. In this run a new electromagnetic lead/quartz fiber calorimeter (QFC) with improved acceptance and resolution <sup>6</sup> was used.

## 2 Results and discussion

For the kaon measurement we modified the 1998 set up of NA52 <sup>7</sup> by placing the target 0.6 m upstream of the calorimeter. The results have been corrected for empty target contributions. The number of participant nucleons  $N$  has been estimated from the energy measured with the calorimeter (figure 1) in the way described in <sup>4</sup>. Particle identifica-




Figure 1. Preliminary energy distribution in arbitrary units in minimum bias Pb+Pb collisions at 158 A GeV, from the 1998 NA52 run.

tion is described in <sup>4</sup> and references there. The positive kaon yield divided by  $N$  is independent of  $N$ , for  $N > 100$  (figure 2) in agreement with previous NA52 results <sup>4</sup>. Assuming that  $N$  is proportional to the volume of the particle source, figure 2 shows that the kaon number density exhibit a discontinuity, saturating above  $N=100$ .

This indicates a transition to a phase characterized by a high degree of chemical equilibrium and enhancement <sup>4</sup> of kaons from the point  $N=100$  on, corresponding to energy density  $\epsilon \sim 1.3 \text{ GeV}/\text{fm}^3$  <sup>8,9,1</sup>, near the critical energy density  $\epsilon_c \sim 1-2 \text{ GeV}/\text{fm}^3$  <sup>2,10</sup>. This change marks the onset of strangeness enhancement seen in kaons in an equilibrated phase, which may be suggestive for a QCD phase transition, depending on the simultaneous appearance of thresholds in other signatures like e.g. the  $J/\Psi$  suppression at the relevant  $\epsilon$  values and their theoretical understanding <sup>10,3,8,11</sup>.

### Acknowledgments

We thank Prof. P. Minkowski for fruitful discussions and the Schweizerischer Nationalfonds for their support.




Figure 2. Preliminary  $K^+$  yield in arbitrary units per participant nucleon  $N$ , as a function of  $N$  from Pb+Pb collisions at 158 A GeV, measured in the 1998 NA52 run. For comparison the 1995 NA52 data <sup>4</sup> are also shown. The 1998 data are scaled by 2.4 with respect to the 1995 data.

### References

1. S. Kabana, proceedings of the conference ICHEP2000 and Univ. of Bern preprint BUHE-00-08.
2. E. Laermann, Nucl. Phys. A610 (96) 1c.
3. M.C. Abreu et al (NA50 coll.), CERN-EP-2000-013.
4. G. Ambrosini et al. (NA52 Coll.), New J. of Phys. (99) 1, 22, G. Ambrosini et al. (NA52 Coll.), New J. of Phys. (99) 1, 23, S. Kabana et al. (NA52 Coll.), Nucl. Phys. A 661 (99) 370c.
5. S. Kabana et al (NA52 coll.), contribution to the conference SQM2000 and Univ. of Bern preprint BUHE-00-05.
6. M. Weber et al (NA52 Coll.), CALOR97, World Scientific (98) 151.
7. M. Weber et al (NA52 coll.), contribution to SQM2000 and Univ. of Bern preprint BUHE-00-04.
8. S. Kabana, hep-ph-0004138.
9. S. Kabana, contrib. to SQM2000 and Univ. of Bern preprint BUHE-00-06.
10. H. Satz, hep-ph/0007069.
11. P.B. Munzinger et al, nucl-th/0007059.