




XJ0300124



ОБЪЕДИНЕННЫЙ  
ИНСТИТУТ  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ

Дубна

E2-2002-251

M. V. Tokarev<sup>1</sup>, Yu. A. Panebratsev<sup>2</sup>, G. P. Škoro<sup>3</sup>,  
I. Zborovský<sup>4</sup>

SELF-SIMILARITY OF HIGH- $p_T$  HADRON  
PRODUCTION IN  $\pi^-p$  AND  $\pi^-A$  COLLISIONS

Presented at the XVI International Conference on Ultra Relativistic  
Nucleus-Nucleus Collisions «Quark Matter'2002»,  
July 18–24, 2002, Nantes, France

<sup>1</sup>E-mail: tokarev@sunhe.jinr.ru

<sup>2</sup>E-mail: panebrat@sunhe.jinr.ru

<sup>3</sup>Institute of Nuclear Science «Vinča», Faculty of Physics,  
University of Belgrade, Belgrade, Yugoslavia; E-mail: goran@ff.bg.ac.yu

<sup>4</sup>Nuclear Physics Institute, Academy of Sciences of the Czech Republic,  
Řež, Czech Republic; E-mail: zborovsk@uif.cas.cz

# 1 Introduction

Particle production with high transverse momenta is traditionally connected with local character of hadron interactions. Locality of the interaction is expressed in terms of the hadron constituents. Scaling features of high- $p_T$  hadron spectra reflects self-similarity of constituent interactions. Therefore, search for scaling regularities in high energy collisions of hadrons and nuclei is of physical interest. Up to this date, the investigation of hadron properties in the high energy collisions has revealed widely known scaling laws. From the most popular and famous, let us mention the Feynman scaling [1] observed in inclusive particle production, the Bjorken scaling [2] in deep inelastic scattering (DIS),  $y$ -scaling [3] valid in DIS on nuclei, limiting fragmentation established for nuclei [4], scaling behaviour of the cumulative particle production [5, 6, 7], the Koba-Nielsen-Olesen (KNO) scaling [8], and others. Another expression of self-similarity in hadronic interactions are described by quark counting rules [9].

One of the methods to study the properties of particle structure, constituent interactions and particle formation is  $z$ -scaling [10]. The scaling was observed in proton-(anti)proton and proton-nucleus high energy collisions. In this paper we develop the concept of the scaling for  $\pi^-p$  and  $\pi^-A$  collisions.

## 2 Basic principles of $z$ -scaling

The idea of  $z$ -scaling is based on the principles of locality, self-similarity and fractality of hadronic interactions. The properties are assumed to be revealed in high energy collisions of hadrons and nuclei. Their most pronounced manifestation is expected in the inclusive production of particles with large transverse momenta.

### 2.1 Locality principle

Following the ideas suggested in Ref. [6], we assume that gross features of the inclusive particle distributions for the reaction

$$M_1 + M_2 \rightarrow m_1 + X \quad (1)$$

can be described in terms of the constituent sub-process

$$(x_1 M_1) + (x_2 M_2) \rightarrow m_1 + (x_1 M_1 + x_2 M_2 + m_2) \quad (2)$$

satisfying to the condition

$$(x_1 P_1 + x_2 P_2 - p)^2 = (x_1 M_1 + x_2 M_2 + m_2)^2. \quad (3)$$

The assumption is expression of locality of the hadronic interactions at constituent level. The  $x_1$  and  $x_2$  are fractions of the incoming four-momenta  $P_1$  and  $P_2$  of the colliding objects with the masses  $M_1$  and  $M_2$ . The  $p$  is four-momentum of the inclusive particle  $m_1$ . The parameter  $m_2$  is introduced to satisfy the internal conservation laws (for isospin, baryon number, strangeness, and so on).

### 2.2 Self-similarity principle

Self-similarity is a scale-invariant property connected with dropping of certain dimensional quantities out of physical picture of the interaction. It means that for description

of physical processes dimensionless quantities are used. One of the approaches to description of production cross sections complying with this requirement is the  $z$ -scaling. The scaling function  $\psi(z)$  depends in a self-similar manner on a single variable  $z$ . The scaling function is expressed via the invariant differential cross section  $E d^3\sigma/dp^3$  as follows

$$\psi(z) = -\frac{\pi s}{\rho\sigma_{in}} J^{-1} E \frac{d^3\sigma}{dp^3}. \quad (4)$$

Here  $s$  is the collision center-of-mass energy squared,  $\sigma_{in}$  is the inelastic cross section,  $J$  is the corresponding Jacobian, and  $\rho = dN/d\eta$  is the particle multiplicity density. The function  $\psi(z)$  is normalized as

$$\int_{z_{min}}^{\infty} \psi(z) dz = 1. \quad (5)$$

The relation allows us to give the physical meaning of the scaling function  $\psi(z)$  as a probability density to form a particle with the corresponding value of the variable  $z$ .

### 2.3 Fractality principle

Principle of fractality states that variables used in the description of the process diverge in terms of the resolution [11, 12]. This property is characteristic for the scaling variable

$$z = z_0 \Omega^{-1}, \quad (6)$$

where

$$\Omega(x_1, x_2) = (1 - x_1)^{\delta_1} (1 - x_2)^{\delta_2}. \quad (7)$$

The variable  $z$  has character of a fractal measure. For a given production process (1), its finite part  $z_0$  is the ratio of the transverse energy released in the binary collision of constituents (2) and the average multiplicity density  $dN/d\eta|_{\eta=0}$ . The divergent part  $\Omega^{-1}$  describes the resolution at which the collision of the constituents can be singled out of this process. The  $\Omega(x_1, x_2)$  represents relative number of all initial configurations containing the constituents which carry fractions  $x_1$  and  $x_2$  of the incoming momenta. The  $\delta_1$  and  $\delta_2$  are the anomalous fractal dimensions of the colliding objects (hadrons or nuclei). The momentum fractions  $x_1$  and  $x_2$  are determined in a way to minimize the resolution  $\Omega^{-1}(x_1, x_2)$  of the fractal measure  $z$  with respect to all possible sub-processes (2) subjected to the condition (3). The variable  $z$  was interpreted [10] as a particle formation length.

## 3 $Z$ -scaling in $\pi^-p$ collisions

In this section we study the properties of data  $z$ -presentation for hadron production in  $\pi^-p$  collisions. One of the ingredients in the definition of the variable  $z$  is the particle multiplicity density  $\rho(s) = dN/d\eta|_{\eta=0}$ . We have simulated the multiplicity density [13] using the PYTHIA [14] in the energy range  $\sqrt{s} = 10 - 200$  GeV. The generated values were parameterized in the form  $\rho(s) = as^b$ , where  $a = 0.59 \pm 0.08$ ,  $b = 0.126 \pm 0.017$ . The energy dependence of  $\rho(s)$  in  $\pi^-p$  collisions is found to be similar as for the experimental measured multiplicity density in the nucleon-nucleon interactions. Therefore we use the experimental values in our analysis.

The second ingredient in the variable  $z$  is the anomalous fractal dimension of the colliding objects. For the nucleon-nucleon collisions  $\delta_1 = \delta_2 \equiv \delta_N$ . The value of  $\delta_N$  was determined in Ref. [10]. For pion-nucleon collisions, we deal with the asymmetric case  $\delta_1 \equiv \delta_\pi \neq \delta_N$ .

In our analysis we verify the hypothesis of the energy independence of data  $z$ -presentation for hadron production in  $\pi^-p$  collisions to determine  $\delta_\pi$ . We use the experimental data on cross sections for  $\pi^{\pm,0}, K^{\pm}$ , and  $\bar{p}$  hadrons produced in  $\pi^-p$  collisions obtained in Refs. [15, 16, 17, 18]. The measurements were made at pion momenta  $p_{lab} = 40, 200$ , and 300 GeV/c over the high transverse momentum range  $0.8 < p_T < 10$ . GeV/c. Note that the data demonstrate strong energy dependence of cross sections as function of the transverse momentum. The difference between the hadron yields increases with  $p_T$ . The energy dependence of the cross sections is contrasted with  $z$ -presentation of the data obtained for  $\delta_\pi = 0.1$ . The result confirms the hypothesis of existence of the energy independence of the  $z$ -scaling in pion-nucleon interactions. The results are illustrated in Figs. 1-3.

Nevertheless, we would like to note that last points of the spectrum for the incident pion momentum  $p_{lab} = 40$  GeV/c show some abundance of the hadron yields. From our point of view it means existence of considerable systematic errors of measurements at high transverse momentum in the  $\pi^-p$  collisions at this energy [18]. We do not observe such anomalies in  $pp$  interactions (Fig. 4).

## 4 Z-scaling in $\pi^-A$ collisions

Using the concept of  $z$ -scaling, we extend our study to hadron production in  $\pi^-A$  collisions. In this case we specify the scaling variable in analogy with its construction for the elementary pion-nucleon interaction. We determine the scale of the variable  $z$  in terms of the average multiplicity density of particles produced in the corresponding pion-nucleon interaction. The anomalous fractal dimensions of nuclei  $\delta_A$  are expressed via the nucleon fractal dimension  $\delta_N$  as follows [10]

$$\delta_A = A \cdot \delta_N. \quad (8)$$


The value of pion  $\delta_\pi$  is used the same as determined from the pion-nucleon interactions.

The experimental data on cross sections [15, 19, 20] have been analysed. The data are shown as function of  $p_T$  and  $z$  in Figs. 5-7.  $Z$ -presentation of the data manifests clear energy independence in the whole region of the transverse momentum considered. The scaling functions for different nuclei  $Be, Cu$ , and  $W$  are within the errors the same. The result is a confirmation of validity of the  $z$ -scaling in particle production in pion-nucleus collisions. The similar properties are observed for hadron production in proton-nucleus collisions, as well (Fig.8).


As seen from Fig.6, the scaling function manifests a power behaviour in the region of large values of  $z$

$$\psi(z) \sim z^{-\beta}. \quad (9)$$

From our point of view, such asymptotic regime reflects fractal character of particle formation. The fractality is connected with definition of the scaling variable  $z$  which has character of a fractal measure. The measure tends to infinity as the resolution increases and does not violate the power law for any value of  $z$ .




a)




b)

**Figure 1.** (a) Dependence of the inclusive cross section of  $\pi^+$ -meson production on transverse momentum  $q_T$  at  $p_{lab} = 40, 200,$  and  $300$  GeV/c and  $\theta_{cm}^{\pi p} \simeq 90^\circ$  in  $\pi^-p$  collisions. Experimental data are taken from [15, 18]. (b) The corresponding scaling function  $\psi(z)$ .




a)




b)

**Figure 2.** (a) Dependence of the inclusive cross section of  $\pi^0$ -meson production on transverse momentum  $q_T$  at  $p_{lab} = 100, 200,$  and  $300$  GeV/c and  $\theta_{cm}^{\pi p} \simeq 90^\circ$  in  $\pi^-p$  collisions. Experimental data are taken from [16, 17]. (b) The corresponding scaling function  $\psi(z)$ .




a)




b)

**Figure 3.** (a) Dependence of the inclusive cross section of  $K^+$ -meson production on transverse momentum  $q_T$  at  $p_{lab} = 40, 200,$  and  $300$  GeV/c and  $\theta_{cm}^{\pi p} \simeq 90^\circ$  in  $\pi^- p$  collisions. Experimental data are taken from [15, 18]. (b) The corresponding scaling function  $\psi(z)$ .




a)




b)

**Figure 4.** (a) Dependence of the inclusive cross section of  $K^+$ -meson production on transverse momentum  $q_T$  at  $p_{lab} = 70, 200, 300,$  and  $400$  GeV/c and  $\theta_{cm}^{pp} \simeq 90^\circ$  in  $pp$  collisions. Experimental data are taken from [21, 22]. The solid lines are obtained by fitting of the data at  $p_{lab} = 70$  and  $400$  GeV/c. (b) The corresponding scaling function  $\psi(z)$ .




a)




b)

**Figure 5.** (a) Dependence of the inclusive cross section of  $\pi^+$ -meson production on transverse momentum  $q_T$  in  $\pi^-A$  collisions at  $p_{lab} = 200$  and  $300$  GeV/c. Experimental data are taken from [15]. (b) The corresponding scaling function  $\psi(z)$ .




a)




b)

**Figure 6.** (a) Dependence of the inclusive cross section of  $\pi^0$ -meson production on transverse momentum  $q_T$  in  $\pi^-A$  collisions at  $\sqrt{s} = 31$  GeV. Experimental data are taken from [19, 20]. (b) The corresponding scaling function  $\psi(z)$ .




a)




b)

**Figure 7.** (a) Dependence of the inclusive cross section of  $K^+$ -meson production on transverse momentum  $q_T$  in  $\pi^- A$  collisions at  $p_{lab} = 200$  and  $300$  GeV/c. Experimental data are taken from [15]. (b) The corresponding scaling function  $\psi(z)$ .



a)



b)

**Figure 8.** (a) Dependence of the inclusive cross section of  $K^+$ -meson production on transverse momentum  $q_T$  in  $pA$  collisions at  $p_{lab} = 200, 300$  and  $400$  GeV/c. Experimental data are taken from [21, 23]. (b) The corresponding scaling function  $\psi(z)$ .



## 5 Conclusions

The  $z$ -presentation of experimental data on cross sections of hadron production in  $\pi^-p$  and  $\pi^-A$  collisions was constructed. The anomalous fractal dimension of pion,  $\delta_\pi = 0.1$ , allows to reproduce the general properties of the  $z$ -scaling established in  $pp$  and  $pA$  interactions. The scaling function  $\psi(z)$  demonstrates energy independence in a wide region of the colliding energy and the transverse momentum. Universal character of  $\psi(z)$  for different nuclei was found. For sufficient large values of  $z$ , the scaling function is governed by a power law.

The  $z$ -scaling confirmed in  $\pi^-p$  and  $\pi^-A$  collisions reflects general properties of particle production such as self-similarity, locality, and fractality. The scaling function  $\psi(z)$  is interpreted as the probability density to form a particle with the formation length  $z$ . The scaling can serve as an effective tool in searching for new physical phenomena in pion induced reactions at small scales.

**Acknowledgments.** This work has been partially supported by the grant No. 020475 of the Czech Ministry of Education, Youth and Physical Training.

## References

- [1] R.P. Feynman, *Phys. Rev. Lett.* **23**, 1415 (1969).
- [2] J.D. Bjorken, *Phys. Rev.* **179**, 1547 (1969); J.D. Bjorken and E.A. Paschos, *Phys. Rev.* **185**, 1975 (1969).
- [3] P. Bosted *et al.*, *Phys. Rev. Lett.* **49**, 1380 (1972).
- [4] J. Benecke *et al.*, *Phys. Rev.* **188**, 2159 (1969).
- [5] A.M. Baldin, *Part. Nuclei* **8**, 429 (1977).
- [6] V.S. Stavinsky, *Physics of Elementary Particles and Atomic Nuclei* **10**, 949 (1979).
- [7] G.A. Leksin, Report No. ITEF-147, 1976; G.A. Leksin, in *Proceedings of the XVIII International Conference on High Energy Physics*, Tbilisi, Georgia, 1976, edited by N.N. Bogolubov *et al.*, (JINR Report No. D1,2-10400, Tbilisi, 1977), p. A6-3.
- [8] Z. Koba, H.B. Nielsen, and P. Olesen, *Nucl. Phys.* **B40**, 317 (1972).
- [9] V.A. Matveev, R.M. Muradyan, and A.N. Tavkhelidze, *Part. Nuclei* **2**, 7 (1971); *Lett. Nuovo Cimento* **5**, 907 (1972); *Lett. Nuovo Cimento* **7**, 719 (1973); S. Brodsky, and G. Farrar, *Phys. Rev. Lett.* **31**, 1153 (1973); *Phys. Rev.* **D11**, 1309 (1975).
- [10] I.Zborovský, M.V.Tokarev, Yu.A.Panebratsev, and G.P.Škoro, *Phys. Rev.* **C59**, 2227 (1999); M.V.Tokarev, T.G.Dedovich, *Int. J. Mod. Phys.* **A15** (2000) 3495; M.V.Tokarev, O.V.Rogachevski, T.G.Dedovich, *J. Phys. G: Nucl. Part. Phys.* **26** (2000) 1671; M.Tokarev, I.Zborovský, Yu.Panebratsev, G.Skoro, *Int. J. Mod. Phys.* **A16** (2001) 1281.
- [11] B. Mandelbrot, *The Fractal Geometry of Nature* (Freeman, San Francisco, 1982).

- [12] L.Nottale, *Fractal Space-Time and Microphysics*. (World Scientific Publishing Co.Pte. Ltd. 1993).
- [13] G. Skoro, private communication.
- [14] T. Sjostrand, *Computer Physics Communications*, **82**, 74 (1994).
- [15] N. D. Giokaris et al., *Phys. Rev. Lett.* **47**, 1690 (1981);  
H. J. Frisch et al., *Phys. Rev.* **D27**, 1001 (1983).
- [16] G. J. Donaldson et al., *Phys. Rev. Lett.* **36**, 1110 (1976); *Phys. Rev. Lett.* **40**, 917 (1978);  
*Phys. Lett.* **B73**, 375 (1978).
- [17] C. DeMarzo et al., *Phys. Rev.* **D36**, 16 (1987).
- [18] L. K. Turchanovich et al., *Yad.Fiz.* **56(10)**, 116 (1993).
- [19] G. Alverson et al., *Phys. Rev.* **D48**, 5 (1993).
- [20] L. Apanasevich et al. *Phys. Rev. Lett.* **81**, 2642 (1998).
- [21] J.W. Cronin et al., *Phys. Rev.* **D11**, 3105 (1975); D. Antreasyan et al., *Phys. Rev.* **D19**,  
764 (1979).
- [22] V.V. Abramov et al., *Sov. J. Nucl. Phys.* **41**, 700 (1985).
- [23] V.V. Abramov et al., *Sov. J. Nucl. Phys.* **41**, 357 (1985).

Received on November 4, 2002.

Токарев М. В. и др.

E2-2002-251

Самоподобие рождения адронов с большими поперечными импульсами в  $\pi^-p$ - и  $\pi^-A$ -взаимодействиях

Изучаются свойства самоподобия рождения адронов с большими поперечными импульсами в  $\pi^-p$ - и  $\pi^-A$ -взаимодействиях. Проведен анализ экспериментальных данных в рамках концепции  $z$ -скейлинга. Определена аномальная фрактальная размерность для налетающего пиона  $\delta_\pi \approx 0,1$ . Установлена энергетическая независимость  $z$ -представления экспериментальных данных. Анализ  $A$ -зависимости  $z$ -скейлинга показал, что формирование адронов в  $\pi^-A$ -взаимодействиях обладает свойством самоподобия.

Работа выполнена в Лаборатории высоких энергий им. В. И. Векслера и А. М. Балдина ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 2002

Tokarev M. V. et al.

E2-2002-251

Self-Similarity of High- $p_T$  Hadron Production  
in  $\pi^-p$  and  $\pi^-A$  Collisions

Self-similar properties of hadron production in  $\pi^-p$  and  $\pi^-A$  collisions over a high- $p_T$  region are studied. The analysis of experimental data is performed in the framework of  $z$ -scaling. The scaling variable depends on the anomalous fractal dimension of the incoming pion. Its value is found to be  $\delta_\pi = 0.1$ . Independence of the scaling function  $\psi(z)$  on the collision energy is shown.  $A$ -dependence of data  $z$ -presentation confirms self-similarity of particle formation in  $\pi^-A$  collisions.

The investigation has been performed at the Veksler-Baldin Laboratory of High Energies, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 2002

Макет *Т. Е. Понeko*

Подписано в печать 24.12.2002.

Формат 60 × 90/16. Бумага офсетная. Печать офсетная.

Усл. печ. л. 0,81. Уч.-изд. л. 1,41. Тираж 415 экз. Заказ № 53678.

Издательский отдел Объединенного института ядерных исследований  
141980, г. Дубна, Московская обл., ул. Жолио-Кюри, 6.

E-mail: [publish@pds.jinr.ru](mailto:publish@pds.jinr.ru)

[www.jinr.ru/publish/](http://www.jinr.ru/publish/)