




XJ0300070



ОБЪЕДИНЕННЫЙ  
ИНСТИТУТ  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ

Дубна

E6-2002-201

K. Ya. Gromov, S. A. Kudrya\*, Yu. V. Narseev,  
Zh. K. Samatov, V. A. Sergienko\*, V. I. Fominykh,  
V. G. Chumin

ON THE  $^{221}\text{Rn} \rightarrow ^{221}\text{Fr}$  DECAY SCHEME

Submitted to «Известия РАН, серия физическая»

---

\*St. Petersburg State University, Russia

Experimental information on the structure of the  $^{221}\text{Fr}$  levels can be gained from investigation of either the  $^{225}\text{Ac}$  ( $T_{1/2} = 10$  d)  $\alpha$ -decay or the  $^{221}\text{Rn}$  ( $T_{1/2} = 25$  min.)  $\beta^-$ -decay. In the latter case there arise difficulties with preparing  $^{221}\text{Rn}$  sources: radon is a noble gas and the only way to produce sufficiently strong  $^{221}\text{Rn}$  sources is via the reaction of spallation of thorium by protons yielding not only  $^{221}\text{Rn}$  but also other radon isotopes. The most comprehensive investigation of  $^{221}\text{Rn}$  sources was performed by Vylov et al. [1]. Data on the  $\gamma$ -spectrum, conversion electron spectrum, and  $\alpha$ -spectrum were gained. A decay scheme was proposed. However, coincidence experiments could not be carried out at that time.

In [2] the  $\alpha$ -decay  $^{225}\text{Ac} \rightarrow ^{221}\text{Fr}$  was investigated. Qualitative analysis was given to  $(\alpha-\gamma)$ -coincidences to prove belonging of  $\gamma$ -transitions to the  $^{225}\text{Ac}$  -decay. The  $^{225}\text{Ac}$  -decay scheme was proposed. The results of the investigation also allow  $^{221}\text{Rn}$  decay scheme proposed in [1] to be somewhat refined relying on, first, the proof [2] of existence of the relevant levels in  $^{221}\text{Fr}$  and, secondly, the relative intensities of  $\gamma$ -rays at the de-excitation of these levels. The  $^{221}\text{Rn} \rightarrow ^{221}\text{Fr}$  decay scheme proposed by us on the basis of an analysis like that is displayed in Fig. 1. It shows all  $\gamma$ -transitions found in [2] at de-excitation of a level. Transitions observed in [1] are marked with an asterisk. A total intensity, including conversion, is given for each  $\gamma$ -transition. Intensities of  $\gamma$ -transitions that were not observed in [1] are calculated with relative intensities of  $\gamma$ -rays from each level given in [2].

Vylov et al. [1] assume that the  $^{221}\text{Rn}$  decay may excite the 393.2-keV level in  $^{221}\text{Fr}$ , which is then de-excited by the 119.9, 168.9, and 197.8-keV transitions. Seven transition from this level are found in the  $^{225}\text{Ac}$  decay (see Fig. 1). Intensity ratios of the 168.9-keV and 197.8-keV  $\gamma$ -rays in [1] and [2] coincide within the error. Therefore, we believe that the 393.2-keV level in  $^{221}\text{Fr}$  is excited at the  $^{221}\text{Rn}$  decay. Intensities of the 114, 139.6, 243.2, 284.5, and 354.9-keV  $\gamma$ -rays are lower than the intensity of the 197.8-keV  $\gamma$ -transition, which explains their absence in [1]. Placing the 119.9-keV  $\gamma$ -transition between the 393.9 and 273.5-keV levels [1] is in conflict with the results [2]. The 273.5-keV level is introduced in [2] on the basis of coincidence of 236-keV  $\gamma$ -rays with  $E_\alpha = 5563$  keV  $\alpha$ -particles. No 273.5-keV  $\gamma$ -rays were observed in [2, 3]. Therefore we believe that

273.5-keV  $\gamma$ -rays in [1] are due to presence of other isotopes in the  $^{221}\text{Rn}$  source. Thus, we think that in [1] they did not have grounds for introducing the 273.5-keV level at the  $^{221}\text{Rn}$  decay and thus for placing the 119.9-keV  $\gamma$ -transition between the 393.9 and 273.5-keV levels.

Intensity ratios of  $\gamma$ -rays from de-excitation of the 294.6, 279.2, 253.5, 234.5, 224.6, 195.8, and 150 keV in [1] and [2] coincide within the tolerable error. Therefore, we believe that excitation of the above levels at the  $^{221}\text{Rn}$  decay is justified. The  $^{221}\text{Fr}$  level at 288.1 keV excited at the  $^{225}\text{Ac}$   $\alpha$ -decay is not found in [1] at the  $^{221}\text{Rn}$  decay.

Gromov et al. [4] and Ardisson et al. [3] assumed excitation of the  $^{221}\text{Fr}$  level at 145.9 keV at the  $^{225}\text{Ac}$  decay. Investigation of the ( $\alpha$ - $\gamma$ )-coincidences [2] confirm existence of a  $^{221}\text{Fr}$  level at this energy. The spectrum of  $\alpha$ -particles coinciding with the 119.9-keV  $\gamma$ -rays is displayed in Fig. 2. Peaks with  $E_\alpha = (5686 \pm 15)$ ,  $(5609 \pm 15)$ , and  $(5443 \pm 15)$  keV are observed in this spectrum. Energies of these lines coincide with energies of  $\alpha$ -particles populating levels at 145.9, 224.6, and 393.2 keV. Coincidences with  $E_\alpha = (5609 \pm 15)$  keV are due to the fact that the 145.9-keV level is populated by 78.8-keV  $\gamma$ -rays at de-excitation of the 224.6-keV level more intensely than by the  $\alpha$ -decay to the 145.9-keV level. The area of the  $E_\alpha = (5443 \pm 15)$  keV peak is small, about 2.5% area of all peaks in Fig. 2. Coincidences of 119.9-keV  $\gamma$ -quanta with  $(5443 \pm 15)$ -keV  $\alpha$ -particles is due to population of the 145.9-keV level by a cascade of the 78.8 and 169.2-keV  $\gamma$ -transitions from the 393.2-keV level through the 224.6-keV level. The intensity ratio of the 119.9 and 46.2-keV transitions is 17 in [2]; that is why 46.2-keV  $\gamma$ -rays were not observed in [1]. Thus, we consider introduction of the 145.9-keV level in the  $^{221}\text{Rn}$  decay scheme to be justified.

Intensity ratios of  $\gamma$ -rays at de-excitation of levels at energies ranging from 25.9 to 108.3 keV in [1] agree with the data in [2] within the tolerable error, i.e., all these levels are excited at the  $^{221}\text{Rn}$  decay and there is no contradiction in their properties.

Intensities of  $\beta^-$ -transitions to  $^{221}\text{Fr}$  level are found from the intensities of  $\gamma$ -transitions at the  $^{221}\text{Rn}$  decay and reduced probabilities  $\text{lgft}$  of these  $\beta^-$ -transitions are calculated [5]. The value  $Q_{\beta^-}$  ( $^{221}\text{Rn}$ ) = 1130(100) keV [6] was used in this calculation. The values of  $I_{\beta^-}$  and  $\text{lgft}$  are shown in Fig. 1.

Levels at 108.3 keV  $(7/2)^-$  and 294.6 keV  $(9/2)^+$  are most intensely populated at the  $^{221}\text{Rn}$  decay, the respective intensities being  $I_{\beta^-} = 10.3(34)$  and  $25(2)$ . To the 108.3-keV level there proceeds a once-forbidden





Fig. 1.  $^{221}\text{Rn}$  decay scheme. Above  $\gamma$ -transitions there are their energies, total (including conversion) intensities (in parentheses), and experimental multipolarity data. An asterisk (\*) marks  $\gamma$ -transitions observed in [1]. Next to the level energies are spins and parities [2], intensities and reduced  $\beta$ -decay probabilities.



**Fig. 2.** Intensity of 119.9-keV  $\gamma$ -ray in  $(\alpha\text{-}\gamma)$ -coincidence experiment as a function of the  $\alpha$ -particle energy.

$\beta^-$ -transition. A  $\beta^-$ -transition to the 294.6-keV level is allowed. The  $^{221}\text{Rn}$  ground state spin is  $7/2$  [7]. Reduced probabilities for transitions to these levels allows a conclusion that the  $^{221}\text{Rn}$  ground state parity is most probably positive. The reduced probabilities for transitions to other  $^{221}\text{Fr}$  levels are not in contradiction with the assumption of the positive parity of the  $^{221}\text{Rn}$  ground state.

Thus, in addition to those in [1], the 393.2-keV  $(7/2, 5/2)^+$  and 145.9-keV  $(1/2)^+$  levels are introduced in the  $^{221}\text{Fr}$  level scheme. The assumption [1] of excitation of the 273.5-keV level at the  $^{221}\text{Rn}$  decay is shown to contradict the results of investigation of  $(\alpha-\gamma)$ -coincidences at the  $^{225}\text{Ac}$  decay [2]. The  $^{221}\text{Rn}$  ground state parity appears to be positive.

#### Reference

1. Ts. Vylov, N.A. Golovkov, B.S. Dzhelepov et al. // *Izv. AN SSSR, ser.fiz.* 1977. V.41. P.1634.
2. S.A. Kudrya, V.M. Gorozhankin, K.Ya. Gromov et al. // *Izv. RAN, ser.fiz.* 2003. V.67. Nom. 1
3. G. Ardisson, R.K. Sheline et al. *Phys. Rev. C*, //V.62, 064306. 2000.
4. K.Ya. Gromov, M.Ya. Kuznetsova, Yu.V. Norseev et al. // *Izv. RAN, ser.fiz.* 1994. V.58. P.35.
5. B.S. Dzhelepov, L.N. Zyryanova, Yu.P. Suslov. // *Beta-processy. Nauka. M.-L.* 1972.
6. G. Audi, O. Bersillou, J. Blachot et al.// *Nucl. Phys.* A624. 1977. P.1
7. AIP Conf. Proceedings 164 , 5-th International Conference, "Nuclei far from stability", Rosseau Lake, Ontario, Canada. 1987. P.126.

Received on August 28, 2002.

Громов К. Я. и др.  
О схеме распада  $^{221}\text{Rn} \rightarrow ^{221}\text{Fr}$

E6-2002-201

Сравниваются результаты исследований  $\beta^-$ -распада  $^{221}\text{Rn}$  и  $\alpha$ -распада  $^{225}\text{Ac}$ . Показано, что при распаде  $^{221}\text{Rn}$  возбуждаются уровни  $^{221}\text{Fr}$  с энергиями 145,9 и 393,2 кэВ. Определены интенсивности и приведенные вероятности  $\beta^-$ -распада на уровни  $^{221}\text{Fr}$ . Сделано заключение о положительной четности основного состояния  $^{221}\text{Rn}$ .

Работа выполнена в Лаборатории ядерных проблем им. В. П. Дзелепова ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 2002

Gromov K. Ya. et al.  
On the  $^{221}\text{Rn} \rightarrow ^{221}\text{Fr}$  Decay Scheme

E6-2002-201

The results of investigating the  $^{221}\text{Rn}$   $\beta^-$ -decay and the  $^{225}\text{Ac}$   $\alpha$ -decay are compared. It is shown that  $^{221}\text{Fr}$  levels at 145.9 and 393.2 keV are excited at the  $^{221}\text{Rn}$  decay. Intensities and reduced probabilities of the  $\beta^-$ -decay to the  $^{221}\text{Fr}$  levels are determined. A conclusion is drawn that the parity of the  $^{221}\text{Rn}$  ground state is positive.

The investigation has been performed at the Dzhelepov Laboratory of Nuclear Problems, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 2002

*Макет Т. Е. Попеко*

Подписано в печать 04.10.2002.

Формат 60 × 90/16. Бумага офсетная. Печать офсетная.

Усл. печ. л. 0,5. Уч.-изд. л. 0,57. Тираж 310 экз. Заказ № 53547.

Издательский отдел Объединенного института ядерных исследований  
141980, г. Дубна, Московская обл., ул. Жолио-Кюри, 6.