

# TABLE OF CONTENTS

|  |        |
|--|--------|
| Foreword   | xxvii  |
| List of Acronyms   | xxix   |
| List of Affiliations   | xxxii  |
| Acknowledgements   | xxxiii |
| <b>PART I: INTRODUCTION</b>  |        |
| Chapter 1 Health Systems Performance Assessment: Goals, Framework and Overview<br><i>Christopher J.L. Murray, David B. Evans</i>                           | 3      |
| <b>PART II: REGIONAL PERSPECTIVES ON PERFORMANCE ASSESSMENT</b>  |        |
| Chapter 2 African Regional Consultation on Health Systems Performance Assessment<br><i>WHO Regional Office for Africa</i>                                  | 23     |
| Chapter 3 American Regional Consultation on Health Systems Performance Assessment<br><i>WHO Regional Office for the Americas</i>                           | 27     |
| Chapter 4 Eastern Mediterranean Regional Consultation on Health Systems Performance Assessment<br><i>WHO Regional Office for the Eastern Mediterranean</i> | 35     |
| Chapter 5 European Regional Consultation on Health Systems Performance Assessment<br><i>WHO Regional Office for Europe</i>                                 | 55     |
| Chapter 6 South-East Asian Regional Consultation on Health Systems Performance Assessment<br><i>WHO Regional Office for South-East Asia</i>                | 69     |
| Chapter 7 Western Pacific Regional Consultation on Health Systems Performance Assessment<br><i>WHO Regional Office for the Western Pacific</i>             | 95     |
| <b>PART III: EXPERT CONSULTATION REPORTS</b>   |        |
| Chapter 8 Technical Consultation on Measurement of the Efficiency of Health Systems  | 109    |
| Chapter 9 Technical Consultation on Concepts and Methods for Measuring the Responsiveness of Health Systems  | 115    |
| Chapter 10 Technical Consultation on Effective Coverage in Health Systems  | 125    |
| Chapter 11 Technical Consultation on Stewardship   | 135    |
| Chapter 12 Technical Consultation on Statistical Strategies for Cross-Population Comparability   | 141    |
| Chapter 13 Technical Consultation on Fairness in Financial Contribution  | 147    |
| Chapter 14 Technical Consultation on the Measurement of Health Inequalities  | 155    |
| Chapter 15 WHO Meetings of Experts on Measuring and Summarizing Health   | 163    |

## PART IV: METHODS AND EMPIRICISM

|                        |   |     |
|------------------------|---|-----|
| INPUTS                 |   |     |
| Chapter 16             | National Health Accounts: Concepts, Data Sources, and Methodology<br><i>Jean-Pierre Poullier, Patricia Hernandez, Kei Kawabata</i>  | 185 |
| Chapter 17             | Patterns of Global Health Expenditures: Results for 191 Countries<br><i>Jean-Pierre Poullier, Patricia Hernandez, Kei Kawabata, William D. Savedoff</i>   | 195 |
| FINANCING              |   |     |
| Chapter 18             | Monitoring the Health Financing Function<br><i>William D. Savedoff, Guy Carrin, Kei Kawabata, Abdelhay Mechbal</i>  | 205 |
| Chapter 19             | Developing Health Financing Policies<br><i>William D. Savedoff, Guy Carrin</i>  | 211 |
| PROVISION AND COVERAGE |   |     |
| Chapter 20             | Beyond Access and Utilization: Defining and Measuring Health System Coverage<br><i>Bakhuti Shengelia, Christopher J.L. Murray, Orvill B. Adams</i>  | 221 |
| Chapter 21             | Provision of Personal and Non-personal Health Services: Proposal for Monitoring<br><i>Orvill B. Adams, Bakhuti Shengelia, Barbara Stilwell, Itziar Larizgoitia, Andrei Issakov, Sylvester Y. Kwankam, Ferdinand Siem Tjam</i>           | 235 |
| Chapter 22             | Inequalities in Coverage: Valid DTP3 and Measles Vaccination in 40 Countries<br><i>Saba Moussavi, Bakhuti Shengelia, Ajay Tandon, Neeru Gupta, Christopher J.L. Murray</i>  | 251 |
| Chapter 23             | Validity of Reported Vaccination Coverage in 45 Countries<br><i>Christopher J.L. Murray, Bakhuti Shengelia, Neeru Gupta, Saba Moussavi, Ajay Tandon, Michel Thieren</i>   | 265 |
| RESOURCE GENERATION    |   |     |
| Chapter 24             | Human, Physical, and Intellectual Resource Generation: Proposals for Monitoring<br><i>Orvill B. Adams, Mario R. Dal Poz, Bakhuti Shengelia, Sylvester Y. Kwankam, Andrei Issakov, Barbara Stilwell, Pascal Zurn, Alexandre Goubarev</i> | 273 |
| STEWARDSHIP            |   |     |
| Chapter 25             | Towards Better Stewardship: Concepts and Critical Issues<br><i>Phyllida Travis, Dominique Egger, Philip Davies, Abdelhay Mechbal</i>  | 289 |
| POPULATION HEALTH      |   |     |
| Chapter 26             | Quantifying Individual Levels of Health: Definitions, Concepts, and Measurement Issues<br><i>Joshua A. Salomon, Colin D. Mathers, Somnath Chatterji, Ritu Sadana, T. Bedirhan Üstün, Christopher J.L. Murray</i>                        | 301 |
| Chapter 27             | Alternative Summary Measures of Average Population Health<br><i>Colin D. Mathers, Joshua A. Salomon, Christopher J.L. Murray, Alan D. Lopez</i>   | 319 |
| Chapter 28             | Life Tables for 191 Countries for 2000: Data, Methods, Results<br><i>Alan D. Lopez, Omar B. Ahmad, Michel Guillot, Brodie D. Ferguson, Joshua A. Salomon, Christopher J.L. Murray, Kenneth H. Hill</i>                                  | 335 |

|                                    |   |     |
|------------------------------------|---|-----|
| Chapter 29                         | Modified Logit Life Table System: Principles, Empirical Validation, and Application<br><i>Christopher J.L. Murray, Brodie D. Ferguson, Alan D. Lopez, Michel Guillot, Joshua A. Salomon, Omar B. Ahmad</i>  | 355 |
| Chapter 30                         | Empirical Evaluation of the Anchoring Vignette Approach in Health Surveys<br><i>Christopher J.L. Murray, Emre Özaltın, Ajay Tandon, Joshua A. Salomon, Ritu Sadana, Somnath Chatterji</i>   | 369 |
| Chapter 31                         | Unpacking Health Perceptions Using Anchoring Vignettes<br><i>Joshua A. Salomon, Ajay Tandon, Christopher J.L. Murray, World Health Survey Pilot Study Collaborating Groups</i>  | 401 |
| Chapter 32                         | Health State Valuations in Summary Measures of Population Health<br><i>Joshua A. Salomon, Christopher J.L. Murray, T. Bedirhan Üstün, Somnath Chatterji</i>   | 409 |
| Chapter 33                         | Methods for Measuring Healthy Life Expectancy<br><i>Colin D. Mathers, Christopher J.L. Murray, Joshua A. Salomon</i>  | 437 |
| HEALTH INEQUALITY                  |   |     |
| Chapter 34                         | A Framework for Measuring Health Inequality<br><i>Emmanuela Gakidou, Christopher J.L. Murray, Julio Frenk</i>   | 471 |
| Chapter 35                         | Measuring Total Health Inequality: Adding Individual Variation to Group-Level Differences<br><i>Emmanuela Gakidou, Gary King</i>  | 485 |
| Chapter 36                         | Determinants of Inequality in Child Survival: Results from 39 Countries<br><i>Emmanuela Gakidou, Gary King</i>  | 497 |
| Chapter 37                         | Measurement Methods for Inequality in the Risk of Adult Mortality<br><i>Emmanuela Gakidou, Ajay Tandon</i>  | 503 |
| FAIRNESS IN FINANCIAL CONTRIBUTION |   |     |
| Chapter 38                         | Assessing the Distribution of Household Financial Contributions to the Health System: Concepts and Empirical Application<br><i>Christopher J.L. Murray, Ke Xu, Jan Klavus, Kei Kawabata, Piya Hanvoravongchai, Riadh Zeramdini, Ana Mylena Aguilar-Rivera, David B. Evans</i> | 513 |
| Chapter 39                         | Household Health System Contributions and Capacity to Pay: Definitional, Empirical, and Technical Challenges<br><i>Ke Xu, Jan Klavus, Kei Kawabata, David B. Evans, Piya Hanvoravongchai, Juan Pablo Ortiz, Riadh Zeramdini, Christopher J.L. Murray</i>                      | 533 |
| Chapter 40                         | Summary Measures of the Distribution of Household Financial Contributions to Health<br><i>Ke Xu, Jan Klavus, Ana Mylena Aguilar-Rivera, Guy Carrin, Riadh Zeramdini, Christopher J.L. Murray</i>  | 543 |
| Chapter 41                         | The Impact of Vertical and Horizontal Inequality on the Fairness in Financial Contribution Index<br><i>Ke Xu, Jan Klavus, David B. Evans, Piya Hanvoravongchai, Riadh Zeramdini, Christopher J.L. Murray</i>  | 557 |

|                        |   |     |
|------------------------|---|-----|
| Chapter 42             | Understanding Household Catastrophic Health Expenditures: a Multi-country Analysis<br><i>Ke Xu, David B. Evans, Kei Kawabata, Riadh Zeramdini, Jan Klavus, Christopher J.L. Murray</i>  | 565 |
| RESPONSIVENESS         |   |     |
| Chapter 43             | Health System Responsiveness: Concepts, Domains and Operationalization<br><i>Nicole B. Valentine, Amala de Silva, Kei Kawabata, Charles Darby, Christopher J.L. Murray, David B. Evans</i>  | 573 |
| Chapter 44             | Classical Psychometric Assessment of the Responsiveness Instrument in the WHO Multi-country Survey Study on Health and Responsiveness 2000–2001<br><i>Nicole B. Valentine, René Lavallée, Bao Liu, Gouke J. Bonsel, Christopher J.L. Murray</i> | 597 |
| Chapter 45             | Weights for Responsiveness Domains: Analysis of Country Variation in 65 National Sample Surveys<br><i>Nicole B. Valentine, Joshua A. Salomon</i>  | 631 |
| Chapter 46             | Patient Experiences with Health Services: Population Surveys from 16 OECD Countries<br><i>Nicole B. Valentine, Juan Pablo Ortiz, Ajay Tandon, Kei Kawabata, David B. Evans, Christopher J.L. Murray</i>   | 643 |
| Chapter 47             | Inequality in Responsiveness: Population Surveys from 16 OECD Countries<br><i>Juan Pablo Ortiz, Nicole B. Valentine, Emmanuela Gakidou, Ajay Tandon, Kei Kawabata, David B. Evans, Christopher J.L. Murray</i>                                  | 653 |
| COMPOSITE ATTAINMENT   |   |     |
| Chapter 48             | Quality and Equity: Preferences for Health System Outcomes<br><i>Emmanuela Gakidou, Christopher J.L. Murray, David B. Evans</i>   | 667 |
| Chapter 49             | Measuring Health System Attainment: the Impact of Variability in the Importance of Social Goals<br><i>Jeremy A. Lauer, David B. Evans, Christopher J.L. Murray</i>  | 677 |
| EFFICIENCY             |   |     |
| Chapter 50             | Health System Efficiency: Concepts<br><i>Ajay Tandon, Jeremy A. Lauer, David B. Evans, Christopher J.L. Murray</i>  | 683 |
| Chapter 51             | Determinants of Health System Performance: Second-Stage Efficiency Analysis<br><i>David B. Evans, Jeremy A. Lauer, Ajay Tandon, Christopher J.L. Murray</i>   | 693 |
| Chapter 52             | Health System Efficiency: Time, Attribution, and Multiple Indicators<br><i>David B. Evans, Christopher J.L. Murray, Ajay Tandon</i>   | 699 |
| MEASUREMENT CHALLENGES |   |     |
| Chapter 53             | Cross-Population Comparability of Evidence for Health Policy<br><i>Christopher J.L. Murray, Ajay Tandon, Joshua A. Salomon, Colin D. Mathers, Ritu Sadana</i>   | 705 |
| Chapter 54             | Towards Evidence-Based Public Health<br><i>Christopher J.L. Murray, Colin D. Mathers, Joshua A. Salomon</i>   | 715 |

|   |   |     |
|---|---|-----|
| Chapter 55  | Statistical Models for Enhancing Cross-Population Comparability<br><i>Ajay Tandon, Christopher J.L. Murray, Joshua A. Salomon, Gary King</i>  | 727 |
| Chapter 56  | Estimating Permanent Income Using Indicator Variables<br><i>Brodie D. Ferguson, Ajay Tandon, Emmanuela Gakidou, Christopher J.L. Murray</i>   | 747 |
| Chapter 57  | WHO Multi-country Survey Study on Health and Responsiveness 2000–2001<br><i>T. Bedirhan Üstün, Somnath Chatterji, Maria Villanueva, Lydia Bendib, Can Çelik, Ritu Sadana, Nicole B. Valentine, Juan Pablo Ortiz, Ajay Tandon, Joshua A. Salomon, Yang Cao, Wan Jun Xie, Emre Özaltın, Colin D. Mathers, Christopher J.L. Murray</i> | 761 |
| Chapter 58  | The World Health Surveys<br><i>T. Bedirhan Üstün, Somnath Chatterji, Abdelhay Mechbal, Christopher J.L. Murray, WHS Collaborating Groups</i>  | 797 |
| INCREASING POLICY RELEVANCE   |   |     |
| Chapter 59  | Subnational Health Systems Performance Assessment: Objectives, Challenges and Strategies<br><i>Phyllida Travis, Abdelhay Mechbal, Ajay Tandon, Brodie D. Ferguson, Michel Thieren, Christopher J.L. Murray</i>  | 809 |
| Chapter 60  | WHO-CHOICE: CHOosing Interventions That Are Cost-Effective<br><i>Raymond C.W. Hutubessy, Rob M.P.M. Baltussen, Tessa Tan Torres–Edejer, David B. Evans, WHO-CHOICE Working Group</i>  | 823 |
| PART V: REPORT OF THE SCIENTIFIC PEER REVIEW GROUP ON HEALTH SYSTEMS PERFORMANCE ASSESSMENT |   |     |
| Chapter 61  | Report of the Scientific Peer Review Group on Health Systems Performance Assessment<br><i>Sudhir Anand, Walid Ammar, Timothy Evans, Toshihiko Hasegawa, Katarzyna Kissimova-Skarbek, Ana Langer, Adetokunbo O. Lucas, Lindiwe Makubalo, Alireza Marandi, Gregg Meyer, Andrew Podger, Peter Smith, Suwit Wibulpolprasert</i>         | 839 |
| Index   |   | 917 |

## LIST OF TABLES

|             |   |     |
|-------------|---|-----|
| Table 1.1   | Intrinsic goals to which the health system contributes  | 8   |
| Table 11.1  | Tasks of the stewardship function   | 136 |
| Table 13.1  | The dimensions of health system contributions   | 150 |
| Table 17.1  | Health spending by income groups and regions  | 196 |
| Table 17.2  | Share of total spending on health by type and region, 2000 (%)  | 198 |
| Table 17.3  | External resources, consumption, and health spending, selected countries, 2000 (I\$ per capita)                         | 201 |
| Table 18.1  | Indicators for the financing function   | 207 |
| Table 21.1  | Expenditure shares to salary and drugs and supplies in eight LAC countries  | 237 |
| Table 21.2  | Incentives  | 244 |
| Table 21.3  | Matrix for the assessment and monitoring of health service provision function   | 246 |
| Table 22.1  | Summary table of the four inequality indices for DTP3 and measles   | 255 |
| Table 22.2  | Correlation of inequalities indices for probability of being covered for DTP3 and measles (Spearman's rho)              | 257 |
| Table 22.3A | Measles: estimated covariates and random effect from probit model   | 258 |
| Table 22.3B | DTP3: estimated covariates and random effect from probit model  | 260 |
| Table 24.1  | Classification of human resources for health systems  | 277 |
| Table 24.2  | Flow of workers' remittances and its share in imports and exports of goods in selected labour exporting countries       | 280 |
| Table 24.3  | Admissions of non-UK qualified nurses to the nursing register, 1990–1998  | 280 |
| Table 27.1  | SMPH criteria met by various forms of health expectancies   | 323 |
| Table 27.2  | Different questions and time perspectives for describing individual health  | 326 |
| Table 28.1  | Mortality data sources (number of countries) for WHO subregions, 2000   | 335 |
| Table 28.2  | Mortality data sources (% of deaths covered) for WHO subregions, 2000   | 336 |
| Table 28.3  | Life expectancy at birth (years), both sexes combined, top 10 and bottom 10 countries, 2000                             | 343 |
| Table 28.4  | Countries with male life expectancy greater than 75.0 years, 2000   | 343 |
| Table 28.5  | Countries with female life expectancy greater than 80.0 years, 2000   | 343 |
| Table 28.6  | Difference in life expectancy at birth for all possible causes of death and causes excluding AIDS, by sex, 2000 (years) | 344 |
| Table 28.7  | Total deaths by sex, age and WHO subregion, 2000  | 346 |
| Table 28.8  | Life expectancy at birth (years) by WHO subregion, 2000   | 348 |
| Table 28.9  | Availability of vital registration data on mortality in the WHO database, 1980–2000 (as of 15 September 2001)           | 350 |
| Table 29.1  | Life tables comprising the empirical dataset  | 360 |

|             |   |     |
|-------------|---|-----|
| Table 29.2  | Characteristics of life tables comprising the empirical dataset   | 360 |
| Table 29.3  | Values of $\theta_x$ , $\gamma_x$ and $l_x$ standard, by sex  | 361 |
| Table 29.4  | Comparison of root mean square error of $\ln(m_x)$ of the Coale-Demeny, Brass and modified logit systems using the 30% life table subset                          | 366 |
| Table 29.5A | Results of regression of selected observed life table parameters on those predicted by the modified logit system (n=541)  | 367 |
| Table 29.5B | Results of regression of selected observed life table parameters on those predicted by the modified logit system, using 25th percentile values for males ages 65+ | 367 |
| Table 30.1  | Number of respondents for each set of vignettes by domain in 69 MCSS surveys  | 372 |
| Table 30.2  | Mobility vignettes as included in the WHO Multi-country Survey Study on Health and Responsiveness 2000–2001   | 372 |
| Table 30.3  | Consistent and near consistent orderings of vignettes by domain in 69 MCSS surveys  | 373 |
| Table 30.4  | Per cent consistent and near consistent orderings by domain and survey  | 375 |
| Table 30.5  | Benefit of the doubt rank order correlation coefficients for mobility and cognition, 69 surveys   | 377 |
| Table 30.6  | Average BDROCC (benefit of the doubt rank order correlation coefficient) by domain and survey   | 380 |
| Table 30.7  | Average BDROCC (benefit of the doubt rank order correlation coefficient) by education group for all domains   | 382 |
| Table 30.8  | Regression analysis of BDROCC (benefit of the doubt rank order correlation coefficient) as a function of age, sex, educational attainment, and survey mode        | 382 |
| Table 30.9  | Eighteen most commonly used vignette response patterns in 69 surveys among consistent raters  | 383 |
| Table 30.10 | Modal mobility vignette response patterns for consistent raters, pooled and across 69 surveys   | 384 |
| Table 30.11 | Distribution of respondents in 69 multi-country survey studies by 28 self-response and vignette response patterns   | 386 |
| Table 30.12 | Distribution of vignette-adjusted raw score (VARS) and self-response categories for mobility in 69 multi-country survey studies                                   | 387 |
| Table 30.13 | Vignette thetas and sigmas for mobility for full dataset and for BDROCC > 0.8 dataset   | 390 |
| Table 31.1  | Distribution of sample used in pilot study of health module for the World Health Survey by country, age, sex, and years of schooling                              | 403 |
| Table 32.1  | Characteristics of responses to different valuation methods   | 415 |
| Table 32.2  | Regression models for visual analogue scale values  | 417 |
| Table 32.3  | Characteristics of the study population   | 418 |
| Table 32.4  | Median categorical descriptions of each condition by domain   | 419 |
| Table 32.5  | Correlation coefficients for mean VAS scores across countries   | 420 |

|             |   |     |
|-------------|---|-----|
| Table 32.6  | Characteristics of the multi-method study population  | 421 |
| Table 32.7  | Spearman's rank order correlation coefficients for different valuation methods, within and between countries  | 422 |
| Table 32.8  | Mean, median and standard deviation of valuation results by method and condition  | 423 |
| Table 32.9  | Results from multi-method analyses  | 425 |
| Table 32.10 | Estimation of valuation function  | 427 |
| Table 32.11 | Implications of estimated valuation functions: changes in overall health decrements associated with changes in levels on individual domains   | 429 |
| Table 32.12 | Predictive validity of regression models  | 430 |
| Table 33.1  | Life expectancy (LE), healthy life expectancy (HALE), and lost healthy years as per cent of total LE (LHE%), at birth and at age 60, by sex and total, WHO regions and world, 2000        | 438 |
| Table 33.2  | Life expectancy (LE), healthy life expectancy (HALE), and lost healthy years (LHE) as per cent of total LE (HLE%), at birth and at age 60, by sex and total, by mortality subregion, 1999 | 440 |
| Table 33.3  | Overall prevalence YLD rates (%) for WHO epidemiological subregions for the year 2000   | 446 |
| Table 33.4  | Cause-specific methods used for estimation of country-level prevalence YLD  | 448 |
| Table 33.5  | Average per cent of total prevalence YLD estimated by every method, for WHO Member States within each epidemiological subregion, GBD 2000 Version 1                                       | 449 |
| Table 33.6  | Major cause groups for which independent comorbidity assumed  | 450 |
| Table 33.7  | Estimation of uncertainty in GBD prior estimates of comorbidity-adjusted prevalences by age and sex for WHO Member States in the year 2000  | 452 |
| Table 33.8  | Estimated correlation matrix for age-specific uncertainty distributions for prevalence YLD estimates for the AFRO and SEARO D regions, year 2000 estimates                                | 453 |
| Table 33.9  | Population surveys conducted using WHO survey instrument 1999–2000  | 454 |
| Table 35.1  | DHS survey year and sample size   | 486 |
| Table 35.2  | Child survival inequality index for 50 countries, estimates and 95% confidence intervals  | 489 |
| Table 35.3  | Relative ranks of child survival inequality by four measures of inequality  | 490 |
| Table 36.1  | Demographic and Health Surveys used in this analysis: country name, three-letter acronym, survey year, and sample size  | 498 |
| Table 37.1  | Some existing individual-level datasets   | 504 |
| Table 37.2  | Predicted mean and median hazard rate, interquartile range, standard deviation, and coefficient of variation, by age and sex, USA   | 507 |
| Table 37.3  | Expected death rates and estimated standard deviation of the expected death rate across small areas in the USA  | 509 |
| Table 37.4  | Standard deviation of estimated risk of death across individuals from two analyses: full survival analysis model with random effect and small-area analysis                               | 510 |



|             |  |     |
|-------------|--|-----|
| Table 38.1  | Main indicators used in the income and burden approaches to analysing the consequences of household health system payments                               | 513 |
| Table 38.2  | Redistributive effect (RE) and poverty headcount difference (DH)   | 521 |
| Table 38.3  | Fairness in financial contribution and catastrophic payments   | 525 |
| Table 38.4  | Correlation coefficients between measures in the income and burden space   | 527 |
| Table 40.1  | Standard deviation of HFC and “augmented standard deviation” based on equal burden HFC ( $HFC_o$ )   | 548 |
| Table 40.2  | The FFC index based on equal burden HFC ( $HFC_o$ ) and HFC mean ( $\overline{HFC}$ )  | 549 |
| Table 40.3  | Theil’s index and the mean logarithmic deviation (MLD)   | 551 |
| Table 40.4  | Inequality in HFC implied by Atkinson’s index  | 552 |
| Table 40.5  | Rank order using different inequality measures   | 553 |
| Table 40.7  | Comparison of different summary measures   | 554 |
| Table 40.6  | The rank correlation coefficient of different inequality measures  | 554 |
| Table 42.1  | Data sources and country codes   | 567 |
| Table 42.2  | Percentage of households with catastrophic health expenditures due to out-of-pocket payments, 59 countries   | 568 |
| Table 42.3  | The determinants of catastrophic health expenditure from cross-country analysis  | 570 |
| Table 43.1  | Existing questionnaires that incorporate domains of responsiveness   | 576 |
| Table 43.2  | Operationalization of the domains in the World Health Survey 2002  | 588 |
| Table 44.1  | Criteria and approaches for constructing an instrument   | 598 |
| Table 44.2  | The operationalization of responsiveness domains in the WHO Multi-country Survey Study   | 600 |
| Table 44.3  | The number of items in the responsiveness module of the Multi-country Survey Study with a short description of the sections and the targeted respondents | 601 |
| Table 44.4  | Average item missing rates for the responsiveness module across 65 surveys   | 603 |
| Table 44.5  | Average item missing rates by survey for 65 surveys  | 604 |
| Table 44.6  | Size of samples for retests in eight countries   | 605 |
| Table 44.7  | Kappa rates for sections of the responsiveness module, calculated from retests in eight countries  | 606 |
| Table 44.8  | Item correlations and alpha coefficients for domain questions on the level of responsiveness   | 607 |
| Table 44.9  | Confirmatory factor analysis standardized coefficients—outpatients   | 609 |
| Table 44.10 | Confirmatory factor analysis standardized coefficients—inpatients  | 611 |
| Table 45.1  | List of 65 surveys analysed, survey modes, and respondent numbers  | 632 |
| Table 45.2  | Domain coefficients from the ordered probit model for 65 surveys   | 636 |
| Table 45.3  | Domain weights for eight domains across 65 surveys   | 638 |

|            |   |     |
|------------|---|-----|
| Table 45.4 | Comparison of responsiveness domain weights from 65 national sample surveys with <i>The World Health Report 2000</i> key informant survey results   | 640 |
| Table 45.5 | Correlation of responsiveness domain weights with GDP per capita and total health expenditure per capita for 65 surveys   | 641 |
| Table 46.1 | Descriptive statistics of the responsiveness module for 16 OECD countries: survey mode, response rates, the number of respondents, and the percentage of respondents using health services  | 644 |
| Table 46.2 | Wording of responsiveness module items and response options for inpatient and outpatient services in the Multi-country Survey Study for the domains of prompt attention, dignity, communication, autonomy, confidentiality, choice, quality of basic amenities, and support | 645 |
| Table 46.3 | Responsiveness domain and overall results for inpatient services in 16 OECD countries: means and standard errors, standardized by country, age and sex  | 648 |
| Table 46.4 | Responsiveness domain and overall results for outpatient services in 16 OECD countries: means and standard errors, standardized by country, age and sex   | 648 |
| Table 46.5 | Countries with high and low performance for two of the most important responsiveness domains: dignity and prompt attention  | 649 |
| Table 46.6 | Correlation of the percentage of the population “very satisfied or satisfied with the way health care runs in their country” and responsiveness domain-specific and overall inpatient and outpatient results  | 650 |
| Table 46.7 | Responsiveness inpatient and outpatient results compared with satisfaction results for the poor and wealthy in 16 OECD countries  | 651 |
| Table 47.1 | Information on the countries analysed   | 654 |
| Table 47.2 | Overall outpatient and inpatient level of responsiveness for 16 OECD countries  | 654 |
| Table 47.3 | Exercise based on key informants  | 656 |
| Table 47.4 | Overall, outpatient, and inpatient inequality in responsiveness (coefficient of variation)  | 657 |
| Table 47.5 | Inequality in responsiveness by domain  | 659 |
| Table 47.6 | A comparison of different inequality measures (coefficient of variation, relative mean deviation, standard deviation of logs, Gini coefficient, Theil index, and mean)  | 660 |
| Table 47.7 | Rank correlation between different inequality measures  | 660 |
| Table 47.8 | Inequality of responsiveness by sex   | 661 |
| Table 48.1 | Sample size and characteristics of surveys used   | 669 |
| Table 48.2 | Relative weights assigned to the three main health system goals   | 670 |
| Table 48.3 | Relative weights and sample standard deviations assigned to system quality vs. equity   | 672 |
| Table 48.4 | Results from the seemingly unrelated regression model   | 673 |
| Table 49.1 | Original weights and benefit of the doubt weights with summary statistics   | 678 |
| Table 51.1 | Variables used in the second-stage analysis   | 695 |
| Table 51.2 | Multivariate analysis for determinants of efficiency on health  | 696 |

|             |  |     |
|-------------|--|-----|
| Table 51.3  | Multivariate analysis for determinants of overall efficiency   | 697 |
| Table 54.1  | A taxonomy of evidence relating to population health and health systems  | 720 |
| Table 54.2  | Example of taxonomy for WHO estimates of life expectancy at birth for Member States for year 2000  | 722 |
| Table 54.3  | Average uncertainty range for life expectancy at birth ( $e_0$ ) estimates   | 722 |
| Table 54.4  | Example of taxonomy for UNICEF/WHO joint estimates of maternal mortality ratios (MMRs) per 100 000 for Member States for year 2000   | 722 |
| Table 54.5  | Average uncertainty ranges for estimates of maternal mortality ratio (MMR) per 100 000   | 722 |
| Table 55.1  | Descriptive statistics (simulated data)  | 729 |
| Table 55.2  | Estimation results: ordered probit   | 730 |
| Table 55.3  | Estimation results: two-stage partial credit   | 732 |
| Table 55.4  | Estimation results: partial credit with covariates   | 733 |
| Table 55.5  | Estimation results: HOPIT  | 742 |
| Table 55.6  | Estimation results: HOPIT $\tau^1$   | 742 |
| Table 55.7  | Estimation results: HOPIT $\tau^2$   | 743 |
| Table 55.8  | Estimation results: HOPIT $\tau^3$   | 743 |
| Table 55.9  | Estimation results: HOPIT $\tau^4$   | 744 |
| Table 55.10 | Estimation results: hierarchical partial credit  | 744 |
| Table 55.11 | Estimation results: hierarchical partial credit $\delta^1$   | 745 |
| Table 55.12 | Estimation results: hierarchical partial credit $\delta^2$   | 745 |
| Table 55.13 | Estimation results: hierarchical partial credit $\delta^3$   | 746 |
| Table 55.14 | Estimation results: hierarchical partial credit $\delta^4$   | 746 |
| Table 56.1  | Variables used in the estimation and validation of permanent income using DIHOPIT  | 753 |
| Table 56.2  | Results of application of random-effect DIHOPIT to Greece ECHP, 1995   | 754 |
| Table 56.3  | Correlation of estimated permanent income with reported income measures, Greece ECHP, 1995   | 754 |
| Table 56.4  | Results of application of random-effect DIHOPIT to Pakistan IHS, 1991  | 755 |
| Table 56.5  | Correlation of permanent income estimates with reported household income and expenditure, Pakistan IHS, 1991   | 756 |
| Table 56.6  | Results of application of random-effect DIHOPIT to Peru LSMS, 2000   | 757 |
| Table 56.7  | Correlation of permanent income estimates with reported household income and expenditure, Peru LSMS, 2000  | 757 |
| Table 56.8  | Item reduction subsets, Peru LSMS, 2000  | 758 |
| Table 56.9  | Spearman rank correlation of permanent income estimated from indicator subsets with full-set permanent income, household income and household expenditure, Peru LSMS, 2000 | 758 |

|              |  |     |
|--------------|--|-----|
| Table 57.1   | Assessment instrument domains  | 766 |
| Table 57.2   | Distribution of survey countries by WHO regions  | 778 |
| Table 57.3   | Countries participating in the survey study  | 778 |
| Table 57.4   | Summary survey diagnostics for different modes   | 779 |
| Table 57.5   | Sex ratio (male/female) in the UN Population Database and in the household survey country samples  | 779 |
| Table 57.6   | Timing of individual sections of full-length household interviews, pilot phase (data from 10 countries)  | 780 |
| Table 57.7   | Mean interview duration (minutes) of full-length household interviews, main phase (data from 10 countries)                                       | 781 |
| Table 57.8A  | Reliability statistics for major sections of the health component of the survey (data from 10 countries in full-length household interview mode) | 781 |
| Table 57.8B  | Reliability statistics for major sections of the responsiveness component of the survey  | 782 |
| Table 57.9   | Summary results of calibration tests   | 782 |
| Table 57.10A | Survey metrics for the household survey, health and responsiveness components  | 783 |
| Table 57.10B | Survey metrics for the brief face-to-face survey, health and responsiveness components   | 784 |
| Table 57.10C | Survey metrics for the postal survey, health and responsiveness components   | 785 |
| Table 57.11  | Average cost per completed interview   | 790 |
| Table 58.1   | Modules of the WHS instrument in 2002–2003   | 799 |
| Table 58.2   | Countries participating in the WHS 2002–2003   | 803 |
| Table 59.1   | Assessment of health system outcomes   | 813 |
| Table 59.2   | Illustration of types of indicators  | 814 |
| Table 60.1   | Leading 10 selected risk factors as per cent of disease burden measured in DALYs   | 827 |
| Table 60.2   | Interventions evaluated  | 828 |
| Table 60.3   | Annual costs, effects, and cost-effectiveness of interventions for AmrA  | 830 |
| Table 61.1   | Matrix for the assessment of human resource generation   | 856 |
| Table 61.2   | Matrix for the assessment of human resource maintenance and utilization  | 856 |
| Table 61.3   | Indicators discussed with the Scientific Peer Review Group, Dec. 2001  | 862 |
| Table 61.4   | Three classifications of tasks for stewardship   | 866 |

## LIST OF FIGURES

|             |   |     |
|-------------|---|-----|
| Figure 1.1  | Boundaries of the health system   | 7   |
| Figure 1.2  | From functions to outcomes  | 9   |
| Figure 1.3  | Defining health system efficiency   | 10  |
| Figure 12.1 | Mapping from unobserved latent variable to observed categorical response categories   | 141 |
| Figure 16.1 | Private and public health expenditures by function, Sri Lanka   | 190 |
| Figure 16.2 | Per capita expenditure on health by region in Mexico, 1995  | 191 |
| Figure 17.1 | Health spending around the world as share of GDP (%), 2000  | 196 |
| Figure 17.2 | Inequality in health spending and income by region, 2000  | 197 |
| Figure 17.3 | Healthy life expectancy and health spending, 2000   | 198 |
| Figure 17.4 | Composition of world spending on health, 2000   | 199 |
| Figure 17.5 | Public share of health spending by income group   | 199 |
| Figure 17.6 | Out-of-pocket share of health spending by income  | 200 |
| Figure 18.1 | The process of developing indicators for the financing function   | 206 |
| Figure 21.1 | Health service provider   | 235 |
| Figure 21.2 | Assessment of health service provision  | 236 |
| Figure 21.3 | Physicians, nurses, and dentists per 1 000 population, in selected OECD countries, 1998   | 238 |
| Figure 21.4 | Matrix of decision autonomy   | 241 |
| Figure 22.1 | Distribution of the probability of valid DTP3   | 254 |
| Figure 22.2 | Comparison of DTP3 and measles inequality variance index  | 256 |
| Figure 23.1 | DTP3 immunization coverage rates by birth cohort (1985–1998) from the DHS, 45 countries (sorted by increasing total valid immunization rates) | 267 |
| Figure 23.2 | DTP3 vaccination rates by birth cohort (1985–1998), according to source of data, 41 countries   | 268 |
| Figure 23.3 | Differences in DTP3 vaccination rates by birth cohort (1985–1998), according to source of data, 45 countries                                  | 268 |
| Figure 23.4 | Comparison of total valid DTP3 vaccination rates for overlapping birth cohorts across two successive DHS surveys in three countries           | 269 |
| Figure 23.5 | Quinquennial change in coverage rates for DTP3 by birth cohort (1986–1998), according to source of data, 16 countries                         | 269 |
| Figure 24.1 | Availability of health technology. 1997   | 274 |
| Figure 24.2 | Total health employment per 1 000 population, 1997  | 275 |
| Figure 24.3 | Comparison of the total health expenditure (THE) and total investment in medical facilities, 1998   | 275 |

|              |  |     |
|--------------|--|-----|
| Figure 24.4  | Comparison of total health expenditure (THE) and total investment in medical facilities, 1998                          | 276 |
| Figure 24.5  | The link between resource generation and service provision functions   | 276 |
| Figure 24.6A | Distribution of countries by physicians per 100 000 inhabitants  | 278 |
| Figure 24.6B | Distribution of countries by nurses per 100 000 inhabitants  | 278 |
| Figure 24.7  | Comparison of the number of health care personnel in selected African countries, 1996                                  | 278 |
| Figure 24.8  | Comparison of the number of health care personnel in selected European countries, 1998                                 | 279 |
| Figure 24.9  | Comparison of the number of nurses to the number of doctors in Europe, 1998  | 279 |
| Figure 24.10 | Trends in total per capita investments in medical facilities in PPP terms  | 283 |
| Figure 26.1  | Health and disability in a single health domain  | 306 |
| Figure 26.2  | Domains of health  | 309 |
| Figure 26.3  | Multidimensional health profile, WHO Health and Responsiveness Surveys 2000–2001, selected surveys and countries       | 311 |
| Figure 26.4  | Latent variable for a health domain: an illustration for mobility  | 312 |
| Figure 27.1  | Survivorship function for a population   | 321 |
| Figure 27.2  | Survivorship function for four health states   | 322 |
| Figure 28.1  | Trends in ${}_{45}q_{15}$ from vital registration data—China, 1990–1998  | 338 |
| Figure 28.2  | Trends in ${}_{45}q_{15}$ (unadjusted) based on estimates from the Sample Survey of Population Change—China, 1991–1998 | 339 |
| Figure 28.3  | Age pattern of HIV mortality   | 342 |
| Figure 28.4  | Distribution of male–female difference in life expectancy at birth, WHO Member States, 2000                            | 344 |
| Figure 28.5  | Chances of dying in childhood (0–4 years) and adulthood (15–59 years), by subregion, 2000                              | 345 |
| Figure 28.6  | Adult mortality versus child mortality for 191 WHO Member States for the year 2000                                     | 347 |
| Figure 28.7  | Average age of death and population, 2000, 191 countries   | 348 |
| Figure 29.1  | Annual logit life table values (1900–95) vs. 1900 logit values (US males)  | 358 |
| Figure 29.2  | Values of $\theta_x$ , $\gamma_x$ by age and sex   | 361 |
| Figure 29.3  | Deviations between observed and predicted logits by age, selected countries  | 362 |
| Figure 29.4  | Isoclines of $e_0$ , ${}_{45}q_{15}$ and ${}_{20}q_{60}$ , selected values, males                                      | 362 |
| Figure 29.5  | Log Mx for four populations with male $e_0 = 65$ years   | 363 |
| Figure 29.6  | Comparison of observed patterns of ${}_5q_0$ and $e_0$ vs. Coale-Demeny model values, males                            | 365 |
| Figure 29.7  | Comparison of observed patterns of ${}_5q_0$ and ${}_{45}q_{15}$ vs. Coale-Demeny model values, males                  | 365 |

|              |  |     |
|--------------|--|-----|
| Figure 29.8  | Comparison of observed patterns of ${}_{45}q_{15}$ and ${}_{20}q_{60}$ vs. Coale-Demeny model values, males  | 365 |
| Figure 29.9  | Predicted vs. observed male ${}_{45}q_{15}$ using the Coale-Demeny and modified logit systems, selecting on the basis of ${}_5q_0$ and $e_0$       | 366 |
| Figure 29.10 | Predicted vs. observed male $e_0$ using the Coale-Demeny and modified logit systems, selecting on the basis of ${}_5q_0$ and ${}_{45}q_{15}$       | 366 |
| Figure 30.1  | Latent mobility scale and cut-point variation in categorical responses   | 370 |
| Figure 30.2  | Stacked-bar diagram of vignette responses in China and the Netherlands   | 373 |
| Figure 30.3  | Three examples of consistent vignette orderings  | 376 |
| Figure 30.4  | Median BDROCC across domains   | 379 |
| Figure 30.5  | Median BDROCC across countries   | 379 |
| Figure 30.6  | 28 self-response and vignette response patterns  | 385 |
| Figure 30.7  | Cumulative mobility distributions in Latvia, Estonia, Romania, and the Czech Republic  | 388 |
| Figure 30.8  | Cut-points and vignette ratings for all respondents  | 391 |
| Figure 30.9  | Average cut-points for each survey and vignette ratings, 55 survey full run  | 392 |
| Figure 30.10 | Cut-point 1 by age category in six countries, y-axis is normalized to vignette 1 equals 100  | 392 |
| Figure 30.11 | Cut-points by educational attainment in Austria Postal Survey, y-axis is normalized to vignette 1 equals 100                                       | 392 |
| Figure 30.12 | Cut-points by sex in eight countries   | 392 |
| Figure 30.13 | Posterior estimates of mobility versus expected values of mobility for the entire dataset  | 393 |
| Figure 30.14 | Comparison of posterior estimates of mobility for full run and BDROCC > 0.8 run  | 394 |
| Figure 30.15 | Comparison of age-standardized raw score with age-standardized posteriors in 55 surveys  | 394 |
| Figure 31.1  | Response category cut-point shift  | 401 |
| Figure 31.2  | Mobility items in the World Health Survey Pilot Study  | 402 |
| Figure 31.3  | Mobility ratings for self-assessment and selected vignettes, China and Sri Lanka (N=1 061 for self-ratings, N=151 for vignettes)                   | 404 |
| Figure 31.4  | Vignette ratings for two mobility questions, pooled results from six countries (China, Myanmar, Pakistan, Sri Lanka, Turkey, United Arab Emirates) | 404 |
| Figure 31.5  | Variation in vignette ratings across age groups in three countries, Myanmar, Pakistan and Turkey   | 405 |
| Figure 32.1  | Measurement framework for health state valuations  | 411 |
| Figure 32.3  | Average VAS score by condition and country   | 418 |
| Figure 32.2  | Distribution of categorical ratings on difficulties performing usual activities for the condition of drug dependence                               | 418 |

|              |  |     |
|--------------|--|-----|
| Figure 32.4  | Box-and-whisker plots of VAS scores by country for four states: own health state, infertility, major depression, and quadriplegia  | 420 |
| Figure 32.5  | Standard deviation of domain descriptions and VAS scores   | 421 |
| Figure 32.6  | Valuation results by method and condition  | 424 |
| Figure 33.1  | Average HALE at birth (males and females combined), 191 Member States, 2000  | 441 |
| Figure 33.2  | Average HALE at birth (males and females combined), 191 Member States, 2000  | 442 |
| Figure 33.3  | Healthy life expectancy at birth versus total life expectancy at birth, by sex, WHO Member States, 2000  | 442 |
| Figure 33.4  | Healthy life expectancy at birth: males versus females, WHO Member States, 2000  | 442 |
| Figure 33.5  | Female-male difference in HALE at birth, 191 Member States, 2000   | 443 |
| Figure 33.6  | Overall prevalence YLD rates (%) for WHO Regions, 2000   | 445 |
| Figure 33.7  | Estimated age-standardized prevalence YLD rate versus life expectancy at birth, by sex, WHO Member States, 2000  | 451 |
| Figure 33.8  | Estimated age-standardized prevalence YLD: country versus regional rate, by sex, WHO Member States, 2000   | 451 |
| Figure 33.9  | Comparison of severity-weighted average prevalences from surveys and GBD priors, developed and developing countries  | 455 |
| Figure 33.10 | Comparison of severity-weighted average prevalences from surveys for developed and developing countries and for A regions versus Euro B and Euro C regions                           | 456 |
| Figure 33.11 | Age-standardized average severity-weighted prevalences versus per capita Gross Domestic Product (PPP 1998), 63 surveys in the WHO 2000–2001 household survey study                   | 456 |
| Figure 33.12 | Age-standardized average severity-weighted prevalences for females versus males, 63 surveys in the WHO 2000–2001 household survey study  | 456 |
| Figure 33.13 | Survey weights by age and sex, 63 surveys in 55 countries, 2000–2001   | 458 |
| Figure 33.14 | Estimated age-standardized posterior prevalence YLD rate versus life expectancy at birth, by sex, WHO Member States, 2000  | 458 |
| Figure 33.15 | Estimated age-standardized posterior prevalence YLD: country versus regional rate, by sex, WHO Member States, 2000   | 459 |
| Figure 33.16 | Uncertainty in average healthy life expectancy at birth (males and females combined) for the year 2000 versus average health expenditure per capita (1998) for 191 WHO Member States | 461 |
| Figure 33.17 | HALE at birth with 95% uncertainty intervals, by sex, 17 epidemiological regions, 2000   | 462 |
| Figure 34.1  | Healthy life span for an individual  | 472 |
| Figure 34.2A | Health survivorship function   | 472 |
| Figure 34.2B | Distribution of healthy life span  | 472 |
| Figure 34.3A | Health risk by age   | 473 |
| Figure 34.3B | Distribution of realized healthy life span   | 473 |



|              |   |     |
|--------------|---|-----|
| Figure 34.4  | Mortality rates by age, UK females, 1901–1995   | 474 |
| Figure 34.5A | Health risk by age  | 475 |
| Figure 34.5B | Distribution of healthy life span   | 475 |
| Figure 34.6  | Distribution of healthy life expectancy   | 477 |
| Figure 34.7  | Transfer of healthy life expectancy   | 478 |
| Figure 34.8  | Transfer of healthy life expectancy   | 479 |
| Figure 34.9  | Transfer of healthy life expectancy   | 479 |
| Figure 35.1  | Distribution of probability of death between birth and age two ( ${}_2q_0$ ) for Benin and the Central African Republic   | 488 |
| Figure 35.2  | Child survival inequality index and 95% confidence intervals for 50 countries   | 491 |
| Figure 35.3  | Child survival inequality index, plotted against five economic and demographic indicators by type of government   | 492 |
| Figure 35.4  | Country rankings of child survival inequality: comparing the individual-level inequality index with existing indices of income- and ethnicity-related inequalities in child survival. A rank of 1 on all scales indicates the highest levels of inequality. | 493 |
| Figure 36.1  | Relationship of total child survival inequality with average level of child mortality   | 500 |
| Figure 36.2  | Within- vs. between-group inequality for 39 countries   | 500 |
| Figure 36.3  | Per cent reduction in total inequality resulting from removing income inequality versus removing health services access inequality  | 500 |
| Figure 36.4  | Per cent reduction in total inequality resulting from raising income to the mean versus raising health services access to the mean for those below the mean   | 501 |
| Figure 37.1  | Predicted distributions of risk of death for males and females aged 45–54 years, USA  | 506 |
| Figure 37.2  | Coefficient of variation in expected death rate, males and females, USA   | 509 |
| Figure 37.3  | Coefficient of variation in expected death rate, males and females, United Kingdom  | 509 |
| Figure 37.4  | Coefficient of variation of risk of death from survival analysis model and small area analysis, USA, males and females  | 510 |
| Figure 38.1  | Illustration of the progressivity index   | 515 |
| Figure 38.2  | The redistributive effect (RE)  | 516 |
| Figure 38.3  | Conceptual framework for the calculation of redistributive effect (RE)  | 516 |
| Figure 38.4  | Distribution of income and the poverty line   | 517 |
| Figure 38.5  | Distribution of household financial contribution (HFC)  | 519 |
| Figure 38.6  | Redistributive effect (RE) and its components (V, H and R)  | 523 |
| Figure 38.7  | Redistributive effect (RE) and out-of-pocket (OOP) share  | 523 |
| Figure 38.8  | Level of poverty and impoverishment due to health payments  | 524 |
| Figure 38.9  | The fairness in financial contribution index (FFC) and the prepayment share of total health expenditure   | 524 |

|              |   |     |
|--------------|---|-----|
| Figure 38.10 | The distribution of household financial contribution by income deciles  | 527 |
| Figure 39.1  | Current income, permanent income, and effective income  | 534 |
| Figure 39.2  | Subsistence expenditure as a share of total consumption expenditure, Bangladesh   | 539 |
| Figure 40.1  | Distribution of household financial contributions (HFC), Spain and Azerbaijan   | 543 |
| Figure 40.2  | The FFC index based on equal burden HFC ( $HFC_o$ ) and HFC mean ( $\overline{HFC}$ )   | 548 |
| Figure 40.3  | HFC distributions, Viet Nam and Zambia  | 553 |
| Figure 41.1  | Decomposing the FFC index: removing the vertical effect   | 559 |
| Figure 41.2  | Concentration index of HFC (CI_HFC) vs. percentage increase in FFC after removing the vertical effect   | 560 |
| Figure 41.3  | Decomposing the FFC index: removing both vertical and extreme horizontal effect   | 560 |
| Figure 41.4  | Sources of unfairness (1-FFC)   | 561 |
| Figure 41.5  | Redistributive effect (RE) vs. FFC  | 562 |
| Figure 42.1  | Proportion of households with catastrophic expenditures vs. share of out-of-pocket payment in total health expenditures (OOP%THE)                             | 569 |
| Figure 44.1  | Grouping of respondents (completed questionnaires) to responsiveness module   | 602 |
| Figure 45.1  | Question on the importance of responsiveness domains asked to respondents in the responsiveness module of the Multi-country Survey Study                      | 633 |
| Figure 45.2  | Frequency of respondents in a country rating a domain as least important or most important  | 635 |
| Figure 45.3  | Domain weights under alternative anchor values  | 637 |
| Figure 45.4  | Comparisons of eight domain weights across 65 surveys. Surveys in descending order based on prompt attention weights  | 640 |
| Figure 46.1  | National health expenditure compared with inpatient and outpatient responsiveness results for 16 OECD countries   | 649 |
| Figure 47.1  | Distribution of responsiveness for 16 countries   | 658 |
| Figure 47.2  | Distribution of responsiveness for Belgium, Portugal, and Finland for males and females   | 661 |
| Figure 47.3  | Inequality in responsiveness vs. total health expenditure as percentage of GDP  | 661 |
| Figure 47.4  | Regression analysis using basic variables (inequality in responsiveness vs. Gini, total health expenditure as percentage of GDP and government effectiveness) | 662 |
| Figure 47.A1 | Inequality scenario 1   | 664 |
| Figure 47.A2 | Inequality scenario 2   | 664 |
| Figure 47.A3 | Inequality scenario 3   | 665 |
| Figure 48.1  | Health system goals as part of the performance assessment framework   | 668 |
| Figure 48.2  | Relative weights assigned to the five health system goals, by country and survey mode   | 671 |
| Figure 49.1  | Alternative (maximum) attainment scores versus original ( <i>World Health Report 2000</i> ) scores, showing 45-degree line                                    | 679 |

|              |   |     |
|--------------|---|-----|
| Figure 49.2  | Histogram showing absolute differences in two sets of attainment scores: alternative (maximum) versus original ( <i>World Health Report 2000</i> )      | 679 |
| Figure 49.3  | Density plot of the changes induced under benefit of the doubt, showing the standard normal distribution  | 679 |
| Figure 49.4  | Alternative ranks (based on maximum scores) versus original ( <i>World Health Report 2000</i> ) ranks   | 679 |
| Figure 49.5  | Histogram of absolute value of differences in two sets of ranks: based on alternative (maximum) and original ( <i>World Health Report 2000</i> ) scores | 680 |
| Figure 50.1  | The health system as a production unit  | 683 |
| Figure 50.2  | Efficiency: free disposal hull (FDH)  | 685 |
| Figure 50.3  | Efficiency: data envelopment analysis (DEA)   | 685 |
| Figure 50.4  | Efficiency: corrected ordinary least squares (COLS)   | 685 |
| Figure 50.5  | Efficiency: stochastic frontier truncated-normal model  | 685 |
| Figure 50.6  | Efficiency: fixed-effect model  | 686 |
| Figure 50.7  | Causal web relating income to health  | 687 |
| Figure 50.8  | Literacy rate versus GDP per capita, PPP  | 687 |
| Figure 50.9  | Educational attainment versus GDP per capita, PPP   | 687 |
| Figure 50.10 | Overall attainment versus health expenditure per capita, 1997 international dollars   | 688 |
| Figure 50.11 | Overall attainment versus educational attainment  | 688 |
| Figure 50.12 | Overall attainment versus proxy “Other”   | 688 |
| Figure 51.1  | Distributions of estimated efficiency for health and overall attainment   | 694 |
| Figure 51.2  | Overall efficiency versus efficiency for health: country means and confidence intervals   | 694 |
| Figure 51.3  | Kernel density estimate of the distribution of logistic transformation of estimated efficiencies  | 696 |
| Figure 52.1  | Defining health system efficiency   | 699 |
| Figure 52.2  | Efficiency and effective coverage of critical interventions   | 702 |
| Figure 53.1  | Proportion of population $\geq 16$ years of age, reporting bad and very bad general health, 12 European countries, 1994                                 | 706 |
| Figure 53.2  | Comparison of average health levels by age and sex, China and the United States of America  | 707 |
| Figure 53.3  | Mapping from latent mobility variable to categories   | 707 |
| Figure 55.1  | Mapping from unobserved latent variable to observed response categories   | 727 |
| Figure 55.2  | Distribution of responses for three self-report questions in countries A and B  | 729 |
| Figure 55.3  | Predicted versus true cut-points: ordered probit for main question  | 730 |
| Figure 55.4  | Predicted versus true mobility: ordered probit for main question  | 730 |
| Figure 55.5  | Predicted probabilities: ordered probit for main question   | 731 |

|              |   |     |
|--------------|---|-----|
| Figure 55.6  | Predicted versus true mobility: two-stage partial credit                                | 732 |
| Figure 55.7  | Predicted probabilities: two-stage partial credit for main question                     | 732 |
| Figure 55.8  | Predicted versus true mobility: partial credit with covariates                          | 733 |
| Figure 55.9  | Predicted versus true mobility: partial credit with covariates (Bayesian)               | 734 |
| Figure 55.10 | Distribution of vignette responses for country A  | 735 |
| Figure 55.11 | Distribution of vignette responses for country B  | 736 |
| Figure 55.12 | Predicted versus true mobility: HOPIT   | 737 |
| Figure 55.13 | Predicted versus true cut-points: HOPIT main question                                   | 737 |
| Figure 55.14 | Predicted versus true cut-points: HOPIT auxiliary question 1                            | 737 |
| Figure 55.15 | Predicted versus true cut-points: HOPIT auxiliary question 2                            | 738 |
| Figure 55.16 | Predicted versus true mobility: HOPIT (Bayesian)  | 738 |
| Figure 55.17 | Predicted versus true mobility: partial credit model                                    | 739 |
| Figure 55.18 | Predicted versus true mobility: partial credit model (Bayesian)                         | 739 |
| Figure 56.1  | Hypothetical indicator cut-points on the permanent income latent variable               | 750 |
| Figure 56.2  | Indicator variable ladder for 23 indicators, Greece ECHP, 1995                          | 754 |
| Figure 56.3  | Indicator variable ladder for 30 indicators, Pakistan IHS, 1991                         | 756 |
| Figure 56.4  | Indicator variable ladder for 24 indicators, Peru LSMS, 2000                            | 757 |
| Figure 57.1  | Response category shift — different rulers  | 764 |
| Figure 57.2  | Quality assurance steps for data  | 776 |
| Figure 57.3  | Different survey modes in the WHO Multi-country Survey Study, 2000–2001                 | 777 |
| Figure 57.4  | WHO Multi-country Survey Study, 2000–2001   | 777 |
| Figure 57.5  | Example of a sample population deviation index  | 780 |
| Figure 57.6  | Sample population representativeness by survey mode                                     | 786 |
| Figure 57.7A | Comparison of household vs. postal surveys for representativeness                       | 787 |
| Figure 57.7B | Comparison of brief face-to-face vs. postal surveys for representativeness              | 787 |
| Figure 57.8A | Summary quality assessment of household surveys by reliability and missing value        | 787 |
| Figure 57.8B | Summary quality assessment of household surveys by reliability and representativeness   | 787 |
| Figure 57.8C | Summary quality assessment of household surveys by representativeness and missing value | 788 |
| Figure 57.8D | Summary quality assessment of WHO surveys by representativeness and missing value       | 788 |
| Figure 57.8E | Summary quality assessment of WHO surveys by response rate and missing value            | 788 |
| Figure 57.8F | Summary quality assessment of WHO surveys by representativeness and response rate       | 788 |
| Figure 57.9A | Representativeness of DHS surveys by country  | 789 |

|              |   |     |
|--------------|---|-----|
| Figure 57.9B | Relationship between missing data and representativeness for DHS surveys  | 789 |
| Figure 57.10 | Comparison of representativeness across different surveys   | 789 |
| Figure 57.11 | Comparison of cost by mode  | 790 |
| Figure 57.12 | Adjusted and unadjusted mobility by age   | 790 |
| Figure 57.13 | Adjusted and unadjusted self-care by age  | 791 |
| Figure 57.14 | Adjusted and unadjusted dignity by age  | 791 |
| Figure 57.15 | Plot of weighted and smoothed log of death rates from survey, compared with corresponding WHO 2000 estimates for females in Nigeria | 791 |
| Figure 58.1  | Different information collection modes for health information systems   | 798 |
| Figure 58.2  | The WHS 2002–2003 geographical distribution of participating countries  | 804 |
| Figure 58.3  | WHS quality assurance procedures  | 805 |
| Figure 58.4  | Example of a sample population deviation index  | 806 |
| Figure 59.1  | Effect of increasing sample size on accuracy of sample and posterior estimates for a given prior estimate                           | 818 |
| Figure 59.2  | Proportion of a provincial population estimated to be poor  | 819 |
| Figure 59.3  | Effects of item reduction: deriving estimates of economic status from asset questions   | 819 |
| Figure 60.1  | Health system efficiency  | 823 |
| Figure 60.2  | Annual costs and effectiveness for CVD risk factor interventions, AmrA  | 831 |
| Figure 60.3A | Probability of inclusion in the optimal mix at “low levels” of resources availability in subregion AmrA                             | 832 |
| Figure 60.3B | Probability of inclusion in the optimal mix at “high levels” of resources availability in subregion AmrA                            | 832 |
| Figure 60.4  | Maximum possible health gains from selected CVD risk factor interventions, subregion AmrA   | 833 |
| Figure 61.1  | A framework for stewardship   | 869 |
| Figure 61.2  | Response category cut-point shift   | 900 |