

RESEARCH COMMUNICATION

PPAR γ knockdown by engineered transcription factors: exogenous PPAR γ 2 but not PPAR γ 1 reactivates adipogenesis

Delin Ren,¹ Trevor N. Collingwood,²
Edward J. Rebar,² Alan P. Wolffe,²
and Heidi S. Camp^{1,3}

¹Department of Molecular Science and Technology, Pfizer Global and Research Development, Ann Arbor, Michigan 48105, USA; ²Sangamo BioSciences Inc., Pt. Richmond, California 94804, USA

To determine functional differences between the two splice variants of PPAR γ (γ 1 and γ 2), we sought to selectively repress γ 2 expression by targeting engineered zinc finger repressor proteins (ZFPs) to the γ 2-specific promoter, P2. In 3T3-L1 cells, expression of ZFP55 resulted in >50% reduction in γ 2 expression but had no effect on γ 1, whereas adipogenesis was similarly reduced by 50%. However, ZFP54 virtually abolished both γ 2 and γ 1 expression, and completely blocked adipogenesis. Overexpression of exogenous γ 2 in the ZFP54-expressing cells completely restored adipogenesis, whereas overexpression of γ 1 had no effect. This finding clearly identifies a unique role for the PPAR γ 2 isoform.

Received October 12, 2001; revised version accepted November 15, 2001.

The nuclear hormone receptor PPAR γ is essential for cellular differentiation and lipid accumulation during adipogenesis (Barak et al. 1999; Kubota et al. 1999; Rosen et al. 1999). The adipocyte-specific γ 2 isoform differs from the more widely expressed γ 1 in that it contains additionally 30 amino acid residues at the amino terminus (Kliwer et al. 1994; Tontonoz et al. 1994a; Zhu et al. 1995). Evidence suggests these residues contribute to a constitutive transcription activation function that is 5–10-fold greater than in γ 1 (Werman et al. 1997). PPAR γ 2 is selectively expressed in adipose tissue (Fajas et al. 1997) and is strongly up-regulated during adipogenesis (Tontonoz et al. 1994b; Wu et al. 1998), suggesting a specific role for this isoform in fat cell differentiation. Nevertheless, a specific role for γ 2 that could not be substituted by γ 1 has not been clearly determined.

The ability to selectively knock out or knock down the expression of a specific gene provides a powerful approach for understanding its biological function. The targeting of individual mRNA splice variants offers an even greater level of selective control and understanding of

differential isoform function. Rationally engineered transcription factors potentially provide a powerful tool for targeted regulation of endogenous genes by combining a functional transcription regulatory domain with a customized DNA binding domain that can bind to a specific sequence within the target gene. C2H2 zinc finger proteins (ZFPs) can be engineered to bind with high specificity to wide a diversity of DNA sequences (Desjarlais and Berg 1992; Choo and Klug 1994; Jamieson et al. 1994; Rebar and Pabo 1994; Greisman and Pabo 1997). Previous studies have demonstrated the utility of both engineered activator- and repressor-ZFPs in the regulation of endogenous chromosomal loci (Bartsevich and Juliano 2000; Beerli et al. 2000; Zhang et al. 2000; Liu et al. 2001). Our goal for this study was to selectively inhibit expression of the PPAR γ 2 isoform in the adipogenic mouse 3T3-L1 cell line by utilizing engineered zinc finger repressor proteins.

Results and Discussion

The mouse PPAR γ gene spans >105 kb (Zhu et al. 1995). Coding exons 1 to 6 are conserved between the γ 1 and γ 2 isoforms (Fig. 1A) and transcription of these is driven by an upstream promoter (P1) that also drives expression of two untranslated γ 1-specific exons, A1 and A2. The additional amino acids at the amino terminus of γ 2 are encoded by an additional exon, B1, that is uniquely regulated by a separate promoter (P2) lies >63 kb downstream of P1. DNase I digestion of the endogenous PPAR γ gene locus in 3T3-L1 cells revealed two DNase I hypersensitive sites in the vicinity of the proximal P2 promoter that represent regions of accessible chromatin at an endogenous locus (Fig. 1B; DHS1 and DHS2). Two six-finger ZFPs (ZFP54 and ZFP55; Fig. 1C) linked to the KRAB transcriptional repression domain (Margolin et al. 1994) were designed to bind specifically to 18-bp sequences within the DHS1 shown in Figure 1B. Each ZFP bound its cognate site on naked DNA with high affinity (Kd = 20 and 44 pM, respectively).

The ZFP54 and ZFP55 repressor proteins were expressed retrovirally in 3T3-L1 cells to similar levels (Fig. 2A). Nontransduced wild-type cells and stable pools of infected cells expressing each ZFP, as well as control cells retrovirally transduced to express LacZ, were induced to differentiate and initiate the adipogenic pathway. Total PPAR γ mRNA level was determined at 0, 2, and 5 d post induction. Over the 5-d time course PPAR γ expression increased ~5.8- to 6.0-fold in both the wild-type and LacZ control cells (Fig. 2B). Up-regulation of total PPAR γ expression was reduced slightly in the presence of ZFP55 (4.8-fold compared to 6.0-fold in wild type) but completely inhibited by ZFP54. However, a time course analysis of expression of the individual PPAR γ isoforms revealed that although ZFP54 effectively inhibited expression of both isoforms, ZFP55 selectively repressed PPAR γ 2 by ~50% and had little effect on PPAR γ 1 compared with wild-type cells (Fig. 2C). Another ZFP-KRAB fusion protein that binds to a sequence that is absent in the PPAR γ gene has no effect on PPAR γ expression or adipogenesis (data not shown). The effects of these ZFPs are confirmed at the protein level whereby PPAR γ 2 expression is virtually knocked out in the presence of ZFP54, whereas only a low but detectable level of

[Key Words: PPAR γ 1; PPAR γ 2; adipogenesis; zinc finger protein]

³Corresponding author.

E-MAIL heidi.camp@pfizer.com; FAX (734) 622-5668.

Article and publication are at <http://www.genesdev.org/cgi/doi/10.1101/gad.953802>.

Ren et al.




Figure 1. Structure of PPAR γ gene and location of accessible chromatin in P2 promoter. (A) Genomic structure of mPPAR γ showing exon splicing (Zhu et al. 1995). (B) Location of DNase I hypersensitive sites (DHS1 and DHS2) in the proximal P2 promoter, along with positions of ZFPs binding sites. (C) ZFP target sequences and finger designs. (Target sequence) The promoter DNA sequence to which each ZFP was designed. (Finger designs) The residues in each position from -1 to +6 of the recognition helix of each finger targeted against the cognate basepair triplet subsite.

PPAR γ 1 remains (Fig. 2D). In the presence of ZFP55 only PPAR γ 2 expression is reduced substantially by ZFP55. Because P1 and P2 are >63 kb apart, it was predicted that ZFPs targeted to P2 would selectively inhibit PPAR γ 2 expression. This was the case with ZFP55, which gave 50% knockdown of PPAR γ 2 only, however ZFP54 suppressed both isoforms almost completely. ZFP54 and ZFP55 bind to opposite strands of the DNA and have slightly different DNA binding affinities, which may contribute to their differential effects on P2.

The promoter specificity exhibited by the two ZFPs facilitates examination of the functional requirements for each PPAR γ isoform in adipogenesis. In complete concordance with inhibited mRNA and protein expression, we find that ZFP54 totally blocks the cellular lipid accumulation that is a marker of adipogenesis (Fig. 3). Furthermore, the cells in which only PPAR γ 2 is selectively repressed by 50% (ZFP55) also show a correspond-




Figure 2. Ectopic expression of PPAR γ 2-specific ZFPs in 3T3-L1 cells. 3T3-L1 cells were infected with retroviral vectors expressing PPAR γ 2 promoter-specific zinc finger repressors, ZFP54 and ZFP55. The uninfected 3T3-L1 cells (wild type) or cells infected with retroviral vector expressing LacZ gene were used as negative controls. (A) Total RNA was isolated from selected cell lines and determined the levels of ZFP mRNA by Northern blot analysis using a 300 bp C-terminal fragment of ZFP as probe. The equivalence of RNA loading was verified by β -actin. (B) Total RNA was isolated on indicated days postdifferentiation and subjected to RNase protection assay using a PPAR γ -specific riboprobe. The protected PPAR γ and β -actin mRNAs. The fold-induction was calculated by normalization of PPAR γ -specific signal with β -actin signal and presented as fold-change compared to day-0 of wild-type cells. (C) The real time quantitative RT-PCR (TaqMan) analysis of PPAR γ 1 and PPAR γ 2 mRNA expression. 40 ng and 60 ng of reversed transcribed total RNA was used for PPAR γ 2 and PPAR γ 1 analysis, respectively. (D) Western blot analysis with a polyclonal antibody against PPAR γ . The arrows indicate PPAR γ 1 and PPAR γ 2 proteins.




Figure 3. Effect of PPAR γ 2-ZFPs on adipogenesis. Cultured wild-type 3T3-L1 and retrovirally infected cells were induced to differentiate with the standard adipogenic hormones for 10 days. Cells were fixed in 3% formaldehyde and stained with Oil Red O. Stained cells were photographed with Nikon-Diaphot300 microscope (10 \times 20) and 3CCD Vida Camera Systems (Optronics Engineering).

ing 50% loss in adipogenic capacity, supporting the idea of a specific requirement for PPAR γ 2 in this pathway.

A key aspect of engineered ZFPs is the high theoretical specificity of DNA binding and gene targeting. Each ZFP is designed to recognize an 18-bp nucleotide sequence that probability dictates to be unique within the mammalian genome. To further characterize the specificity of PPAR γ targeting, we analyzed the effect of ZFP expression on other genes both upstream and downstream of PPAR γ in the adipogenic gene cascade. The cell cyclin-dependent kinase inhibitors p21 and p27 play key roles in cell cycle progression (Harper et al. 1993; Toyoshima and Hunter 1994). P27 is expressed abundantly in growth-arrested preadipocytes where its expression is decreased transiently on hormonal induction of differentiation but returns to initial levels as the cells exit clonal expansion. p21 is absent in growth-arrested cells but increases during S-phase of mitotic expansion (Morrison and Farmer 1999). Here we show that the expression profile of both p21 and p27 is preserved irrespective of the presence of ZFPs (Fig. 4A).

Transient expression of the transcription factors C/EBP β and C/EBP δ occurs very early during adipocyte differentiation in response to the standard adipogenic hormones (IBMX/DEX/Insulin) (Cao et al. 1991; Wu et al. 1998). The proximal promoter of PPAR γ 2 contains a tandem array of C/EBP binding elements (Zhu et al. 1995). PPAR γ is up-regulated immediately following C/EBP β and C/EBP δ (Wu et al. 1998) and one would predict that expression of these upstream genes would not be affected by targeted repression of PPAR γ . In concordance with this prediction, we found that the expression level of C/EBP β protein was not significantly affected by the presence of either ZFPs (Fig. 4A). C/EBP δ expression

also was not inhibited, but rather an increase in its expression was observed in the presence of ZFP54. This suggests a potential negative feedback mechanism whereby PPAR γ may actively down-regulate C/EBP δ , in keeping with the reduction in C/EBP δ observed during progression of adipogenesis (Wu et al. 1998). Taken together these results support the notion that the inhibitory effects of ZFPs are specific.

Genes that are expressed during adipogenesis after PPAR γ induction include C/EBP α and the adipocyte-specific aP2 (Wu et al. 1998; Hamm et al. 2001). As predicted, the loss of PPAR γ expression in the presence of ZFP54 abrogates both C/EBP α and aP2 expression (Fig. 4A,B), supporting a key role for PPAR γ in the regulatory cascade leading to the expression of these genes. However, in the presence of ZFP55 the expression of C/EBP α was delayed (Fig. 4A, lanes 3,7,15), although wild-type expression levels were achieved by day five. This suggests that although the remaining PPAR γ expression in the presence of ZFP55 is sufficient to fully stimulate C/EBP α expression, it is not, in itself, sufficient to achieve wild-type levels of cellular differentiation and lipid accumulation.

The studies by others described earlier have demonstrated an absolute requirement for PPAR γ expression in adipose tissue development, but they failed to differentiate clearly between the specific regulatory capacity of PPAR γ 1 and γ 2 isoforms (Barak et al. 1999; Kubota et al. 1999; Rosen et al. 1999). In the present study, the abrogation of endogenous expression of both PPAR γ isoforms by ZFP54 provides a unique cell-based model system in which to study the differential functions of each isoform. Selective repression of the endogenous PPAR γ gene would be expected to generate what otherwise should be an adipogenically competent cell line that is devoid of




Figure 4. Effects of ZFP54 and ZFP55 on the expression of cyclin-dependent kinase inhibitors, C/EBPs and aP2. Whole cell extracts and total RNA from the experiment shown in Figure 2 were subjected to the following analysis. (A) Western blot analysis of p21 and p27, and C/EBP gene family members (β , δ , and α). Arrows indicate the 42-kD and 30-kD isoforms of C/EBP α . (B) The aP2 gene expression was analyzed by Northern blot.




Figure 5. Ectopic expression of PPAR γ 2, but not γ 1 restores adipogenesis in ZFP54 cells. The L1/ZFP54 cells were infected with the retroviral constructs expressing either PPAR γ 1 (pBMN/neo- γ 1) or PPAR γ 2 (pBMN/neo- γ 2). The retroviral vector expressing LacZ was used as empty vector control. The doubly infected cell lines were designated as ZFP54/ γ 1, ZFP54/ γ 2, and ZFP54/LacZ, respectively. After the selection with puromycin and geneticin, cells were subjected to the differentiation protocol as described previously. (A) Total RNA was isolated from the indicated cell lines and PPAR γ mRNA level was quantitated by RNase protection assay. (B) PPAR γ 1 and γ 2 protein levels by Western blot analysis. (C) Cells were differentiated with the adipogenic hormones for 10 d and stained with Oil Red O. (D) C/EBP α and β protein expression by Western blot analysis. The membrane was stripped and reblotted with a rabbit polyclonal antibody against TBP.

only PPAR γ . Rescue of these cells by exogenous expression of either PPAR γ 1 or γ 2 might be expected to identify isoform-specific functional differences in the capacity to potentiate adipogenesis. Here we used retrovirus to overexpress each PPAR γ isoform on the background of ZFP54 expressing 3T3-L1 cells. Data in Figure 5A,B demonstrate the comparable levels of expression of exogenous PPAR γ 1 and γ 2 mRNA and proteins at day 0 in the presence of ZFP54. The exogenously expressed PPAR γ 1 isoform was completely incapable of inducing adipogenesis in the presence of ZFP54 (Fig. 5C). In marked contrast, expression of PPAR γ 2 effectively restored cellular differentiation and lipid accumulation. In addition, PPAR γ 2 fully restored C/EBP α expression at 7 d postinduction, whereas PPAR γ 1 was totally ineffective (Fig. 5D). Neither isoform had any effect on the expression of C/EBP β or TBP (Fig. 5D). This result clearly demonstrates for the first time a regulatory function for PPAR γ 2 in adipogenesis that cannot be achieved by PPAR γ 1 in the absence of exogenous ligand.

It is unclear how the 30 amino acid residues unique to the PPAR γ 2 isoform confer additional regulatory function. However, in addition to rendering the constitutive activation function of the PPAR γ 2 amino terminus up to 10-fold greater than that of PPAR γ 1 (Werman et al. 1997), clinical studies have identified a human allelic variant of at least one of these residues (Pro12Ala) that has been variously associated with decreased receptor activity, lower body mass index, obesity, improved insulin sensitivity, and decreased risk of type 2 diabetes (Beamer et al. 1998; Deeb et al. 1998; Altshuler et al. 2000).

The effective restoration of the adipogenic cascade by adding back only the product of the targeted gene, PPAR γ , indicates that in this study the use of engineered repressor ZFPs does not result in the general disruption of cellular metabolic processes. This observation, which demonstrates a precise specificity for ZFP targeting, represents an important advance in the rapid establishment of cell-based gene knockdown models and such an approach may, in future, be readily transferable to whole animal transgenic studies.

Materials and methods

Design, synthesis, and DNA binding affinity of zinc finger proteins

Pairs of 3-finger ZFPs were generated as described previously (Zhang et al. 2000) then linked to form each 6-finger ZFP. The DNA binding affinity of each ZFP was determined using a gel shift technique essentially as described earlier (Zhang et al. 2000) except that 10 μ M zinc was used.

Plasmids and stable cell lines

The retroviral expression vector pBMN (Garry Nolan Laboratory, Stanford University) was constructed by inserting an internal ribosome entry site (IRES) along with either *puro*^r or *neo*^r. The retroviral plasmids pBMN-puro-ZFP54 and pBMN-puro-ZFP55 were constructed by inserting ZFP54 and ZFP55 cDNA fragments into pBMN-puro vector at *Eco*RI and *Xho*I sites. pBMN/neo- γ 1 was generated by inserting the full-length mPPAR γ 1 cDNA (nucleotide positions -6 to +1415) into pBMN-*neo* vector at *Bam*HI and *Not*I sites. pBMN/neo- γ 2 construct was derived from mPPAR γ 2/pSport (a gift from Dr. B. Spiegelman, Harvard Medical School, Boston, MA) by specifically ligating the blunt-ended PPAR γ 2 cDNA into pBMN/neo vector at the blunt-ended *Eco*RI site. To generate infectious recombinant viruses, Phoenix-ECO cells (Garry Nolan Laboratory, Stanford University, CA) were cultured in DMEM medium containing 10% FBS. Cells were transfected with 15 μ g of retroviral plasmids by Lipofectamine 2000 kit (GIBCO BRL) following the protocol as recommended by the manufacturer. Viral supernatant was harvested at 72 h posttransfection and added onto ~75% confluent 3T3-L1 cells in the presence of 8 μ g/mL polybrene (Sigma) for 2 h. Infected cells were selected by adding 2

Functional differences between PPAR γ 1 and PPAR γ 2

μ g/mL puromycin (Sigma) for 5 d. For double infection, L1/ZFP54 cells were infected with either pBMN/neo- γ 1, or pBMN/neo- γ 2, and cells were selected with both 600 μ g/mL geneticin (GIBCO BRL) and 2 μ g/mL puromycin for 14 d. Pools of stably infected cells were used in these experiments.

Cell culture and differentiation

Uninfected 3T3-L1 and virally infected 3T3-L1 cells were cultured and maintained for 2 d postconfluence in DMEM containing 10% CS. Differentiation protocol was conducted as described previously (Camp et al. 2001)

Mapping of DNase I accessible chromatin regions

DNase I mapping was performed essentially as described previously (Liu et al. 2001). Undifferentiated 3T3-L1 cell nuclei were partially digested with DNase I and the genomic DNA was extracted followed by *Xba*I digestion at -1588 bp in the P2 promoter. The probe for Southern blotting of the proximal P2 promoter was an *Xba*I/*Eco*RI fragment spanning from -1588 to -1007.

The real-time quantitative PCR (TaqMan)

Gene-specific primers and probes were designed using the Primer Express software (Perkin Elmer Life Sciences). The real-time quantitative RT-PCR reaction was performed essentially following the manufacturer's protocol. Briefly, reaction mixture contained 5.5 mM MgCl₂, 500 μ M dNTP, 2.5 μ M random hexamers, 200 nM FAM-probe, and 600 nM of both forward and reverse primers in a final volume of 25 μ L, and was analyzed in ABI PRISM 7700 sequence detection system (Perkin Elmer Life Sciences). Relative quantitation of PPAR γ mRNA levels was plotted as fold-change compared to day 0 of wild-type 3T3-L1 cells. 18S ribosomal RNA was used for normalization. TaqMan reverse transcriptase reactions were performed in triplicates and the experiments were repeated independently at least three times.

RNA isolation, Northern blot, and RNase protection assay

Total RNA was isolated from cultured cells using Ultraspec RNA system (Biotecx Laboratories, Inc.). ZFP, aP2, and rat β -actin cDNA were labeled with [α -³²P] dCTP (Amersham) using random primed labeling kit (GIBCO BRL). Northern blot analysis and RNase protection assay (RPA) were performed as described previously (Camp et al. 1999).

Immunoblot analysis

Cells were lysed in HNTG cell lysate buffer (50 mM Hepes, 150 mM NaCl, 10% glycerol, 1% Triton, 1.5 mM MgCl₂, and 1 mM EDTA) at 4°C for 15 min, followed by centrifugation at 15,000 rpm at 4°C for 10 min. The supernatant was collected and the protein concentration was determined by BCA protein assay (Pierce). Western blot analysis was carried out as described previously (Camp et al. 1999). Rabbit polyclonal antibodies against C/EBP α , C/EBP β , C/EBP δ , TBP, and monoclonal antibodies against p21 and p27 were purchased from Santa Cruz.

Acknowledgments

We thank Neelam Srivastava, Katherine Pasquetti, and Hong Qi for technical assistance, and Casey Case, Todd Leff, and Yen Choo for valuable discussion and criticism. We dedicate this work to the memory of our friend and colleague, Alan P. Wolffe.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

Altshuler, D., Hirschhorn, J.N., Klannemark, M., Lindgren, C.M., Vohl, M.C., Nemesh, J., Lane, C.R., Schaffner, S.F., Bolk, S., Brewer, C., et al. 2000. The common PPAR γ Pro12Ala polymorphism is associated with decreased risk of type 2 diabetes. *Nat. Genet.* **26**: 76–80.

Barak, Y., Nelson, M.C., Ong, E.S., Jones, Y.Z., Ruiz-Lozano, P., Chien, K.R., Koder, A., and Evans, R.M. 1999. PPAR γ is required for placental, cardiac, and adipose tissue development. *Mol. Cell* **4**: 585–595.

Bartsevich, V.V. and Juliano, R.L. 2000. Regulation of the MDR1 gene by transcriptional repressors selected using peptide combinatorial libraries.

Mol. Pharmacol. **58**: 1–10.

Beamer, B.A., Yen, C.J., Andersen, R.E., Muller, D., Elahi, D., Cheskin, L.J., Andres, R., Roth, J., and Shuldiner, A.R. 1998. Association of the Pro12Ala variant in the peroxisome proliferator-activated receptor- γ 2 gene with obesity in two Caucasian populations. *Diabetes* **47**: 1806–1808.

Beerli, R.R., Dreier, B., and Barbas, C.F., III. 2000. Positive and negative regulation of endogenous genes by designed transcription factors. *Proc. Natl. Acad. Sci.* **97**: 1495–1500.

Camp, H.S., Whitton, A.L., and Tafuri, S.R. 1999. PPAR γ activators down-regulate the expression of PPAR γ in 3T3-L1 adipocytes. *FEBS Lett.* **447**: 186–190.

Camp, H.S., Chaudhry, A., and Leff, T. 2001. A novel potent antagonist of peroxisome proliferator-activated receptor gamma blocks adipocyte differentiation but does not revert the phenotype of terminally differentiated adipocytes. *Endocrinology* **142**: 3207–3213.

Cao, Z., Umek, R.M., and McKnight, S.L. 1991. Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes & Dev.* **5**: 1538–1552.

Choo, Y. and Klug, A. 1994. Selection of DNA binding sites for zinc fingers using rationally randomized DNA reveals coded interactions. *Proc. Natl. Acad. Sci.* **91**: 11168–11172.

Deeb, S.S., Fajas, L., Nemoto, M., Pihlajamaki, J., Mykkanen, L., Kuusisto, J., Laakso, M., Fujimoto, W., and Auwerx, J. 1998. A Pro12Ala substitution in PPAR γ 2 associated with decreased receptor activity, lower body mass index and improved insulin sensitivity. *Nat. Genet.* **20**: 284–287.

Desjarlais, J.R. and Berg, J.M. 1992. Redesigning the DNA-binding specificity of a zinc finger protein: A data base-guided approach. *Proteins* **12**: 101–104.

Fajas, L., Auboeuf, D., Raspe, E., Schoonjans, K., Lefebvre, A.M., Saladin, R., Najib, J., Laville, M., Fruchart, J.C., Deeb, S., et al. 1997. The organization, promoter analysis, and expression of the human PPAR γ gene. *J. Biol. Chem.* **272**: 18779–18789.

Greisman, H.A. and Pabo, C.O. 1997. A general strategy for selecting high-affinity zinc finger proteins for diverse DNA target sites. *Science* **275**: 657–661.

Hamm, J.K., Park, B.H., and Farmer, S.R. 2001. A role for C/EBP β in regulating peroxisome proliferator-activated receptor gamma activity during adipogenesis in 3T3-L1 preadipocytes. *J. Biol. Chem.* **276**: 18464–18471.

Harper, J.W., Adami, G.R., Wei, N., Keyomarsi, K., and Elledge, S.J. 1993. The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G₁ cyclin-dependent kinases. *Cell* **75**: 805–816.

Jamieson, A.C., Kim, S.H., and Wells, J.A. 1994. In vitro selection of zinc fingers with altered DNA-binding specificity. *Biochemistry* **33**: 5689–5695.

Kliwer, S.A., Forman, B.M., Blumberg, B., Ong, E.S., Borgmeyer, U., Mangelsdorf, D.J., Umehono, K., and Evans, R.M. 1994. Differential expression and activation of a family of murine peroxisome proliferator-activated receptors. *Proc. Natl. Acad. Sci.* **91**: 7355–7359.

Kubota, N., Terauchi, Y., Miki, H., Tamemoto, H., Yamauchi, T., Komeda, K., Satoh, S., Nakano, R., Ishii, C., Sugiyama, T., et al. 1999. PPAR γ mediates high-fat diet-induced adipocyte hypertrophy and insulin resistance. *Mol. Cell* **4**: 597–609.

Liu, P.Q., Rebar, E.J., Zhang, L., Liu, Q., Jamieson, A.C., Liang, Y., Qi, H., Li, P.X., Chen, B., Mendel, M.C., et al. 2001. Regulation of an endogenous locus using a panel of designed zinc finger proteins targeted to accessible chromatin regions. Activation of vascular endothelial growth factor A. *J. Biol. Chem.* **276**: 11323–11334.

Margolin, J.F., Friedman, J.R., Meyer, W.K., Vissing, H., Thiesen, H.J., and Rauscher, F.J., III. 1994. Kruppel-associated boxes are potent transcriptional repression domains. *Proc. Natl. Acad. Sci.* **91**: 4509–4513.

Morrison, R.F. and Farmer, S.R. 1999. Role of PPAR γ in regulating a cascade expression of cyclin-dependent kinase inhibitors, p18^(INK4c) and p21^(Waf1/Cip1), during adipogenesis. *J. Biol. Chem.* **274**: 17088–17097.

Rebar, E.J. and Pabo, C.O. 1994. Zinc finger phage: Affinity selection of fingers with new DNA-binding specificities. *Science* **263**: 671–673.

Rosen, E.D., Sarraf, P., Troy, A.E., Bradwin, G., Moore, K., Milstone, D.S., Spiegelman, B.M., and Mortensen, R.M. 1999. PPAR γ is required for the differentiation of adipose tissue in vivo and in vitro. *Mol. Cell* **4**: 611–617.

Ren et al.

- Tontonoz, P., Hu, E., Graves, R.A., Budavari, A.I., and Spiegelman, B.M. 1994a. mPPAR γ 2: Tissue-specific regulator of an adipocyte enhancer. *Genes & Dev.* **8**: 1224–1234.
- Tontonoz, P., Hu, E., and Spiegelman, B.M. 1994b. Stimulation of adipogenesis in fibroblasts by PPAR γ 2, a lipid-activated transcription factor. *Cell* **79**: 1147–1156.
- Toyoshima, H. and Hunter, T. 1994. p27, a novel inhibitor of G₁ cyclin-Cdk protein kinase activity, is related to p21. *Cell* **78**: 67–74.
- Werman, A., Hollenberg, A., Solanes, G., Bjorbaek, C., Vidal-Puig, A.J., and Flier, J.S. 1997. Ligand-independent activation domain in the N terminus of peroxisome proliferator-activated receptor gamma (PPAR γ). Differential activity of PPAR γ -1 and -2 isoforms and influence of insulin. *J. Biol. Chem.* **272**: 20230–20235.
- Wu, Z., Xie, Y., Morrison, R.F., Bucher, N.L., and Farmer, S.R. 1998. PPAR γ induces the insulin-dependent glucose transporter GLUT4 in the absence of C/EBP α during the conversion of 3T3 fibroblasts into adipocytes. *J. Clin. Invest.* **101**: 22–32.
- Zhang, L., Spratt, S.K., Liu, Q., Johnstone, B., Qi, H., Raschke, E.E., Jamieson, A.C., Rebar, E.J., Wolffe, A.P., and Case, C.C. 2000. Synthetic zinc finger transcription factor action at an endogenous chromosomal site. Activation of the human erythropoietin gene. *J. Biol. Chem.* **275**: 33850–33860.
- Zhu, Y., Qi, C., Korenberg, J.R., Chen, X.N., Noya, D., Rao, M.S., and Reddy, J.K. 1995. Structural organization of mouse peroxisome proliferator-activated receptor gamma (mPPAR γ) gene: Alternative promoter use and different splicing yield two mPPAR γ isoforms. *Proc. Natl. Acad. Sci.* **92**: 7921–7925.



PPAR γ knockdown by engineered transcription factors: exogenous PPAR γ 2 but not PPAR γ 1 reactivates adipogenesis

Delin Ren, Trevor N. Collingwood, Edward J. Rebar, et al.

Genes Dev. 2002, **16**:

Access the most recent version at doi:[10.1101/gad.953802](https://doi.org/10.1101/gad.953802)

References

This article cites 30 articles, 16 of which can be accessed free at:
<https://genesdev.cshlp.org/content/16/1/27.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

