

Toward a Geometry of Common Sense: A Semantics and a Complete Axiomatization of Mereotopology

Nicholas Asher

Dpt of Philosophy, University of Texas at Austin
316 Waggener Hall
Austin, Texas, USA
nasher@bertie.la.utexas.edu

Laure Vieu

IRTT, University Paul Sabatier
118 route de Narbonne
31062 Toulouse cedex, France
vieu@irit.fr

Abstract

Mereological and topological notions of connection, part, interior and complement are central to spatial reasoning and to the semantics of natural language expressions concerning locations and relative positions. While several authors have proposed axioms for these notions, no one with the exception of Tarski [18], who based his axiomatization of mereological notions on a Euclidean metric, has attempted to give them a semantics. We offer an alternative to Tarski, starting with mereotopological notions that have proved useful in the semantic analysis of spatial expressions. We also give a complete axiomatization of this account of mereotopological reasoning.

1 Introduction

Mereological and topological notions of connection, part, interior and complement are central to spatial reasoning and to the Natural Language (NL) semantics of expressions concerning locations and relative positions. For example, reasoning about objects inside other objects or on them may involve complex inferences based on a semantics for the prepositions *in* and *on* that exploits these mereotopological notions properly [4]. Reasoning about these concepts is also present in many commonsense reasoning tasks about spatial position and navigation [13, 16]. We propose here a semantics and complete axiomatization of these notions that takes the linguistic semantics of spatial expressions to be a fundamental guide to commonsense spatial and geometrical reasoning. We do this for two reasons. First we are interested in reasoning about spatial situations from linguistically given information, in particular the spatial prepositions, movement verbs, and noun phrases referring to portions of space. In such a task we must understand exactly what sort of information spatial expressions in NL convey. Second, we are interested in how the narrative structure of discourse can convey spatial information [2].

When examining commonsense reasoning from this linguistic perspective, two observations become apparent. The first is the absence in NL of a natural way of referring to points without measure. Such points are fundamental to the mathematical conception of topological spaces, geometry, and analysis. But they are foreign to space as it is usually expressed in narrative texts, newspaper articles and texts about how to get to a certain place, which are the sort of texts with which we have been concerned. At the very least, mathematical points are not needed to express spatial relationships within NL. Furthermore, the NL expressions that

we do use to refer to "points," "borders," and "surfaces" do not refer in the following examples to the mathematical notions:

- (1) The point of this pencil is actually an irregular surface with several peaks.
- (2) There is a scratch in the surface of the table.

Mathematical points, unlike pencil points, can't be surfaces, nor can mathematical surfaces have scratches (which presuppose a depth). Even the points, surfaces and boundaries of our commonsense world are three dimensionally extended objects.

The other observation is that agents use spatial information contained in texts even though this information does not contain any system of universal coordinates by means of which we may spatially situate the objects talked about in the text in the way we might use a coordinate system (e.g. longitude, latitude and altitude) to situate the position of a robot. NL strongly suggests that space is a *relational* concept as Leibniz argued, not an "absolute" concept as Newton thought. Our conception of space is constructed from or dependent upon the relative positions of objects to each other.'

This is not to say that our commonsense conception of space has nothing to do with mathematical conceptions of space. Commonsense reasoning approximates but simplifies the mathematical concepts, since the full power of mathematical topology, geometry or analysis, is not needed in commonsense spatial reasoning tasks. That there should be this compatibility between mathematics and commonsense seems evident to us in that, for example, we can express mathematically precise locations in NL relative to a system of coordinates-*north*, *south*, *east*, *west*, *longitude* and *latitude*, together with numerical expressions within these dimensions and the dimension of altitude. We can also when needed introduce the language of analysis. Indeed, one of the accomplishments of the theory we propose below is that we make this notion of approximation between commonsense and mathematics itself formally precise. But it is clear that mathematical expressions are not always present in NL when spatially useful information is present. Thus, the use of mathematically precise notions of space is not an essential part of our conception of space.

Our aim here is to develop the foundations of a commonsense geometry. In section 2, we present a language in which

¹Like time which may be construed out of events only, not taking states into account [12], space may be constructed from material objects only. Indeed, immaterial portions of space such as *the space beneath the table* are entities dependent of objects.

we cannot refer to points without extension and in which there are no expressions for referring to a coordinate system. We also present in section 2 a semantics for this language. In sections 3 and 4, we present a new sound and complete theory for the topological aspects of this qualitative, relational conception of space. In section 5, we develop a modal analysis of the approximation between this commonsense conception of space and the mathematical one.

2 Structures for Relational Mereotopology

Our attempt to build topology and geometry out of a domain of individuals or "bodies" instead of points is inspired by Whitehead [21], Tarski [18] and Clarke [6,7], as well as work in mereology [14]. Tarski is the only one we know who has presented an axiomatized and complete mereotopological theory besides ours, but he classified his domain of "bodies" into "spheres" and non-spheres, thus recovering topological concepts from metrical ones. Our topological theory most directly exploits Clarke's work which corrects a mistake of Whitehead's, leading to an inconsistency. However, since Clarke presents (in the Russellian philosophical style) only an axiomatic calculus, we will begin by describing the formal language and the semantic structures we have found necessary for the commonsense conception of space embodied in NL.

To state our relational, topological theory, we use a first order language, L_{RT} with one non logical two place relation symbol, C , and a denumerable infinity of individual nonlogical constants (a, a_0, a^*, b, b_1 , etc.) and variables (x, y, z, u, v, w, x_i , etc.). C represents intuitively the notion of connection between individuals. There are intuitively two sorts, EC (external connection), in which two objects share a boundary and O (overlap) in which two objects have a common part which is an object. A further kind of contact is also needed, $WCont$ (weak contact), in which two objects are not connected but are in some sense "vanishingly close" to each other. Examples of weak contact are the relation between a glass and the table on which it is standing and the relation between the glass and the wine it contains, while an example of external connection or "strong contact" is the relation between the stem of the glass and the cup of the glass. To express this notion of vanishing closeness, it is natural to suppose that each object has surrounding it a smallest neighborhood. While $WCont$ may be defined in terms of C it seems natural to us to think of its semantics using neighborhood. Further, neighborhoods may be useful if we wish to impose constraints on smallest neighborhoods that we cannot state within L_{RT} (for example, that for some fixed real number 5, the neighborhood of x is within 6 of x). The presence of neighborhoods is also a useful feature if we wish to incorporate principles about how properties persist over spatial regions (see for instance [3]).

Besides these notions, NL semantics requires us to capture the more classical topological notions of openness and closure (for details see [4]). To show that the right topological notions of open and closed properties are grasped, it suffices to show that our cognitive spaces are structures built over topological spaces. The fact that our structures are based on set-theory and have points as primary entities is not in contradiction with our aims. It doesn't really matter how one chooses to express the models of a theory: once complete-

ness is proven, they are all equivalent in the sense that our language and inferential mechanisms cannot distinguish between them. Moreover, there are two clear advantages of expressing our structures in terms of classical topological spaces. First, it's a practical way to compare the concepts we introduced with better-known ones, and second, it will give us the opportunity to show in section 5 that classical topology may be seen as a limit of mereotopology.

The domain of our structures is a subset of the power set of a topological space. Every subset corresponding to the space occupied by a physical object must clearly have a non-empty interior. However, few people working on mereotopology have noticed an additional important feature of cognitive space, emphasized in qualitative spatial reasoning by [10] and [8]: the subsets of a topological space corresponding to physical objects are three dimensional throughout or "regular". Regularity has two defining features: for any n -dimensional regular space, (1) every part of the objects in that space must be also n -dimensional (smooth boundaries); (2) these objects must have no "holes" of a lower dimension (full interiors). This means that subsets including a single isolated point, as well as subsets with a single interior point deleted are not in the structure. This feature has a further consequence: we must modify the union and intersection operators to ensure that they are internal composition laws on our domain of objects. To make this clear, consider this example. Suppose we have the one-dimensional space, $\langle R^+, \{ \text{any union of open intervals} \} \rangle$. Then, clearly any open interval is "regular" but the union of $(0,1)$ and $(1,2)$ has a deleted point, 1. The correct union operator of two regular subsets preserving "regularness" is, not ordinary u , but a new operator u^* , introduced below. Similarly its dual n^* replaces set intersection.

This discussion leads naturally to our definition of the models

2.1 Definition of the models

Let $T = \langle X, \mathbf{X} \rangle$ be a non-empty topological space (X being the set of points, \mathbf{X} the set of open subsets of X , int and cl the interior and closure operators, $open$ and $closed$ the open and closed properties, and \sim the relative complement wrt X). Let $RT_T = \langle Y, f, \{ \cdot \} \rangle$ be a structure such that

- 1) Y is a set with the following properties
 - (i) $Y \subseteq \wp(X)$ and $X \in Y$
 - (ii) $\forall x \in Y (int(x) \in Y \ \& \ int(x) \neq \emptyset \ \& \ int(x) = int(cl(x)))$
(full interiors)
 - (iii) $\forall x \in Y (cl(x) \in Y \ \& \ cl(x) = cl(int(x)))$
(smooth boundaries)
 - (iv) $\forall x \in Y (int(\sim x) \neq \emptyset \rightarrow \sim x \in Y)$
 - (v) $\forall x \in Y \ \forall y \in Y (int(x \cap y) \neq \emptyset \rightarrow x \cap^* y \in Y)$
 - (vi) $\forall x \in Y \ \forall y \in Y \ x \cup^* y \in Y^2$
where $x \cap^* y = x \cap y \cap cl(int(x \cap y))$ and $x \cup^* y = x \cup y \cup int(cl(x \cup y))$
 - (vii) $\exists x \in Y \ \exists y \in Y (x \cap y \neq \emptyset \ \& \ int(x \cap y) = \emptyset)$
 - (viii) $\exists x \in Y \ \exists y \in Y (x \cap y = \emptyset \ \& \ \forall z \in Y ((open(z) \ \& \ x \subseteq z) \rightarrow y \cap cl(z) \neq \emptyset))$
- 2) f is a function over $Y (f: Y \rightarrow Y)$ such that:

$$x \subseteq f(x) \ \& \ open(f(x)) \ \& \ (cl(f(x)) \cap y \neq \emptyset \rightarrow x \cap cl(f(y)) \neq \emptyset) \ \& \ ((open(x) \ \& \ y \subseteq x) \rightarrow f(y) \subseteq x)$$

²Note that $x \cap^* y \subseteq x \cap y$, $x \cup y \subseteq x \cup^* y$, $x \cap^* x = x$, $x \cup^* x = x$

$f(x)$ is called the (smallest) neighborhood of x , a straightforward extension of the topological notion of neighborhood of a point. It is important to note that X cannot be dense (in the sense that $(\text{open}(x) \ \& \ \text{open}(y) \ \& \ x \subset y \ \& \ x \neq y) \rightarrow \exists z (\text{open}(z) \ \& \ z \neq x \ \& \ z \neq y \ \& \ x \subset z \ \& \ z \subset y)$ because the notion of a minimal open set is vacuous then, since the intersection of any set of open sets may not be an open set.

3) $\llbracket \cdot \rrbracket$ is a function assigning individual terms a denotation in Y . The interpretation of C will be given below. (We assume the usual extension of the interpretation of constants 0 to an interpretation of terms $\llbracket \cdot \rrbracket_g$, where g is an assignment of objects in the domain to the variables occurring free in the term.)

Of all these constraints only conditions (vii) and (viii) above have not yet been motivated. They are needed to ensure that our structure is nontrivial in the sense that it does have instances of weak contact and external connection.

2.2 Semantics for L

$RT_T \models_g C(x,y)$ iff $\llbracket x \rrbracket_g \cap \llbracket y \rrbracket_g \neq \emptyset$
 $RT_T \models_g WCont(x,y)$ iff not $RT_T \models_g C(x,y)$ and $f(\llbracket x \rrbracket_g) \cap \llbracket y \rrbracket_g \neq \emptyset$

$RT = \{ RT_T : T \text{ is a non-empty topological space} \}$

Validity in every RT_T model is written \vdash_{RT}

2.3 Axiomatization

Our axiomatization extends and corrects that of [6]. Classical mereology [14], as well as [6], contains a fusion operator for summing up any collection of individuals into a new individual. This general fusion operator is in fact unnecessary. In addition, removing Clarke's axiom stating the existence of the fusion of any collection of individuals is a simple way of making the theory first-order, and gives a response to a criticism often given of mereology, that it is neither cognitively nor ontologically acceptable to assume the existence of individuals having as parts any collection of individuals. This change is visible in axioms (A4-8). The next change is in the definition of connectedness (D10). Clarke's definition makes it impossible to have connected spaces as soon as it is split into two externally connected parts, since the sum of two interiors equals the entire space and thus is not connected according to his definition₃. Finally, we added $WCont$, and axioms that ensure that the theory is not trivially verified in any topology (because, e.g., there is no external connection).

Theory RT_0

- (A1) $C(x,x)$
- (A2) $C(x,y) \rightarrow C(y,x)$
- (A3) $(C(z,x) \ \& \ C(z,y)) \rightarrow x=y$
- (D1) $P(x,y) \stackrel{\text{def}}{=} \forall z (C(z,x) \rightarrow C(z,y))$
- (D2) $PP(x,y) \stackrel{\text{def}}{=} P(x,y) \ \& \ \neg P(y,x)$
- (D3) $O(x,y) \stackrel{\text{def}}{=} \exists z (P(z,x) \ \& \ P(z,y))$
- (D4) $EC(x,y) \stackrel{\text{def}}{=} C(x,y) \ \& \ \neg O(x,y)$
- (D5) $TP(x,y) \stackrel{\text{def}}{=} P(x,y) \ \& \ \exists z (EC(z,x) \ \& \ EC(z,y))$
- (D6) $NTP(x,y) \stackrel{\text{def}}{=} P(x,y) \ \& \ \neg \exists z (EC(z,x) \ \& \ EC(z,y))$
- (A4) $\exists x \forall u C(u,x)$

(A4) and (A3) entail the existence of a unique universe a^* .

(A5) $\forall x \forall y \exists z \forall u (C(u,z) \rightarrow (C(u,x) \vee C(u,y)))$

(A5) and (A3) entail the existence of a unique sum of x and y , $x+y$ for every x and y .

³ We owe this observation to Carola Eschenbach.

(A6) $\forall x \forall y (O(x,y) \rightarrow \exists z \forall u (C(u,z) \rightarrow \exists v (P(v,x) \ \& \ P(v,y) \ \& \ C(v,u))))$

(A6) and (A3) entail the existence of a unique nonempty intersection, $x*y$ for every x and y such that $O(x,y)$.

(A7) $\forall x (\exists y \neg C(y,x) \rightarrow \exists z \forall u (C(u,z) \rightarrow \exists v (\neg C(v,x) \ \& \ C(v,u))))$

(A7) and (A3) entail the existence of a unique complement of x in a^* , $\neg x$, for every $x \neq a^*$.

(A8) $\forall x \exists y \forall u (C(u,y) \rightarrow \exists v (NTP(v,x) \ \& \ C(v,u)))$

(A8) and (A3) entail the existence of a unique interior, ix , for every x .

(D7) $cx \stackrel{\text{def}}{=} \neg i(\neg x)$

Since the complement is involved in this definition, $c(x)$ exists only if $x \neq a^*$. We make the c operator a function by adding axiom (A9):

(A9) $c(a^*) = a^*$

(D8) $OP(x) \stackrel{\text{def}}{=} x = ix$

(D9) $CL(x) \stackrel{\text{def}}{=} cx = cx$

(A10) $(OP(x) \ \& \ OP(y) \ \& \ O(x,y)) \rightarrow OP(x*y)$

(D10) $Con(x) \stackrel{\text{def}}{=} \neg \exists y \exists z (x=y+z \ \& \ \neg C(y,z))$

(D11) $WCont(x,y) \stackrel{\text{def}}{=} \neg C(x,y) \ \& \ \forall z ((P(x,z) \ \& \ OP(z)) \rightarrow C(cz,y))$

(A11) $\exists x \exists y EC(x,y)$

(A12) $\exists x \exists y WCont(x,y)$

(A13) $\forall x \exists y (P(x,y) \ \& \ OP(y) \ \& \ \forall z (P(x,z) \ \& \ OP(z)) \rightarrow P(y,z))$

(A13) and (A3) entail the existence of a unique neighborhood nx for each x ; if x is open, $nx=x$. Fact 3 shows that neighborhoods and $WCont$ interact in the right way.

Fact 1: i) $\neg x=x$; ii) $icx=ix$; iii) $cix=cx$. [20]

Fact 2: $WCont(x,y) \rightarrow WCont(y,x)$ [20]

Fact 3: $(\neg C(x,y) \ \& \ C(x,c(ny))) \rightarrow WCont(x,y)$

Proof: \rightarrow : By (A13), $\forall z ((P(x,z) \ \& \ OP(z)) \rightarrow P(nx,z))$.

Since $P(nx,z) \rightarrow P(c(nx),cz)$, $(C(c(nx),y) \ \& \ P(nx,z)) \rightarrow C(cz,y)$; so $\neg C(x,y) \ \& \ \forall z ((P(x,z) \ \& \ OP(z)) \rightarrow C(cz,y))$.

\leftarrow : Directly from (D11) since by (A13), $P(x,nx) \ \& \ OP(nx)$

3 The Soundness Proof

When added to the axioms and rules for first order logic, the set of axioms we have set up for relational mereotopology (RT_0) form the basis of a proof relation \vdash_{RT_0} . A proof in RT_0 is as usual a finite sequence of sentences of L, each one of which is either an axiom or derivable from the other lines using Modus Ponens or Universal Generalization. We now provide a class of models for RT_0 to show that it is sound. Clarke did not do this, nor have any of the people following his work done this. This has led researchers, including Clarke, to miss certain important features of the theory~e.g., the models are not just the power sets of topological spaces with the subsets having an empty interior removed, and the difficulty of defining points within this theory (attempted in [7], but proved in [20] to be incorrect).

The question arises as to whether the constraints we have imposed on Y and hence on RT_T , though intuitively motivated, are consistent. It will suffice to show that, under the hypotheses that $Y \neq \emptyset$ and $Y \neq \{X\}$ (needed to satisfy the constraints corresponding to (A1) and (A12)):

Fact 4: Let X and Y be as defined above for RT_T and suppose that $Y \neq \emptyset$ and $Y \neq \{X\}$; then, the elements asserted to exist in Y in clauses (i) and (iv)-(vi) verify the constraints on their interiors and closures imposed by clauses (ii) and (iii).

The proof though long is not hard. Our conclusion is that indeed the constraints are satisfiable by a substructure of a classical topological space.

Theorem 5: $\vdash RT_0 \phi \rightarrow \vdash_{RT} \phi$

The proof is by induction on the complexity of a proof in RT_0 . We look here only at the basis case to show for an arbitrary model $RT_T = \langle Y, f, I \rangle$ that $RT_T \vdash (A1)$ to (A13).

That a structure RT_T verifies (A1) and (A2) is trivial. That RT_T verifies (A4) follows immediately from the fact that $X \in Y$. To show (A3), it suffices to show that $\forall x \in Y \forall y \in Y (\forall z \in Y (z \cap x \neq \emptyset \rightarrow z \cap y \neq \emptyset) \rightarrow x \subseteq y)$. If $y = X$, $x \subseteq y$ is always true; if $y \neq X$, suppose $\forall z \in Y (z \cap x \neq \emptyset \rightarrow z \cap y \neq \emptyset)$ and $\neg x \subseteq y$, we have $y \neq X \rightarrow \sim y \in Y$ and $\neg x \subseteq y \rightarrow x \cap \sim y \neq \emptyset$ then by hypothesis, $y \cap \sim y \neq \emptyset$, which is impossible. \square

More complex are (A5 - A8). Let's first consider (A5): to show $RT_T \vdash \forall x \forall y \exists z \forall u (C(u, z) \leftrightarrow (C(u, x) \vee C(u, y)))$ where $z = x * y$, we show that $\llbracket x + y \rrbracket_g = \llbracket x \rrbracket_g \cup \llbracket y \rrbracket_g$ for some arbitrary assignment g . This amounts to showing: $\forall x \in Y \forall y \in Y \forall u \in Y (u \cap (x \cup y) \neq \emptyset \leftrightarrow (u \cap x \neq \emptyset \vee u \cap y \neq \emptyset))$

\leftarrow : Since $(u \cap x \neq \emptyset \vee u \cap y \neq \emptyset) \rightarrow u \cap (x \cup y) \neq \emptyset$ and $x \cup y \subseteq x \cup^* y$, $u \cap (x \cup y) \neq \emptyset$
 \rightarrow : By definition, $x \cup^* y = (x \cup y) \cup \text{int}(\text{cl}(x \cup y))$. So $u \cap (x \cup^* y) \neq \emptyset \rightarrow u \cap (x \cup y) \neq \emptyset \vee u \cap \text{int}(\text{cl}(x \cup y)) \neq \emptyset$. If $u \cap (x \cup y) \neq \emptyset$, then the conclusion follows trivially. If $u \cap \text{int}(\text{cl}(x \cup y)) \neq \emptyset$, then $\text{cl}(u) = \text{cl}(\text{int}(u))$ since $u \in Y$, and so $\text{cl}(\text{int}(u)) \cap \text{int}(\text{cl}(x \cup y)) \neq \emptyset$, and $\text{cl}(a) \cap \text{int}(b) \neq \emptyset \rightarrow a \cap \text{int}(b) \neq \emptyset$. So $\text{int}(u) \cap \text{int}(\text{cl}(x \cup y)) \neq \emptyset$. Then, $\text{int}(u) \cap \text{cl}(x \cup y) \neq \emptyset$, and so $\text{int}(u) \cap (\text{cl}(x) \cup \text{cl}(y)) \neq \emptyset$, or equivalently $\text{int}(u) \cap \text{cl}(x) \neq \emptyset \vee \text{int}(u) \cap \text{cl}(y) \neq \emptyset$. So $u \cap x \neq \emptyset \vee u \cap y \neq \emptyset$. \square

Now let's consider the proof for (A6): $RT_T \vdash \forall x \forall y (O(x, y) \rightarrow \exists z \forall u (C(u, z) \leftrightarrow \exists v (P(v, x) \& P(v, y) \& C(v, u))))$ where $z = x * y$. As with (A5), we will show that $\llbracket x * y \rrbracket_g = \llbracket x \rrbracket_g \cap \llbracket y \rrbracket_g$, and this amounts to showing:

$$(*) \quad \forall x \in Y \forall y \in Y (\exists z \in Y (z \subseteq x \& z \subseteq y) \rightarrow \forall u \in Y (u \cap (x \cap y) \neq \emptyset \leftrightarrow \exists v \in Y (v \subseteq x \& v \subseteq y \& v \cap u \neq \emptyset)))$$

To show (*) we note the following fact of our models (which follows from the fact that $x \cap^* y \subseteq x \cap y$ and that $x \cap^* y \in Y$, if $\text{int}(x \cap y) \neq \emptyset$): $\forall x \in Y \forall y \in Y (\exists z \in Y (z \subseteq x \& z \subseteq y) \leftrightarrow \text{int}(x \cap y) \neq \emptyset)$. So to prove (*), assume $z \subseteq x \& z \subseteq y$. By the fact above $\text{int}(x \cap y) \neq \emptyset$. So now we must show: $u \cap (x \cap y) \neq \emptyset \leftrightarrow \exists v \in Y (v \subseteq x \& v \subseteq y \& v \cap u \neq \emptyset)$.

\rightarrow : Take $v = x \cap y$
 \leftarrow : By hypothesis $v \subseteq x \cap y$. So $\text{cl}(\text{int}(v)) \subseteq \text{cl}(\text{int}(x \cap y))$; $\text{cl}(v) = \text{cl}(\text{int}(v))$ since $v \in Y$; so $v \subseteq \text{cl}(\text{int}(x \cap y))$; so, $v \subseteq (x \cap y) \cap \text{cl}(\text{int}(x \cap y))$. So $v \subseteq x \cap^* y$. Since $v \cap u \neq \emptyset$ by hypothesis, $u \cap (x \cap^* y) \neq \emptyset$. \square

The proofs that $RT_T \vdash (A7)$ and $RT_T \vdash (A8)$ are similar to the proof of (A6). These proofs establish that the interpretation of the operators i and \sim are the operators int and \sim .

Proof for (A9): Since the interpretation of the operators i and \sim are the operators int and \sim , the definition in RT_0 of the operator c for all variables not equal to a^* and the topological theorem: $\text{cl}(x) = \sim \text{int}(\sim x)$ yield that $\llbracket cx \rrbracket_g$ for $g(x) \neq X$ is $\text{cl}(g(x))$. The interpretation of c as cl can be extended to all of Y , so that $RT_T \vdash c(a^*) = a^*$. \square

Proof for (A10): $RT_T \vdash (\text{OP}(x) \& \text{OP}(y) \& O(x, y)) \rightarrow \text{OP}(x * y)$. It suffices to show $\forall x \in Y \forall y \in Y [(\text{open}(x) \&$

$\text{open}(y) \& \exists z \in Y (z \subseteq x \& z \subseteq y)) \rightarrow \text{open}(x \cap^* y)]$. Assume $(\text{open}(x) \& \text{open}(y) \& \exists z \in Y (z \subseteq x \& z \subseteq y))$; in view of the fact used to verify (A6) about the existence of $x \cap^* y$, $\text{int}(x \cap y) \neq \emptyset$. But since $(\text{open}(x) \& \text{open}(y)) \rightarrow \text{open}(x \cap y)$, then $\text{int}(x \cap y) = x \cap y$ and since $x \cap y \subseteq \text{cl}(x \cap y)$, $x \cap^* y = x \cap y \cup \text{cl}(x \cap y) = x \cap y$. \square

That RT_T verifies (A11) and (A12) comes directly from conditions (vii) and (viii) on Y , and that RT_T verifies (A13) follows easily from the definition of the function f . \square

4 Completeness

Since our axioms for our mereotopology are all 1st order, completeness of RT_0 amounts to showing completeness for a particular 1st order theory. This means that we need to show that if ϕ is not a theorem then we can construct a counter model of the appropriate sort. We do this by means of the usual Henkin method, in which three lemmas are crucial:

Lindenbaum Lemma: Every RT_0 consistent set of sentences can be extended to a maximal consistent set.

Saturation or Witness Lemma: Every RT_0 consistent set of sentences Σ can be extended to a saturated set Σ' in the extension of L , $L(a_0, a_1, \dots)$, such that $\Sigma' \vdash \exists x A \rightarrow A(a_n/x)$, for every formula with one free variable A and a_n is a witness for x .

Henkin Lemma: Every RT_0 maximal consistent saturated set Σ yields an MT_0 model \mathcal{M}_Σ such that $\mathcal{M}_\Sigma \vdash \phi$ iff $\phi \in \Sigma$.

The first two lemmas have a standard proof. But for the proof of the Henkin Lemma, we must spell out in detail the construction of \mathcal{M}_Σ .

We assume that every consistent RT_0 set can be extended to a maximal, RT_0 consistent saturated set. Given a maximal RT_0 -consistent, saturated set Σ , we have a collection of constants occurring in Σ , Σ_C , that will form the basis of our mereotopology; equivalence classes of these constants represent our objects. But since we must model C by non-empty intersection, we must in the model \mathcal{M}_Σ , constructed from Σ , represent these objects by sets of points. We will in fact define points by means of our basic objects, by appealing to an ultrafilter-like construction. It is similar to that used e.g., by [12] to construct temporal instants from temporally extended intervals and states. Clarke [7] also suggests that the ultrafilter construction can be used to reconstruct points, but he uses just one construction adapted from Russell and Wiener that actually makes the system inconsistent as soon as there is one external connection [20]. We use two sorts of ultrafilter constructions, one for interior points (IP) and one for boundary points (BP).

$$IP(\alpha) =_{\text{def}} \alpha \subseteq \Sigma_C \& \alpha \neq \emptyset \& \quad (a)$$

$$\forall x \forall y ((x \in \alpha \& y \in \alpha) \rightarrow (O(x, y) \& x * y \in \alpha)) \& \quad (b)$$

$$\forall x \forall y ((x \in \alpha \& P(x, y)) \rightarrow y \in \alpha) \& \quad (c)$$

$$\alpha \text{ maximal-i.e., } \forall \beta (\beta \text{ verifies (a), (b) and (c)}) \& \quad (d)$$

$$BP(\alpha) =_{\text{def}} \alpha \subseteq \Sigma_C \& \exists x \exists y (x \in \alpha \& y \in \alpha \& EC(x, y)) \& \forall x$$

$$\forall y [(x \in \alpha \& y \in \alpha) \rightarrow ((O(x, y) \& x * y \in \alpha) \vee \exists z \exists t$$

$$(z \in \alpha \& t \in \alpha \& P(z, x) \& P(t, y) \& EC(z, t)))] \& \forall x$$

$$\forall y ((x \in \alpha \& P(x, y)) \rightarrow y \in \alpha) \& \alpha \text{ maximal}$$

We are now in a position to construct the model \mathcal{M}_Σ . We associate with each equivalence class of constants $[c_n] = \{c : \text{HRT} \vdash c = c_n\}$ for each c_n in Σ_C , the set of points $\Omega_{[c_n]} = \{\alpha : P(\alpha) \vee BP(\alpha)\}$ & $\{c_n \in \alpha\}$. The domain in \mathcal{M}_Σ , $D_{\mathcal{M}_\Sigma}$, is

⁴Since $\forall x \in Y \forall y \in Y (x \subseteq y \rightarrow \forall z \in Y (z \cap x \neq \emptyset \rightarrow z \cap y \neq \emptyset))$, we have $RT_T \vdash P(x, y)$ iff $\llbracket x \rrbracket \subseteq \llbracket y \rrbracket$, an intuitive consequence.

such that $D_{\mathcal{M}_\Sigma} = \{\Omega_{[c_n]} : c_n \in \Sigma_C\}$, $\Pi_{\mathcal{M}_\Sigma}$ is an interpretation function of constants in Σ_C such that $\{c_n\}_{\mathcal{M}_\Sigma} = \Omega_{[c_n]}$, and $f_{\mathcal{M}_\Sigma}$ is a function from $D_{\mathcal{M}_\Sigma}$ to $D_{\mathcal{M}_\Sigma}$ such that $f_{\mathcal{M}_\Sigma}(\Omega_{[c_n]}) = \Omega_{[c_n]}$ where $\Sigma \vdash RT_0 c_m = n(c_n)$. We set $\mathcal{M}_\Sigma = \langle D_{\mathcal{M}_\Sigma}, f_{\mathcal{M}_\Sigma}, \Pi_{\mathcal{M}_\Sigma} \rangle$.

Lemma 6: (Henkin Lemma) $\mathcal{M}_\Sigma \models \phi$ iff $\phi \in \Sigma$.

The proof by induction that $\mathcal{M}_\Sigma \models \phi$ iff $\phi \in \Sigma$ is standard except for the base clause. Our axioms concern only one primitive predicate C and so we need only to check atomic formulas of the form $C(x,y)$. What we need to show is:

$C(a,b) \in \Sigma$ iff $\mathcal{M}_\Sigma \models C(a,b)$

\rightarrow : Assume that $C(a,b) \in \Sigma$. Then $O(a,b) \in \Sigma \vee EC(a,b) \in \Sigma$ by (D3). By the "ultrafilter" constructions, this assures either that an interior point $c \in [a] \cap [b]$ or a boundary point $c \in [a] \cap [b]$. In either case we have $[a] \cap [b] \neq \emptyset$. So we have $\mathcal{M}_\Sigma \models C(a,b)$.

\leftarrow : Now assume $\mathcal{M}_\Sigma \models C(a,b)$. So $[a] \cap [b] \neq \emptyset$ in \mathcal{M}_Σ . But the common points must all either be boundary points or at least some must be interior points. By the "ultrafilter" construction then we conclude $EC(a,b) \in \Sigma$ or $O(a,b) \in \Sigma$. By (D3) and (D4) then, $C(a,b) \in \Sigma$.

The final step in proving completeness is that we need to show that \mathcal{M}_Σ is a model in RT . To show this we must show that every one of the conditions on models in RT is fulfilled. To this end, we must embed the open elements of the domain of \mathcal{M}_Σ in a classical topological space. We have already the set of points that forms the space; it is just $\cup\{\Omega_{[c_n]} : c_n \in \Sigma_C\} =_{\text{def}} \Sigma_U$. Now we must find an embedding of $\{\Omega_{[c_n]} : c_n \in \Sigma_C \ \& \ \Sigma \vdash OP(c_n)\}$ in some set of open sets in $\wp(\Sigma_U)$, to form the topology. We will simply take $\{\Omega_{[c_n]} : c_n \in \Sigma_C \ \& \ \Sigma \vdash OP(c_n)\} \cup \{\emptyset\} \cup \{UX : X \subseteq \{\Omega_{[c_n]} : c_n \in \Sigma_C \ \& \ \Sigma \vdash OP(c_n)\}\} =_{\text{def}} \Sigma_U$ to be the relevant set. $\langle \Sigma_U, \Sigma_U \rangle$ is a structure in which: (i) $\emptyset \in \Sigma_U$; (ii) $x, y \in \Sigma_U \rightarrow x \cap y \in \Sigma_U$; (iii) $X \subseteq \Sigma_U \rightarrow UX \in \Sigma_U$; (iv) $\Sigma_U \in \Sigma_U$. Thus, $\langle \Sigma_U, \Sigma_U \rangle$ is a non-empty topological space (Σ_U is assured to be non-empty because of (A11), (A12) and the "ultrafilter" construction of points). Further, fact 7 now follows immediately from our axioms and definitions.

Fact 7: There is an embedding of $D_{\mathcal{M}_\Sigma}$ (the identity function on $D_{\mathcal{M}_\Sigma}$) into $\langle \Sigma_U, \Sigma_U \rangle$ such that:

- (i) $[a^*]_{\mathcal{M}_\Sigma} = \Sigma_U$; (ii) $[x+y]_{\mathcal{M}_\Sigma} = [x]_{\mathcal{M}_\Sigma} \cup [y]_{\mathcal{M}_\Sigma}$;
- (iii) $[x \cdot y]_{\mathcal{M}_\Sigma} = [x]_{\mathcal{M}_\Sigma} \cap [y]_{\mathcal{M}_\Sigma}$; (iv) $[-x]_{\mathcal{M}_\Sigma} = \neg [x]_{\mathcal{M}_\Sigma}$;
- (v) $[ix]_{\mathcal{M}_\Sigma} = \text{int}([x]_{\mathcal{M}_\Sigma})$; (vi) $[cx]_{\mathcal{M}_\Sigma} = \text{cl}([x]_{\mathcal{M}_\Sigma})$.

We can now verify that all the conditions of RT hold of \mathcal{M}_Σ relative to the embedding of the previous lemma.

Lemma 8:

- (i) $D_{\mathcal{M}_\Sigma} \subseteq \wp(\Sigma_U) \ \& \ \Sigma_U \in D_{\mathcal{M}_\Sigma}$;
- (ii) $\forall x \in D_{\mathcal{M}_\Sigma} (\text{int}(x) \in D_{\mathcal{M}_\Sigma} \ \& \ \text{int}(x) \neq \emptyset \ \& \ \text{int}(x) = \text{int}(\text{cl}(x)))$;
- (iii) $\forall x \in D_{\mathcal{M}_\Sigma} (\text{cl}(x) \in D_{\mathcal{M}_\Sigma} \ \& \ \text{cl}(x) = \text{cl}(\text{int}(x)))$;
- (iv) $\forall x \in D_{\mathcal{M}_\Sigma} (\text{int}(\sim x) \neq \emptyset \rightarrow \sim x \in D_{\mathcal{M}_\Sigma})$;
- (v) $\forall x \in D_{\mathcal{M}_\Sigma} \ \forall y \in D_{\mathcal{M}_\Sigma} (\text{int}(x \cap y) \neq \emptyset \rightarrow x \cap y \in D_{\mathcal{M}_\Sigma})$;
- (vi) $\forall x \in D_{\mathcal{M}_\Sigma} \ \forall y \in D_{\mathcal{M}_\Sigma} \ x \cup y \in D_{\mathcal{M}_\Sigma}$;
- (vii) $\exists x \in D_{\mathcal{M}_\Sigma} \ \exists y \in D_{\mathcal{M}_\Sigma} (x \cap y \neq \emptyset \ \& \ \text{int}(x \cap y) = \emptyset)$;
- (viii) $\exists x \in D_{\mathcal{M}_\Sigma} \ \exists y \in D_{\mathcal{M}_\Sigma} (x \cap y = \emptyset \ \& \ \forall z \in D_{\mathcal{M}_\Sigma} ((\text{open}(z) \ \& \ x \subseteq z) \rightarrow y \cap \text{cl}(z) \neq \emptyset))$;
- (ix) $\forall x \in D_{\mathcal{M}_\Sigma} (f(x) \in D_{\mathcal{M}_\Sigma} \ \& \ x \subseteq f(x) \ \& \ \text{open}(f(x)) \ \& \ \text{cl}(f(x)) \cap y \neq \emptyset \rightarrow x \cap \text{cl}(f(y)) \neq \emptyset \ \& \ ((\text{open}(u) \ \& \ x \subseteq u) \rightarrow f(x) \subseteq u))$

The proof of (i) follows directly from the definition of $D_{\mathcal{M}_\Sigma}$ and (A4), (v), (vi), (vii) and (viii) follow directly from the axioms (A6, A5, A11 and A12) of RT_0 respectively.

To show (ii), we note that by (A7), for each $a \in \Sigma_C$, $ia \in \Sigma_C$ so $\text{iala} \in D_{\mathcal{M}_\Sigma}$; and by (A1) $C(ia, ia) \in \Sigma$. So by the "ultrafilter" construction and Fact 7, there is a point in iala , i.e., $\text{iala} = \text{int}(\text{iala}) \neq \emptyset$.

We now show that $ia = \text{ica} \in \Sigma$, so that $\text{int}(\text{iala}) = \text{int}(\text{cl}(\text{iala}))$ holds of all objects $D_{\mathcal{M}_\Sigma}$, insofar as they are elements of $\langle \Sigma_U, \Sigma_U \rangle$ and where int and cl are the topological operators in $\langle \Sigma_U, \Sigma_U \rangle$. By axiom (A3) this amounts to showing that $C(b, ia) \rightarrow C(b, ica)$. But $C(b, ia) \rightarrow O(b, a)$ and $C(b, ica) \rightarrow O(b, ca)$ by (A8), (D6) and (D3). So we need to show that $O(b, a) \rightarrow O(b, ca)$. To this end we note: $\vdash RT_0 P(b, ca) \rightarrow P(ib, a)$.

Now to the proof of $O(b, a) \rightarrow O(b, ca)$:

\rightarrow : $O(b, a) \rightarrow \exists z (P(z, b) \ \& \ P(z, a))$ (D3) and $P(a, ca)$ by (D1), (D7), (A7), (A8) and (A9). By the transitivity of P , $O(b, a) \rightarrow \exists z (P(z, b) \ \& \ P(z, ca))$ and so $O(b, ca)$.

\leftarrow : $O(b, ca) \rightarrow \exists z (P(z, b) \ \& \ P(iz, a))$ by (D3) and the fact we noted above. Since $P(\text{id}, d)$ (proof from (A8) and (D1)) and the transitivity of P , $O(b, ca) \rightarrow \exists z (P(iz, b) \ \& \ P(iz, a))$. So $O(b, ca) \rightarrow \exists z' (P(z', b) \ \& \ P(z', a))$ and so $O(b, ca) \rightarrow O(b, a)$.

To prove (iii), we see that by (A8) and (A9), $\forall a \in \Sigma_C$, $ca \in \Sigma_C$ so $\text{ical} \in D_{\mathcal{M}_\Sigma}$. Let's now show that $cx = \text{cix} \in \Sigma$, so that by Fact 7, $\text{cl}(\text{ix}) = \text{cl}(\text{int}(\text{ix}))$ in \mathcal{M}_Σ . If $a = a^*$, then since $ca^* = a^*$ by (A9) and $ia^* = a^*$ (since $\vdash RT_0 \forall x \text{NTP}(x, a^*)$), $\text{cl}(x) = \text{cl}(\text{int}(x))$ in \mathcal{M}_Σ . If $a \neq a^*$, $ca = \text{cia} \rightarrow \neg i(-a) = \neg i(-ia)$ by (D7). Since $\neg a = a$, $ca = \text{cia} \rightarrow i(-a) = i(-(-a))$, then $ca = \text{cia} \rightarrow i(-a) = i(\text{c}(-a))$, but $i(-a) = i(\text{c}(-a))$ is true by the previous result exploiting again the correspondence in Fact 7.

To show (iv), we note that by (A7), $\forall a \in \Sigma_C (a \neq a^* \rightarrow \neg a \in \Sigma_C)$, so if $\text{iala} \in \Sigma_U$ then $\sim \text{iala} \in D_{\mathcal{M}_\Sigma}$. But $\forall a \in D_{\mathcal{M}_\Sigma} (a \neq \Sigma_U \rightarrow \text{int}(\sim ax) \neq \emptyset)$ since $\text{int}(\sim a) \neq \emptyset \rightarrow \text{cl}(a) \neq \Sigma_U$ and $\text{cl}(a) \neq \Sigma_U \rightarrow a \neq \Sigma_U$ (cf lemma in the proof of $RT_T \vdash (A7)$); so if $\text{int}(\sim \text{iala}) \neq \emptyset$ then $\sim \text{iala} \in D_{\mathcal{M}_\Sigma}$.

Finally, to show (ix), we note that by (A13) $\forall a \in \Sigma_C \ \forall b \in \Sigma_C$ so $\text{inal} \in D_{\mathcal{M}_\Sigma}$ and $\text{ial} \subseteq \text{inal} \ \& \ \text{open}(\text{inal}) \ \& \ \forall b \in \Sigma_C ((\text{open}(\text{ib}) \ \& \ \text{ial} \subseteq \text{ib}) \rightarrow \text{inal} \subseteq \text{ib})$. It suffices now to show that $\forall b \in \Sigma_C (\text{cl}(\text{inal}) \cap \text{ib} \neq \emptyset \rightarrow \text{ial} \cap \text{cl}(\text{ib}) \neq \emptyset)$. We have $\neg C(x, y) \rightarrow (C(x, c(ny)) \rightarrow C(c(nx), y)) \in \Sigma$, directly from Fact 3 and Fact 2, and $C(x, y) \rightarrow (C(x, c(ny)) \ \& \ C(c(nx), y)) \in \Sigma$ since $P(x, cnx)$ and $P(y, cny)$ are trivial facts in RT_0 .

From the Lindenbaum, Witness and Henkin lemmas, the desired result now follows:

Theorem 9: Every RT_0 consistent set of sentences has a model in the class RT ; or equivalently, $\vdash_{RT} \phi \rightarrow \vdash_{RT_0} \phi$.

5 Extension : Microscopization

In this section, we explore one refinement of commonsense mereotopology. RT_0 meets our requirements for a topology exemplifying a relational conception of space without mathematical points. For many reasoning tasks and for the semantics of NL expressions this information is sufficient (See e.g. [4] for details). When, for instance, a cup is on top of a table, the proper semantics for the relation between the cup and the table is weak contact. However, looking much closer we may see that there are objects or a space between the cup and the table. As we refine the granularity of our space (that is, as we consider smaller and smaller size objects), we eliminate weak contact relations. Note that this

revision never happens for an external connection, for we cannot find any object between the stem and the cup of the glass, or between one's hand and one's wrist.

This sort of refinement is also present in changes of granularity signified by the use of certain modals in NL. Consider again, for example, the sentences (1) and (2) above in which there is arguably a shift in granularity-passing, e.g., from the description of an object as a point to one where it is described as a surface. Superficially, these descriptions are inconsistent⁵, so we suppose that there is some sort of shift from one model or world to another. We call this sort of shift of granularity *microscopization*. We can suppose that this process of refinement of granularity continues to a "limit" as follows. We suppose that we can find parts of each object in the space. Next, we suppose we could find parts of those parts, and so on forming a sequence of objects decreasing in size. These sequences of objects are, mathematically speaking Cauchy sequences; and if we suppose that the limits of these sequences exist, then we will have constructed dense spaces of points by partitioning the original objects of our commonsense conception of space based on middle-sized objects.

The use of modal adverbs like *actually* (in sentence (1) above) suggests that we should model this process of reflection *modally*, each world will reflect an increasingly fine-grained view of what the parts of objects are. At the end, we pass from the notion of a physical part to a mathematical part. We can axiomatize the idea that there are worlds of commonsense mereotopology and there are refined worlds with traditional, mathematical conceptions of topology.

The modal extension for L_{RT} that we envision will include a modal operator $[m]$. To the usual clauses for a first order language, we add: if ϕ is a formula of L_{RT} , then $[m]\phi$ is also a formula. A modal RT_0 model is a quintuple, $\langle W, D, R, c, [m] \rangle$, where W is a nonempty set (of worlds), D a function from W into a discrete, "pseudotopological" structure of the sort outlined in the semantics for L_{RT} in section 1 above; R a transitive and asymmetric alternativeness relation in W ; c an injective counterpart function from $D(w)$ into $D(w')$ for each w and w' such that wRw' such that if $a \cap b \neq \emptyset$ in w then for all w' such that wRw' $c(a) \cap c(b) \neq \emptyset$ in w' ; and ll an assignment to each nonlogical constant of an appropriate intension (function from worlds to extensions).

The domain of each accessible world is strictly larger than its R -predecessors (it is this that allows us to capture the asymmetry of R), reflecting the refinement of the granularity size of objects. We further suppose that for each object a in w and wRw' , w' contains additional objects that are parts of a (or strictly speaking the counterpart of a). The collections of points in one world need not be the same collection in another, but an object (collection of points in one world) will always have a counterpart in each accessible world. Finally, we suppose a limit to the R chains in which every object is the sum of points without interiors. This falsifies certain axioms of RT_0 at the limit worlds—namely the axiom that every object has a nonempty interior. These limit worlds are clas-

sical topologies of the sort we used to build the models for RT_0 . We insist that every world other than a limit world is in effect a model in RT , and the limit worlds verify all except (A1), (A8), (A11), (A12) and (A13). We will call the class of all modal models MRT . Because MRT is a modal semantics with variable domains, we must change the notion of truth and validity. Following standard techniques (see [11]) we will say that a formula ϕ is either verified (1) falsified (0) or undefined at a world. An atomic formula ϕ is undefined at w if it contains a constant that denotes an object $\notin D(w)$ or a variable whose assignment is not in $D(w)$. The recursive satisfaction definition follows the strong Kleene semantics. We say that ϕ is *MRT valid* iff in every MRT model M and every world w in M if ϕ is defined at w in M , then ϕ is true at w in M .

Axiomatization: In addition to the axioms (A2-7) and (A9-10) of RT_0 , we now have the axioms below for MRT_0 (modal RT_0). We define **LIMIT** as:

$$\text{LIMIT} =_{\text{def}} \exists w \forall u (P(w, u) \ \& \ \neg C(w, u) \ \& \ (u \neq w \rightarrow C(u, u))) \ \& \ \forall x \forall y (C(y, x) \rightarrow O^*(y, x)) \ \& \ \neg(A12) \ \& \ \neg(A13)$$

This formula describes these limit worlds; it asserts the existence of \emptyset which is unique by extensionality. Below we will refer to this element as Null and its existence if only in limit worlds prompts us to add a new notion of overlap (O^*). Of interest are the axioms (B1), (B2), and (B7)-(B10) and Nec^* . (B2) says that once two objects are connected they are always connected under microscopization. (B8) captures the process of microscopization, while (B9) says that a microscopization of the current world always exists or else the current world is already a limit world. (B7) asserts that such limit worlds exist. (B10) asserts that the whole space is connected, while (B1) and (Nec^*) say that the normal axioms of RT_0 hold except at limit worlds.

- (B1) $((A1) \ \& \ (A8) \ \& \ (A11) \ \& \ (A12) \ \& \ (A13)) \vee \text{LIMIT}$
- (B2) $\forall x \forall y (C(x, y) \rightarrow [m]C(x, y))$
- (B3) $[m]\phi \rightarrow [m][m]\phi$
- (B4) $[m]\forall x \phi \rightarrow \forall x [m]\phi$
- (B5) $\exists x [m]\forall y P(y, x)$
- (B6) $\phi \rightarrow \psi \rightarrow [m](\phi \rightarrow \psi)$
- (B7) $[m] \rightarrow [m] \rightarrow \text{LIMIT}$
- (B8) $[m]\forall x [m]\exists y, y' ((P(y, x) \ \& \ P(y', x) \ \& \ \neg O(y, y'))$
- (B9) $([m]\phi \rightarrow \neg [m]\neg \phi) \vee \text{LIMIT}$
- (B10) $\text{Con}(a^*)$
- (D3') $O^*(x, y) =_{\text{def}} \exists z (z \neq \text{Null} \ \& \ P(z, x) \ \& \ P(z, y))$

Nec^* : For all instances of the axioms of FOL with identity, (A2-7), (A9-10), (B1-10) $\phi, \vdash_{MRT_0} [m]\phi$

6 Related Work

Qualitative spatial reasoning has in the past focussed more on reasoning about orientations than on topology. A number of authors have defined topological relations from orientational primitives, extending Allen's interval calculus [1] to 3 dimensions (for instance [15]), but it is easily shown that mereotopological concepts such as overlap or external connection can be correctly grasped in this case only for a quite limited domain of parallelepipedic objects [20].

Others such as [8] use a domain of points and all the power of classical topology (and Euclidean geometry), on the grounds that it was the only sound theory modelling these concepts. Our work shows that an alternative based on a naive conception of space as it is expressed in NL can be de-

⁵The "points" referred to here are in effect extended objects, i.e. they have nothing to do with the abstract points in the models, but they are subject to constraints that are incompatible with those constraints that define surfaces (see [4] for the definitions of NL notions of point, border and surface objects in RT_0).

veloped rigorously at least as far as the topology is concerned. Furthermore, the full topology of R^3 is a higher order concept and we have proposed a first-order and axiomatizable alternative.

As far as we know, the only other mereotopological approach in qualitative reasoning apart from ours has been taken by the AI group in Leeds, the last version of their theory being presented in [16]. While also based on Clarke's proposal, they remove the topological definitions of interior and closure. They also change the complement definition so that it is equivalent to the two following axioms $\forall x \exists y \forall u (C(u,y) \rightarrow \neg NTPP(u,x))$ and $P(x,y) \rightarrow \forall u \exists v (C(u,v) \& (C(x,v) \rightarrow NTPP(v,y)))$. As shown in [16], these imply that there can be no atoms, and that it is now impossible to assert the existence of interiors-otherwise the theory is inconsistent. This last feature is one of the motivations for the change in the theory, since for them, differentiating between an individual, its closure and its interior has no cognitive support. On our point of view, it is on the contrary cognitively important to be able to view material objects as closed individuals and their complements as open ones, so that their interpretations don't share any point. Indeed, we don't want the air around the glass to have a "glass boundary" belonging to it, that is why in RT_0 , the glass and the air are in weak contact. But in [16], an object and its complement are externally connected, and so in that theory the unintuitive consequence about glass boundaries seems to follow unless one does a lot of fiddling with the way we refer to objects in NL. We have two strong motivations for wanting topological operators in our theory, one being the possibility of defining strong contact and weak contact, the other being that we think it ontologically important to be able to recover classical topology as a "limit" model, which we do in the process of microscopization. Note that the authors of [16] don't provide any soundness or completeness proofs for their theory.

Recently, in formal ontology there has been much work done on the ontological relation between mereological concepts and topological ones. [19] provides a nice description and a classification of this work. One active trend in this field is to show that mereology alone supports topology, at the cost of having domains containing both extended individuals and boundaries [9, 5, 17]. It seems to us important to guarantee the ontological homogeneity of the domain, in order to avoid the need to classify *a priori* the spatial entities we will represent in the theory, which even if metaphysically justifiable does not seem to us relevant to spatial reasoning tasks in AI. Besides, we have already stressed the fact that no linguistically described entity corresponds to an infinitely thin mathematical object, so that considering boundaries as constructs more abstract than extended individuals, as we do, is cognitively grounded. In addition, we note that none of these authors provide a soundness or completeness proofs to support their intuitions about the kind of models their theories are supposed to have.

References

1. J. Allen. Maintaining knowledge about temporal intervals. *Communications of the ACM* 26(11):832-843, 1983
2. N. Asher, M. Aurnague, M. Bras, P. Sablayrolles & L. Vieu. Computing the spatiotemporal structure of discourse. In *Proceedings of IWCS'94*, Tilburg, 1994.
3. N. Asher & J. Lang: When nonmonotonicity comes from distances. In *Proceedings of KI'94*, pp. 308-318, Berlin: Springer, 1994.
4. M. Aurnague, L. Vieu. A three-level approach to the semantics of space. In: C. Zelinsky-Wibbelt (ed.) *The Semantics of Prepositions - From Mental Processing to Natural Language Processing.*, pp. 393-439, Berlin: Mouton de Gruyter, 1993.
5. R. Chisholm. Spatial continuity and the theory of part and whole - A Brentano study. *Brentano Studien*, 4:11-23, 1992/93.
6. B. Clarke. A calculus of individuals based on "connection". *Notre Dame Journal of Formal Logic* 22(3):204-218, 1981.
7. B. Clarke. Individuals and points. *Notre Dame Journal of Formal Logic* 26(1):61-75, 1985.
8. E. Davis. *Representations of commonsense knowledge*. Palo Alto: Morgan Kaufmann, 1990.
9. C. Eschenbach. A mereotopological definition of point. In: C. Eschenbach, Ch. Habel & B. Smith (eds.): *Topological foundations of Cognitive Science*, Papers from the Workshop at the FISCI-CS, Buffalo, July 1994.
10. P. Hayes. The second naive physics manifesto. In: J.R. Hobbs & R.C. Moore (eds.) *Formal theories of the commonsense world*, pp. 1-36, Norwood: Ablex, 1985.
11. G. Hughes & M. Cresswell. *An introduction to modal logic*. London: Routledge, 1968.
12. H. Kamp. Events, instants and temporal reference - In R. Bauerle, U. Egli, A. von Stechow (eds.) *Meaning, use and interpretation of language*, pp. 376-417, Berlin: de Gruyter, 1979.
13. B. Kuipers. Modelling spatial knowledge. *Cognitive Science*, 2(2): 129-154, 1978.
14. S. Lesniewski. O podstawach matematyki. *Przegląd filozoficzny*, 30-34, 1927-1931.
15. A. Mukerjee, G. Joe. A Qualitative Model for Space. In *Proceedings of AAAI'90*, pp. 721-727, Cambridge (MA): MIT Press, 1990.
16. D. Randell, Z. Cui & A. Cohn. A spatial logic based on regions and connection. In *Proceedings of KR'92*, pp 394-398, Los Altos: Morgan Kaufmann, 1992.
17. B. Smith. The formal ontology of space. In: L. Hahn (ed.) *Roderick Chisholm*, Library of living philosophers. La Salle: Open Court, to appear.
18. A. Tarski. Les fondements de la geometrie des corps. In: A. Tarski *Logique, Semantique, Metamathematique*. Paris: Armand Colin, 1972.
19. A. Varzi. On the boundary between mereology and topology, Technical Report #9308-04, IRST Trento, 1993.
20. L. Vieu. Semantique des relations spatiales et inferences spatio-temporelles -Ph.D. dissertation. Toulouse: Universite Paul Sabatier, 1991.
21. A.N. Whitehead. *Process and Reality*, New York: Macmillan, 1929.