

## Article (refereed)

---

Thomas, Philippe J.; Mineau, Pierre; **Shore, Richard F.**; Champoux, Louise; Martin, Pamela A.; Wilson, Laurie K.; Fitzgerald, Guy; Elliott, John E.. 2011 Second generation anticoagulant rodenticides in predatory birds: Probabilistic characterisation of toxic liver concentrations and implications for predatory bird populations in Canada. *Environment International*, 37 (5). 914-920. [10.1016/j.envint.2011.03.010](https://doi.org/10.1016/j.envint.2011.03.010)

Copyright © 2011 Elsevier Ltd.

This version available <http://nora.nerc.ac.uk/14168/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the authors and/or other rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

**This document is the author's final manuscript version of the journal article prior to the peer review process. Some differences between this and the publisher's version may remain. You are advised to consult the publisher's version if you wish to cite from this article.**

[www.elsevier.com/](http://www.elsevier.com/)

Contact CEH NORA team at  
[noraceh@ceh.ac.uk](mailto:noraceh@ceh.ac.uk)

**Title:** Second generation anticoagulant rodenticides in predatory birds: probabilistic characterisation of toxic liver concentrations and implications for predatory bird populations in Canada.

**Authors:** Philippe J. Thomas<sup>1</sup>, Pierre Mineau<sup>1</sup>, Richard F. Shore<sup>2</sup>, Louise Champoux<sup>3</sup>, Pamela A. Martin<sup>4</sup>, Laurie K. Wilson<sup>5</sup>, Guy Fitzgerald<sup>6</sup>, John E. Elliott<sup>7</sup>

<sup>1</sup> National Wildlife Research Centre, Science & Technology Branch, Environment Canada, Ottawa, Ontario, K1A 0H3, Canada, [philippe.thomas@ec.gc.ca](mailto:philippe.thomas@ec.gc.ca) (corresponding author), [pierre.mineau@ec.gc.ca](mailto:pierre.mineau@ec.gc.ca),

Tel: (613) 998-0511 Fax: (613) 998-0458

<sup>2</sup> Centre for Ecology & Hydrology, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster LA1 4AP, UK, [rfs@ceh.ac.uk](mailto:rfs@ceh.ac.uk)

<sup>3</sup> Science & Technology Branch, Environment Canada, Ste-Foy, QC, G1V 4H5, Canada, [louise.champoux@ec.gc.ca](mailto:louise.champoux@ec.gc.ca)

<sup>4</sup> Science & Technology Branch, Environment Canada, Burlington, ON, L7R 4A6, Canada, [pamela.martin@ec.gc.ca](mailto:pamela.martin@ec.gc.ca)

<sup>5</sup> Canadian Wildlife Service, Environment Canada, Delta, BC, V4K 3N2, Canada, [laurie.wilson@ec.gc.ca](mailto:laurie.wilson@ec.gc.ca)

<sup>6</sup> Université de Montréal, Saint-Hyacinthe, QC, J2S 7C6, Canada, [guy.fitzgerald@umontreal.ca](mailto:guy.fitzgerald@umontreal.ca)

<sup>7</sup> Science & Technology Branch, Environment Canada, Delta, BC, V4K 3N2, Canada, [john.elliott@ec.gc.ca](mailto:john.elliott@ec.gc.ca)

**Abstract:**

Second-generation anticoagulant rodenticides (SGARs) are widely used to control rodent pests but exposure and poisonings occur in non-target species, such as birds of prey. Liver residues are often analyzed to detect exposure in birds found dead but their use to assess toxicity of SGARs is problematic. We analyzed published data on hepatic rodenticide residues and associated symptoms of anticoagulant poisoning from 270 birds of prey using logistic regression to estimate the probability of toxicosis associated with different liver SGAR residues. We also evaluated exposure to SGARs on a national level in Canada by analyzing 196 livers from great horned owls (*Bubo virginianus*) and red-tailed hawks (*Buteo jamaicensis*) found dead at locations across the country. Analysis of a broader sample of raptor species from Quebec also helped define the taxonomic breadth of contamination. Calculated probability curves suggest significant species differences in sensitivity to SGARs and significant likelihood of toxicosis below previously suggested concentrations of concern (<0.1 mg/kg). Analysis of birds from Quebec showed that a broad range of raptor species are exposed to SGARs, indicating that generalized terrestrial food chains could be contaminated in the vicinity of the sampled areas. Of the two species for which we had samples from across Canada, great horned owls are exposed to SGARs to a greater extent than red-tailed hawks and liver residue levels were also higher. Using our probability estimates of effect, we estimate that a minimum of 11% of the sampled great horned owl population is at risk of being directly killed by SGARs. This is the first time the potential mortality impact of SGARs on a raptor population has been estimated.

**Keywords:** rodent, exposure, liver residues, toxicity threshold, anticoagulant rodenticide

SGAR – second generation anticoagulant rodenticide

GHOW – great horned owl

RTHA – red-tailed hawk

BAOW – barred owl

BNOW – barn owl

ALL – all bird species pooled

EIIS – ecological incident information system

EPA – environmental protection agency

1 **1. Introduction**

2 Introduced in the 1970s, second-generation anticoagulant rodenticides (SGARs) were  
3 developed to combat the reported development of rodent resistance to first-generation  
4 compounds (Buckle et al. 1994). These newer anticoagulant poisons differ from their  
5 first-generation counterparts in that they are more acutely toxic at lower doses (often  
6 allowing a lethal dose to be obtained in a single feeding), and are more persistent in  
7 vertebrate livers (Parmar et al. 1987, Stone et al. 1999, Newton et al. 1999, Erickson and  
8 Urban 2004). Greater acute toxicity increases the potential for primary poisoning  
9 amongst non-target species while the longer tissue half-lives of SGARs enhance the  
10 potential for bioaccumulation in non-target predators in particular, and so may increase  
11 the risk of secondary poisoning. Furthermore, rodents survive for several days after  
12 consuming a lethal dose of SGARs and often will continue feeding on the bait (Cox and  
13 Smith 1992). That increases the likelihood that the body burden in poisoned rodents may  
14 significantly exceed the LD50 or even LD100 dose, and poisoned animals may remain  
15 active and available for capture by predators for some period after ingestion of the  
16 rodenticide. Additionally, poisoned rodents exhibit an altered state of behaviour, such as  
17 spending more time in open areas in a lethargic state, and this may further predispose  
18 them to predation (Cox and Smith 1992).

19 SGARs bind and inhibit vitamin K epoxide reductase and persist for at least six  
20 months in organs and tissues containing this enzyme such as the liver (Stone et al. 1999,  
21 Eason et al. 2002). In an attempt to monitor exposure in non-target wildlife, the presence  
22 of detectable SGAR residues as well as the magnitude of concentrations has been  
23 measured in the livers of some Canadian, American and European predatory birds and

24 scavengers (Albert et al. 2010, Newton et al. 1990, Shore et al. 1999, Shore et al. 2006).  
25 There was a common trend among those studies for most SGARs, namely brodifacoum,  
26 bromadiolone, difenacoum and difethialone being detected at an increasing frequency in  
27 numerous predators and scavengers. Species most commonly monitored in North  
28 America are great horned owls (*Bubo virginianus*) and red-tailed hawks (*Buteo*  
29 *jamaicensis*) (Albert et al. 2010, Erickson and Urban 2004).

30 It is still uncertain what SGAR liver concentration is diagnostic of a potentially lethal  
31 dose and, indeed Erickson and Urban (2004) have questioned whether such a cause-effect  
32 relationship is appropriate. A sometimes cited –toxicity threshold is given as –greater  
33 than 0.1 – 0.2 mg/kg wet weight (Newton et al. 1998, Newton et al. 1999). This was, in  
34 fact, described as a –potentially lethal range and was derived for a single species, the  
35 barn owl (*Tyto alba*); it stems from two sets of observations (Shore et al. 2001). Firstly,  
36 barn owls diagnosed post-mortem as having died from rodenticides had liver  
37 concentrations > 0.1 mg/kg. Those owls exhibited classical toxicosis signs such as  
38 haemorrhaging from organs such as the heart, lungs, liver, brain and/or subcutaneous  
39 areas (Newton et al. 1998). Secondly, owls that were experimentally poisoned had liver  
40 residues in the range of 0.2 – 1.72 mg/kg (Newton et al. 1999). However, it is uncertain  
41 whether these barn owl criteria would apply to other species. Liver residues associated  
42 with SGAR poisonings in various species typically range over two orders of magnitude  
43 and were reported to be as low as 0.01 mg/kg wet wt in one great horned owl that was  
44 examined (Stone et al. 1999). Thus, liver SGAR concentrations associated with toxicity  
45 vary markedly among both individuals and species. This suggests a probabilistic  
46 approach; which we adopt to review the evidence pertaining to how liver residues are

47 related to toxicity. Our principal objectives are: i) to determine SGAR liver  
48 concentrations that may be associated with mortality in birds (ie- to quantify the -toxicity  
49 threshold) and ii) using the threshold values, assess the extent and severity of exposure  
50 in Canadian birds of prey.

51

## 52 **2. Methods**

### 53 **2.1. Toxicity Threshold**

#### 54 2.1.1. Literature Search

55 Recently published (~ last 10 years) peer-reviewed publications as well as the United  
56 States Environmental Protection Agency (EPA)'s Ecological Incident Information  
57 System (EIS) were surveyed in order to locate liver residue data sets for birds of prey.  
58 The EIS is the EPA's database managing information on incidents linked to the exposure  
59 of non target plants and animals to pesticides. It is currently managed by the Office of  
60 Pesticide Programs (Mastrota 2007). Data were retained for our assessment if they met a  
61 set of pre-determined conditions. These conditions included:

- 62 i) SGAR detection limits in liver were under 0.02 mg/kg wet wt;
- 63 ii) post-mortem evaluations were conducted prior to liver extraction and analysis;  
64 pathophysiological signs of rodenticide poisoning were included.
- 65 iii) post-mortem evaluations were conducted by a reputable professional such as a  
66 doctor of veterinary medicine (DVM); and
- 67 iv) adequate sample sizes were available ( $n > 15$ ) for any given species (in order to  
68 have greater statistical power).

#### 69 2.1.2. Data Analysis

70 Raptor necropsies with attending SGAR liver analyses were collected and compiled  
71 in database software, and each case was given a binary code as positive (1) or negative  
72 (0) for pathophysiological signs of poisoning. A positive coding meant that, after a  
73 detailed post-mortem evaluation, an anticoagulant was diagnosed as being the cause of  
74 death or a significant contributory factor (ie- when necropsies showed hemorrhage or  
75 anemia in the absence of traumatic injury or infectious or parasitic diseases and an  
76 anticoagulant residue was detected in the liver). A negative coding represented cases  
77 where the cause of death was deemed to be natural or accidental (for example incidental  
78 take, hunting, motor vehicle collisions, starvation).

79 The binary dataset was imported into SAS/STAT (version 9.2 TS2M0). Residue  
80 concentrations of all SGAR compounds were summed for the logistic regression.  
81 Concentrations were log transformed to meet the assumption of normality and re-tested.  
82 The PROC LOGISTIC macro was invoked to determine how liver residues affected  
83 presence or absence of poisoning symptoms. An effects plot was generated to illustrate  
84 the relationship and equations were built for every species with sufficient data ( $n \geq 15$ ).  
85 Using these equations, liver residue levels (in mg/kg wet weight (ww)) were determined  
86 for probabilities of 5%, 10%, 15% and 20% of exhibiting pathologies consistent with  
87 rodenticides exposure. Species comparisons were completed using analysis of variance  
88 (ANOVA) in conjunction with Tukey's Studentized Range test. Because all birds were  
89 found dead or moribund, there was a logical inference that those pathologies  
90 (haemorrhaging of the heart, lungs, liver, brain and/or subcutaneous areas) were  
91 responsible for, or strongly contributed to, the mortality of the individual.

## 92 **2.2. Exposure extent in Canada**

93 2.2.1. Sample Collection

94 To obtain a cross-Canada survey of residue levels, liver samples of birds were  
95 selected, irrespective of the cause of death, from British Columbia, the prairie provinces,  
96 Ontario and Quebec. The birds were collected near agricultural and urban areas of the  
97 country where SGAR use was thought to be common. They were typically submitted to  
98 rehabilitation or veterinary centres either dead or in a moribund state. Initial diagnosis  
99 frequently involved car strike or other obvious ‘mishap’. They were not chosen because  
100 they showed signs of anticoagulant poisoning, but rather reflect the population of  
101 reported birds of prey dying from a multitude of causes. The subsequent liver samples  
102 were harvested initially as part of previous investigations of exposure to heavy metals or  
103 other toxicants, and then rodenticides residues were determined in later years. Three main  
104 collections were sampled. These included an Ontario/prairie sample of red-tailed hawks  
105 and great-horned owls, two common species known to scavenge; a broader phylogenetic  
106 collection from Quebec and a collection of three owl species from British Columbia (barn  
107 owl, barred owl [*Strix varia*] and great-horned owl). Those owl species are less mobile  
108 than most of the hawk species and were chosen to help identify geographical patterns of  
109 contamination and hence, potential sources of rodenticides residues. Results from the  
110 latter have already been reported (Albert et al. 2010).

111 2.2.2. Chemical Analysis

112 Chemical analysis was conducted at the National Wildlife Research Center in Ottawa,  
113 Ontario, Canada. Methods were similar to those reported by Albert et al. (2010). 50 mg of  
114 liver was ground in a mortar with about 5 g anhydrous sodium sulphate (Fisher no. S420-  
115 3). The resulting mixture was transferred to an amber glass septum bottle and acetonitrile

116 (EMD Omnisolv, AX0142-1, HPLC grade; 1 x 7 mL and 2 x 5 mL) was used for  
117 extraction. The extract was shaken for 2 minutes by hand and 15 minutes mechanically.  
118 After centrifuging for 15 minutes at 1000 rpm, the supernatant was removed and  
119 transferred into a 40 mL conical tube. The supernatant of the two subsequent extractions  
120 were combined with the first supernatant. The total product was evaporated to 10 mL  
121 under a stream of nitrogen in a water bath kept at 40°C.

122 In order to clean up liver extract, a 2 mL portion was transferred into a test tube and  
123 heated to dryness. The sample was reconstituted in acetonitrile and cleaned by solid-  
124 phase extraction. After the introduction of the sample into the SPE cartridge, the tube  
125 containing the sample was rinsed with acetonitrile and added to the SPE cartridge  
126 solution. The eluate was then evaporated to dryness and reconstituted in MeOH and  
127 filtered through an Acrodisc® syringe filter with a polyvinylidene fluoride (PVDF)  
128 membrane. A volume of 10 µL of the diluted filtered extract was analyzed by liquid  
129 chromatography-mass spectrometry (LC-MSMS). Some of the owl samples analysed  
130 (mainly from British Columbia) were not cleaned using an SPE cartridge. However,  
131 limits of detection were calculated for the procedure with and without an SPE sample  
132 cleaning phase and were found to be identical. For this reason, both SPE-cleaned data and  
133 non-SPE data were pooled for our analysis.

134 Brodifacoum, bromadiolone and difethialone were detected with a triple quadrupole  
135 mass Quatro-Ultima (Waters) with negative electrospray ionization (ESI) in multiple  
136 reaction monitoring scanning mode (MRM). LC-MSMS, MRM parameters and triple  
137 quadrupole settings were identical as the ones reported in Albert et al. 2010.



138 The method's detection limit was 0.005 mg/kg for difethialone and 0.002 mg/kg for  
139 brodifacoum and bromadiolone. The standards were all analytical grade (>98% purity). A  
140 calibration curve was built with five levels of concentrations ranging from 2.5 to 80 pg  
141 with an  $r^2 > 0.99$ . Samples were diluted in order to fit within the limits of the calibration  
142 curve. Recoveries at low and high level were >70% for all compounds. Known amounts  
143 of coumatetralyl (5 pg/L; transition 291.00>140.90) and flocoumafen (1 pg/L; transition  
144 541.40>382.00) were added to each sample prior to the injection allowing ion  
145 suppression monitoring. Methanol was injected between each sample to monitor any  
146 possible contamination.

### 147 2.2.3. Statistical Analysis

148 Since great horned owls and red-tailed hawks represented the two species consistently  
149 found across Canada (no red-tailed hawk samples were submitted from British Columbia,  
150 however) and for which we had a large enough sample size to warrant a meaningful  
151 analysis, cumulative frequency distribution graphs were constructed for these species.  
152 The graphs were generated through a bootstrapping procedure (501 samples) using  
153 BurrliOZ (version 1.0.14, © Commonwealth Scientific and Industrial Research  
154 Organisation, Australia 2000). Using the values identified in our toxicity threshold  
155 analysis, it was possible to identify the percentage of the sampled population exposed to  
156 SGARs belonging to a certain risk category (5%, 10%, 15% and 20% risk of becoming  
157 symptomatic).

158

## 159 **3. Results**

### 160 **3.1. Toxicity Threshold**

161 Five sources of data matched our criteria and were used in the analysis. Data  
162 published by Newton et al. (1990, 1998, 2000; n=45), Albert et al. (2010; n=164) as well  
163 as data from the Ecological Incident Information System (EIIS; n=61). All but four of the  
164 EIIS cases were submitted by the State of New York and several of the values were  
165 published in Stone et al. (1999, 2003). Barn owl samples were collected from localized  
166 areas across Canada and the United Kingdom (UK) with a few individuals from the  
167 United States (USA). Barred owl samples were mostly collected in Canada with only one  
168 from the USA while red-tailed hawk samples were obtained from the USA only. Great  
169 horned owl samples were collected from across both Canada and the USA. Samples were  
170 often collected from relatively developed areas or areas where the public was likely to  
171 report and submit carcasses.

172 There were significant differences between species in liver SGAR concentrations  
173 ( $F_{(4,535)}=12.68$ ,  $p<0.0001$ ). Post hoc-tests (Tukey's Studentized Range test,  $\alpha = 0.05$ )  
174 revealed that, on average, red-tailed hawks (n=32) were the species with the highest liver  
175 concentrations of SGARs (Figure 1). All three owl species (great horned owl [n=86],  
176 barred owl [n=26] and barn owl [n=126]) had SGAR liver residues that were comparable.

177 Logistic regression plots were calculated to predict the probability of a bird being  
178 symptomatic as a function of SGAR liver residues (Figure 2). This was done for each  
179 species separately and for all species combined (total of 270 individuals). Only the  
180 predicted probability curve for the great horned owl (GHOW) was located inside the 95%  
181 confidence limits for the pooled data and the estimated probability of becoming  
182 symptomatic differed significantly between species ( $F_{(1,4)} = 82.9$ ,  $p<0.0001$ ). The curve  
183 for the red-tailed hawk curve differed from those of the three owl species and the curves

184 for the great horned owl and the barn owl also differed from each other (Tukey's  
185 Studentized Range post-hoc test,  $P < 0.05$ ).

186 Using the probability curves, we calculated the predicted SGAR liver residue levels  
187 for different probability risk thresholds for different species (Table 1), although this was  
188 not possible for red-tailed hawks, as the data for this species could not be significantly  
189 modeled by a logistic regression. The majority of the calculated values are under the  
190  $>0.1$ - $0.2$  mg/kg threshold suggested by Newton et al. (1999) and all are below  $0.2$  mg/kg.  
191 If the lower range of  $0.1$  mg/kg and  $0.2$  mg/kg from the potentially lethal range suggested  
192 for barn owls is applied to the barn owl probability curve, they correspond to toxicity  
193 probabilities of  $11\%$  and  $22\%$ , respectively. The higher  $0.7$  mg/kg level proposed by the  
194 Rodenticide Registrants Task Force (Erickson and Urban 2004) corresponds to a  $54\%$   
195 probability of effect in barn owls.

196 Although the differences among the species curves indicate that probabilities of  
197 toxicity should be considered on a species-by-species basis, that is not possible where  
198 data for species are lacking. In such cases, it may be necessary to estimate toxicity  
199 probabilities on the basis of pooled data for other species. The probability curve for the  
200 pooled data in our study predicts that one in 20 birds with detectable residues would  
201 become symptomatic with SGAR liver residues of  $0.02$  mg/kg and one in five when  
202 residue levels reach  $0.08$  mg/kg.

### 203 **3.2. The extent of SGAR exposure in Canada**

204 Of the two species sampled over a relatively broad area of Canada (great horned owl,  
205 red-tailed hawk), great horned owls were most consistently exposed to SGARs (Figure  
206 3). Roughly  $65\%$  of great horned owls across Canada had detectable levels of SGARs in

207 their liver (detection limit of 0.005 mg/kg ww). Frequency of exposure in red-tailed  
208 hawks seemed to increase eastward from the Prairie Provinces to Ontario and Quebec.  
209 The frequency of exposed birds was the lowest (~20%) in the Prairie and Northern  
210 provinces (and territories), increased to ~70% in Ontario and reached the highest in  
211 Quebec (~90% of red-tailed hawks found with detectable SGAR liver residues), although  
212 the sample size in Quebec was smaller than in the other regions. However, as sampling  
213 was fortuitous and sampling effort was not uniform, these spatial comparisons must be  
214 considered preliminary.

215 Great horned owls and red-tailed hawks were exposed to a number of SGARs (Figure  
216 4). The majority of great-horned owls had multiple compounds in the liver; it was the  
217 only species with detectable levels of all three registered compounds. Sixty percent of  
218 red-tailed hawks had detectable liver residues of one or two compounds (Figure 4).

219 Although the proportion of great horned owls with detectable residues was greater than  
220 for red-tailed hawks, this difference was not significant when data were compared for  
221 those provinces from which carcasses of both species were collected (Prairie Provinces,  
222 Ontario and Quebec; (paired t-test,  $t_{(2)} = -0.78$ ,  $p = 0.26$ ; Figure 4). Brodifacoum and  
223 bromadiolone were both detected in great horned owls and red-tailed hawks.  
224 Difethialone was only ever detected in great horned owls (Table 2) but has only been  
225 registered in Canada relatively recently.

226 When the liver SGAR concentrations in great horned owls measured in the present  
227 study were plotted as a cumulative frequency graph (Figure 5; birds with detectable  
228 residues only), it was apparent that approximately 25% had liver SGARs that exceeded  
229 the 20% probability level for effect (0.07mg/kg; Table 1). The lack of a probability curve

230 for red-tailed hawks precludes making a similar calculation for that species, but it is  
231 evident that liver residue levels were much lower than for great-horned owls (Figure 5  
232 and 6). For-example, 50% of great horned owls with detectable residues had liver  
233 concentrations greater than 0.05 mg/kg ww compared with only 10% of red-tailed hawks.  
234 Comparison of liver concentrations in the two species in which birds were matched by  
235 province confirmed that liver residues were significantly higher in the owls than in the  
236 hawks (paired t-test;  $t_{(2)} = -4.0$ ,  $p=0.03$ ). This finding is in contrast to the previously  
237 published literature (Figure 1) where liver residues were higher in red-tailed hawks than  
238 in great-horned owls.

239 Of the small number of individuals from 13 other species analyzed from Quebec,  
240 eight of those had at least one individual with detectable liver SGAR residues (Figure 7).  
241 That indicates that a wide breadth of species is probably also exposed to these  
242 compounds elsewhere in Canada.

243

## 244 **4. Discussion**

### 245 **4.1. Toxicity Threshold**

246 Critical SGAR liver concentrations associated with adverse effects and/or mortality  
247 have not been defined for most raptor species (Walker et al. 2008a), and establishing liver  
248 -toxicity thresholds for SGARs is problematic (Stone et al. 2003). This is partly  
249 because there are a number of factors that contribute uncertainty. For instance, the limit  
250 of quantification used to measure the liver SGAR residues can vary widely with the  
251 analytical method. That can lead to underestimates of the extent of contamination but,  
252 conversely, inflation of residue magnitude if residues which were detected but were

253 below the level of quantification using older analytical methodology were assigned an  
254 inflated limit value (Taylor et al. 2009). Species also vary markedly in their sensitivity to  
255 SGARs. This is known for laboratory mammals (World Health Organisation 1995) but  
256 almost nothing is known about the relative sensitivity of different avian species (Walker  
257 et al. 2008a). Our risk probability curves strongly suggest significant differences exist  
258 among raptor species.

259 To date, the only residue toxicity threshold for SGARs in raptors that has been  
260 suggested is the >0.1-0.2 mg/kg –potentially lethal range for barn owls (Newton et al.  
261 1998, 1999). At best, that provides a range of concern for potential toxicity, and gives no  
262 indication of likelihood of effects. The approach described in the current study offers a  
263 major advance in our ability to assess risk from SGAR residues in that it proposes  
264 quantitative toxicity thresholds for different probability levels of dying from SGAR  
265 intoxication for three species, including the barn owl. If sufficient data were available, it  
266 should be possible to extend this approach to other species. That, in turn, would help to  
267 identify raptor species that may be more sensitive to SGAR toxicity. Overall, on the basis  
268 of the probability curves defined so far, it would seem that the >0.1-0.2 mg/kg level for  
269 barn owls already carries considerable risk of acute intoxication (> 10-20% of barn owls  
270 with this residue being likely to suffer mortality). Clearly, the probability of acute  
271 poisoning associated with the 0.7 mg/kg residue level proposed by the Rodenticide  
272 Registrants Task Force (Erickson and Urban 2004) is worse still.

273 The probabilistic methods described here are, as with all predictive methods, subject  
274 to biases and uncertainties. Of these, perhaps two of the most important are likely to be  
275 underestimation of non-lethal residues, because birds characterised as –zeros in the

276 probabilistic plot may have metabolised some of their non-lethal SGAR residues before  
277 dying [from non-SGAR related causes], and over-estimation of residues associated with  
278 mortality because birds ingest more than a lethal dose before they die; animals typically  
279 die some 5-7 days after ingestion of a lethal dose (Meehan 1984). Both biases would  
280 have the effect of flattening the probability curve.

## 281 **4.2. Exposure extent in Canada**

### 282 4.2.1. Spatial extent

283 Stone et al. (2003) stated that, at the time, SGARs appeared to be present in the  
284 majority of great horned owls and in roughly half of the red-tailed hawks from the  
285 sampled areas of the State of New York. That conclusion can be directly applied to our  
286 situation in Canada. Furthermore, a substantial fraction of a number of other raptors in  
287 Quebec (from the western half of the province including areas surrounding Gatineau,  
288 Montreal, Sherbrooke, Quebec and as far north as Obedjiwan) were also exposed to  
289 SGARs (43% – or 13 of 30 birds tested), supporting the notion that other avian species  
290 are also being impacted by SGAR use. This wider exposure in Quebec suggests a broad  
291 contamination of terrestrial food chains as Accipiters, such as the Cooper’s hawk, as well  
292 as other species such as the merlin and the American kestrel, feed predominantly on small  
293 birds and occasionally on insects (Ehrlich et al. 1988). Small birds, if the source of  
294 rodenticides, are most likely being exposed to SGARs from insects or other invertebrates,  
295 and possibly through direct uptake of grain-based baits.

296 In our study, great horned owls were consistently exposed to SGARs across the  
297 country. In apparent contrast, their daytime ecological counterpart, the red-tailed hawk,  
298 showed an increasing frequency of exposure eastward from the Prairie Provinces. This

299 difference could be explained by the lower dietary diversity of owls than hawks. Marti  
300 and Kochert (1995) showed that, on a finer scale, food-niche breadth became narrower  
301 along an eastward transect from the west coast of North America. This may reflect  
302 greater diversity of available prey in the west that could permit local populations of those  
303 two raptors to increase their diet segregation in western regions (Marti and Kochert  
304 1995). Houston et al. (1998) lists the main prey of great horned owls as including rabbits  
305 and hares, coots and other waterfowl and mice. While snowshoe hares (*Lepus*  
306 *americanus*), black-tailed jackrabbits (*Lepus californicus*), and ground squirrels  
307 (*Spermophilus spp.*) dominate the hawk's diet in western and northern parts of North  
308 America (Preston and Beane 2009). The bulk of their diet in eastern and midwestern  
309 North America includes voles (*Microtus*), mice (*Peromyscus spp.*, *Reithrodontomys spp.*,  
310 *Mus musculus*), rats (*Sigmodon hispidus*, *Oryzomys palustris*), and cottontails (*Sylvilagus*  
311 *spp.*) (Preston and Beane, 2009). Thus, it may be that in eastern areas that are more  
312 agricultural and urban (and subject to a higher degree of SGAR use), red-tailed hawks are  
313 exposed more frequently to SGARs through their increased feeding on rodents and  
314 reduced predation on other prey.

315 To obtain a more reliable estimate on actual exposure in Canada, we examined the  
316 livers of birds found dead from all causes. Our data indicate that, despite a smaller human  
317 population and the harsher climate in Canada (albeit some south-western regions of the  
318 country are characterised by milder weather), both of which should limit the need for  
319 rodenticides, the scale of exposure reported in our study are comparable to those in  
320 Europe. In the French Department of Loire Atlantique, 73% of a sample consisting of  
321 common kestrels (*Falco tinnunculus*), common buzzards (*Buteo buteo*), barn owls and



322 tawny owls (*Strix aluco*) had detectable SGAR liver residues (Lambert et al. 2007). In the  
323 UK, between 40% and 74% of barn owls, kestrels, and avian scavengers such as buzzards  
324 and red kites (*Milvus milvus*) found dead from various causes had detectable liver SGAR  
325 residues (Newton et al. 1999; Shore et al., 1999, 2006; Walker 2008b). However, it  
326 should be noted that the sampled areas of Canada were those with higher population  
327 densities and where landscape features are not greatly dissimilar from Europe. That may  
328 at least in part account for the apparent similarity in the frequency of contamination.

329       The widespread exposure in Canada in part most likely reflects the increase in sales  
330 and use of SGARs in the last few decades (Albert et al. 2010), and the use of persistent  
331 compounds that remain detectable in the liver long after the exposure event (Laas et al.  
332 1985). However, it is also clear from our data that multiple exposures, as detected by the  
333 presence of multiple compounds in the liver, are common. Although SGARs cannot be  
334 used legally on crops or orchards in Canada and are labelled for ‘\_indoor uses’ only,  
335 ‘\_indoor’ is defined to include use of baits outside farms and food establishments. This is  
336 likely to increase the exposure of non-target organisms. SGARs in Canada are currently  
337 labelled for domestic use although this is likely to change soon. Proposed regulatory  
338 actions relating to exposure risks for wildlife includes (amongst others), prohibiting use  
339 of SGAR compounds in residential settings or outdoor areas where wildlife may be  
340 exposed. In the case of commercial applications, bait stations would be required where  
341 wildlife could be exposed. Furthermore, labels of commercial class products would be  
342 amended to state that those products could be used only by certified operators, farmers  
343 and persons authorized in government-approved pest control programs (Pest Management

344 Regulatory Agency 2009). Those risk mitigation measures should have an overall  
345 positive impact on reducing unnecessary exposure risks to wildlife.

346       Regarding the impact of SGARs, we must be cautious in extrapolating from our data  
347 to predict likely mortality. However, if the probability of mortality is applied to each  
348 residue value in our dataset for great horned owls, this equates to an estimated predicted  
349 mortality of 11% (calculated by multiplying the probability of being exposed to SGARs  
350 [65% in GHOW] by the mean probability of exhibiting signs of intoxication [17% in  
351 GHOW]). This is the first time that the scale of potential mortality from SGARs has been  
352 estimated for any wild raptor population. That estimate may well be too low, as some  
353 proportion of poisoned birds likely die out of sight (Shore et al. 2005) and so be under-  
354 represented in our sample. Furthermore, our estimates of the scale of mortality do not  
355 account for any indirect effects that SGARs may have. Sub-lethal exposures may  
356 indirectly increase mortality associated with natural or accidental events. For instance,  
357 SGARs may hinder the recovery of birds from non-fatal collisions or accidents. They  
358 may also impair hunting ability through behavioural changes such as lethargy, thus  
359 increasing the probability of starvation. Intoxication with rodenticides has been shown to  
360 alter behaviour in rodents (Cox and Smith 1992) but there is no evidence to date of  
361 indirect effects in free-ranging raptors (Shore et al. 2005).

362       The lack of a probability plot for red-tailed hawks means that a comparable estimate  
363 for SGAR-induced mortality in Canada cannot be made for this species. The available  
364 data suggest that red-tailed hawks may be more sensitive to SGARs than great horned  
365 owls (Figure 2) but red-tailed hawks generally had lower liver SGAR concentrations in  
366 Canada, and, it is notable that in New York, great horned owls are poisoned more

367 frequently than red-tailed hawks (Stone et al. 1999, 2003). Additional studies and  
368 monitoring of red-tailed hawk SGAR residues would strengthen our ability to estimate  
369 the risk of toxicosis following exposure to SGARs.

#### 370 4.2.2. Future directions

371 Most studies that investigate exposure of non-target species to SGARs have focused  
372 on the uptake of poisoned rodents by various predators (Newton et al.1990, 1999; Berny  
373 et al.1997; McDonald et al.1998; Howald et al. 1999; Shore et al. 1999, 2003). The  
374 finding that falcons and accipiters were also exposed in Quebec suggests that terrestrial  
375 food chains are broadly contaminated by SGARs despite their very restricted use.  
376 Invertebrates represent another route of exposure, especially in insectivorous avian  
377 species (Dowding et al. 2006). Some potential routes of exposure to aerial insectivores  
378 include the consumption of invertebrates that previously fed on rodent faeces or carcasses  
379 and even the consumption of ground-dwelling earthworms and beetles that ingested  
380 residues or actual rodent bait (Spurr and Drew 1999; Dunlevy et al. 2000). Clearly, given  
381 the fact that many ecosystems contain a larger proportion of insectivorous vertebrates  
382 relative to higher trophic predators, exposure could even be greater in those taxa  
383 (Dowding et al. 2010). Developing probability curves or even metabolism studies for a  
384 wider range of species would provide us with insight into the relative sensitivities and  
385 risks to other species (Watanabe et al 2010). Finally, researching further indirect effects  
386 of SGARs on survival would refine current risk assessments of direct and indirect  
387 mortalities in wildlife.

#### 388 **4.3. Conclusion**

389           Our results continue to support recommendations that persistent SGARs such as  
390 brodifacoum, bromadiolone and difethialone should be used with caution (or not at all in  
391 some circumstances) given that it appears difficult to eliminate the risk of exposure to  
392 non-target wildlife. The results presented will hopefully aid policy-makers in refining  
393 risk-assessments of SGARs on non-target wildlife.

394           Our results can also help regulatory agencies worldwide provide guidance on both  
395 commercial and residential use of SGARs and enforce appropriate risk mitigation as  
396 needed. In this context, the extent of non-target exposure to SGARs may not always  
397 depend on the amount of bait used, but also on the way it is used (Shore et al. 2006).  
398 Focusing on improving application methods, such as baiting in areas of high rat activity  
399 only, conducting periodic and frequent searches for dead or dying rodents, enclosing the  
400 bait in a fashion that reduces invertebrate uptake may help reduce exposure of SGARs to  
401 predatory birds and other non-target species. Whether or not rodenticide resistance is  
402 common, an Integrated Pest Management (IPM) approach, that seeks to combine  
403 mechanical, biological and chemical controls, should be favoured as opposed to relying  
404 on a purely chemical mode of control.

405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416

417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460

### **Acknowledgements**

We wish to thank all the wildlife agencies, provincial ministries, Conservation Officers, wildlife rehabilitation centers and shelters and the various wildlife rescue associations who submitted raptor carcasses for this study. National Wildlife Research Centre staff are thanked for specimen bank archiving and rodenticide residue analysis. Thanks are extended to Nick Mastrota for access to the US EPA EIIS, Dr. Malcolm McAdie and Dr. Craig Stephens and staff at the Centre for Coastal Health for conducting post mortem examinations and Dr. Ward Stone for his unflagging and inspirational work on contaminant and pesticide issues in New York State. Funding was from the Canadian Wildlife Service and the Pesticide Science Fund of Environment Canada.

### **References:**

- Albert, C.A., Wilson, L.K., Mineau, P., Trudeau, S. and Elliott, J.E. 2010. Anticoagulant rodenticides in three owl species from western Canada, 1998-2003. Arch. Environ. Contam. Toxicol. 163,451-459.
- Berny, P.J., Buronfosse, T., Buronfosse, F., Lamarque, F. and Lorgue, G. 1997. Field evidence of secondary poisoning of foxes (*Vulpes vulpes*) and buzzards (*Buteo buteo*) by bromadiolone, a 4-year survey. Chemosphere. 35,1817–1829.
- Buckle, A.P., Prescott, C.V. and Ward, K.J. 1994. Resistance to the first and second generation anticoagulant rodenticides – a new perspective. In: Halverson, W.S., Marsh, R.E. (eds) Proc. 16<sup>th</sup> Vert. Pest Conf. pp 138-144.
- Cox, P. and Smith, R.H. 1992. Rodenticide ecotoxicology: pre-lethal effects of anticoagulants on rat behaviour. In: Borreco JE, Marsh RE (eds) Proc. 15<sup>th</sup> Vert. Pest Conf. University of California, Davis, California, USA. pp 165-170.
- Dowding, J.E., Lovegrove, T.G., Ritchie, J., Kast, S.N. and Puckett, M. 2006. Mortality of northern New Zealand dotterels (*Charadrius obscurus aquilonius*) following an aerial poisoning operation. Notornis. 53,235–239.
- Dowding, C.V., Shore, R.F., Worgan, A., Baker, P.J. and Harris, S. 2010. Accumulation of anticoagulant rodenticides in a non-target insectivore, the European hedgehog (*Erinaceus europaeus*). Environ. Pollut. 158,161-66.
- Dunlevy, P.A., Campbell, E.W. and Lindsey, G.D. 2000. Broadcast application of a placebo rodenticide bait in a native Hawaiian forest. Int. Biodeterior. Biodegradation. 45,199–208.
- Eason, C.T., Murphy, E.C., Wright, G.R.G., Spurr, E.B. 2002. Assessment of risks of brodifacoum to non-target birds and mammals in New Zealand. Ecotoxicology. 11,35-48.

461 Ehrlich, P.R., Dobkin, D.S. and Wheye, D. 1988. The birder's handbook – a field guide  
462 to the natural history of North American birds. Simon & Schuster Inc. New York, New  
463 York.  
464  
465 Erickson, W. and Urban, D. 2004. Potential risks of nine rodenticides to birds and  
466 nontarget mammals: a comparative approach. United States Environmental Protection  
467 Agency, Washington, DC, USA.  
468  
469 Hedgal, P.L. and Colvin, B.A. 1988. Potential hazard to eastern screech-owls and other  
470 raptors of brodifacoum bait used for vole control in orchards. Environ. Toxicol. Chem.  
471 7,245-260.  
472  
473 Houston, C.S., Dwight, G.S. and Christoph, R. 1998. Great Horned Owl (*Bubo*  
474 *virginianus*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of  
475 Ornithology; Retrieved from the Birds of North America Online:  
476 <http://bna.birds.cornell.edu/bna/species/372>  
477  
478 Howald, G.R., Mineau, P., Elliott, J.E., and Cheng, K.M. 1999. Brodifacoum poisoning  
479 of avian scavengers during rat control on a seabird colony. Ecotoxicology 8,429-446.  
480  
481 Laas, F.Y., Forss, D.A., Godfrey, M.E.R. 1985. Retention of brodifacoum in sheep  
482 tissues and excretion in faeces. N.Z. J. Agric. Res. 28,357–359.  
483  
484 Lambert, O., Pouliquen, H., Larhantec, M., Thorin, C. and L'Hostis, M. 2007. Exposure  
485 of raptors and waterbirds to anticoagulant rodenticides (difenacoum, bromadiolone,  
486 coumatetralyl, coumafen, brodifacoum): epidemiological survey in Loire Atlantique  
487 (France). Bull. Environ. Contam. Toxicol. 79,91-94.  
488  
489 McDonald, R.A., Harris, S., Turnbull, G., Brown, P. and Fletcher, M. 1998.  
490 Anticoagulant rodenticides in stoats (*Mustela erminea*) and weasels (*Mustela nivalis*) in  
491 England. Environ. Pollut. 103,17–23.  
492  
493 Marti, C.D. and Kochert, M.N. 1995. Are Red-tailed Hawks and Great Horned Owls  
494 diurnal-nocturnal dietary counterparts? Wilson Bull. 107,615-628.  
495  
496 Mastrota, N.F. 2007. EIIS Version 2.0 – User Manual. EFED Information and Support  
497 Branch, Environmental Fate and Effects Division, Office of Pesticide Programs, USA.  
498  
499 Meehan, A.P. 1984. Rats and Mice: their Biology and Control. Rentokil Limited, East  
500 Grinstead, Sussex, UK.  
501  
502 Newton, I., Wyllie, I. and Freestone, P. 1990. Rodenticides in British barn owls. Environ.  
503 Pollut. 68,101-117.  
504  
505 Newton, I., Dale, L., Finnie, J.K., Freestone, P., Wright, J., Wyatt, C. and Wyllie, I. 1998.  
506 Wildlife and Pollution: 1997/98 Annual Report. JNCC Report No. 285

507  
508 Newton, I., Shore, R.F., Wyllie, I., Birks, J.D.S., Dale, L. 1999. Empirical evidence of  
509 side-effects of rodenticides on some predatory birds and mammals. In: Cowan DP, Feare  
510 CJ (eds) Advances in vertebrate pest management. Filander Verlag, Furth, pp 347–367.  
511  
512 Newton, I., Afsar, A., Dale, L., Finnie, J., Shore, R.F., Wright, J., Wyatt, C. and Wyllie, I.  
513 2000. Wildlife and Pollution: 1998/99 Annual Report. JNCC Report No.305  
514  
515 Parmar, G., Bratt, H., Moore, R. and Batten, P.L. 1987. Evidence for a common binding-  
516 site in vivo for the retention of anticoagulants in rat liver. Hum. Toxicol. 6,431–432.  
517  
518 Pest Management Regulatory Agency. 2009. Proposed Risk Mitigation Measures for  
519 Eight Rodenticides. Re-evaluation Note REV2009-05. Health Canada. Ottawa, Ontario.  
520 22 June 2009. 23 pages.  
521  
522 Preston, C.R. and Beane, R.D. 2009. Red-tailed Hawk (*Buteo jamaicensis*), The Birds of  
523 North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved  
524 from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/052>  
525  
526 Shore, R.F., Birks, J.D.S. and Freestone, P. 1999. Exposure of non-target vertebrates to  
527 second-generation rodenticides in Britain, with particular reference to the polecat *Mustela*  
528 *putorius*. N.Z. J. Ecol. 23,199-206.  
529  
530 Shore, R.F., Malcolm, H.M., Wienburg, C.L., Turk, A., Horne, J.A., Dale, L., Wyllie, I.  
531 and Newton, I. 2001. Wildlife and pollution: 1999/2000 Annual report. JNCC Report No.  
532 321  
533  
534 Shore, R.F., Birks, J.D.S., Afsar, A., Wienburg, C.L. and Kitchener, A.C. 2003. Spatial  
535 and temporal analysis of second-generation anticoagulant rodenticide residues in polecats  
536 (*Mustela putorius*) from throughout their range in Britain, 1992–1999. Environ. Pollut.  
537 122,183–193.  
538  
539 Shore, R.F., Malcolm, H.M., Wienburg, C.L., Walker, L.A., Turk, A. and Horne, J.A.  
540 2005. Wildlife and Pollution: 2001/02 Annual Report. JNCC Report No. 352  
541  
542 Shore, R.F., Malcolm, H.M., McLennan, D., Turk, A., Walker, L.A., Wienburg, C.L. and  
543 Burn, A.J. 2006. Did foot-and-mouth disease-control operations affect rodenticide  
544 exposure in raptors? J. Wildl. Manage. 70,588-593.  
545  
546 Spurr, E.B. and Drew, K.W. 1999. Invertebrates feeding on baits used for vertebrate pest  
547 control in New Zealand. N.Z. J. Ecol. 23,167–173.  
548  
549 Stone, W.B., Okoniewski, J.C. and Stedelin, J.R. 1999. Poisoning of wildlife with  
550 anticoagulant rodenticides in New York. J. Wildl. Dis. 35,187-193.  
551

552 Stone, W.B., Okoniewski, J.C. and Stedelin, J.R. 2003. Anticoagulant rodenticides and  
553 raptors: recent findings from New York, 1998-2001. Bull. Environ. Contam. Toxicol.  
554 70,34-40.  
555  
556 Taylor, M.J., Sharp, E.A., Melton, L.M., Giela, A. and Watson, J.E. 2009. Walk on the  
557 wild side: an investigation into secondary poisoning of wild birds using LC-MS. British  
558 Mass Spectrometry Society. 60,12-14.  
559  
560 Walker, L.A., Turk, A., Long, S.M., Wienburg, C.L., Best, J. and Shore, R.F. 2008a.  
561 Second generation anticoagulant rodenticides in tawny owls (*Strix aluco*) from Great  
562 Britain. Sci. Total Environ. 392,93-98.  
563  
564 Walker, L.A., Shore, R.F., Turk, A., Pereira, M.G., and Best, J. 2008b. The Predatory  
565 Bird Monitoring Scheme (PBMS) — identifying chemical risks to top predators in  
566 Britain. Ambio. 37,466-471.  
567  
568 Walker, L.A., Lawlor, A.J., Llewellyn, N., Peréira, M.G., Potter, E., Townsend, J., Turk,  
569 A., and Shore, R.F. 2010. The Predatory Bird Monitoring Scheme (PBMS) Report 2006-  
570 7. A contract report from the Centre for Ecology & Hydrology to Natural England, 65  
571 pages. Retrieved from the website at  
572 [http://pbms.ceh.ac.uk/docs/AnnualReports/PBMSreport2006\\_2007\\_FINAL.pdf](http://pbms.ceh.ac.uk/docs/AnnualReports/PBMSreport2006_2007_FINAL.pdf)  
573  
574 Watanabe, K.P., Saengtienchai, A., Tanaka, K.D., Ikenaka, Y. and Ishizuka, M. 2010.  
575 Comparison of warfarin sensitivity between rat and bird species. Comp. Biochem. Phys.  
576 C. 152,114-119.  
577  
578 World Health Organisation. 1995. Environmental Health Criteria 175: Anticoagulant  
579 Rodenticides. World Health Organisation (WHO), Geneva.  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597



598 **Figure captions:**

599

600 **Figure 1:** Published liver SGAR residues (combined concentrations of bromadiolone,  
601 brodifacoum and difethialone) in barred owl (BAOW), barn owl (BNOW), great horned  
602 owl (GHOW) and red-tailed hawk (RTHA). Total number of birds = 270 and do not  
603 include birds with non-detected residues. Diamond in the center of the box represents  
604 average, line is the median, box is the upper and lower quartiles and the whiskers are the  
605 standard deviation. Sources of the data are: Newton et al. 1990, 1998, 2000; Stone et al.  
606 1999, 2003; Albert et al. 2010; EIIS 2010 download.

607

608 **Figure 2:** Effect plot of the probability of becoming symptomatic (0,1) as a function of  
609  $\log_{10}$  [mg/kg]. ALL represents pooled data (n=270), BAOW represents barred owls  
610 (n=26), BNOW represents barn owls (n=126), GHOW represents great horned owls  
611 (n=86) and RTHA represents red-tailed hawks (n=32). Shading represents 95%  
612 confidence limits for ALL birds. Curves were drawn using the formula  $y(\text{probability}) =$   
613  $1/(1+\exp(-(\text{int} + b*x)))$  where int is the intercept and b is the parameter estimate for X  
614 (concentration).

615

616 **Figure 3:** Percentage of great horned owls (GHOW) and red-tailed hawks (RTHA)  
617 across Canada sampled in our study that had detectable ( $\geq 0.005$  mg/kg ww) liver SGAR  
618 residues. No RTHA samples were collected from PYR. PYR stands for the Pacific and  
619 Yukon region of Canada and PNR is the Prairie and Northern Region.

620

621 **Figure 4:** Percentage of great horned owls (GHOW) and red-tailed hawks (RTHA) with  
622 0, 1, 2 and 3 different SGARs detected in the liver. Tested compounds were brodifacoum,  
623 bromadiolone and difethialone.

624

625 **Figure 5:** Cumulative frequency graph for liver SGAR residues in 79 great horned owls.  
626 Red line represents the 20% probability level for effect (0.07 mg/kg; Table 1).

627

628 **Figure 6:** Cumulative frequency graph for liver SGAR residues in 42 red-tailed hawks.

629

630 **Figure 7:** Numbers of birds of prey from Québec that contained detectable and non-  
631 detectable liver SGAR residues (13/30 samples tested positive or 43%).

632

633 **Table 1:** Toxicity threshold values (mg/kg ww) for 5%, 10%, 15% and 20% probability  
 634 risk levels. For-example, in barred owls (BAOW), an owl with 0.06mg/kg SGAR  
 635 residues in the liver would have a 5% chance of showing signs of toxicosis. Sample sizes  
 636 (n) as well as the number of positive (1) and negative (0) cases are presented. P value  
 637 representing binary logit model fit is also showed. BNOW stands for barn owl, GHOW is  
 638 the great horned owl, RTHA the red-tailed hawk and ALL represents the pooled data for  
 639 all birds.  
 640

Probability	BAOW n=26 0=22 1=4 p=0.008	BNOW n=126 0=114 1=12 p=<0.0001	GHOW n=86 0=62 1=24 p=<0.0001	RTHA n=32 0=3 1=29 p=0.37	ALL n=270 0=201 1=69 p=<0.0001
0.05	0.06	0.05	0.02	---	0.02
0.10	0.09	0.09	0.03	---	0.04
0.15	0.13	0.13	0.05	---	0.06
0.20	0.16	0.18	0.07	---	0.08

641 --- - values not presented if binary logit model fit was not statistically significant  
 642  
 643  
 644  
 645  
 646  
 647  
 648  
 649  
 650  
 651  
 652  
 653  
 654  
 655  
 656  
 657  
 658  
 659  
 660  
 661  
 662  
 663  
 664  
 665  
 666  
 667  
 668  
 669


670 **Table 2:** Geometric mean (range) liver SGAR concentrations [mg/kg ww] for great  
 671 horned owls (GHOW) and red-tailed hawks (RTHA) from the Pacific and Yukon region  
 672 of Canada (PYR), the prairie and northern region (PNR), Ontario and Quebec.  
 673

	PYR		PNR		Ontario		Quebec		Pooled – all provinces	
	GHOW	RTHA	GHOW	RTHA	GHOW	RTHA	GHOW	RTHA	GHOW	RTHA
<b>Brodifacoum</b>	0.04 (0.003- 0.61) n=28	N/A	0.008 (0.001- 0.016) n=6	0.004 (0.001- 0.02) n=3	0.007 (0.001- 0.05) n=17	0.006 (0.001- 0.17) n=18	0.013 (0.003- 0.08) n=7	0.01 (0.008- 0.04) n=5	0.017 (0.001- 0.61) n=58	0.006 (0.001- 0.17) n=26
<b>Bromadiolone</b>	0.03 (0.005- 0.57) n=33	N/A	0.007 (0.001- 0.07) n=7	0.004 (0.001- 0.008) n=3	0.01 (0.001- 0.07) n=15	0.004 (0.001- 0.06) n=25	0.01 (0.003- 0.14) n=6	0.003 (0.002- 0.006) n=4	0.018 (0.001- 0.57) n=61	0.004 (0.001- 0.064) n=32
<b>Difethialone</b>	0.02 (0.013- 0.03) n=3	N/A	ND	ND	0.003 (0.003- 0.003) n=1	ND	ND	ND	0.013 (0.003- 0.03) n=4	0
<b>Pooled - all compounds</b>	0.03 (0.003- 0.609) n=64	N/A	0.007 (0.001- 0.07) n=13	0.004 (0.001- 0.017) n=6	0.008 (0.001- 0.07) n=33	0.005 (0.001- 0.17) n=43	0.012 (0.003- 0.14) n=13	0.006 (0.002- 0.04) n=9	0.016 (0.001 - 0.61) n=123	0.005 (0.001 - 0.064) n=58


674 N/A = no samples obtained; ND = no detectable residue in any livers; n= number of birds with detectable  
 675 residues.

676  
 677  
 678  
 679

680  
681  
682



683  
684 **Figure 1**  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696



697  
 698  
 699  
 700  
 701  
 702  
 703  
 704  
 705  
 706  
 707  
 708  
 709  
 710  
 711

**Figure 2**

712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757




Figure 3

758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803




Figure 4

804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849




Figure 5







Figure 6

850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890



891  
 892  
 893  
 894  
 895  
 896  
 897  
 898  
 899  
 900  
 901  
 902  
 903  
 904  
 905  
 906  
 907  
 908  
 909  
 910  
 911  
 912  
 913  
 914  
 915

**Figure 7**

916  
917  
918  
919  
920