

Packing Plane Spanning Trees into a Point Set

Ahmad Biniáz*

Alfredo García†

Abstract

Let P be a set of n points in the plane in general position. We show that at least $\lfloor n/3 \rfloor$ plane spanning trees can be packed into the complete geometric graph on P . This improves the previous best known lower bound $\Omega(\sqrt{n})$. Towards our proof of this lower bound we show that the center of a set of points, in the d -dimensional space in general position, is of dimension either 0 or d .

1 Introduction

In the two-dimensional space, a *geometric graph* G is a graph whose vertices are points in the plane and whose edges are straight-line segments connecting the points. A subgraph S of G is *plane* if no pair of its edges cross each other. Two subgraphs S_1 and S_2 of G are *edge-disjoint* if they do not share any edge.

Let P be a set of n points in the plane. The *complete geometric graph* $K(P)$ is the geometric graph with vertex set P that has a straight-line edge between every pair of points in P . We say that a sequence S_1, S_2, S_3, \dots of subgraphs of $K(P)$ is *packed into* $K(P)$, if the subgraphs in this sequence are pairwise edge-disjoint. In a packing problem, we ask for the largest number of subgraphs of a given type that can be packed into $K(P)$. Among all subgraphs, plane spanning trees, plane Hamiltonian paths, and plane perfect matchings are of interest. Since $K(P)$ has $n(n-1)/2$ edges, at most $\lfloor n/2 \rfloor$ spanning trees, at most $\lfloor n/2 \rfloor$ Hamiltonian paths, and at most $n-1$ perfect matchings can be packed into it.

A long-standing open question is to determine whether or not it is possible to pack $\lfloor n/2 \rfloor$ plane spanning trees into $K(P)$. If P is in convex position, the answer in the affirmative follows from the result of Bernhart and Kanien [3], and a characterization of such plane spanning trees is given by Bose et al. [5]. In CCCG 2014, Aichholzer et al. [1] showed that if P is in general position (no three points on a line), then $\Omega(\sqrt{n})$ plane spanning trees can be packed into $K(P)$; this bound is obtained by a clever combination of crossing family (a set of pairwise crossing edges) [2] and double-stars

(trees with only two interior nodes) [5]. Schnider [12] showed that it is not always possible to pack $\lfloor n/2 \rfloor$ plane spanning double stars into $K(P)$, and gave a necessary and sufficient condition for the existence of such a packing. As for packing other spanning structures into $K(P)$, Aichholzer et al. [1] and Biniáz et al. [4] showed a packing of 2 plane Hamiltonian cycles and a packing of $\lceil \log_2 n \rceil - 2$ plane perfect matchings, respectively.

The problem of packing spanning trees into (abstract) graphs is studied by Nash-Williams [11] and Tutte [13] who independently obtained necessary and sufficient conditions to pack k spanning trees into a graph. Kundu [10] showed that at least $\lceil (k-1)/2 \rceil$ spanning trees can be packed into any k -edge-connected graph.

In this paper we show how to pack $\lfloor n/3 \rfloor$ plane spanning trees into $K(P)$ when P is in general position. This improves the previous $\Omega(\sqrt{n})$ lower bound.

2 Packing Plane Spanning Trees

In this section we show how to pack $\lfloor n/3 \rfloor$ plane spanning tree into $K(P)$, where P is a set of $n \geq 3$ points in the plane in general position (no three points on a line). If $n \in \{3, 4, 5\}$ then one can easily find a plane spanning tree on P . Thus, we may assume that $n \geq 6$.

The *center* of P is a subset C of the plane such that any closed halfplane intersecting C contains at least $\lfloor n/3 \rfloor$ points of P . A *centerpoint* of P is a member of C , which does not necessarily belong to P . Thus, any halfplane that contains a centerpoint, has at least $\lfloor n/3 \rfloor$ points of P . It is well known that every point set in the plane has a centerpoint; see e.g. [7, Chapter 4]. We use the following corollary and lemma in our proof of the $\lfloor n/3 \rfloor$ lower bound; the corollary follows from Theorem 4 that we will prove later in Section 3.

Corollary 1 *Let P be a set of $n \geq 6$ points in the plane in general position, and let C be the center of P . Then, C is either 2-dimensional or 0-dimensional. If C is 0-dimensional, then it consists of one point that belongs to P , moreover n is of the form $3k+1$ for some integer $k \geq 2$.*

Lemma 1 *Let P be a set of n points in the plane in general position, and let c be a centerpoint of P . Then, for every point $p \in P$, each of the two closed halfplanes, that are determined by the line through c and p , contains at least $\lfloor n/3 \rfloor + 1$ points of P .*

*University of Waterloo, Canada. Supported by NSERC Postdoctoral Fellowship. ahmad.biniiaz@gmail.com

†Universidad de Zaragoza, Spain. Partially supported by H2020-MSCA-RISE project 734922 - CONNECT and MINECO project MTM2015-63791-R. olaverri@unizar.es




Figure 1: Illustration of the proof of Lemma 1.

Proof. For the sake of contradiction assume that a closed halfplane \bar{H} , that is determined by the line through c and p , contains less than $\lceil n/3 \rceil + 1$ points of P . By symmetry assume that \bar{H} is to the left side of this line oriented from c to p as depicted in Figure 1. Since c is a centerpoint and \bar{H} contains c , the definition of centerpoint implies that \bar{H} contains exactly $\lceil n/3 \rceil$ points of P (including p and any other point of P that may lie on the boundary of \bar{H}). By slightly rotating \bar{H} counterclockwise around c , while keeping c on the boundary of \bar{H} , we obtain a new closed halfplane that contains c but misses p . This new halfplane contains less than $\lceil n/3 \rceil$ points of P ; this contradicts c being a centerpoint of P . \square

Now we proceed with our proof of the lower bound. We distinguish between two cases depending on whether the center C of P is 2-dimensional or 0-dimensional. First suppose that C is 2-dimensional. Then, C contains a centerpoint, say c , that does not belong to P . Let p_1, \dots, p_n be a counter-clockwise radial ordering of points in P around c . For two points p and q in the plane, we denote by \vec{pq} , the ray emanating from p that passes through q .

Since every integer $n \geq 3$ has one of the forms $3k$, $3k + 1$, and $3k + 2$, for some $k \geq 1$, we will consider three cases. In each case, we show how to construct k plane spanning directed graphs G_1, \dots, G_k that are edge-disjoint. Then, for every $i \in \{1, \dots, k\}$, we obtain a plane spanning tree T_i from G_i . First assume that $n = 3k$. To build G_i , connect p_i by outgoing edges to $p_{i+1}, p_{i+2}, \dots, p_{i+k}$, then connect p_{i+k} by outgoing edges to $p_{i+k+1}, p_{i+k+2}, \dots, p_{i+2k}$, and then connect p_{i+2k} by outgoing edges to $p_{i+2k+1}, p_{i+2k+2}, \dots, p_{i+3k}$, where all the indices are modulo n , and thus $p_{i+3k} = p_i$. The graph G_i , that is obtained this way, has one cycle $(p_i, p_{i+k}, p_{i+2k}, p_i)$; see Figure 2. By Lemma 1, every closed halfplane, that is determined by the line through c and a point of P , contains at least $k + 1$ points of P . Thus, all points $p_i, p_{i+1}, \dots, p_{i+k}$ lie in the closed halfplane to the left of the line through c and p_i that is oriented from c to p_i . Similarly, the points p_{i+k}, \dots, p_{i+2k} lie in the closed halfplane to the left of the oriented line from c to p_{i+k} , and the points $p_{i+2k}, \dots, p_{i+3k}$ lie in the closed halfplane to the left of the oriented line

from c to p_{i+2k} . Thus, all the k edges outgoing from p_i are in the convex wedge bounded by the rays $\vec{cp_i}$ and $\vec{cp_{i+k}}$, all the edges outgoing from p_{i+k} are in the convex wedge bounded by $\vec{cp_{i+k}}$ and $\vec{c_{i+2k}}$, and all the edges from p_{i+2k} are in the convex wedge bounded by $\vec{cp_{i+2k}}$ and $\vec{c_{i+3k}}$. Therefore, the spanning directed graph G_i is plane. As depicted in Figure 2, by removing the edge (p_{i+2k}, p_i) from G_i we obtain a plane spanning (directed) tree T_i . This is the end of our construction of k plane spanning trees.




Figure 2: The plane spanning trees T_1 (the top) and T_2 (the bottom) are obtained by removing the edges (p_{1+2k}, p_1) and (p_{2+2k}, p_2) from G_1 and G_2 , respectively.

To verify that the k spanning trees obtained above are edge-disjoint, we show that two trees T_i and T_j , with $i \neq j$, do not share any edge. Notice that the tail of every edge in T_i belongs to the set $I = \{p_i, p_{i+k}, p_{i+2k}\}$, and the tail of every edge in T_j belongs to the set $J = \{p_j, p_{j+k}, p_{j+2k}\}$, and $I \cap J = \emptyset$. For contrary, suppose that some edge (p_r, p_s) belongs to both T_i and T_j , and without loss of generality assume that in T_i this edge is oriented from p_r to p_s while in T_j it is oriented from p_s to p_r . Then $p_r \in I$ and $p_s \in J$. Since $(p_r, p_s) \in T_i$ and the largest index of the head of every outgoing edge from p_r is $r + k$, we have that $s \leq (r + k) \pmod n$. Similarly, since $(p_s, p_r) \in T_j$ and the largest index of the head of every outgoing edge from p_s is $s + k$, we have that $r \leq (s + k) \pmod n$. However, these two inequalities cannot hold together; this contradicts our assumption




Figure 3: The dimension of a point set in the plane, that is not in general position, can be any number in $\{0, 1, 2\}$.

that (p_r, p_s) belongs to both trees. Thus, our claim, that T_1, \dots, T_k are edge-disjoint, follows. This finishes our proof for the case where $n = 3k$.

If $n = 3k + 1$, then by Lemma 1, every closed halfplane that is determined by the line through c and a point of P contains at least $k + 2$ points of P . In this case, we construct G_i by connecting p_i to its following $k + 1$ points, i.e., $p_{i+1}, \dots, p_{i+k+1}$, and then connecting each of p_{i+k+1} and p_{i+2k+1} to their following k points. If $n = 3k + 2$, then we construct G_i by connecting each of p_i and p_{i+k+1} to their following $k + 1$ points, and then connecting p_{i+2k+2} to its following k points. This is the end of our proof for the case where C is 2-dimensional.

Now we consider the case where C is 0-dimensional. By Corollary 1, C consists of one point that belongs to P , and moreover $n = 3k + 1$ for some $k \geq 2$. Let $p \in P$ be the only point of C , and let p_1, \dots, p_{n-1} be a counter-clockwise radial ordering of points in $P \setminus \{p\}$ around p . As in our first case (where C was 2-dimensional, c was not in P , and n was of the form $3k$) we construct k edge-disjoint plane spanning trees T_1, \dots, T_k on $P \setminus \{p\}$ where p playing the role of c . Then, for every $i \in \{1, \dots, k\}$, by connecting p to p_i , we obtain a plane spanning tree for P . These plane spanning trees are edge-disjoint. This is the end of our proof. In this section we have proved the following theorem.

Theorem 2 *Every complete geometric graph, on a set of n points in the plane in general position, contains at least $\lfloor n/3 \rfloor$ edge-disjoint plane spanning trees.*

3 The Dimension of the Center of a Point Set

The *center* of a set P of $n \geq d + 1$ points in \mathbb{R}^d is a subset C of \mathbb{R}^d such that any closed halfspace intersecting C contains at least $\alpha = \lceil n/(d + 1) \rceil$ points of P . Based on this definition, one can characterize C as the intersection of all closed halfspaces such that their complementary open halfspaces contain less than α points of P . More precisely (see [7, Chapter 4]) C is the intersection of a finite set of closed halfspaces $\overline{H_1}, \overline{H_2}, \dots, \overline{H_m}$ such that for each $\overline{H_i}$

1. the boundary of $\overline{H_i}$ contains at least d affinely independent points of P , and
2. the complementary open halfspace H_i contains at

most $\alpha - 1$ points of P , and the closure of H_i contains at least α points of P .

Being the intersection of closed halfspaces, C is a convex polyhedron. A *centerpoint* of P is a member of C , which does not necessarily belong to P . It follows, from the definition of the center, that any halfspace containing a centerpoint has at least α points of P . It is well known that every point set in the plane has a centerpoint [7, Chapter 4]. In dimensions 2 and 3, a centerpoint can be computed in $O(n)$ time [9] and in $O(n^2)$ expected time [6], respectively.

A set of points in \mathbb{R}^d , with $d \geq 2$, is said to be in *general position* if no $k + 2$ of them lie in a k -dimensional flat for every $k \in \{1, \dots, d - 1\}$.¹ Alternatively, for a set of points in \mathbb{R}^d to be in general position, it suffices that no $d + 1$ of them lie on the same hyperplane. In this section we prove that if a point set P in \mathbb{R}^d is in general position, then the center of P is of dimension either 0 or d . Our proof of this claim uses the following result of Grünbaum.

Theorem 3 (Grünbaum, 1962 [8]) *Let \mathcal{F} be a finite family of convex polyhedra in \mathbb{R}^d , let I be their intersection, and let s be an integer in $\{1, \dots, d\}$. If every intersection of s members of \mathcal{F} is of dimension d , but I is $(d - s)$ -dimensional, then there exist $s + 1$ members of \mathcal{F} such that their intersection is $(d - s)$ -dimensional.*

Before proceeding to our proof, we note that if P is not in general position, then the dimension of C can be any number in $\{0, 1, \dots, d\}$; see e.g. Figure 3 for the case where $d = 2$.

Observation 1 *For every $k \in \{1, \dots, d + 1\}$ the dimension of the intersection of every k closed halfspaces in \mathbb{R}^d is in the range $[d - k + 1, d]$.*

Theorem 4 *Let P be a set of $n \geq d + 1$ points in \mathbb{R}^d , and let C be the center of P . Then, C is either d -dimensional, or contained in a $(d - s)$ -dimensional polyhedron that has at least $n - (s + 1)(\alpha - 1)$ points of P for some $s \in \{1, \dots, d\}$ and $\alpha = \lceil n/(d + 1) \rceil$. In the latter case if P is in general position and $n \geq d + 3$, then C consists of one point that belongs to P , and n is of the form $k(d + 1) + 1$ for some integer $k \geq 2$.*

¹A flat is a subset of d -dimensional space that is congruent to a Euclidean space of lower dimension. The flats in 2-dimensional space are points and lines, which have dimensions 0 and 1.

Proof. The center C is a convex polyhedron that is the intersection of a finite family \mathcal{H} of closed halfspaces such that each of their complementary open halfspaces contains at most $\alpha - 1$ points of P [7, Chapter 4]. Since C is a convex polyhedron in \mathbb{R}^d , its dimension is in the range $[0, d]$. For the rest of the proof we consider the following two cases.

- (a) The intersection of every $d + 1$ members of \mathcal{H} is of dimension d .
- (b) The intersection of some $d + 1$ members of \mathcal{H} is of dimension less than d .

First assume that we are in case (a). We prove that C is d -dimensional. Our proof follows from Theorem 3 and a contrary argument. Assume that C is not d -dimensional. Then, C is $(d - s)$ -dimensional for some $s \in \{1, \dots, d\}$. Since the intersection of every s members of \mathcal{H} is d -dimensional, by Theorem 3 there exist $s + 1$ members of \mathcal{H} whose intersection is $(d - s)$ -dimensional. This contradicts the assumption of case (a) that the intersection of every $d + 1$ members of \mathcal{H} is d -dimensional. Therefore, C is d -dimensional in this case.

Now assume that we are in case (b). Let s be the largest integer in $\{1, \dots, d\}$ such that every intersection of s members of \mathcal{H} is d -dimensional; notice that such an integer exists because every single halfspace in \mathcal{H} is d -dimensional. Our choice of s implies the existence of a subfamily \mathcal{H}' of $s + 1$ members of \mathcal{H} whose intersection is d' -dimensional for some $d' < d$. Let s' be an integer such that $d' = d - s'$. By Observation 1, we have that $d' \geq d - s$, and equivalently $d - s' \geq d - s$; this implies $s' \leq s$. To this end we have a family \mathcal{H}' with $s + 1$ members for which every intersection of s' members is d -dimensional (because $s' \leq s$ and $\mathcal{H}' \subseteq \mathcal{H}$), but the intersection of all members of \mathcal{H}' is $(d - s')$ -dimensional. Applying Theorem 3 on \mathcal{H}' implies the existence of $s' + 1$ members of \mathcal{H}' whose intersection is $(d - s')$ -dimensional. If $s' < s$, then this implies the existence of $s' + 1 \leq s$ members of $\mathcal{H}' \subseteq \mathcal{H}$, whose intersection is of dimension $d - s' < d$. This contradicts the fact that the intersection of every s members of \mathcal{H} is d -dimensional. Thus, $s' = s$, and consequently, $d' = d - s' = d - s$. Therefore C is contained in a $(d - s)$ -dimensional polyhedron I which is the intersection of the $s + 1$ closed halfspaces of \mathcal{H}' . Let H_1, \dots, H_{s+1} be the complementary open halfspaces of members of \mathcal{H}' , and recall that each H_i contains at most $\alpha - 1$ points of P . Let \bar{I} be the complement of I . Then,

$$\begin{aligned} n &= |I \cup \bar{I}| = |I \cup H_1 \cup \dots \cup H_{s+1}| \\ &\leq |I| + |H_1| + \dots + |H_{s+1}| \leq |I| + (s + 1)(\alpha - 1), \end{aligned}$$

where we abuse the notations I , \bar{I} , and H_i to refer to the subset of points of P that they contain. This inequality implies that I contains at least $n - (s + 1)(\alpha - 1)$ points

of P . This finishes the proof of the theorem except for the part that P is in general position.

Now, assume that P is in general position and $n \geq d + 3$. By the definition of general position, the number of points of P in a $(d - s)$ -dimensional flat is not more than $d - s + 1$. Since I is $(d - s)$ -dimensional, this implies that

$$n - (s + 1)(\alpha - 1) \leq d - s + 1.$$

Notice that n is of the form $k(d + 1) + i$ for some integer $k \geq 1$ and some $i \in \{0, 1, \dots, d\}$. Moreover, if i is 0 or 1, then $k \geq 2$ because $n \geq d + 3$. Now we consider two cases depending on whether or not i is 0. If $i = 0$, then $\alpha = k$. In this case, the above inequality simplifies to $k(d - s) \leq d - 2s$, which is not possible because $k \geq 2$ and $d \geq s \geq 1$. If $i \in \{1, \dots, d\}$, then $\alpha = k + 1$. In this case, the above inequality simplifies to $(k - 1)(d - s) + i \leq 1$, which is not possible unless $d = s$ and $i = 1$. Thus, for the above inequality to hold we should have $d = s$ and $i = 1$. These two assertions imply that $n = k(d + 1) + 1$, and that I is 0-dimensional and consists of one point of P . Since $C \subseteq I$ and C is not empty, C also consists of one point of P . \square

References

- [1] O. Aichholzer, T. Hackl, M. Korman, M. J. van Kreveld, M. Löffler, A. Pilz, B. Speckmann, and E. Welzl. Packing plane spanning trees and paths in complete geometric graphs. *Information Processing Letters*, 124:35–41, 2017. Also in CCCG'14, pages 233–238.
- [2] B. Aronov, P. Erdős, W. Goddard, D. J. Kleitman, M. Klugerman, J. Pach, and L. J. Schulman. Crossing families. *Combinatorica*, 14(2):127–134, 1994. Also in SoCG'91, pages 351–356.
- [3] F. Bernhart and P. C. Kainen. The book thickness of a graph. *Journal of Combinatorial Theory, Series B*, 27(3):320–331, 1979.
- [4] A. Biniiaz, P. Bose, A. Maheshwari, and M. H. M. Smid. Packing plane perfect matchings into a point set. *Discrete Mathematics & Theoretical Computer Science*, 17(2):119–142, 2015.
- [5] P. Bose, F. Hurtado, E. Rivera-Campo, and D. R. Wood. Partitions of complete geometric graphs into plane trees. *Computational Geometry: Theory and Applications*, 34(2):116–125, 2006.
- [6] T. M. Chan. An optimal randomized algorithm for maximum tukey depth. In *Proceedings of the 15th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 430–436, 2004.
- [7] H. Edelsbrunner. *Algorithms in Combinatorial Geometry*. Springer, 1987.
- [8] B. Grünbaum. The dimension of intersections of convex sets. *Pacific Journal of Mathematics*, 12(1):197–202, 1962.

- [9] S. Jadhav and A. Mukhopadhyay. Computing a centerpoint of a finite planar set of points in linear time. *Discrete & Computational Geometry*, 12:291–312, 1994.
- [10] S. Kundu. Bounds on the number of disjoint spanning trees. *Journal of Combinatorial Theory, Series B*, 17(2):199–203, 1974.
- [11] C. St. J. A. Nash-Williams. Edge-disjoint spanning trees of finite graphs. *Journal of the London Mathematical Society*, 36(1):445–450, 1961.
- [12] P. Schnider. Packing plane spanning double stars into complete geometric graphs. In *Proceedings of the 32nd European Workshop on Computational Geometry, EuroCG*, pages 91–94, 2016.
- [13] W. T. Tutte. On the problem of decomposing a graph into n connected factors. *Journal of the London Mathematical Society*, 36(1):221–230, 1961.