



# The impact of phytogetic feed additives on ruminant production: A review

N.A. Nastoh\*, M. Waqas, A.A. Çınar and M. Salman

Ondokuz Mayıs University, Faculty of Veterinary Medicine, Department of Animal Nutrition and Nutritional Diseases, Kurupelit Campus/Atakum, 55139, Samsun, Turkey

**KEY WORDS:** antioxidant, blood profile, meat and milk, performance, phytogetic feed additive, ruminant

Received: 25 April 2024

Revised: 14 July 2024

Accepted: 22 July 2024

**ABSTRACT.** Ruminant livestock, a vital source of high-value proteins such as red meat and milk, requires efficient and sustainable farming practices. To enhance production and quality, researchers have turned to various feed additives, with phytogetic feed additives (PFA) gaining significant attention following the European Union's ban on antibiotics in 2006. PFA, which are plant-derived products, are also often referred to as phytonutrients, phytochemical, plant natural compounds, plant bioactive compounds, or herbal feed additives, and are natural and safe alternatives to synthetic antibiotics. The review highlights the diverse benefits of PFA in ruminant livestock production. These compounds can improve ruminant performance, rumen fermentation, milk production, reproductive health, blood profiles, antioxidant and antimicrobials properties, immunity and meat quality. Specific bioactive compounds in PFA, such as essential oils, polyphenols and flavonoids, exhibit significant antimicrobial, antifungal, antioxidant and antiparasitic properties. They also modulate ruminal fermentation, reduce methane emission, and enhance feed intake and palatability. The mechanisms through which PFA function include improving feed utilisation and growth performance, as well as mitigating oxidative stress and inflammation. The mechanism behind this effect is based on donating hydrogen atoms, activating detoxifying enzymes and quenching reactive oxygen species. The review underscores the need for further research to fully understand these processes. Considering the rising global demand for meat and dairy products, PFA present a viable solution for sustainable livestock farming. They offer a natural effective and safe alternative to antibiotics, promising improved productivity, animal health, and food quality. This review aims to provide a comprehensive characterization of PFA and their potential in ruminant nutrition and production.

\* Corresponding author:  
e-mail: nastohnoor1919@gmail.com

## Introduction

The production of ruminant livestock is crucial to meet the growing demand for high-value proteins and to ensure food security and safety, both of which are essentials for human health (Caprarulo et al., 2022). Animal products such as meat, milk and eggs provide proteins of high biological value, which are easily digestible and contain all essential amino

acids (Pulina et al., 2017). These products are also rich in critical nutrients like heme-iron, vitamin B<sub>12</sub>, D<sub>3</sub>, zinc and calcium. The main objective of advancing dairy farming is to enhance the sustainability and efficiency of agricultural practices by producing ecological products free from hormones, antibiotics, and other xenobiotics. Utilising biologically active compounds (BAS) as feed additives is a practical, cost-effective, and scientifically robust method of

achieving this goal (Boyko et al., 2021). Feed additives are intentionally incorporated into livestock diets or drinking water to improve feed quality, enhance growth performance, aid digestion of anti-nutritive factors, prevent nutrient depletion, and positively impact animal production. These additives also increase animal resistance to diseases and parasites and can optimise energy consumption (Singh, 2015; Silveira et al., 2021). One of the most important challenges in animal husbandry is formulating a balanced diet that meets all the nutritional needs of animals (Pandey et al., 2019). Modern intensive farming systems have yielded excellent results in terms of obtaining high-quality animal products. However, even greater benefits related to fertility, daily weight gain, and disease resistance can be achieved by enriching diets with appropriate supplements (Singh, 2015). Due to these positive effects, Mihaylova et al. (2020) have predicted that global meat production would increase from 258 mln t in 2005–2007 to 374 mln t by 2030. Animal feed additives are categorised as technological, sensory, nutritional, zootechnical, and coccidiostats/histomonostats (Karásková et al., 2015; Pandey et al., 2019; Purba et al., 2020). They also include antioxidants, antimicrobials, immunomodulators, enzymes, prebiotics, probiotics, hormones, organic acids, mycotoxin binders, emulsifiers, metabolic modifiers, essential oils, phytochemical herbs, and pellet binders (Singh, 2015). Historically, humans used herbs to care for and feed domestic animals, with evidence of plant-based treatments dating back to Mesopotamia around 2600 BC (Hussein et al., 2015; Franz et al., 2020). In modern animal husbandry, the focus remains on increasing productivity, profitability, and farm efficiency to meet the rising nutrient demand driven by the growing global population. Various growth stimulants, including antibiotic growth promoters (AGP), have been employed to boost production. However, concerns about human health risks led the European Commission to ban in-feed antibiotics in 2006 (Hussein et al., 2015; Karásková et al., 2015; Steiner and Syed, 2015; Lillehoj et al., 2018; Stevanović et al., 2018; Pandey et al., 2019; Franz et al., 2020; Skoufos et al., 2020). As a result, livestock producers and researchers have sought safe, effective, and affordable alternatives to antibiotics. Fragrant plants, herbs, plant secondary metabolites, and aromatic vegetables, along with their essential oils (EO), have emerged as promising growth enhancers (Caroprese et al., 2020; Purba et al., 2020). These plant-derived compounds, known as phytobiotics, phytochemical feed additives (PFA), phytochemical compounds (PC), plant secondary metabolites (PSM), and phytochemical

compounds (PCC), can offer significant nutraceutical and nutritional benefits for animal health and nutrition (Yang et al., 2015; Stevanović et al., 2018).

Phytochemicals from herbs and vegetables serve as natural protection against pathogens and spoilage organisms. Compounds such as polyphenols, quinines, flavonoids, alkaloids, and polypeptides exhibit key bioactive properties essential for maintaining animal health. Feed additives help manage environmental stressors and improve food utilisation (Caroprese et al., 2020). Essential oils and other plant-based additives play a significant role in increasing livestock productivity by improving feed efficiency and performance indices such as milk yield, digestibility, and growth rate (Vasta et al., 2019; Purba et al., 2020). Phytobiotic compounds offer antimicrobial, antifungal, antioxidant, and antiparasitic benefits. They can alter rumen fermentation and microflora, reducing methane emissions and protein degradation, while improving feed intake and palatability in ruminants (Stevanović et al., 2018; Artuso-Ponte et al., 2020). Although the exact mechanisms of action of PFA are not yet fully elucidated, they are considered safe for animal nutrition. For example, the phytonutrient (PN) is known for its antimicrobial properties, which are effective against protozoa, fungi, and bacteria, as well as for its ability to modulate rumen fermentation and decrease ammonia levels (Oh et al., 2017; Purba et al., 2020). PFA contain essential oils and various plant secondary compounds. These volatile aromatic metabolites, including terpenoids and flavonoids, possess antimicrobial properties, enhance feed utilisation, modify rumen fermentation, inhibit methanogenesis, and improve overall animal performance (Singh and Gaikwad, 2020). For instance, the inclusion of limonene and carvacrol in the diets of lactating ewes has been shown to significantly enhance weight gain of weaned lambs, likely due to improved energy utilisation by the ewes (Varga-Visi et al., 2023). Additionally, incorporating winery waste products and citrus into ruminant diets as feed additives has been found to increase dry matter intake, improve nutrient digestibility, rumen fermentation, growth performance, and both health status and meat quality (Tayengwa and Mapiye, 2018). The use of *Coleus amboinicus* Lour in lamb diets was shown to reduce methane production, improve rumen environment, and meat quality due to increased n-3 PUFA content (Yanza et al., 2022). The latter authors also showed that a composite plant extract (CPE) containing horse gram seed, Shatavari root, Rohitaka bark, and pomegranate

peel could enhance rumen fermentation, lactation, and nutrient utilisation efficiency, all while reducing methane emissions. Considering these substantially beneficial properties, phytogetic feed additives are an excellent alternative to synthetic antibiotic growth promoters in increasing animal performance (Steiner and Syed, 2015). The objective of this review is to provide a comprehensive summary of the impact and applicability of PFA as alternatives to AGP in ruminant nutrition. The article discusses PFA effects on growth performance, antioxidant capacity, antimicrobial activity, milk production and composition, immunomodulation, blood profile, carcass characteristics, methane and ammonia emissions, rumen fermentation and volatile fatty acid production, gut microbiota reproduction, palatability, and feed intake. Given the lack of comprehensive reviews covering all these parameters in a single article, this study aims to fill this gap. The data collected will provide a compendium of knowledge on the use of various types of PFA in ruminant nutrition and how to optimise dietary strategies to improve productivity while reducing environmental pollution from methane, nitrogen, and other greenhouse gases.

## Literature search methods

To gather comprehensive data regarding the effects of dietary supplementation with PFA on various aspects of ruminant health and performance, we conducted an extensive literature review, and eventually included 180 studies from different sources. Our aim was to capture a detailed picture of how these natural additives influence parameters such as rumen fermentation, methane and ammonia emissions, milk production, reproduction, blood profile, antioxidant and antimicrobial properties, immunity, as well as meat composition and quality. We utilised several major databases, including ScienceDirect, PubMed, Scopus, Google Scholar, and Web of Science, ensuring a thorough query. Searches were carefully developed using specific keywords tailored to each database. The terms used included 'phytogetic feed additives', 'ruminants', 'cow', 'sheep', 'goat', 'buffalo', 'growth performance', 'carcass traits', 'ruminal fermentation', 'antioxidant', 'immunity', 'anti-microbial', 'methane and ammonia emission', 'milk composition', 'blood metabolites', and 'meat quality'. This approach to data collection has allowed us to compile a robust dataset of recent studies, providing valuable insights into the multifaceted effects of PFA on ruminant livestock.

## Brief definition, classification, and application of phytogetic feed additives

Medicinal and aromatic plants have contributed significantly to the development of human societies by improving living conditions from ancient times. Historical manuscripts, such as the Hippocratic Corpus by Hippocrates and *De Causis Plantarum* and *Historia Plantarum* by Theophrastus, dating from 300 to 500 BC, highlight their historical importance (Inoue et al., 2017). Phytogetic compositions are plant-derived bioactive substances that positively affect animal health and growth. These substances are typically utilised in the form of essential oils (EO) and extracts derived from herbs. Some are considered to possess antioxidant, antimicrobial, antifungal, and antiviral properties (Yang et al., 2015). Plant-derived additives are also added to animal's diets to improve their production and can generally be classified as spices, herbs, essential oils, and oleoresins (Hussein et al., 2015; Franz et al., 2020; Pashtetsky et al., 2020). Phytogetic, phytobiotic, and phytochemicals are natural bioactive compounds originating from plants, supplemented into livestock diets to improve performance and profitability (Lillehoj et al., 2018). These compounds are sourced from spices, herbs, and essential oils such as anise, cloves, thyme, or melissa (Steiner and Syed, 2015). Phytobiotics, also known as PN, are organic compounds produced by plant as secondary metabolites that have a wide range of properties protecting against bacteria, fungi, and yeasts. They are also used as rumen environment modifiers and methane mitigators in ruminants (Oh et al., 2017; Purba et al., 2020). These additives are studied in four main areas: nutritional supplementing (minerals, vitamins, herbal enzymes), animal husbandry supplementation (immunomodifiers, digestive stimulants, growth enhancers), technological supplementation (coccidiostats, antioxidants), and sensory replacements (Karásková et al., 2015; Stevanović et al., 2018). PFA include EO and many other plant-secondary compounds (PSC). The antimicrobial properties of the plants they are extracted from are often attributed to their EO, which contain terpenoids and phenylpropanoids (Huang et al., 2018). Herbs produce a wide array of metabolites crucial for their survival and growth. These metabolites are primarily categorised into two parts: primary and secondary metabolites (PPM and PSM, respectively). The former are necessary

for the growth and development of plants, while the latter protect plants from threats such as predators, stress, and environmental factors. In other words, secondary metabolites provide defence against insects, microorganisms, and other harmful animals (Karásková et al., 2015; Valenzuela-Grijalva et al., 2017; Pashtetsky et al., 2020). PSM are divided into three main classes: terpenes, phenolics, and nitrogen-containing compounds. PSM include various biologically active compounds that enhance plant defence against diseases and are used as flavouring agents (Singh and Singh, 2018; Nawrot-Chorabik et al., 2022). Classification of plant secondary metabolites is illustrated in Figure 1.

toxic and deleterious effects, PFA can be considered safe and may serve as the primary alternative to AGP. A fundamental restriction in using PFA is the lack of understanding of the optimal dose necessary for improving the growth performance of animals and the adaptation of ruminal microbes; additionally, the chemical structure of feed additives can influence their dosage requirements (Kholif and Olafadehan, 2021). Similarly, Steiner and Syed (2015) referenced European regulations highlighting improvements in digestibility, animal production, and animal health status by PFA application. Phytochemical compounds are usually applied in dry, solid, and ground form, or as extracts





Figure 1. Classification of plant secondary metabolites (Nawrot-Chorabik et al., 2022)

Many studies have documented the inhibitory role of PN in deaminating amino acids and methane production in the rumen, as well as enhancing the production of propionate and butyrate through fermentation. Active components such as cinnamaldehyde, eugenol, capsaicin, and allicin in some phytochemicals can activate transient receptor potentials in mammals, leading to changes in neurons, intestines, and pancreas (Oh et al., 2017). Interest in using PFA as natural alternatives for synthetic additives has recently increased (Singh, 2015). However, Yang et al. (2015) expressed some concerns regarding their use, including cost-related-effects, side effects, and regulatory obstacles. According to Stevanović et al. (2018), to align the market for animal production with consumer expectations, the 'CGE' concept must be applied, where 'C' stands for clean, 'G' for green, and 'E' for ethical. These authors also believe that despite some

(Lillehoj et al., 2018; Franz et al., 2020). Recently, encapsulation techniques have been developed to protect active ingredients against heat to prevent the release of odours until they reach the digestive tract (Steiner and Syed, 2015). Figure 2 demonstrates the reasons for using PFA.

Medicinal and aromatic plants (MAP) have a wide global distribution, including America, Africa, Europe, Australia, India, and China. Reports indicate the presence of 7 500 species in India, 6 000 species in China, 5 000 species in Africa, and 2 000 species in Europe (Pandey et al., 2020). According to estimates, 70 000 herbal species are currently utilised in traditional medicine (Inoue et al., 2017). These plants are used directly or in the form of PSM, with approximately 200 000 PSM identified worldwide (Singh and Singh, 2018). In addition, Steiner and Syed (2015) reported the presence of 320 000 PSM worldwide and noted that 1 mln



**Figure 2.** Primary reasons for using phytogetic feed additives (PFA, %) (Pandey et al., 2019)

AGP/AGPs – antimicrobial growth promoters, FCR – feed conversion ratio

compounds are produced in accordance with European legislation (1831/2003). According to Gessner et al. (2015), regulations define the use of plants in animal diets. Currently, the most applicable plant families in livestock feed include *Lamiaceae*, *Apiaceae*, *Rosaceae*, *Asteraceae*; other prevalent plant families are *Alliaceae*, *Brassicaceae*, *Cannabaceae*, *Fabaceae*, *Gentianaceae*, *Lauraceae*, *Myrtaceae*, *Papaveraceae*, *Rosaceae*, *Solanaceae*, *Zingiberaceae*, and some species with limited distribution (Franz et al., 2020).

## PFA effects on ruminant performance parameters

Medications containing antibiotics have increasingly been replaced by aromatic plants and the EO they produce. These plants are gaining significance due to their antibacterial properties and stimulating effects on animal digestive systems (Al-Kassie, 2009). Recently, PFA have gained attention as a substitute dietary strategy to AGP. Feed additives that promote growth primarily function by maintaining feed hygiene (e.g., through organic acids) and, more importantly, by positively influencing the ecology of the gastrointestinal microbiota by managing possible infections (Hussein et al., 2015). Herbs and their extracts, as MAP, offer innovative methods to enhance ruminant production and feed utilisation, while improving the quality of their products, such as milk and meat, possibly by affecting rumen digestion (Skoufos et al., 2020). The precise mechanisms of PFA as AGP are not yet fully elucidated. These mechanisms depend on factors such as pharmacokinetics, structure, dosage, animal species, production stage, and administration time (Skoufos et al., 2020). Phytogetic compounds, especially when combined with EO, can potentially increase feed intake by improving palatability through enhanced flavour and aroma

(Yang et al., 2015). The addition of 44.1 ppm of a prebiotic (cobalt lactate and arabinogalactons) and EO (carvacrol, caryophyllene, p-cymene, cineole, terpinene, and thymol) to the starter of newborn Holstein calves resulted in a linear increase in feed conversion ratio (FCR), dry matter intake (DMI), and average daily gain (ADG), along with an increase in body frame measurements (Liu et al., 2020). Similar results were obtained by Hashemzadeh et al. (2022), who observed that supplementing heat-stressed lambs with a mixture of cinnamon, turmeric, rosemary, and clove buds at 1% to 2% led to improvements in DMI, ADG, and body weight (BW). Adding 3 g/head/day of red propolis extract to the late pregnancy diet of Santa Ines ewes increased ( $P = 0.02$ ) crude protein and organic matter digestibility, as well as nitrogen intake ( $P = 0.03$ ) (Morsy et al., 2021). Conversely, increased consumption of a quebracho and chestnut tannin extract mixture exerted a negative impact on milk output, actual protein content, DMI, fat and protein-corrected milk (FPCM), and apparent nutritional digestibility, primarily when supplied at 0.90 and 1.80% diet DM (Aguerre et al., 2016). The growth performance of Sewa sheep was improved by adding 300 mg/kg of oregano essential oil (OEO) concentrate to their diets, which modified the structure and composition of their intestinal flora (Sun et al., 2022). The effects of PFA on ruminant performance are presented in Table 1.

## PFA effect on rumen fermentation and volatile fatty acid profile

The production of high-quality proteins for ruminants is mainly associated with ruminal microbial activity. However, this process can lead to significant protein and energy losses due to microbial fermentation, resulting in the release of ammonia and methane – two major environmental contaminants

**Table 1.** Effects of phytogetic feed additives (PFA) on ruminant performance

PFA source	Dosage	Study type	Animal species	Diet	Result	References
<i>Foeniculum vulgare</i> seed powder	0.75 and 1.5% of diet	<i>in vivo</i>	male lambs	forage-concentrate 40:60	DMI, LDG, FBW, and FCR ↑ ( $P < 0.05$ ) at 1.5%	Hajalazadeh et al., 2019
<i>Lippia alba</i> hay	30-60-90 g/kg of total DMI	<i>in vivo</i>	Alpine goat	TMR	DMI, acid detergent fibre, feed efficiency ↑ ( $P < 0.05$ ); the intake of total carbohydrates, NDF, and NFC ↓ ( $P < 0.001$ )	Da Silva et al., 2022
<i>Echinacea purpurea</i> leaf (EP)	4–8-12 g/head/day	<i>in vivo</i>	Ossimi lambs	TMR	BW ↑ 11.82, 13.80, and 12.26% for 4, 8 and 12 g EP, respectively; WG and growth rate ↑ (36.99 and 37.48%), respectively by 8 g EP	Tantawi et al., 2023
Blend of PFA (Digestarom, DSM) and Monensin M	0.5 g/head/day PFA, 0.3 g/head/day monensin	<i>in vivo</i>	Angus steers	TMR	similar DMI and ADG, but ↓ FI as monensin in the beginning phase but performance in the end	Yang et al., 2023
Grape pomace (GP)	5–10% of diet	<i>in vivo</i>	Churra ewe	TMR forage-concentrate 40:60	BBW-FBW and ADG ↓ not significant	Gómez-Cortés et al., 2018
<i>Woodfordia fruticosa</i> , <i>Solanum nigrum</i> , <i>Trigonella foenum-graecum</i>	2–3% of total DMI	<i>in vivo</i>	Jamunapari goat	concentrate and roughage	all nutrients digestibility ↑ ( $P < 0.05$ ); DMI and BWG ↓	Choubey et al., 2015
Anise, clove, and thyme EO	2 ml/head/day	<i>in vivo</i>	Shame goat	TMR	digestibility of ADF, OM, and ether extract ↑ ( $P < 0.05$ )	El-Essawy et al., 2021
Thyme leaf and celery seeds mixture 1:1	20 g/head/day	<i>in vivo</i>	Barki ewes	concentrate and fodder 60:40	digestibility and nutrient intake ↑ ( $P < 0.05$ )	Khattab et al., 2020
Rosemary and lemongrass	10 g of each separately/day/h	<i>in vivo</i>	Damascus goat	concentrate and berseem clover	NI ↓ ( $P > 0.05$ ); OM and fibre digestion ↑ ( $P < 0.05$ )	Kholif et al., 2017
Oregano and carvacrol	50 mg/kg of DM each separately	<i>in vivo</i>	dairy cow	forage and concentrate 60:40	digestibility, N utilisation ↑	Benchaar, 2020
Garlic extract, coriander seed, mesquite pods and oregano	6 ml/head/day orally	<i>in vivo</i>	sheep	TMR	ingestive behaviour and NI ↓ ( $P > 0.05$ ); ND ↑ ( $P < 0.05$ )	Da Silva et al., 2017
Black rice-purple maize extract 80:20	2-4-6% of diet	<i>in vivo</i>	male dairy cattle	TMR	intake of DM, OM, CP, NDF, ADG, and ADF ↑; EEI ↑ ( $P < 0.05$ )	Prommachart et al., 2021
Blend of <i>Echinacea</i> , garlic, thyme, caraway and liquorice	35, 20, 15, 15, and 15%, respectively	<i>in vivo</i>	Limousine bulls	roughage and concentrate	daily gain ↑ by 212 g/h/day while feed intake was ↓; protein and fat digestibility ↑ by 1.3 and 2.5%, respectively	Klebaniuk et al., 2016

EO – essential oil, TMR – total mixed ration, DM – dry matter, DMI – dry matter intake, LDG – live daily gain, FBW – final body weight, FCR – feed conversion ratio, NDF – neutral detergent fibre, NFC – non-fibre carbohydrate, BW – body weight, WG – weight gain, BBW – birth body weight, ADG – average daily gain, BWG – body weight gain, OM – organic matter, NI – nutrient intake, ND – nutrient digestibility, ADF – acid detergent fibre, EEI – ether extract intake;  $P < 0.05$  indicates that there is significant effect of PFAs on the given parameters

(Simitzis, 2017). Supplementation of Holstein cow diets with tannin extract from *Acacia mearnsii* at up to 0.43% of DMI did not affect acetate levels. However, it caused a linear increase in the molar fraction of butyrate and a corresponding decrease in propionate in the ruminal fluid (Oliveira et al., 2023). Similarly, the addition of citral oil at 0.08, 0.16, or 0.24 ml per kg BW to Saanen goat diets resulted in a linear increase in butyrate proportion in the ruminal fluid. The ingestion of citral oil was shown to positively affect ruminal propionate and butyrate levels; nonetheless, it had a negative impact on the acetate ratio in goats (Canaes et al., 2017). Meanwhile, Muñoz-Cuautle et al. (2022) did not observe

significant changes in ruminal parameters with the inclusion of 0.02 and 0.04% oregano essential oils in the lamb diet. Soltan and Patra (2020) reported that propolis improved rumen nitrogen metabolism and the formation of individual and total short-chain fatty acids, which benefits rumen microbial fermentation. Additionally, propolis can reduce methane production during ruminal fermentation. Similarly, Min and Solaiman (2018) found that diets rich in phytonutrient tannins could modify the microbiota and mitigate methane gas emissions. Studies have indicated that tannins can affect rumen fermentation and microbiota in both beneficial and detrimental ways. The effectiveness of tannins varies depending on their

quantity, source, and the diet in which they are included. Rivera-Chacon et al. (2022) demonstrated that a phytogetic combination of clove powder (*Syzygium aromaticum*), mint oil (*Mentha arvensis*), thymol, and eugenol influenced rumen microbial populations and fermentation patterns. In an *in vitro* study, Faniyi et al. (2021) found that neem, drumstick, and scent (*Ocimum gratissimum*), along with spices like garlic, ginger, and onion leaves, could influence rumen fermentation and decrease sheep methane generation (Faniyi et al., 2021). Overall, depending on the additive and its concentration, different PFA, such as tannins and essential oils, can greatly improve fermentation and volatile fatty acid (VFA) production, while lowering methane emissions, although their efficacy varies greatly. Additional research is necessary

to maximise these benefits for both ruminant health and environmental sustainability. PFA effects on rumen fermentation and VFA profile are presented in Table 2.

## PFA effect on methane and ammonia emission

In addition to causing environmental pollution and intensifying the greenhouse effect, ruminal methane (CH<sub>4</sub>) emissions also contribute to low production efficiency and animal energy losses. Therefore, it is crucial to develop strategies for reducing methane emissions from ruminants. Research has shown that feed additives such as plant extracts and nitrogen-containing compounds can significantly

**Table 2.** Effects of phytogetic feed additives (PFA) on rumen fermentation and volatile fatty acid (VFA) profile

PFA sources	Dosage	Study type	Animal species	Diet	Result	References
Bee propolis extract	3 g/head/day	<i>in vivo</i>	Santa Ines ewes	basal diet	total short-chain fatty acids and acetate/propionate proportion ↑ ( $P < 0.05$ )	Morsy et al., 2021
<i>Boswellia sacra</i> resin (BS)	2 and 4 g/head/day	<i>in vivo</i>	Nubian goats	basal diet	linear ↓ ( $P < 0.01$ ) in acetate molar ratio. Linear ↓ ( $P = 0.01$ ) in total short-chain fatty acids. linear ↑ ( $P < 0.01$ ) in propionate molar proportions; and quadratically ↓ ( $P < 0.05$ ) in branched-chain volatile fatty acids	Soltan et al., 2021
Oregano and cinnamon EO, turmeric extract, and tannic acid	300 mg – encapsulated oils/kg and 12.5 mg of curcumin/kg	<i>in vivo</i>	Holstein calves	concentrate	butyric acid, acetic acid, and propionic acid concentrations ↑	Brunetto et al., 2023
Red osier dogwood	0.3, 0.7 and 0.10% of diet	<i>in vivo</i>	beef heifers	high-grain diet	↑ rumen pH and total volatile fatty acid	Wei et al., 2019
CFE	50-100-150 g/day	<i>in vivo</i>	Holstein dairy calves	TMR	↑ total volatile fatty acid	Zhao et al., 2023
Oregano EO	0, 130 mg, and 260mg/day/h	<i>in vivo</i>	Pingliang red cattle	TMR	↑ propionate and butyrate's relative abundance in the high group	Zhang et al., 2021
TA	0.3, 0.9 and 2.7% dry matter	<i>in vivo</i>	Holstein bull	TMR	irrespective of dose TA ↓ ( $P < 0.05$ ) valerate and isobutyrate; 0.9 and 2.7 % ↓ ( $P < 0.05$ ) butyrate; acetate and acetate to propionate ratio ↑ ( $P < 0.05$ )	Wang et al., 2022
Rosemary and lemongrass	10 g of each separately/ day/h	<i>in vivo</i>	Damascus goat	concentrate and berseem clover	the ruminal concentration of SCFA, propionate, total UFA, and linoleic acid ↑ ( $P < 0.05$ ); total SFA ↓ ( $P < 0.05$ )	Kholif et al., 2017
PFA mixture	3 and 6 g/h/day	<i>in vivo</i>	Friesian cows	TMR	rumen pH, total VFA, propionate and acetate ↑ ( $P < 0.05$ )	Kholif et al., 2021
Oregano and carvacrol	50 mg/kg oDM each separately	<i>in vivo</i>	dairy cow	forage and concentrate 60:40	ammonia, pH, VFA, and protozoa count ↓	Benchaar, 2020
<i>Dolichos biflorus</i> seed, <i>Asparagus racemosus</i> root, <i>Amoora rohituka</i> bark Homogeneous blend	10, 20, 30, and 40 g/kg substrate	<i>in vitro</i>	Beetal goat rumen fluids	green bajra + concentrate	VFA and ratio of propionate ↑ ( $P < 0.001$ ); acetate's percentage and the ratio of acetate to propionate ↓	Shilwant et al., 2023

EO – essential oil, CFE – citrus flavonoid extract, TMR – total mixed ration, SCFA – short-chain fatty acids, UFA – unsaturated fatty acids, SFA – saturated fatty acids, TA – tannic acid;  $P < 0.01$  indicates significant effect of PFA on the given parameters

reduce ruminant methane emissions (Sun et al., 2021). These phytochemical additives reduce hyperammonia ( $\text{NH}_3\text{-N}$ ) levels, suppress methanogenic and other harmful bacterial populations, increase VFA concentrations, and lower  $\text{NH}_3\text{-N}$  concentrations, leading to favourable alterations of rumen fermentation (Kholif and Olafadehan, 2021). Different combinations of phytogenic materials can lower the amount of  $\text{CH}_4$  released by ruminants into the atmosphere. Methane is the second most significant greenhouse gas, and ruminants are a major source of its production (Karásková et al., 2015). Research indicates that hops or oak tannin extracts, used alone or in combination, do not affect the daily intestinal  $\text{CH}_4$  production of cows (Focant et al., 2019). However, feeding Friesian lactating cows with 4% coriander per head per day significantly reduced ( $P < 0.001$ ) ruminal ammonia-N content (Matloup et al., 2017). Legumes, which are high in tannins, can be used to increase dietary nitrogen content and improve pasture productivity. Tropical legumes like *Desmanthus* and *Leucaena leucocephala* have strong antimethanogenic properties and are considered a viable method for reducing intestinal  $\text{CH}_4$  emission in cattle production (Almeida et al., 2021). In a study on Segurena sheep, olive cake treated with alkali was used to assess its effect on enteric methane emissions. These authors found that adding 400 g/kg of olive cake to the basal diet significantly decreased the production of enteric  $\text{CH}_4$  (Aguilera et al., 2021). An *in vitro* work revealed that adding *Moringa oleifera* oil changed fermentation kinetics from acetate to propionate, thereby markedly reducing methane production in all mixed rations. Additionally, low and medium roughage feeds (30–50%) were shown to reduce the population of protozoa and methanogens (Ebeid et al., 2020). Buffalo calves were fed three distinct plant mixtures: Mix-1 (ajwain oil and lemon grass oil at a 1:1 ratio at 0.05% DMI), Mix-2 (garlic and soapnut at a 2:1 ratio at 2% of DMI), and Mix-3 (garlic, soapnut, harad, and ajwain at a 2:1 ratio at 1% of DMI). The result showed that feed conversion ratio was high (10.2%) in Mix-3, while methane emission was low up to 17.8% in Mix-1 (l/kg DMI and l/kg digestible DMI) (Samal et al., 2016). Methane production and ammonia emissions of beetal goats decreased linearly ( $P < 0.05$ ) when a composite plant extract (*Dolichos biflorus* seed, *Asparagus racemosus* root, *Amoora rohituka* bark) was homogenously mixed and applied at levels of 10, 20, 30 and 40 g/kg free moisture substrate. The outcomes improved with increasing doses of the composite (Shilwant et al., 2023).

In both *in vivo* and *in vitro* studies, polyphenols from *Coleus amboinicus* lour (CAL) included at 20% in the lambs' diet significantly reduced ruminal methane production by suppressing the growth of methanogen bacterial communities. Additionally, CAL supplementation favourably affected the ruminal environment by balancing the development of ruminal microorganisms responsible for fatty acid fermentation and biohydrogenation (Yanza et al., 2022). Additionally, an *in vitro* study by Hundal et al. (2019) investigated the effects of various herbal feed additives on the nutritional value. The study found that a TMR supplemented with *Anethum sowa* at 2% of the substrate resulted in more effective fermentation compared to TMR with other herbal feed additives. This supplementation increased the production of total and individual VFA while reducing methane and ammonia emissions. Another *in vitro* research has shown that the combination of a low dose of eucalyptus oil (5  $\mu\text{l}$ ) with mulethi root extract (15  $\mu\text{l}$ ) can significantly reduce the rate of ruminal ammonia production ( $P < 0.001$ ) without compromising feed digestion (Chanu et al., 2020). Feeding lactating cows a diet consisting of immature herbage and vine leaves has been shown to decrease methane emissions, as demonstrated for a ration comprising 69% legume-grass herbage, 13% grass hay, and 13% vine leaves (Birkinshaw et al., 2022). Adding green tea and oregano extract to dairy cow diets at concentrations of 0.056% and 0.028%, respectively, enhanced the digestible fraction of ingested DM and reduced methane emissions, measured in g per kg of digestible DMI (Kolling et al., 2018). The  $\text{CH}_4/\text{VFA}$  molar ratio decreased when hop pellets at the highest dose (475 mg/l) were combined with oak extracts, whereas the same additives applied separately exerted no significant effect. Additionally, oak extract, whether used alone at the highest dose (309 mg/l) or in combination with hop pellets, significantly reduced ammonia nitrogen concentrations (Quynh et al., 2018).

### PFA effect on milk production and composition

The production of high-quality milk by dairy cattle is closely associated with optimal animal health (Yang et al., 2019). Understanding the nutritional value, technological properties, and traceability of milk requires a thorough examination of potential correlations between feeding systems and milk composition. Such insights could lead to



the identification of new biomarker molecules (Li et al., 2020). Consequently, enhancing dairy performance may be achieved by incorporating appropriate feed additives into the diets of lactating animals (Focant et al., 2019). For instance, the addition of *Acacia mearnsii* tannin extract at up to 0.43% on a DM basis did not significantly affect milk yield or composition in Holstein cows, although it did result in a slight reduction in milk urea nitrogen (Oliveira et al., 2023). Canaes et al. (2017) reported no significant effects on milk production, fat-corrected milk (FCM), and fatty acid profile except for a linear reduction in milk fat production when citral oil was included in the diets of Saanen goats. Nubian goats receiving 2–4 g/day/animal of *Boswellia sacra* resin showed a linear ( $P < 0.05$ ) increase in milk yield, energy-corrected milk (ECM), and net energy for lactation (Soltan et al., 2021). Adding a blend of green tea and curcuma extract (95:5) to the diet of lactating cows at 0.175 g resulted in a significant increase ( $P < 0.05$ ) in energy-corrected milk yield (Winkler et al., 2015). Furthermore, including 10% ensiled olive cake in the diet of dairy cows (concentrate-to-roughage ratio of 64:36) led to an increase in milk fat yield, although the percentages of fat and protein, as well as protein yield, remained unchanged (Neofytou et al., 2020). Similarly, replacing maize silage with 60, 120, and 180 g of pomegranate by-product silage in a TMR diet for dairy cows resulted in improved digestibility of NDF and ADF at the levels of 60 and 120 g. While total milk production and ECM increased, milk protein and lactose content decreased at the 120 g level (Khorsandi et al., 2019). In conclusion, the impact of various PFA on milk production and composition varies considerably. Pomegranate by-product silage, green tea, curcuma extract, and *Boswellia sacra* resin have all shown promising effects, whereas tannin extract and citral oil have provided only modest benefits. Further research is necessary to fully realize the benefits of these supplements in enhancing dairy animal health and milk production. The effects of PFA on milk production and composition are summarised in Table 3.

## PFA effect on reproductive performance

The unit cost of production in the cow-calf industry is determined by reproductive performance (Rasby and Funston, 2016). Increasing reproductive efficiency is critical for both commercial success and the long-term viability of livestock operations (Swelum et al., 2021). While, modern reproductive technologies have expanded the reproductive

performance in various domestic cattle breeds, conventional procedures are also commonly applied to improve reproduction outcomes. Advances in biotechnologies have enabled detailed analysis and modification of reproductive processes both *in vitro* and *in vivo* (Mihaylova et al., 2020). Phytochemicals, used as feed additives, serve multiple purposes, including reducing methane emissions and substituting antibiotics (Swelum et al., 2021). Among these, phytoestrogens – phenolic compounds found in plants – are particularly noteworthy for their estrogenic activity, either as agonists or antagonists in mammals (Hashem and Soltan, 2016). According to Swelum et al. (2021), PFA in fresh and conserved semen affected hormone profiles, sperm quality, and animal sexual behaviour. Azimi et al. (2020) studied the effects of purslane aqueous extract (PAE), purslane methanolic extract (PME), and purslane ethanolic extract (PEE) at concentrations of 25, 50, and 100 µg/ml on the quality of frozen-thawed goat spermatozoa. The results demonstrated that PAE at 50 µg/ml, PME at 50 µg/ml, and PEE at 50 µg/ml significantly reduced malondialdehyde (MDA) levels and improved total motility, viability, and mitochondrial activity ( $P < 0.05$ ) compared to the control. Of all extracts studied, PME at 50 µg/ml produced the best results. Another study reported the effect of quercetin in tris citric acid extender at concentrations of 50, 100, 150, and 200 µM on the post-thaw quality and *in vivo* fertility of buffalo spermatozoa. The results showed a significant ( $P < 0.05$ ) improvement in progressive motility (PM%), plasma membrane integrity (PMI%), supra vital plasma membrane integrity (SVPMI%), and acrosome integrity (ACR-1%) following treatment. After thawing, there was a significant increase in sperm SVPMI at 100 and 150 µM, as well as sperm PM and ACR-I at 150 and 200 µM (Ahmed et al., 2019). Similar results were obtained when 1.0% green tea extract (GTE) was added to the tris-citric acid extender. This addition enhanced total antioxidant capacity, increased *in vitro* longevity (%), and *in vivo* fertility, while reducing lipid peroxidation and DNA fragmentation in buffalo bull spermatozoa, following a freezing and thawing procedure (Ahmed et al., 2020). An *in vitro* study demonstrated that the addition of up to 100 µg/ml of *Moringa oleifera* leaf extract, along with reproductive hormones, increased the maturation rate of sheep oocytes. This treatment likely stimulated mRNA expression and the synthesis of essential proteins, such as maturation-promoting factor (MPF) (Barakat et al., 2015). In another *in vitro* study, the application of

**Table 3.** Effects of phytochemical feed additives (PFA) on milk production and composition

PFA sources	Dosage	Study type	Animal species	Diet	Result	References
<i>Quercus robur</i> 'oak tannin' (OT) <i>Humulus lupulus</i> 'hops' (HP)	169 g of OT; 56 g of HP.	<i>in vivo</i>	Holstein lactating cow	TMR	hops (HP) did not affect milk production and composition; OT-HP mixture enhanced the milk's fatty acid composition; $\alpha$ linoleic acid was increased by 17.5%	Focant et al., 2018
<i>Lippia alba</i> hay	30-60-90 g/kg of total DMI	<i>in vivo</i>	Alpine goat	TMR	MY, corrected MY <sup>3.5%</sup> , corrected solids, defatted dry extract, and milk contents $\uparrow$ ( $P < 0.05$ )	Da Silva et al., 2022
Anise, clove, and thyme EO	2 ml/head/day	<i>in vivo</i>	Shame goat	TMR	MY, protein, and lactose $\downarrow$ . Fat yield and fat composition $\uparrow$ ( $P < 0.05$ ); milk C12 and C8 FA $\downarrow$ ( $P < 0.05$ ); composition of UFA and MUFA $\uparrow$ ( $P < 0.05$ ). PUFA $\downarrow$ ( $P < 0.05$ )	El-Essawy et al., 2021
Thyme leaf and celery seeds mixture 1:1	20 g/head/day	<i>in vivo</i>	Barki ewes	concentrate and fodder 60:40	WG, ADG, MY, milk component yield, and feed efficiency $\uparrow$ ( $P < 0.05$ ); milk composition $\downarrow$	Khattab et al., 2020
<i>Capsicum oleoresin</i>	250-500-1000 g/head	<i>in vivo</i>	Holstein cow	TMR	MY and energy corrected MY $\uparrow$ ; MF $\downarrow$	Oh et al., 2015
Rosemary and lemongrass	10 g of each separately/d/head	<i>in vivo</i>	Damascus goat	concentrate and berseem clover	MF and milk production $\uparrow$ ( $P < 0.05$ )	Kholif et al., 2017
PFA mixture	3 g and 6 g/head/d	<i>in vivo</i>	Friesian cows	TMR	milk rate, composition, total solids, fat, protein, and lactose $\uparrow$ at 3 g	Kholif et al., 2021
Coriander oil	4 g/head/d	<i>in vivo</i>	Friesian cow	concentrate and berseem clover	milk production, ECM, MF, milk efficiency $\uparrow$ ( $P < 0.05$ )	Matloup et al., 2017
Propolis	5 g/kg diet	<i>in vivo</i>	Barki ewes	concentrate and berseem hay	MY, MF, and milk total solid $\uparrow$ ( $P < 0.05$ )	Shedeed et al., 2019
<i>Dolichos biflorus</i> seed, <i>Asparagus racemosus</i> root, <i>Amoora rohituka</i> bark Homogeneous blend	20 g/kg of diet	<i>in vivo</i>	Beetal goat	Bajra and concentrate 60:40	MY $\uparrow$ ( $P = 0.017$ ), lactose, and protein $\uparrow$ ( $P = 0.045$ ); MF and solid $\downarrow$ ( $P > 0.10$ ); the number of somatic cells $\downarrow$ ( $P = 0.045$ )	Shilwant et al., 2023
Mulberry leaf (flavonoid)	15, 30,45 g/day	<i>in vivo</i>	Murrah buffalo	TMR	daily MY $\uparrow$ ( $P = 0.038$ ) in all levels; T3 and T4 in serum $\uparrow$ ( $P < 0.05$ ), FCM, and milk protein $\uparrow$ by 45 g	Li et al., 2020
Grape seed and grape mark extract	1% of diet	<i>in vivo</i>	Holstein cow	TMR	MY and milk protein yield $\uparrow$	Gessner et al., 2015

TMR – total mixed ration, DMI – dry matter intake, FA – fatty acid, UFA – unsaturated fatty acid, MUFA – monounsaturated fatty acid, PUFA – polyunsaturated fatty acid, EO – essential oil, WG – weight gain, ADG – average daily gain, MY – milk yield, MF – milk fat, ECM – energy-corrected milk, FCM – fat-corrected milk, T3 – triiodothyronine, T4 – thyroxine;  $P < 0.05$  indicates significant effect of PFA on the given parameters

quercetin at concentrations of 1, 10 and 100 ng/ml to cattle ovarian cells resulted in a reduction in proliferating cell nuclear antigen (PCNA) and Bax accumulation. Additionally, the release of progesterone (P4) from granulosa cells, and insulin-like growth factor I (IGF-I) secretion from the cattle cells were increased at the low dose (10 ng), but decreased at the high dose (100 ng) (Sirotkin et al., 2019). Similarly, Ardeshirnia et al. (2017) reported that quercetin concentrations of 5 and 10  $\mu$ g/ml improved the quality of frozen-thawed ram epididymal spermatozoa.

## PFA effect on blood profile

Blood metabolites are commonly applied to diagnose and predict disorders, as well as to assess normal physiology and well-being of animals. Waqas et al. (2023) reported increased blood urea nitrogen levels, and elevated serum globulin and total protein levels when the cows were fed a plant-based additive mixture. These results indicated optimal kidney function, improved nutritional status, increased microbial protein synthesis, and minimal protein catabolism (Waqas et al., 2023).

Weaned dairy heifer calves exposed to heat stress showed a negligible effect of PFA supplementation (Digestaron, DSM Nutritional Products Inc.) on feed intake, average daily gain (ADG), blood O<sub>2</sub> partial pressure, and blood levels of inflammation markers such as haptoglobin and lipopolysaccharide-binding protein (LBP). In all measurements, serum haptoglobin and plasma LBP concentrations were 44% and 38% lower, respectively, in PFA calves compared control calves (Wickramasinghe et al., 2023). Another study supplemented a mixture of cinnamon, turmeric, rosemary, and clove buds to the diet of heat-stress lambs, which resulted in a 1–2% increase in cholesterol but reduced creatinine levels (Hashemzadeh et al., 2022). Moreover, the addition of 250 g/head/day of anise seed powder to the diet of Holstein calves for 60 days did not significantly affect the content of total protein, albumin, glucose, cholesterol, aspartate aminotransferase (AST), gamma-glutamyl transferase (GGT), total antioxidant status (TAS), and total oxidant status (TOS) compared to control (Demirhan and Karafakioğlu, 2022). The effects of PFA on blood profile is presented in Table 4.

## Effect of PFA as antioxidants in ruminants

Excessive levels of reactive oxygen species (ROS), such as hydroxyl radical (OH•), superoxide anion radical, or superoxide anion (O<sub>2</sub>•<sup>-</sup>), are associated with various diseases, such as cancer and cardiovascular disorders (Aqeel et al., 2019). The antioxidant properties of a plant are typically related to its content of phenolic compounds, which generally act by inhibiting oxidation-reduction reactions or scavenging oxygen by donating hydrogen atoms (Amarowicz and Pegg, 2019). In line with this, Piao et al. (2023) have noted that antioxidant compounds in plants can suppress oxidative reactions by donating electrons to metal ions, with polyphenolic compounds demonstrating greater activity and effectiveness compared to vitamins E and C. Secondly, plant extracts, rich in hydroxyl groups similar to flavonoids, can provide more electrons, thereby enhancing antioxidant capacities. This process involves suppressing oxygen atoms and stimulating the production of antioxidant enzymes like glutathione peroxidase (GPx) and superoxide dismutase (SOD) (Purba and Paengkoum, 2022). Furthermore, Hrelia and Angeloni (2020)

**Table 4.** Effects of phytogetic feed additives (PFA) on blood profile

PFA sources	Dosage	Study type	Animal species	Diet	Result	References
Bee propolis extract	3 g/head/day	<i>in vivo</i>	Santa Ines ewes	basal diet	total protein, globulin, and glucose ↑ ( $P < 0.05$ ); cortisol, triiodothyronine, and thyroxine ↓ ( $P < 0.05$ )	Morsy et al., 2021
<i>Boswellia sacra</i> resin (BS)	2 and 3 g/day/head	<i>in vivo</i>	Nubian goats	basal diet	blood plasma glucose, nonesterified free fatty acid, and β-hydroxybutyrate ↓ ( $P < 0.01$ )	Soltan et al., 2021
Oregano and cinnamon EO, turmeric EO, and tannic acid	300 mg EO/kg and 12.5 mg of turmeric/kg of diet	<i>in vivo</i>	Holstein's calves	concentrate	serum glucose ↓, globulin, and total protein ↑	Brunetto et al., 2023
<i>Echinacea purpurea</i> leaf	4-8-12 g/head/day	<i>in vivo</i>	Ossimi lambs	TMR	triglycerides, cholesterol and AST ↓	Tantawi et al., 2023
Red osier dogwood	0.3,0.7 and 0.10% of diet	<i>in vivo</i>	beef heifers	high-grain diet	serum amyloid and haptoglobin of plasma ↑	Wei et al., 2019
<i>Capsicum oleoresin</i>	250-500-1000 g/head	<i>in vivo</i>	Holstein cow	TMR	blood serum β-hydroxybutyrate, neutrophils, eosinophils, and neutrophil-eosinophil ratio ↑	Oh et al., 2015
Coriander oil	4 g/head/day	<i>in vivo</i>	Friesian cow	concentrate and berseem clover	glucose concentration ↑ ( $P = 0.013$ ); short-chain fatty acid ↓ ( $P < 0,05$ )	Matloup et al., 2017
Carvacrol and limonene	0.5 g/head/day	<i>in vivo</i>	Tzigai sheep	grass hay + concentrate	β-hydroxybutyrate ↓ ( $P < 0.001$ ), triglycerides ↓ ( $P = 0.014$ ); nonesterified fatty acid ↓ ( $P = 0.021$ ); fructosamine ↓ ( $P = 0.002$ )	Varga-Visi et al., 2023
Black rice- purple corn extract 80:20	2, 4, 6% of diet	<i>in vivo</i>	male dairy cattle	TMR	glucose, urea, total cholesterol, AST, ALT, and protein carbonyl ↑, but MDA concentration ↑ ( $P < 0.05$ )	Prommachart et al., 2021

TMR – total mixed ration, EO – essential oil, AST – aspartate aminotransferase, ALT – alanine aminotransferase,  $P < 0.05$  indicates significant effect of PFA on the given parameters

have indicated that the mitochondria and chloroplasts of plant cells are associated with natural antioxidants, which can be transported to the mitochondria of animal cells when plants are ingested. This transfer can alter oxidative phosphorylation and electron transportation chain processes. The process of oxidation is a vital issue to consider from the first seconds of birth due to the sensitivity of animals to oxidative stress (Piao et al., 2023). This adverse conditions may occur when there is a deficiency of antioxidants, which can help prevent the spread of reactive oxygen species and mitigate their effects (Singh, 2015; Muñoz-Cuautle et al., 2022). Various intrinsic free radicals, and those originating from feed components can react with lipids in the animal body, making them unstable, reducing their nutritional value, and shortening the shelf life of meat (Singh, 2015). Plants and spices are well known as sources of antioxidants, and volatile oils extracted from plants of the family *Labiatae* are most attractive in this regard. Due to having phenolic and terpenoid compounds, rosemary, oregano, and thyme of the family *Labiatae*, as well as spices from the family *Zingiberaceae*, such as anise and coriander, are recognised for their significant antioxidant properties (Hussein et al., 2015). Plant polysaccharides also have scavenging properties that neutralise free radicals in the animal body and can prevent lipid peroxidation by altering the functional length chain of the lipid peroxidation reaction. Additionally, they stimulate the activation of antioxidant enzymes such as GPx and SOD (Hussein et al., 2015). Herbal extracts (HE) are widely used in animal nutrition to improve health status and immunity, and thus their incorporation into animal diets is recommended particularly in addressing oxidative stress (Piao et al., 2023). The findings of Manuelian et al. (2021) have shown that PFA can control oxidative reactions as effectively as synthetic antioxidants such as vitamins E and C. The latter authors argued that while research on PFA has primarily focused on poultry rather than ruminants, there is potential for PFA to serve as a unique alternative to synthetic growth promoters. Similarly, Yang et al. (2015) have argued that PFA can also be successfully substituted as a natural antioxidant with synthetic antioxidants like butylated hydroxytoluene and ethoxyquin. According to Pashtetsky et al. (2020), carvacrol and thymol SM may penetrate the cytoplasmic membrane, disrupt it and subsequently depolarise within the cytoplasm. According to Brunetto et al. (2023), incorporating a mixture of oregano and cinnamon EO with turmeric extract and tannin to the diet of Holstein steers led

to a reduction in the levels of thiobarbituric acid reactive substance (TBARS) and ROS; in contrast, there was an increase in the activity of glutathione S-transferase and protein thiol levels. Similarly, Tantawi et al. (2023) reported that the inclusion of 4%, 8%, or 12% *Echinacea purpurea* leaf extract (EP) as a dietary supplement for lambs significantly increased ( $P < 0.01$ ) SOD activity, glutathione levels, and total antioxidant capacity (TAC). Further, a blend of seeds and leaves (75:25) of three Bangladeshi traditional medicinal plants was used as a PFA in the daily feed in post-weaned bull calves, resulting in increased total antioxidant levels and activity of glutathione peroxidase (Bostami et al., 2021). OEO has been shown to act as a potent natural antioxidant, maintaining the stability of lamb meat at a high level (Muñoz-Cuautle et al., 2022). Supplementing lamb diet with a blend of rosemary, cinnamon, turmeric, and clove buds at 2% was shown to enhance total liver antioxidant capacity and reduced MDA levels in the liver (Hashemzadeh et al., 2022). Kumar et al. (2022) found that a mixed leaf meal of eucalyptus and poplar (*Populus deltoides*) (50 g/head/day) containing 0.71 g of condensed tannins, 2.30 g of tannin phenolics, and 3.19 g of phenolics, could positively alter the immune and antioxidant status of buffalo calves. HE have been widely recognised as alternatives to antibiotics due to their active properties. They can reduce oxidative stress in ruminants and positively affect animal meat quality and production (Huang et al., 2018). Supplementing buffalo cattle rations with 1% green tea extract and Tris citric acid extender as feed additives improved total antioxidant capacity (Ahmed et al., 2020). Similarly, a notable increase in antioxidant activities was observed when using PFA from three plant species (*Woodfordia fruticosa*, *Solanum nigrum*, and *Trigonella foenum-graecum*) in the diet of adult goats (Choubey et al., 2015). Adding ethanolic saffron petal extract (SPE) to lamb diets reduced lipid oxidation and enhanced antioxidant status. The antioxidant level in lambs were similarly affected by SPE and vitamin E (Alipour et al., 2019). The antioxidant effects of selected PFA on ruminants are shown in Table 5.

### **PFA effect on the gut microbiota of ruminants**

The antimicrobial properties of PFA have been well-documented in the published research (Skoufos et al., 2020). The active substances responsible for their antimicrobial effects include tannins (Huang et al., 2018) and phenolic compounds

**Table 5.** Antioxidant effects of selected phytogetic feed additives (PFA) on ruminants

Plant species/PSM	Plant part/material	Dosage	Study type	Animal species	Diet/material	Result	References
Condensed tannin	–	1, 1.5, and 2%	<i>in vivo</i>	lamb	FILM and wheat bran	G, SOD, and CAT activity, total thiol, protein thiol ( $P < 0.05$ ) ↑, LPO ↓	Dey et al., 2015
PFAs ( <i>Woodfordia fruticose</i> - <i>Solanum nigrum</i> - <i>Trigonella foenum-graecum</i> )	flowers, shoots, leaves -whole plant, seeds, respectively	2 and 3 %	<i>in vivo</i>	goat	concentrate and roughage	GPx, C, and GST ↑ – non-enzymatic antioxidant ↓	Choubey et al., 2015
Oregano and green tea extract	extract	10 g/day/head-5g/day/head respectively	<i>in vivo</i>	Jersey cow	basal diet	oxidative stress biomarker of blood ↓	Vizzotto et al., 2021
<i>Punica granatum</i> PMG. <i>Caesalpinia spinosa</i> TA. <i>Castanea sativa</i> CH. <i>Uncaria gambir</i> GM.	extract	80 µg/ml	<i>in vitro</i>	Holstein dairy cow	bovin aortic endothelial cell	antioxidant activity was 63% ↓, 45% ↓, 51% ↓, and 27% ↓ for PMG, GM, CH, and TA, respectively;	Ciampi et al., 2020
Pomegranate	peel + seed	400 g seed + 400 g peels	<i>in vivo</i>	Holstein dairy cow	basal diet	PTAA ↑; MDA and blood SOD ↑	Safari et al., 2018
Pomegranate peel	extract	0.1, 1.0 and 10 µg/ml	<i>in vitro</i>	Dairy cattle	bovine mammary epithelial cell	MDA ↓ (31.1%, $P < 0.001$ )	Mastrogiovanni et al., 2020
<i>Astragalus membranaceus</i>	root	1%	<i>in vivo</i>	cashmere goat	basal diet	MDA ↓ ( $P < 0.01$ ) – SOD and CAT ↑ ( $P < 0.001$ )	Luo et al., 2020
Grape	extract	670 mg/head/day	<i>in vivo</i>	Holstein heifer	TMR	Total antioxidant capacity ↑ ( $P < 0.05$ )	Engler et al., 2022
Mulberry	leaf (flavonoid)	15 g, 30, 45 g/day	<i>in vivo</i>	Murrah buffaloes	TMR	MDA, CAT ↓ ( $P = 0.012$ ); GPx, HSP ↑ ( $P < 0.01$ ); T-AOC ↓ ( $P = 0.001$ )	Li et al., 2020
Black rice- purple corn	extract	2-4-6% of diet	<i>in vivo</i>	male dairy cattle	TMR	MDA ↓ ( $P < 0.05$ )	Prommachart et al., 2021
<i>Astragalus membranaceus</i>	root	5, 10, 15, 20, 30 r/kg of diet	<i>in vivo</i>	Ram lambs	basal diet	T-SOD, and T-AOC ↑ ( $P \leq 0.018$ ). CAT and MDA ↑ ( $P \leq 0.001$ )	Hao et al., 2020
Green tea	extract	10 mg/kg/BW/d	<i>in vivo</i>	Holstein calves	colostrum and milk replacer	antioxidative status ↓	Maciej et al., 2016
Oregano	essential oil	0.02–0.04% of diet	<i>in vivo</i>	Hampshire lamb	basal diet (concentrate)	antioxidant activity ↑ ( $P < 0.05$ ).	Muñoz-Cuautle et al., 2022;
<i>Origanum vulgare</i> , <i>Citrus</i> spp., and <i>Echinacea</i> spp. (blend 1) – <i>Capsicum</i> spp., <i>Cinnamomum</i> spp., <i>Eugenia caryophyllata</i> spp. (blend 2) + an emulsifier each	extract	Blend 1 (0.11%), Blend 2 (0.0125%) of diet	<i>in vivo</i>	ewes, rams	concentrate barley-based diet	GPx and T-SOD in blood remained unchanged ↓, TBARs ↓	Passetti et al., 2021
Propolis	extract	150-200-250 µl/kg of BW/day	<i>in vivo</i>	lactating lamb	basal diet	↓ ( $P = 0.01$ ) SOD activity, ↑ ( $P = 0.01$ ) non-protein thiols and protein thiols concentration	Cécere et al., 2021
Propolis	extract	5 g/kg of diet	<i>in vivo</i>	Barki ewes	basal diet	↓ $P < 0.01$ hydrogen peroxide, nitric oxide, and SOD	Shedeed et al., 2019
<i>Curcuma longa</i> 'Curcumin'	extract	100 mg–200 mg/kg for 30 days	<i>in vivo</i>	Nursing lambs	concentrate	↑ antioxidant ability of serum against peroxyl radicals ↓ lipoperoxidation	Molosse et al., 2019
Orange peel	essential oils (OPEO)	150 mg-300 mg-450 mg/kg concentrate	<i>in vivo</i>	Chios ewes	concentrate and alfalfa hay	blood plasma GPx, GR, and SOD – activities of milk lactoperoxidase ↑ ( $P < 0.05$ ).	Kotsampasi et al., 2018
<i>Woodfordia fruticose</i> , <i>Solanum nigrum</i> , and <i>Trigonella</i>	(flowers, shoots, and leaves), (whole plant) and (seeds), respectively	2 and 3% of diet	<i>in vivo</i>	Jamunapari goats	concentrate and roughage	↑ ( $P < 0.05$ ) GPx, CAT, and glutathione-S-transferase at 3%; ↑ non-enzymatic antioxidant indices	Choubey et al., 2015

PSM – plant secondary metabolite, FILM – *Ficus infectoria* leaf meal, G – glutathione, SOD – superoxide dismutase, CAT – catalase, LPO – lipid peroxidation, GPx – glutathione peroxidase, GST – glutathione-S-transferase, PTAA – plasma total antioxidant activity, MDA – malondialdehyde, TMR – total mixed ration, HSP – heat shock protein, T-AOC – total antioxidant capacity, T-SOD – total superoxide dismutase; TBARs – thiobarbituric acid reactive substances; OPEO – orange peel essential oil; GR – glutathione reductase, PMG – pomegranate, GM – gambier, CH – chestnut;  $P < 0.01$  indicates significant effect of PFA on the given parameters

(Skoufos et al., 2020). Plants such as oregano thyme and sage, which belong to the family *Labiatae* are among the most commonly used for this purpose. The mechanism of action of antimicrobials can be associated with hydrophobic essential oils in these plants, which strongly penetrate microbial cell membranes, fragmenting them, leading to ion leakage. *Sanguinaria canadensis* has also been reported to exhibit remarkable antimicrobial activity (Hussein et al., 2015). Plants contain hydrophobic EO that can penetrate the lipid layers of bacterial cell walls and mitochondria, leading to the accumulation of these oils in the lipid layer. This accumulation disrupts ion transport, compromises membrane integrity, and alters the osmotic pressure of the cell. The decrease in ATP synthesis and increased hydrolysis indicate a rapid dissipation of H<sup>+</sup> and K<sup>+</sup> ion gradients, which serve as proton-motive sources. This leads to the depletion of the intracellular ATP pool slowing bacterial growth due to reduced transmembrane electric potential and increased membrane proton permeability. Cell death occurs when the loss of vital molecules and ions surpasses the bacterial tolerance threshold (Simitzis, 2017). The result of Álvarez-Martínez et al. (2021) demonstrated that membrane disruption is the main mechanism of action of terpenes, flavones, flavonols, certain alkaloids, and phenylpropanoids, which showed promising antibacterial activity, either alone or in combination with extracts. The potent antimicrobial properties of EO enable them to combat various food-borne and pathogenic bacterial strains, including *Salmonella typhimurium*, *Escherichia coli*, *Shigella dysenteriae*, *Bacillus cereus*, *Pseudomonas aeruginosa*, *Campylobacter*, *Listeria monocytogenes*, and *L. innocua* (Stevanović et al., 2018). Researchers have reported significant antimicrobial potential of some plants and substances obtained from them. Clove, cinnamon, and caraway have shown considerable activity (Wickramasinghe et al., 2023), as have turmeric, oregano, thyme, and rosemary (Karásková et al., 2015). Additionally, thymol and thyme (Salehi et al., 2018), carvacrol and thymol (Memar et al., 2017), and OEO have also demonstrated substantial antimicrobial effects (Leyva-López et al., 2017). Red-osier dogwood exhibited the potential to penetrate the membranes of the rumen microbiota (Erinle et al., 2023). Furthermore, the phytochemicals in propolis phytochemicals, also known for their antibacterial properties, can improve ruminal fermentation, and thus provide nutritional benefits for ruminants (Soltan and Patra, 2020). Gillig et al. (2019) reported that the active ingredients in lemongrass, cinnamon, and oregano

essential oils – citral, cinnamaldehyde, and carvacrol, respectively – as well as extracts from grape seeds, olive oil, green tea, and allspice buds, can be administered individually, in combination, or in combination with fast-acting antibiotics to produce persistent residues. However, Ye et al. (2018) observed that the addition of monensin or cinnamaldehyde did not significantly reduce protozoan population or methane production under the conditions of their study. Zhou et al. (2020) demonstrated *in vitro* that using different levels of OEO (13 mg/l, 52 mg/l, 91 mg/l, and 130 mg/l) slightly increased the relative abundance of *Prevotella* and *Dialister* bacteria as the concentration of OEO increased. When EO incorporation rates increased, certain bacteria showed distinct polynomial responses, while other reacted in similar patterns. Ashraf et al. (2023) investigated the effectiveness of *Zingiber officinale* (ginger) and *Curcuma longa* (turmeric) against coccidia in sheep. The authors found that administering 300 mg/kg BW of turmeric reduced the number of oocytes by 44.92%, 54.32%, 64.21%, and 61.95% in a 4-week test, respectively. For the same dosage of ginger, the reductions were 39.65%, 54.81%, 57.18%, and 58.22% after 4 weeks, respectively. Additionally, the agar well diffusion method was employed to assess the antimicrobial activity of *Halexylon salicornicum* aqueous and solvent extracts against a range of animal pathogens, including *Salmonella typhimurium*, *Pseudomonas aerogenes*, *Streptococcus pyogenes*, *Escherichia coli*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Bacillus cereus*, *Shigella flexneri*, and *Enterococcus faecium*. The extract demonstrated antimicrobial activity against all tested strains of animal pathogens (El-Desoukey et al., 2022). Chan et al. (2018) evaluated the effectiveness of several plant species – *Cinnamomum burmannii*, *Origanum vulgare* L., *Syzygium aromaticum* L., *Cinnamomum cassia* Presl, *Fallopia japonica* Houtt, and *Punica granatum* L. – against five lactic acid bacteria (LAB) and five food-borne pathogenic bacteria (FBPB). Overall, the findings revealed that phenolic-rich extracts had significant antibacterial activity against FBPB but were less effective against LAB. In another study, 143 different essential oils were tested for their effect on *Staphylococcus aureus* during the stationary phase (Xiao et al., 2020). These authors identified 10 EO as particularly effective against *Staphylococcus aureus* in both the growth and stationary phases (Xiao et al., 2020).

Another study found an effect of residues of 36 plant species on the anaerobic fermentation of alfalfa. Interestingly, species such as *Myristica houtt*,

*Elsholtzia ciliata*, *Agastache rusgosa*, *Citrus aurantium*, *Citrus reticulata*, *Atractylodes lancea* T., *Codonopsis pilosula* F., and *Syzygium aromaticum* L., relatively decreased drug resistance, especially in *Salmonella* and *E. coli* strains. Additionally, a reduction in pathogenic bacterial counts was also observed. Increasing the concentration of citrus flavonoid extract (CFE) positively correlated with higher levels of beneficial bacteria, such as *Bifidobacterium* spp., the *Clostridium coccoides-Eubacterium rectale* group, and *Faecalibacterium* in faeces. Conversely, the count of *Ruminococcus torques* and *Rosburia* was reduced after CFE addition, while increasing counts of *Bacteroides* and *Phascolarctobacterium* were recorded (Li et al., 2022). Kholif and Olafadehan (2021) found that high doses of PSM as feed additives inhibited the growth and activity of cellulolytic bacteria in the rumen, while low and moderate doses did not significantly affect bacterial activity. *Haloxylon salicornicum* (Moq.) Bunge ex Boiss has been identified as a natural prophylactic antimicrobial and growth promoter, especially beneficial for chicken and cattle farms. It serves as a preferable alternative to chemical antimicrobials in animal feed, with no adverse effects or residues in animal products (El-Desoukey et al., 2022). The effects of selected PFA on ruminant microbial species are illustrated in Table 6.

## PFA effects on the immunological status of ruminants

PFA are used in animal nutrition primarily for their potential to enhance the immune system of ruminants (Karásková et al., 2015). According to recent research, plant nutrients exert physiological influence on host animals through receptor-mediated mechanisms, affecting insulin regulation, oxidative stress, and immune responses (Oh et al., 2017). Liu et al. (2020) studied the effects of a combination of essential oils (carvacrol, caryophyllene, p-cymene, cineole, terpinene, and thymol) and prebiotics (arabinogalactans and cobalt lactate at 44.1 ppm) added to the starter diet of newborn Holstein calves. The results demonstrated increased concentrations of IgG, IgM, IgA, and total serum protein in the blood. Similarly, a diet including essential oils of oregano and cinnamon, along with turmeric extract, was fed to Holstein calves. This diet resulted in a reduction in lymphocytes and heavy-chain immunoglobulins, while levels of IgA and ceruloplasmin increased.

Brunetto et al. (2023) administered 1.5 g/head/day of *Tribulus terrestris* extract (TTE) in feed and 0.5–1 µl in peripheral blood mononuclear cells demonstrated both *in vivo* and *in vitro* that TTE may modulate immunological responses and induce lymphocyte transformation in rams. This effect occurs by inhibiting lymphocyte DNA degradation and promoting their blastogenic capacity through antioxidant activity, which implies their ability to produce cytokines (Abdelrazek et al., 2018). In another study, curcumin (0.9 mg/kg BW) and a phytogetic blend of thymol, carvacrol, and cinnamaldehyde (2.7 mg/kg BW) were used as an alternative to monensin and flavomycin antibiotics in the diet of Holstein calves. Results revealed that the curcumin and phytogetic blend combination significantly reduced ( $P = 0.05$ ) monocyte and lymphocyte counts, while higher IgA and IgG titres were observed in the curcumin group (Molosse et al., 2022). *Terminalia bellirica* Gaertn. Roxb., *Emblica officinalis* Gaertn., and *Terminalia chebula* Retz. were included in the diet of post-weaned bull calves at a 75:25 ratio of seed to bark for each plant, with 0.5% of each plant used in the ration. All three treatments resulted in a decrease in IGF level, enhanced ( $P < 0.05$ ) lymphocytic proliferation, and increased IgG levels (Bostami et al., 2021). Moreover, propolis supplementation in Barki ewes at 5 g/kg diet resulted in a significant increase ( $P < 0.05$ ) in white blood cell (WBC) counts and IgA titre, while the value of mean corpuscular haemoglobin (MCH) decreased ( $P < 0.05$ ); however, no effect was observed for IgM and IgG titres (Shedeed et al., 2019). Additionally, supplementation with 4.5% licorice extract to a cottonseed hull-based diet for karakul sheep improved IgA and IgG concentrations and elevated total antioxidant capacity (Guo et al., 2019).

## Meat composition and carcass quality

Meat is vital to a healthy and well-balanced diet as it is a source of protein, iron, zinc, essential fatty acids, and B vitamins (Wood, 2017) reference. The fatty acid (FA) composition has an impact on the nutritional quality of meat. On the other hand, consumption of excessive amounts of meat has been associated with pathophysiology of cancer and heart failure, with saturated fatty acids (SFA) identified as significant contributors. In contrast, polyunsaturated fatty acid (PUFA), particularly n-3 PUFA, has been recognised for their positive impact on health, including reduced risk of thrombosis and coronary heart disease in humans (Smeti et al., 2018).

**Table 6.** Effects of selected phytogenic feed additives (PFA) on ruminant microbial species

Plant species/ PSM	Plant part/material	Dosage	Study type	Animal species	Diet/material	Result	References
Oregano	EO	130 mg/day/head 260 mg/day/head	<i>in vivo</i>	Pingliang red cattle	TMR	↑ <i>Parabacteroides distasonis</i> and <i>Bacteroides thetaiotaomicron</i> counts	Zhang et al., 2021
Oregano	EO	4–7 g/day	<i>in vivo</i>	Merino sheep	basal diet	↑ ruminal fungi count at 7 g ↓ ruminal protozoa count; ↓ total ruminal bacteria count; ↑ <i>R. flavefaciens</i> , <i>R. albus</i> , and <i>F. succinogenes</i> at 4 g	Zhou et al., 2019
Aloe vera	extract	3% of total ration	<i>in vivo</i>	Rathi calve	TMR	↓ protozoal counts and ruminal ammonia-N	Bhati et al., 2017
<i>Enterolobium cyclocarpum</i> (Jacq) Griseb. and <i>Gliricidia sepium</i> (Jacq.) Steud	foliage and pods	15-30-45%	<i>in vivo</i>	heifer	basal diet ( <i>Brachiaria brizantha</i> 'Hochst; ex A; rich'; Stapf)	↓ methanogenic archaea, total protozoa, and total bacteria in the rumen population	Molina-Botero et al., 2019
Propolis	extract	150-200-250 µl/kg of BW/day	<i>in vivo</i>	lactating lamb	basal diet	↓ ( $P = 0.01$ ) counts of <i>Escherichia coli</i> and total coliform counts in faeces	Cécere et al., 2021
<i>Cymbopogon citratus</i> (lemongrass)	EO	–	<i>in vitro</i>	agar well diffusion method	–	generated inhibition zone against <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> and <i>Bacillus cereus</i> , was $32.0 \pm 0.75$ , $48.0 \pm 1.05$ and $21.0 \pm 0.64$ mm respectively; the inhibition zone against <i>Proteus vulgaris</i> was $23.0 \pm 0.73$ mm; and no inhibition zone for <i>Pseudomonas aeruginosa</i> and <i>E. coli</i> were detected	Shendurse et al., 2021
Thymol, cinnamaldehyde, and eugenol	EOM	1 g/kg of diet 2 g/kg of diet	<i>in vitro</i>	strained rumen fluid	–	↑ <i>ruminococcus albus</i> and <i>Selenomonas ruminantium</i> $P < 0.001$ and $P < 0.005$ ↑ <i>Ruminococcus flavefaciens</i> , <i>Butyrivibrio fibrisolvens</i> , and fungi $P < 0.001$	Kim et al., 2019
Oregano (ORO), garlic (GAO), peppermint (PEO)	EO	0.50 g/l	<i>in vitro</i>	rumen fluid	–	ORO and GAO ↓ <i>Firmicutes</i> phylum counts- PEO ↑ <i>Firmicutes</i> phylum counts ORO and PEO ↑ <i>Bacteroidetes</i> phylum counts all EOs ↓ <i>Butyrivibrio</i> genus; 67 single OTUs, (operational taxonomic units), exhibit substantial differences ( $P \leq 0.05$ )	Patra and Yu, 2015
PPC	feed additive	mixture of alfalfa and PPC in a ratio of 40:60 (DM basis)	<i>in vivo</i>	lamb	fresh forage	<i>E. coli</i> O157:H7 faecal shedding ↓	Huang et al., 2015
Carvacrol and limonene	extract	0.5 g/head/day	<i>in vivo</i>	Tzigai sheep	grass hay + concentrate	number of nematode eggs in faeces ↓ ( $P = 0.02$ ); <i>Haemonchus contortus</i> nematodes in abomasum ↓	Varga-Visi et al., 2023
<i>Eucalyptus camaldulensis</i>	leaf meal	40,80 and 120 g/h/day	<i>in vivo</i>	Swamp buffalo	concentrate and rice straw	protozoa and proteolytic bacteria were ↓ ( $P < 0.05$ ); fungal zoospores, total viable bacteria, amylolytic, cellulolytic ↓	Thao et al., 2015

EO – essential oil, EOM – essential oil mixture, PPC – purple prairie clove, TMR – total mixed ration, DM – dry matter;  $P < 0.01$  indicates significant effect of PFA on the given parameters



**Table 7.** Effects of selected phytogetic feed additives (PFA) on meat characteristics

PFA sources	Dosage	Study type	Animal species	Diet	Result	References
Mesquite pods ( <i>Prosopis juliflora</i> ) extract	6 ml/day/each group	<i>in vivo</i>	Santa Ines × Dorper F1 crossbred lambs	pasture and supplementation	performance, exterior carcass measurements, leg weight, femoral muscles, carcass weight, and commercial cuts ↓ ( $P < 0.05$ )	Coelho et al., 2020
<i>Foeniculum vulgare</i> seed powder	0.75 and 1.5% of diet	<i>in vivo</i>	Male lambs	forage-concentrate 40:60	weights of both warm and cold carcass, lean meat, testicles, gallbladder, loin, and eye muscle area ↑ ( $P < 0.05$ ) at 1.5% than control	Hajalizadeh et al., 2019
REO	0.3 and 0.6 ml/head/day	<i>in vivo</i>	lamb	oat-hay and concentrate	acceptability and flavour ↑ ( $P < 0.05$ ). Total PUFA ↑. Omega-6, PUFA/SFA and n-6/n-3 percentages ↑	Smeti et al., 2018
Blend of essential oils of (oregano, garlic, lemon, rosemary, thyme, eucalyptus and sweet orange)	3.5 g and 7 g/head/day	<i>in vivo</i>	crossbred young bull	sugarcane: concentrate 10:90	meat colour, chemical and FA composition and WHC ↓. LO ↓ at 3.5 g. 7g resulted to pro-oxidant effects	Guerrero et al., 2014

REO – rosemary essential oils, PUFA – poly unsaturated fatty acid, SFA – saturated fatty acid, WHC – water holding capacity, LO – lipid oxidation, FA – fatty acids;  $P < 0.05$  indicates significant effect of PFA on the given parameters

A meta-analysis examining the effects of EO as a replacement for monensin in beef cattle diets showed improved parameters of the ribeye area (weighted mean differences (WMD) = 0.82 cm<sup>2</sup>;  $P < 0.0001$ ), subcutaneous fat thickness (WMD = 0.56 mm;  $P < 0.0001$ ), and carcass dressing percentage (WMD = 0.38%;  $P = 0.03$ ) (Torres et al., 2021). Meanwhile, Piran Filho et al. (2021) found that a PFA mixture containing cinnamon, caraway, licorice, and vanilla increased the carcass weight and DMI of F1 Angus × Nellore bulls. Adding daily a blend of oregano oil and cobalt (4 g and 7 g) to the basal diet of Suffolk lambs improved carcass weight and dressing percentages ( $P < 0.01$ ) (7 g) compared to other treatments; head width was also greater ( $P > 0.01$ ) at 7 g, while rump width increased ( $P > 0.01$ ) in lambs fed both doses of the blend (Piran Filho et al., 2021). The addition of rosemary and clove EO and an active principal blend (eugenol, thymol, and vanillin) to a concentrated diet of Nellore heifers (up to 4 g/head/day), alone or in combination, had no effects ( $P > 0.05$ ) on fat thickness, marbling, pH, muscle area, thawing, and drip loss; however, colour, texture lipid oxidation, and cooking loss ( $P < 0.05$ ) were affected by the EO treatments, in comparison to the control diet, EO and the active principle blend decreased lipid oxidation, cooking loss, and color loss (de Oliveira Monteschio et al., 2017). The effects of selected PFA on meat quality parameters are shown in Table 7.

## Conclusions

Continuous efforts are being made to produce high-quality products that benefit human health and are central to animal nutrition science. This review emphasised the benefits, as well as nutritional and medicinal properties of plants used as feed additives in ruminants. Interest in phytogetic feed additives is increasing due to the unique ability of ruminants to efficiently utilise plants and their by-products. Commonly used plants such as oregano, rosemary, thyme, coriander, garlic, eucalyptus, anise, clove, peppermint, cinnamon, turmeric, or drumstick have shown satisfactory results, particularly as antioxidant and growth promoters. Additionally, fruit extracts such as mulberry, pomegranate, and grape seed have been shown to positively impact performance, digestibility, milk production, immunity, and antioxidant status. However, the use of plant extracts faces several challenges, including limited production, underdeveloped extraction technologies, unclear dosing guidelines based on species, weight, and age, high cost and inconsistent effects. Additionally, there have been relatively few animal feeding studies and limited research on the mechanisms of action of many plant extracts. To address these challenges, further research is needed to elucidate the mechanisms of various phytogetic feed additives and their practical applications in ruminant nutrition.

## Conflict of interest

The Authors declare that there is no conflict of interest.

## References

- Abdelrazek H.M., Elgawish R.A., Ahmed E.A., Bahr H.I., 2018. *In vitro* and *in vivo* effects of *Tribulus terrestris* on immunological parameters, lymphocyte proliferation, and DNA integrity in sheep. *Small Rumin. Res.* 169, 67–73, <https://doi.org/10.1016/j.smallrumres.2018.10.014>
- Aguerre M.J., Capozzolo M.C., Lencioni P., Cabral C., Wattiaux M.A., 2016. Effect of quebracho-chestnut tannin extracts at 2 dietary crude protein levels on performance, rumen fermentation, and nitrogen partitioning in dairy cows. *J. Dairy Sci.* 99, 4476–4486, <https://doi.org/10.3168/jds.2015-10745>
- Aguilera J., Molina-Alcaide E., 2021. Enteric methane production in adult sheep of the Segureña breed fed diets containing alkali-treated olive cake. *Small Rumin. Res.* 194, 106295, <https://doi.org/10.1016/j.smallrumres.2020.106295>
- Ahmed H., Jahan S., Khan A., L Khan., Khan B.T., Ullah H., Riaz M., Ullah K., 2020. Supplementation of green tea extract (GTE) in extender improves structural and functional characteristics, total antioxidant capacity and *in vivo* fertility of buffalo (*Bubalus bubalis*) bull spermatozoa. *Theriogenology* 145, 190–197, <https://doi.org/10.1016/j.theriogenology.2019.10.024>
- Ahmed H., Jahan S., Salman M.M., Ullah F., 2019. Stimulating effects of Quercetin (QUE) in tris citric acid extender on post-thaw quality and *in vivo* fertility of buffalo (*Bubalus bubalis*) bull spermatozoa. *Theriogenology* 134, 18–23, <https://doi.org/10.1016/j.theriogenology.2019.05.012>
- Alipour F., Vakili A., Mesgaran M.D., Ebrahimi H., 2019. The effect of adding ethanolic saffron petal extract and vitamin E on growth performance, blood metabolites, and antioxidant status in Baluchi male lambs. *Asian Australas. J. Anim. Sci.* 32, 1695–1704, <https://doi.org/10.5713/ajas.18.0615>
- Al-Kassie G.A.M., 2009. Influence of two plant extracts derived from thyme and cinnamon on broiler performance. *Pak. Vet. J.* 29, 169–173
- Almeida A.K., Hegarty R.S., Cowie A., 2021. Meta-analysis quantifying the potential of dietary additives and rumen modifiers for methane mitigation in ruminant production systems. *Anim. Nutr.* 7, 1219–1230, <http://dx.doi.org/10.1016/j.aninu.2021.09.005>
- Álvarez-Martínez F., Barrajón-Catalán E., Herranz-López M., Micol V., 2021. Antibacterial plant compounds, extracts, and essential oils: An updated review on their effects and putative mechanisms of action. *Phytomedicine* 90, 153626, <https://doi.org/10.1016/j.phymed.2021.153626>
- Amarowicz R., Pegg R.B., 2019. Natural antioxidants of plant origin. In: C.F.R. Ferreira, L. Barros (Editors). *Advances in Food and Nutrition Research*. Elsevier Inc. Amsterdam (Netherlands), pp. 1–81, <https://doi.org/10.1016/bs.afnr.2019.02.011>
- Aqeel M.T., Rahman N.U., Khan A.U., Ahmad A., Ashraf Z., Rasheed U., Mansoor S., 2019. Phenolic derivatives with antioxidant and anti-inflammatory activities: an *in silico*, *in vitro* and *in vivo* study. *Pak. Vet. J.* 39, 598–602, <http://dx.doi.org/10.29261/pakvetj/2019.052>
- Ardeshirnia R., Zandi M., Sanjabi M., 2017. The effect of quercetin on fertility of frozen-thawed ram epididymal spermatozoa. *S. Afr. J. Anim. Sci.* 47, 237–244, <http://dx.doi.org/10.4314/sajas.v47i2.16>
- Artuso-Ponte V., Pastor A., Andratsch M., 2020. The effects of plant extracts on the immune system of livestock: The isoquinoline alkaloids model. In: P. Florou-Paneri, E. Christaki, I. Giannenas. *Feed Additives. Aromatic Plants and Herbs in Animal Nutrition and Health*. Elsevier Inc. Amsterdam (Nederland), pp. 295–310, <http://dx.doi.org/10.1016/B978-0-12-814700-9.00017-0>
- Ashraf A., Shahardar R.A., Wani Z.A., Bulbul K.H., Allaiie I.M., 2023. Comparative efficacy of allopathic and herbal drugs in sheep naturally infected with coccidiosis. *Res. Vet. Sci.* 164, 105001, <https://doi.org/10.1016/j.rvsc.2023.105001>
- Azimi G., Farshad A., Farzinpour A., Rostamzadeh J., Sharafi M., 2020. Evaluation of used Purslane extracts in Tris extenders on cryopreserved goat sperm. *Cryobiology* 94, 40–48, <https://doi.org/10.1016/j.cryobiol.2020.05.001>
- Barakat I.A., Khalil W.K., Al-Himaidi A.R., 2015. *Moringa oleifera* extract modulates the expression of fertility related genes and elevation of calcium ions in sheep oocytes. *Small Rumin. Res.* 130, 67–75, <http://dx.doi.org/10.1016/j.smallrumres.2015.06.011>
- Benchaar C., 2020. Feeding oregano oil and its main component carvacrol does not affect ruminal fermentation, nutrient utilization, methane emissions, milk production, or milk fatty acid composition of dairy cows. *J. Dairy Sci.* 103, 1516–1527, <https://doi.org/10.3168/jds.2019-17230>
- Bhati M., Dhuria R., Sharma T., Meel M., Saini S., 2017. Effect of Aloe vera as herbal feed additive on digestibility of nutrients and rumen fermentation in Rathi calves. *Vet. Pract.* 18, 282–283
- Birkinshaw A., Kreuzer M., Sutter M., Reidy B., Terranova M., 2022. Effects of early herbage cutting and vine leaves on methane emission, urine nitrogen losses, and the milk fatty acid profile of dairy cows. *J. Dairy Sci.* 105, 7416–7431, <https://doi.org/10.3168/jds.2021-21458>
- Bostami A.R., Khan M.R.I., Rabbi A.Z., Siddiqui M.N., Islam M.T., 2021. Boosting animal performance, immune index, and antioxidant status in post-weaned bull calves through dietary augmentation of selective traditional medicinal plants. *Vet. Anim. Sci.* 14, 100197, <https://doi.org/10.1016/j.vas.2021.100197>
- Boyko T., Chaunina E., Buzmakova N., Zharikova E., 2021. Biologically active additives for cows as a factor in the production of environmentally friendly products in animal husbandry. *IOP Conf. Ser.: Earth. Environ. Sci.* 624, <https://doi.org/10.1088/1755-1315/624/1/012063>
- Brunetto A.L., Giacomelli C.M., Favero J.F et al., 2023. Phytogenic blend in the diet of growing Holstein steers: Effects on performance, digestibility, rumen volatile fatty acid profile, and immune and antioxidant responses. *Anim. Feed Sci. Tech.* 297, 115595, <https://doi.org/10.1016/j.anifeedsci.2023.115595>
- Canaes T.S., Zanferari F., Maganhe B.L., Takiya C.S., Silva T.H., Del Valle T.A., Rennó F.P., 2017. Increasing dietary levels of citral oil on nutrient total tract digestibility, ruminal fermentation, and milk composition in Saanen goats. *Anim. Feed Sci. Tech.* 229, 47–56, <https://doi.org/10.1016/j.anifeedsci.2017.05.002>
- Caprarulo V., Ventura V., Amatucci A., Ferronato G., Gilioli G., 2022. Innovations for reducing methane emissions in livestock toward a sustainable system: analysis of feed additive patents in ruminants. *Animals* 12, 2760, <https://doi.org/10.3390/ani12202760>
- Caroprese M., Ciliberti M.G., Albenzio M., 2020. Application of aromatic plants and their extracts in dairy animals. In: P. Florou-Paneri, E. Christaki, I. Giannenas (Editors). *Feed Additives*. Academic Press. Cambridge, MA (USA), pp. 261–277, <http://dx.doi.org/10.1016/B978-0-12-814700-9.00015-7>

- Cécere B.G., da Silva A.S., Molosse V.L. et al., 2021. Addition of propolis to milk improves lactating lamb's growth: Effect on antimicrobial, antioxidant and immune responses in animals. *Small Rumin. Res.* 194, 106265, <http://dx.doi.org/10.1016/j.smallrumres.2020.106265>
- Chan C.-L., Gan R.-Y., Shah N.P., Corke H., 2018. Polyphenols from selected dietary spices and medicinal herbs differentially affect common food-borne pathogenic bacteria and lactic acid bacteria. *Food Control.* 92, 437–443, <http://dx.doi.org/10.1016/j.foodcont.2018.05.032>
- Chanu Y.M., Paul S.S., Dey A., Dahiya S.S., 2020. Reducing ruminal ammonia production with improvement in feed utilization efficiency and performance of Murrah buffalo (*Bubalus bubalis*) through dietary supplementation of plant-based feed additive blend. *Front. Vet. Sci.* 7, 464, <https://doi.org/10.3389/fvets.2020.00464>
- Choubey M., Pattanaik A.K., Baliyan S., Dutta N., Jadhav S.E., Sharma K., 2015. Dietary supplementation of a novel phytogetic feed additive: effects on nutrient metabolism, antioxidant status and immune response of goats. *Anim. Prod. Sci.* 56, 1612–1621, <http://dx.doi.org/10.1071/AN14770>
- Ciampi F., Sordillo L., Gandy J., Caroprese M., Sevi A., Albenzio M., Santillo A., 2020. Evaluation of natural plant extracts as antioxidants in a bovine in vitro model of oxidative stress. *J. Dairy Sci.* 103, 8938–8947, <https://doi.org/10.3168/jds.2020-18182>
- Coelho E., Cunha M., Santos M. et al., 2020. Phytogetic additive to improve nutrient digestibility, carcass traits, and meat quality in sheep finished on rangeland. *Livest. Sci.* 241, 104268, <http://dx.doi.org/10.1016/j.livsci.2020.104268>
- Da Silva C.S., de Souza E.J.O., Pereira G.F.C., Cavalcante E.O., de Lima E.I.M., Torres T.R., da Silva J.R.C., da Silva D.C., 2017. Plant extracts as phytogetic additives considering intake, digestibility, and feeding behavior of sheep. *Trop. Anim. Health Pro.* 49, 353–359, <https://doi.org/10.1007/s11250-016-1199-y>
- Da Silva N.I.S., de Moura J.F.P., de Lucena Nascimento M.E., Machado F.C.G., Costa T.G.P., de Araújo Filho J.M., Cordão M.A., Edvan R.L., Oliveira R.L., Bezerra L.R., 2022. Effect of *Lippia alba* hay as phytogetic feed additive on the lactation performance, milk composition, and rumen and blood parameters of Alpine goats. *Small Rumin. Res.* 215, 106767, <http://dx.doi.org/10.1016/j.smallrumres.2022.106767>
- de Oliveira Monteschio J., de Souza K.A., Vital A.C.P., Guerrero A., Valero M.V., Kempinski E.M. B.C., Barcelos V.C., Nascimento K.F., do Prado I.N., 2017. Clove and rosemary essential oils and encapsuled active principles (eugenol, thymol and vanillin blend) on meat quality of feedlot-finished heifers. *Meat Sci.* 130, 50–57, <https://doi.org/10.1016/j.meatsci.2017.04.002>
- Demirhan S.A., Karafakioğlu Y.S., 2022. Effects of anise supplement to ration of beef cattle on performance and some blood parameters. *Kocatepe Vet. J.* 15, 7–14, <https://doi.org/10.30607/kvj.815532>
- Dey A., Dutta N., Pattanaik A.K., Sharma K., 2015. Antioxidant status, metabolic profile and immune response of lambs supplemented with tannin rich *Ficus infectoria* leaf meal. *Jpn. J. Vet. Res.* 63, 15–24
- Ebeid H.M., Mengwei L., Kholif A.E., Hassan F.-u., Lijuan P., Xin L., Chengjian Y., 2020. *Moringa oleifera* oil modulates rumen microflora to mediate in vitro fermentation kinetics and methanogenesis in total mix rations. *Curr. Microbiol.* 77, 1271–1282, <https://doi.org/10.1007/s00284-020-01935-2>
- El-Desoukey R.M., Albarakaty F.M., Alzamel N.M., AlZain M.N., 2022. Ethnobotanical, phytochemical and antimicrobial activity of *Haloxylon salicornicum* (Ramth) as agraze and promising shrub against selected animal microbes. *Saudi J. Biol. Sci.* 29, 103328, <http://dx.doi.org/10.1016/j.sjbs.2022.103328>
- El-Essawy A.M., Anele U., Abdel-Wahed A., Abdou A.R., Khattab I., 2021. Effects of anise, clove and thyme essential oils supplementation on rumen fermentation, blood metabolites, milk yield and milk composition in lactating goats. *Anim. Feed Sci. Tech.* 271, 114760, <http://dx.doi.org/10.1016/j.anifeedsci.2020.114760>
- Engler P., Desguerets C., Benarbia M.E.A., Mallem Y., 2022. Supplementing young cattle with a rumen-protected grape extract around vaccination increases humoral response and antioxidant defenses. *Vet. Anim. Sci.* 15, 100232, <https://doi.org/10.1016/j.vas.2022.100232>
- Erinle T.J., Boulianne M., Miar Y., Scales R., Adewole D., 2023. Red osier dogwood and its use in animal nutrition: A review. *Anim. Nutr.* 13, 64–77, <https://doi.org/10.1016/j.aninu.2022.11.001>
- Faniyi T.O., Adewumi M.K., Jack A.A., Adegbeye M.J., Elghandour M.M., Barbabosa-Pliego A., Salem A.Z., 2021. Extracts of herbs and spices as feed additives mitigate ruminal methane production and improve fermentation characteristics in West African Dwarf sheep. *Trop. Anim. Health Pro.* 53, 312, <https://doi.org/10.1007/s11250-021-02751-x>
- Focant M., Froidmont E., Archambeau Q., Van Q.C.D., Larondelle Y., 2019. The effect of oak tannin (*Quercus robur*) and hops (*Humulus lupulus*) on dietary nitrogen efficiency, methane emission, and milk fatty acid composition of dairy cows fed a low-protein diet including linseed. *J. Dairy Sci.* 102, 1144–1159, <https://doi.org/10.3168/jds.2018-15479>
- Franz C.M., Baser K., Hahn-Ramssl I., 2020. Herbs and aromatic plants as feed additives: aspects of composition, safety, and registration rules. In: P. Florou-Paneri, E. Christaki, I. Giannenas (Editors). *Feed Additives*. Academic Press. Cambridge, MA (USA), pp. 35–56, <https://doi.org/10.1016/B978-0-12-814700-9.00003-0>
- Gessner D., Koch C., Romberg F.-J., Winkler A., Dusel G., Herzog E., Most E., Eder K., 2015. The effect of grape seed and grape marc meal extract on milk performance and the expression of genes of endoplasmic reticulum stress and inflammation in the liver of dairy cows in early lactation. *J. Dairy Sci.* 98, 8856–8868, <https://doi.org/10.3168/jds.2015-9478>
- Gilling D.H., Ravishankar S., Bright K.R., 2019. Antimicrobial efficacy of plant essential oils and extracts against *Escherichia coli*. *J. Environ. Sci. Health A* 54, 608–616, <https://doi.org/10.1080/10934529.2019.1574153>
- Gómez-Cortés P., Guerra-Rivas C., Gallardo B., Lavín P., Mantecón A., de la Fuente M., Manso T., 2018. Grape pomace in ewes diet: Effects on meat quality and the fatty acid profile of their suckling lambs. *Food Res. Int.* 113, 36–42, <https://doi.org/10.1016/j.foodres.2018.06.052>
- Guerrero A., Rivaroli D.C., Valero M.V., Sañudo C., Jorge A.M., Prado I.N., 2014. Consumers acceptability of beef from animals supplemented with essential oils. 60th International Congress of Meat Science and Technology. Punta Del Este (Uruguay)
- Guo X., Cheng L., Liu J., Zhang S., Sun X., Al-Marashdeh O., 2019. Effects of licorice extract supplementation on feed intake, digestion, rumen function, blood indices and live weight gain of Karakul sheep. *Animals* 9, 279, <https://doi.org/10.3390/ani9050279>
- Hajalizadeh Z., Dayani O., Khezri A., Tahmasbi R., Mohammadabadi M., 2019. The effect of adding fennel (*Foeniculum vulgare*) seed powder to the diet of fattening lambs on performance, carcass characteristics and liver enzymes. *Small Rumin. Res.* 175, 72–77, <https://doi.org/10.1016/j.smallrumres.2019.04.011>

- Hao X., Wang P., Ren Y., Liu G., Zhang J., Leury B., Zhang C., 2020. Effects of *Astragalus membranaceus* roots supplementation on growth performance, serum antioxidant and immune response in finishing lambs. *Asian Australas. J. Anim.* 33, 965–972, <https://doi.org/10.5713/ajas.19.0295>
- Hashem N.M., Soltan Y.A., 2016. Impacts of phytoestrogens on livestock production: A review. *Egypt. J. Nutr. Feeds* 19, 81–89, <https://doi.org/10.21608/ejnf.2016.74871>
- Hashemzadeh F., Rafeie F., Hadipour A., Rezadoust M.H., 2022. Supplementing a phytogenic-rich herbal mixture to heat-stressed lambs: Growth performance, carcass yield, and muscle and liver antioxidant status. *Small Rumin. Res.* 206, 106596, <https://doi.org/10.1016/j.smallrumres.2021.106596>
- Hrelia S., Angeloni C., 2020., New mechanisms of action of natural antioxidants in health and disease. *Antioxidants* 9, 344, <https://doi.org/10.3390/antiox9040344>
- Huang Q.Q., Jin L., Xu Z., Barbieri L.R., Acharya S., Hu T., McAllister T., Stanford K., Wang Y., 2015. Effects of purple prairie clover (*Dalea purpurea* Vent.) on feed intake, nutrient digestibility and faecal shedding of *Escherichia coli* O157: H7 in lambs. *Anim. Feed Sci. Technol.* 207, 51–61, <http://dx.doi.org/10.1016/j.anifeedsci.2015.06.009>
- Huang Q., Liu X., Zhao G., Hu T., Wang Y., 2018. Potential and challenges of tannins as an alternative to in-feed antibiotics for farm animal production. *Anim. Nutr.* 4, 137–150, <https://doi.org/10.1016/j.aninu.2017.09.004>
- Hundal J., Wadhwa M., Bakshi M., 2019. Herbal feed additives containing essential oil: Impact on the nutritional worth of complete feed in vitro. *Trop. Anim. Health Prod.* 51, 1909–1917, <https://doi.org/10.1007/s11250-019-01887-1>
- Hussein S.H., Laz E.S., Salem H.H., Hasb Allah T.I., 2015. Phytogenics as animal feed additive. *Egypt. J. Chem. Environ. Health* 1, 421–434, <https://dx.doi.org/10.21608/ejceh.2015.253590>
- Inoue M., Hayashi S., Craker L., 2017. Culture, history, and applications of medicinal and aromatic plants in Japan. In: H.A. El-Shemy (Editor). *Aromatic and Medicinal Plants-Back to Nature*. IntechOpen Ltd. London (UK), pp. 95–110, <http://dx.doi.org/10.5772/66505>
- Karášková K., Suchý P., Straková E., 2015. Current use of phytogenic feed additives in animal nutrition: a review. *Czech J. Anim. Sci.* 60, 521–530, <https://doi.org/10.17221/8594-CJAS>
- Khattab M., Kholif A., Abd El Tawab A., Shaaban M., Hadhoud F., El-Fouly H., Olafadehan O., 2020. Effect of replacement of antibiotics with thyme and celery seed mixture on the feed intake and digestion, ruminal fermentation, blood chemistry, and milk lactation of lactating Barki ewes. *Food Funct.* 11, 6889–6898, <https://doi.org/10.1039/D0FO00807A>
- Kholif A.E., Olafadehan O.A., 2021. Essential oils and phytogenic feed additives in ruminant diet: Chemistry, ruminal microbiota and fermentation, feed utilization and productive performance. *Phytochem. Rev.* 20, 1087–1108, <https://doi.org/10.1007/s11101-021-09739-3>
- Kholif A., Hassan A.A., El Ashry G.M., Bakr M., El-Zaiat H., Olafadehan O., Matloup O., Sallam S., 2021. Phytogenic feed additives mixture enhances the lactational performance, feed utilization and ruminal fermentation of Friesian cows. *Anim. Biotechnol.* 32, 708–718, <http://dx.doi.org/10.1080/10495398.2020.1746322>
- Kholif A., Matloup O., Morsy T., Abdo M., A. Elella A., Anele U., Swanson K., 2017. Rosemary and lemongrass herbs as phytogenic feed additives to improve efficient feed utilization, manipulate rumen fermentation and elevate milk production of Damascus goats. *Livest. Sci.* 204, 39–46, <http://dx.doi.org/10.1016/j.livsci.2017.08.001>
- Khorsandi S., Riasi A., Khorvash M., Hashemzadeh F., 2019. Nutrients digestibility, metabolic parameters and milk production in postpartum Holstein cows fed pomegranate (*Punica granatum* L.) by-products silage under heat stress condition. *Anim. Feed Sci. Technol.* 255, 114213, <http://dx.doi.org/10.1016/j.anifeedsci.2019.114213>
- Kim H., Jung E., Lee H.G., Kim B., Cho S., Lee S., Kwon I., Seo J., 2019. Essential oil mixture on rumen fermentation and microbial community – an *in vitro* study. *Asian Australas. J. Anim.* 32, 808–814, <http://dx.doi.org/10.5713/ajas.18.0652>
- Klebaniuk R., Bakowski M., Kowalczyk-Vasilev E., Olcha M., Widz J., Zajac M., 2016. Effect of herbal mixture in beef cattle diets on fattening performance and nutrient digestibility. *Ann. Warsaw Uni. Life Sci.* 55, 187–195
- Kolling G., Stivanin S., Gabbi A. et al., 2018. Performance and methane emissions in dairy cows fed oregano and green tea extracts as feed additives. *J. Dairy Sci.* 101, 4221–4234, <https://doi.org/10.3168/jds.2017-13841>
- Kotsampasi B., Tsiplakou E., Christodoulou C. et al., 2018. Effects of dietary orange peel essential oil supplementation on milk yield and composition, and blood and milk antioxidant status of dairy ewes. *Anim. Feed Sci. Technol.* 245, 20–31, <https://doi.org/10.1016/j.anifeedsci.2018.08.007>
- Kumar K., Dey A., Rose M.K., Dahiya S.S., 2022. Impact of dietary phytogenic composite feed additives on immune response, antioxidant status, methane production, growth performance and nutrient utilization of buffalo (*Bubalus bubalis*) calves. *Antioxidants* 11, 325, <http://dx.doi.org/10.3390/antiox11020325>
- Leyva-López N., Gutiérrez-Grijalva E.P., Vazquez-Olivo G., Heredia J.B., 2017. Essential oils of oregano: Biological activity beyond their antimicrobial properties. *Molecules* 22, 989, <http://dx.doi.org/10.3390/molecules22060989>
- Li M., Hassan F.-u., Tang Z., Peng L., Liang X., Li L., Peng K., Xie F., Yang C., 2020. Mulberry leaf flavonoids improve milk production, antioxidant, and metabolic status of water buffaloes. *Front. Vet. Sci.* 7, 599, <https://doi.org/10.3389/fvets.2020.00599>
- Li X., Chen F., Wang X., Xiong Y., Liu Z., Lin Y., Ni K., Yang F., 2022. Innovative utilization of herbal residues: exploring the diversity of mechanisms beneficial to regulate anaerobic fermentation of alfalfa. *Bioresource Technol.* 360, 127429, <https://doi.org/10.1016/j.biortech.2022.127429>
- Lillehoj H., Liu Y., Calsamiglia S., Fernandez-Miyakawa M.E., Chi F., Cravens R.L., Oh S., Gay C. G., 2018. Phytochemicals as antibiotic alternatives to promote growth and enhance host health. *Vet. Res.* 49, 76, <https://doi.org/10.1186/s13567-018-0562-6>
- Liu T., Chen H., Bai Y., Wu J., Cheng S., He B., Casper D.P., 2020. Calf starter containing a blend of essential oils and prebiotics affects the growth performance of Holstein calves. *J. Dairy Sci.* 103, 2315–2323, <http://dx.doi.org/10.3168/jds.2019-16647>
- Luo Y., Su L., Su R., Wang B., Liu C., Wang Z., Zhao L., Jin Y., 2020. Effects of *Astragalus Membranaceus* supplementation on oxidative stability of Cashmere goat. *Food Sci. Nutr.* 8, 5550–5556, <http://dx.doi.org/10.1002/fsn3.1786>
- Maciej J., Schäff C., Kanitz E., Tuchscherer A., Bruckmaier R., Wolfram S., Hammon H., 2016. Effects of oral flavonoid supplementation on the metabolic and antioxidative status of newborn dairy calves. *J. Dairy Sci.* 99, 805–811, <http://dx.doi.org/10.3168/jds.2015-9906>

- Manuelian C.L., Pitino R., Simoni M., Mavrommatis A., De Marchi M., Righi F., Tsiplakou E., 2021. Plant feed additives as natural alternatives to the use of synthetic antioxidant vitamins on livestock mammals' performances, health, and oxidative status: A review of the literature in the last 20 years. *Antioxidants* 10, 1461, <https://doi.org/10.3390/antiox10091461>
- Mastrogiovanni F., Bernini R., Basiricò L., Bernabucci U., Campo M., Romani A., Santi L., Lacetera N., 2020. Antioxidant and anti-inflammatory effects of pomegranate peel extracts on bovine mammary epithelial cells BME-UV1. *Nat. Prod. Res.* 34, 1465–1469, <https://doi.org/10.1080/14786419.2018.1508149>
- Matloup O.H., Abd El Tawab A.M., Hassan A.A., Hadhoud F.I., Khattab M.S.A., Khalel M.S., Sallam S.M.A., Kholif A.E., 2017. Performance of lactating Friesian cows fed a diet supplemented with coriander oil: feed intake, nutrient digestibility, ruminal fermentation, blood chemistry, and milk production. *Anim. Feed Sci. Technol.* 226, 88–97, <http://dx.doi.org/10.1016/j.anifeedsci.2017.02.012>
- Memar M.Y., Raei P., Alizadeh N., Aghdam M.A., Kafil H.S., 2017. Carvacrol and thymol: strong antimicrobial agents against resistant isolates. *Rev. Res. J. Med. Microbiol.* 28, 63–68, <http://dx.doi.org/10.1097/MRM.000000000000100>
- Mihaylova D., Krastanov A., Vasilev N., 2020. Non-hormonal feed additives as an alternative in animal reproduction. *Trakia J. Sci.* 4, 405–411, <http://dx.doi.org/10.15547/tjs.2020.04.016>
- Min B.R., Solaiman S., 2018. Comparative aspects of plant tannins on digestive physiology, nutrition and microbial community changes in sheep and goats: A review. *J. Anim. Physiol. Anim. Nutr.* 102, 1181–1193, <https://doi.org/10.1111/jpn.12938>
- Molina-Botero I.C., Arroyave-Jaramillo J., Valencia-Salazar S., Barahona-Rosales R., Aguilar-Pérez C.F., Burgos A.A., Arango J., Ku-Vera J.C., 2019. Effects of tannins and saponins contained in foliage of *Gliricidia sepium* and pods of *Enterolobium cyclocarpum* on fermentation, methane emissions and rumen microbial population in crossbred heifers. *Anim. Feed Sci. Technol.* 251, 1–11, <https://doi.org/10.1016/j.anifeedsci.2019.01.011>
- Molosse V.L., Deolindo G.L., Glombosky P. et al., 2022. Curcumin or microencapsulated phytochemical blend to replace ionophore and non-ionophore antibiotics in weaned calves: Effects on growth performance and health. *Livest. Sci.* 263, 105029, <https://doi.org/10.1016/j.livsci.2022.105029>
- Molosse V., Souza C.F., Baldissera M.D., Glombosky P., Campigotto G., Cazaratto C.J., Stefani M.L. M., da Silva A.S., 2019. Diet supplemented with curcumin for nursing lambs improves animal growth, energetic metabolism, and performance of the antioxidant and immune systems. *Small Rumin. Res.* 170, 74–81, <http://dx.doi.org/10.1016/j.smallrumres.2018.11.014>
- Morsy A., Soltan Y., El-Zaiat H., Alencar S.M.d., Abdalla A.L., 2021. Bee propolis extract as a phytochemical feed additive to enhance diet digestibility, rumen microbial biosynthesis, mitigating methane formation and health status of late pregnant ewes. *Anim. Feed Sci. Technol.* 273, 114834, <http://dx.doi.org/10.1016/j.anifeedsci.2021.114834>
- Muñoz-Cuautle A., Ortega-Cerrilla M.E., Herrera-Haro J.G., Nava-Cuellar C., Gutiérrez-Olvera C., Ramírez-Bribiesca J.E., Zetina-Córdoba P., 2022. Effect of Oregano (*Lippia graveolens*) Essential oil as a phytochemical feed additive on productive performance, ruminal fermentation, and antioxidant activity in lamb meat. *Agriculture* 12, 973, <https://doi.org/10.3390/agriculture12070973>
- Nawrot-Chorabik K., Sulkowska M., Gumulak N., 2022. Secondary metabolites produced by trees and fungi: achievements so far and challenges remaining. *Forests* 13, 1338, <https://doi.org/10.3390/f13081338>
- Neofytou M., Miltiadou D., Sfakianaki E., Constantinou C., Symeou S., Sparaggis D., Hager-Theodorides A.L., Tzamaloukas O., 2020. The use of ensiled olive cake in the diets of Friesian cows increases beneficial fatty acids in milk and Halloumi cheese and alters the expression of *SREBF1* in adipose tissue. *J. Dairy Sci.* 103, 8998–9011, <https://doi.org/10.3168/jds.2020-18235>
- Oh J., Giallongo F., Frederick T., Pate J., Walusimbi S., Elias R., Wall E., Bravo D., Hristov A., 2015. Effects of dietary *Capsicum* oleoresin on productivity and immune responses in lactating dairy cows. *J. Dairy Sci.* 98, 6327–6339, <https://doi.org/10.3168/jds.2014-9294>
- Oh J., Wall E.H., Bravo D.M., Hristov A.N., 2017. Host-mediated effects of phytonutrients in ruminants: A review. *J. Dairy Sci.* 100, 5974–5983, <https://doi.org/10.3168/jds.2016-12341>
- Oliveira L.N., Pereira M.A.N., Oliveira C.D.S., Oliveira C.C., Silva R.B., Pereira R.A.N., DeVries T.J., Pereira M.N., 2023. Effect of low dietary concentrations of *Acacia mearnsii* tannin extract on chewing, ruminal fermentation, digestibility, nitrogen partition, and performance of dairy cows. *J. Dairy Sci.* 106, 3203–3216, <http://dx.doi.org/10.3168/jds.2022-22521>
- Pandey A.K., Kumar P., Saxena M., 2019. Feed additives in animal health. In: R.C. Gupta, A. Srivastava, R. Lall (Editors). *Nutraceuticals in Veterinary Medicine*. Springer Nature. Cham (Switzerland), pp. 345–362, [http://dx.doi.org/10.1007/978-3-030-04624-8\\_23](http://dx.doi.org/10.1007/978-3-030-04624-8_23)
- Pandey A.K., Kumar P., Saxena M., Maurya P., 2020. Distribution of aromatic plants in the world and their properties. In: P. Florou-Paneri, E. Christaki, I. Giannenas (Editors). *Feed Additives*. Elsevier Inc. Amsterdam (Netherlands), pp. 89–114, <http://dx.doi.org/10.1016/B978-0-12-814700-9.00006-6>
- Pashtetsky V., Ostapchuk P., Kuevda T., Zubochenko D., Yemelianov S., Uppe V., 2020. Use of phytobiotics in animal husbandry and poultry. *International Scientific Conference on Biotechnology and Food Technology*. E3S Web of Conf. 215, <http://dx.doi.org/10.1051/e3sconf/202021502002>
- Passetti L.C., Passetti R.A.C., McAllister T.A., 2021. Effect of essential oil blends and a nonionic surfactant on rumen fermentation, anti-oxidative status, and growth performance of lambs. *Transl. Anim. Sci.* 5, txb118, <https://doi.org/10.1093/tas/txb118>
- Patra A.K., Yu Z., 2015. Essential oils affect populations of some rumen bacteria *in vitro* as revealed by microarray (RumenBactArray) analysis. *Front. Microbiol.* 6, 297, <https://doi.org/10.3389/fmicb.2015.00297>
- Piao M., Tu Y., Zhang N., Diao Q., Bi Y., 2023. Advances in the application of phytochemical extracts as antioxidants and their potential mechanisms in ruminants. *Antioxidants* 12, 879, <https://doi.org/10.3390/antiox12040879>
- Piran Filho F.A., Turner T.D., Mueller I., Daniel J.L.P., 2021. Influence of phytochemical feed additive on performance of feedlot cattle. *Front. Anim. Sci.* 2, 767034, <http://dx.doi.org/10.3389/fanim.2021.767034>
- Prommachart R., Uriyapongson J., Cherdthong A., Uriyapongson S., 2021. Feed intake, nutrient digestibility, antioxidant activity in plasma, and growth performance of male dairy cattle fed black rice and purple corn extracted residue. *Trop. Anim. Sci. J.* 44, 307–315, <https://doi.org/10.5398/tasj.2021.44.3.307>

- Pulina G., Francesconi A.H.D., Stefanon B. et al., 2017. Sustainable ruminant production to help feed the planet. *Ital. J. Anim. Sci.* 16, 140–171, <https://doi.org/10.1080/1828051X.2016.1260500>
- Purba R.A.P., Paengkoum P., Paengkoum S., 2020. The links between supplementary tannin levels and conjugated linoleic acid (CLA) formation in ruminants: A systematic review and meta-analysis. *PLoS One* 15, e0216187, <http://dx.doi.org/10.1371/journal.pone.0216187>
- Purba R.A.P., Paengkoum P., 2022. Exploring the phytochemical profiles and antioxidant, antidiabetic, and antihemolytic properties of *Sauropus androgynus* dried leaf extracts for ruminant health and production. *Molecules* 27, 8580, <https://doi.org/10.3390/molecules27238580>
- Quynh Chau Dang Van Q.C.D.V., Gardin C., Mignolet É., Froidmont É., Focant M., Larondelle Y., 2018. *In vitro* effects of hop pellets and oak extracts in combination on ruminal fermentation parameters. *Biotechnol. Agron. Soc. Environ.* 22, 75–83, <http://dx.doi.org/10.25518/1780-4507.16370>
- Rasby R., Funston R.N., 2016. Invited review: Nutrition and management of cows: Supplementation and feed additives. *Prof. Anim. Sci.* 32, 135–144, <https://doi.org/10.15232/pas.2015-01462>
- Rivera-Chacon R., Castillo-Lopez E., Ricci S., Petri R.M., Reisinger N., Zebeli Q., 2022. Supplementing a phytogenic feed additive modulates the risk of subacute rumen acidosis, rumen fermentation and systemic inflammation in cattle fed acidogenic diets. *Animals* 12, 1201, <https://doi.org/10.3390/ani12091201>
- Safari M., Ghasemi E., Alikhani M., Ansari-Mahyari S., 2018. Supplementation effects of pomegranate by-products on oxidative status, metabolic profile, and performance in transition dairy cows. *J Dairy Sci.* 101, 11297–11309, <https://doi.org/10.3168/jds.2018-14506>
- Salehi B., Mishra A.P., Shukla I., Sharifi-Rad M., Contreras M.d.M., Segura-Carretero A., Fathi H., Nasrabadi N.N., Kobarfard F., Sharifi-Rad J., 2018. Thymol, thyme, and other plant sources: Health and potential uses. *Phytother Res.* 32, 1688–1706, <https://doi.org/10.1002/ptr.6109>
- Samal L., Chaudhary L.C., Agarwal N., Kamra D.N., 2016. Impact of phytogenic feed additives on growth performance, nutrient digestion and methanogenesis in growing buffaloes. *Anim. Prod. Sci.* 58, 1056–1063, <http://dx.doi.org/10.1071/AN15610>
- Shedeed H.A., Farrag B., Elwakeel E.A., Abd El-Hamid I.S., El-Rayes M.A.-H., 2019. Propolis supplementation improved productivity, oxidative status, and immune response of Barki ewes and lambs. *Vet. World* 12, 834–843, <https://doi.org/10.14202/vetworld.2019.834-843>
- Shendurse A.M., Sangwan R.B., Kumar A., Ramesh V., Patel A.C., Gopikrishna G., Roy S., 2021. Phytochemical screening and antibacterial activity of lemongrass (*Cymbopogon citratus*) leaves essential oil. *J. Pharmacogn. Phytochem.* 10, 445–449
- Shilwant S., Hundal J.S., Singla M., Patra A.K., 2023. Ruminal fermentation and methane production in vitro, milk production, nutrient utilization, blood profile, and immune responses of lactating goats fed polyphenolic and saponin-rich plant extracts. *Environ. Sci. Pollut. Res.* 30, 10901–10913, <https://doi.org/10.1007/s11356-022-22931-y>
- Silveira R.F., Roque-Borda C.A., Vicente E.F., 2021. Antimicrobial peptides as a feed additive alternative to animal production, food safety and public health implications: An overview. *Anim. Nutr.* 7, 896–904, <https://doi.org/10.1016/j.aninu.2021.01.004>
- Simitzis P.E., 2017. Enrichment of animal diets with essential oils – a great perspective on improving animal performance and quality characteristics of the derived products. *Medicines* 4, 35, <https://doi.org/10.3390/medicines4020035>
- Singh A., Singh I.K. (Editors), 2018. *Molecular aspects of plant-pathogen interaction*. Springer. (Singapore), <https://doi.org/10.1007/978-981-10-7371-7>
- Singh J., Gaikwad D.S., 2020. Phytogenic feed additives in animal nutrition. In: J. Singh, A.N. Yadav (Editors). *Natural bioactive products in sustainable agriculture*. Springer. (Singapore), pp. 273–289, [http://dx.doi.org/10.1007/978-981-15-3024-1\\_13](http://dx.doi.org/10.1007/978-981-15-3024-1_13)
- Singh P.K., 2015. *An overview of feed additives*. Animal Feed Additives. New India Publ. Age. New Delhi (India), pp. 1–13, <https://doi.org/10.59317/9789389130515>
- Sirotkin A.V., Hrabovszká S., Štochmalová A., Grossmann R., Alwasel S., Harrath A.H., 2019. Effect of quercetin on ovarian cells of pigs and cattle. *Anim. Reprod. Sci.* 205, 44–51, <https://doi.org/10.1016/j.anireprosci.2019.04.002>
- Skoufos I., Bonos E., Anastasiou I., Tsinas A., Tzora A., 2020. Effects of phytobiotics in healthy or disease challenged animals. In: P. Florou-Paneri, E. Christaki, I. Giannenas (Editors). *Feed Additives*. Elsevier Inc. Amsterdam (Netherlands), pp. 311–337, <http://dx.doi.org/10.1016/B978-0-12-814700-9.00018-2>
- Smeti S., Hajji H., Mekki I., Mahouachi M., Atti N., 2018. Effects of dose and administration form of rosemary essential oils on meat quality and fatty acid profile of lamb. *Small Rumin. Res.* 158, 62–68, <http://dx.doi.org/10.1016/j.smallrumres.2017.10.007>
- Soltan Y.A., Patra A.K., 2020. Bee propolis as a natural feed additive: bioactive compounds and effects on ruminal fermentation pattern as well as productivity of ruminants. *Indian J. Anim. Hlth.* 59, 50–61, <http://dx.doi.org/10.36062/ijah.59.2SPL.2020.50-61>
- Soltan Y.A., Morsy A.S., Hashem N.M., Sallam S.M., 2021. *Boswellia sacra* resin as a phytogenic feed supplement to enhance ruminal fermentation, milk yield, and metabolic energy status of early lactating goats. *Anim. Feed Sci. Tech.* 277, 114963, <http://dx.doi.org/10.1016/j.anifeedsci.2021.114963>
- Steiner T., Syed B., 2015. Phytogenic feed additives in animal nutrition. In: A. Mathe (Editor). *Medicinal and Aromatic Plants of the World: Scientific, Production, Commercial and Utilization Aspects*. Springer. Dordrecht (Netherlands), pp. 403–423, [http://dx.doi.org/10.1007/978-94-017-9810-5\\_20](http://dx.doi.org/10.1007/978-94-017-9810-5_20)
- Stevanović Z.D., Bošnjak-Neumüller J., Pajić-Lijaković I., Raj J., Vasiljević M., 2018. Essential oils as feed additives – Future perspectives. *Molecules* 23, 1717, <https://doi.org/10.3390/molecules23071717>
- Sun J., Cheng Z., Zhao Y., Wang Y., Wang H., Ren Z., 2022. Influence of increasing levels of oregano essential oil on intestinal morphology, intestinal flora and performance of Sewa sheep. *Ital. J. Anim. Sci.* 21, 463–472, <http://dx.doi.org/10.1080/1828051X.2022.2048208>
- Sun K., Liu H., Fan H., Liu T., Zheng C., 2021. Research progress on the application of feed additives in ruminal methane emission reduction: a review. *PeerJ* 9, e11151, <https://doi.org/10.7717/peerj.11151>
- Swelum A.A., Hashem N.M., Abdelnour S.A., Taha A.E., Ohran H., Khafaga A.F., El-Tarabily K.A., Abd El-Hack M.E., 2021. Effects of phytogenic feed additives on the reproductive performance of animals. *Saudi J. Biol. Sci.* 28, 5816–5822, <https://doi.org/10.1016/j.sjbs.2021.06.045>
- Tantawi A.A., Imbabi T.A., Abdelhakeam M.A., Hassan H.M., Nasr M.A.F., Elganani S., 2023. Does phytogenic natural compound improve growth, physiological status, antioxidant parameters, digestibility and nutritive value of Ossimi lambs. *Small Rumin. Res.* 229, 107130, <http://dx.doi.org/10.1016/j.smallrumres.2023.107130>
- Tayengwa T., Mapiye C., 2018. Citrus and winery wastes: Promising dietary supplements for sustainable ruminant animal nutrition, health, production, and meat quality. *Sustainability* 10, 3718, <https://doi.org/10.3390/su10103718>

- Thao N.T., Wanapat M., Kang S., Cherdthong A., 2015. Effects of supplementation of eucalyptus (*E. camaldulensis*) leaf meal on feed intake and rumen fermentation efficiency in swamp buffaloes. *Asian Australas. J. Anim. Sci.* 28, 951–957, <https://doi.org/10.5713/ajas.14.0878>
- Torres R.N.S., Paschoaloto J.R., Ezequiel J.M.B., da Silva D., Almeida M.T.C., 2021. Meta-analysis of the effects of essential oil as an alternative to monensin in diets for beef cattle. *Vet. J.* 272, 105659, <https://doi.org/10.1016/j.tvjl.2021.105659>
- Valenzuela-Grijalva N.V., Pinelli-Saavedra A., Muhlia-Almazan A., Domínguez-Díaz D., González-Ríos H., 2017. Dietary inclusion effects of phytochemicals as growth promoters in animal production. *J. Anim. Sci. Technol.* 59, 8, <https://doi.org/10.3390/pathogens4010137>
- Varga-Visi É., Nagy G., Csivincsik Á., Tóth T., 2023. Evaluation of a phytogetic feed supplement containing carvacrol and limonene on sheep performance and parasitological status on a hungarian milking sheep farm. *Vet. Sci.* 10, 369, <https://doi.org/10.3390/vetsci10060369>
- Vasta V., Daghighi M., Cappucci A., Buccioni A., Serra A., Viti C., Mele M., 2019. Invited review: Plant polyphenols and rumen microbiota responsible for fatty acid biohydrogenation, fiber digestion, and methane emission: Experimental evidence and methodological approaches. *J. Dairy Sci.* 102, 3781–3804, <https://doi.org/10.3168/jds.2018-14985>
- Vizzotto E.F., Stivanin S.C.B., de Paris M., Passos L.T., Werncke D., Klein C.P., Stone V., Matté C., Zanela M.B., Fischer V., 2021. Supplementation with green tea and oregano extracts on productive characteristics, blood metabolites, and antioxidant status of Jersey cows during the transition period. *Animal* 15, 100032, <https://doi.org/10.1016/j.animal.2020.100032>
- Wang Z., Zhao Y., Lan X., He J., Wan F., Shen W., Tang S., Zhou C., Tan Z., Yang Y., 2022. Tannic acid supplementation in the diet of Holstein bulls: Impacts on production performance, physiological and immunological characteristics, and ruminal microbiota. *Front. Nutr.* 9, 1066074, <https://doi.org/10.3389/fnut.2022.1066074>
- Waqas M., Salman M., Sharif M.S., 2023. Application of polyphenolic compounds in animal nutrition and their promising effects. *J. Anim. Feed Sci.* 32, 233–256, <https://doi.org/10.22358/jafs/159718/2023>
- Wei L.Y., Goma W.M.S., Ametaj B.N., Alexander T.W., Yang W.Z., 2019. Feeding red osier dogwood (*Cornus sericea*) to beef heifers fed a high-grain diet affected feed intake and total tract digestibility. *Anim. Feed Sci. Technol.* 247, 83–91, <http://dx.doi.org/10.1016/j.anifeedsci.2018.11.006>
- Wickramasinghe H.K.J.P., Stepanchenko N., Oconitrillo M.J., Goetz B.M., Abeyta M.A., Gorden P.J., Baumgard L.H., Appuhamy J.A.D.R.N., 2023. Effects of a phytogetic feed additive on weaned dairy heifer calves subjected to a diurnal heat stress bout. *J. Dairy Sci.* 106, 6114–6127, <https://doi.org/10.3168/jds.2022-22856>
- Winkler A., Gessner D.K., Koch C., Romberg F.-J., Duse G.I., Herzog E., Most E., Eder K., 2015. Effects of a plant product consisting of green tea and curcuma extract on milk production and the expression of hepatic genes involved in endoplasmic stress response and inflammation in dairy cows. *Arch. Anim. Nutr.* 69, 425–441, <http://dx.doi.org/10.1080/1745039X.2015.1093873>
- Wood J.D., 2017. Meat composition and nutritional value. In: F. Toldra (Editor). *Lawrie's Meat Sci.* Elsevier Inc. Amsterdam (Netherlands), pp. 635–659, <https://doi.org/10.1016/B978-0-08-100694-8.00020-0>
- Xiao S., Cui P., Shi W., Zhang Y., 2020. Identification of essential oils with activity against stationary phase *Staphylococcus aureus*. *BMC Complement. Med. Ther.* 20, 99, <https://doi.org/10.1186/s12906-020-02898-4>
- Yang C., Chowdhury M.A.K., Hou Y., Gong J., 2015. Phytogetic compounds as alternatives to in-feed antibiotics: potentials and challenges in application. *Pathogens* 4, 137–156, <https://doi.org/10.3390/pathogens4010137>
- Yang L., He K., Li J., Ma Y., Li R., Hou M., Gao Y., Zhong J., 2019. Correlation of antibody levels with peripheral lymphocyte subsets and routine hematological parameters after vaccination with FMD vaccine in young sires. *Pak. Vet. J.* 39, 251–255, <http://dx.doi.org/10.29261/pakvetj/2019.045>
- Yang W., Lima P., Ramirez S., Schwandt E., McAllister T., 2023. Effects of a phytogetic feed additive on growth performance, feed intake, and carcass traits of beef steers. *Appl. Anim. Behav. Sci.* 39, 423–432, <https://doi.org/10.15232/aas.2023-02421>
- Yanza Y.R., Szumacher-Strabel M., Lechniak D., Ślusarczyk S., Kolodziejski P., Patra A.K., Váradyová Z., Lisiak D., Vazirigohar M., Cieslak A., 2022. Dietary *Coleus amboinicus* Lour. decreases ruminal methanogenesis and biohydrogenation, and improves meat quality and fatty acid composition in longissimus thoracis muscle of lambs. *J. Anim. Sci. Biotechnol.* 13, 5, <https://doi.org/10.1186/s40104-021-00654-3>
- Ye D., Karnati S.K.R., Wagner B., Firkins J.L., Eastridge M.L., Aldrich J.M., 2018. Essential oil and monensin affect ruminal fermentation and the protozoal population in continuous culture. *J. Dairy Sci.* 101, 5069–5081, <https://doi.org/10.3168/jds.2017-13646>
- Zhang R., Wu J., Lei Y. et al., 2021. Oregano essential oils promote rumen digestive ability by modulating epithelial development and microbiota composition in beef cattle. *Front. Nutr.* 8, 722557, <https://doi.org/10.3389/fnut.2021.722557>
- Zhao Y., Yu S., Li L., Zhao H., Li Y., Jiang L., Liu M., 2023. Feeding citrus flavonoid extracts decreases bacterial endotoxin and systemic inflammation and improves immunometabolic status by modulating hindgut microbiome and metabolome in lactating dairy cows. *Anim. Nutr.* 13, 386–400, <https://doi.org/10.1016/j.aninu.2023.03.007>
- Zhou R., Wu J., Lang X., Liu L., Casper, Wang D.P.C., Zhang L., Wei S., 2020. Effects of oregano essential oil on in vitro ruminal fermentation, methane production, and ruminal microbial community. *J. Dairy Sci.* 103, 2303–2314, <https://doi.org/10.3168/jds.2019-16611>
- Zhou R., Wu J., Zhang L.X. et al., 2019. Effects of oregano essential oil on the ruminal pH and microbial population of sheep. *PLoS One* 14, e0217054, <https://doi.org/10.1371/journal.pone.0217054>